

Manuscript Number: SAMES-D-14-00186R1

Title: First evidence of testate amoebae in Lago Fagnano (54° S), Tierra del Fuego (Argentina): Proxies to reconstruct environmental changes

Article Type: Research Paper

Keywords: Tierra del Fuego, Lago Fagnano, testate amoebae, earthquake, environmental changes

Corresponding Author: Dr. Mauro Caffau, Ph.D.

Corresponding Author's Institution: Istituto Nazionale di Oceanografia e di Geofisica Sperimentale - OGS

First Author: Mauro Caffau, Ph.D.

Order of Authors: Mauro Caffau, Ph.D.; Davide Lenaz; Emanuele Lodolo; Massimo Zecchin; Cinzia Comici; Alejandro Tassone

Abstract: We report here the first findings of testate amoebae at high southern latitudes (54° S) from four gravity cores recovered in the Lago Fagnano (Tierra del Fuego, Argentina), where twelve taxa have been recognized. Among them, *Centropyxis constricta* "constricta", *Centropyxis elongata*, *Diffugia globulus*, *Diffugia oblonga* "oblonga", and *Diffugia protaeiformis* "amphoralis" are always present, while other taxa are randomly distributed. According to the sand/silt ratio in the different cores, the Total Organic Carbon content and the Carbon/Nitrogen ratio, as well as the presence/disappearance and abundance of testate amoebae from cluster analysis, we infer a correlation between major textural/granulometrical changes found in the cores and environmental changes. A seismic event occurred on 1949, which substantially modified the morphology of the eastern Lago Fagnano shoreline and the supply pattern from two main eastern tributaries of the lake, is recorded in the studied cores. This event has in part modified the distribution of testate amoebae taxa within the studied cores. Present results show that testate amoebae represent important indicators to detect changes occurring in the environment in which they live.

*Journal of South American Earth Sciences*  
*Editorial Office*

Ms. Ref. No.: SAMES-D-14-00186

Title: *First evidence of testate amoebae in Lago Fagnano (54° S), Tierra del Fuego (Argentina): Proxies to reconstruct environmental changes*

Dear Dr. Audemard,  
please find the revised version of the above-mentioned paper.

We have remarkably changed the text, according to the reviewers requests and comments. In the new text the changes are marked in red. Below you will find the description of how we modified the text.

The main modifications made on the text are:

- 1) Sampling frequency on samples has been improved (smaller sampling range)
- 2) Details on palaeo-earthquakes in the study area have been inserted
- 3) Cluster analyses on testate amoebae have been performed

Here below, detailed answers to the reviewers:

Reviewer #1:

This change should be clearly plotted on concerned core representation. Should it be possible to add a higher frequency sampling of the few cm bounding the change, in order to check the precise position of the change? If any textural characteristics of the sediment also display the abrupt change, please add it on the logs.

**We increased the number of the studied samples (44 samples in the first version, actually 97 samples) and modified the way to display them in the diagrams.**

Nevertheless, as the detected abrupt change is attributed to a major earthquake, a few references on this subject should be added to discuss this interpretation.

**We have added some bibliographic citations about Holocene events related to earthquakes in the study area, as revealed from seismic data acquired in the Lago Fagnano.**

Reviewer #2:

For the minor revisions, please refer to the comments in the attached edited manuscript file. I really hope that the author finds my comments helpful as he continues to revise the manuscript.

**As far as concerns the minor edits in the text we accepted them and corrected accordingly.**

Major Revisions

1- Title and Key Words

\* Tierra del Fuego is divided between Argentina and Chile, and so is Lago Fagnano. In the introduction, the author states that the samples were collected from the Argentine side, so I suggest adding the name of the country to the title and the key words.

**OK, changed accordingly (see the new title).**

2- Introduction

\*a general description of the test (shell) size and morphology is missing in this section. Therefore, the author may wish to add few sentence to describe the size and different types of test (e.g. autogenous vs. xenogenous tests; refer to Patterson and Kumar 2002).

**OK, done.**

\* I think the objectives for this study need to be made clearer. The second paragraph in the introduction (P2 Line 35 to 50) states the main finding of the study without mentioning the main objectives.

**OK, in the Introduction paragraph we have made clearer the aim of the work.**

\* Did the 1949 seismic event modify the morphology of Lago Fagnano in general, or was the impact exclusive to the eastern section of the lake? This needs to be made clearer.

OK, we have discussed a bit more the impact of this event in the morphology of the lake, and introduced more details on the palaeo-earthquake evidence (mass-wasting events recorded in the Lago Fagnano)

### 3- Materials and Methods

\* The author does not provide any rationale for the sampling design, the choice of sampling techniques and details about the sampling process. Therefore, I suggest adding a subsection (e.g. sampling design and field methods) under Materials and Methods to provide these important details.

In the Materials and Methods paragraph we have discussed all the methodology applied on the sampling material, and clearly indicated the adopted techniques.

\* There are some major inconsistencies between measurements or values reported in the methods, results and discussion sections and the tables provided with the paper. Below I list these in more details:

A. P4 Line 57. Subsamples for arcellacean analyses were collected from each core at approximately 4 cm intervals. However, this does not match what is shown in tables 1 to 4. Tables 1, 2 and 3 show that the interval was approximately 8 cm between most of the subsamples, while table 4 revealed that the interval between the analysed subsamples was approximately 16 cm.

We sampled with smaller ranges in correspondence of the critical points (every 2 cm). We have refined the sampling interval in general (every 4 cm).

B. The reported core length for cores CF-1B (76 cm) and CF-F (82.5 cm) does not match the depths shown in figures 3 (80 cm) and figure 5 (83.5 cm).

OK, corrected.

\* A major weakness in the methods section is the lack of any statistical analysis, such as cluster analysis, Detrended Correspondence Analysis (DCA), Principle Component Analysis (PCA), Redundancy Analysis (RDA) and/or Canonical Correspondence Analysis (CCA). The data presented in this study must be explored in more details with the help some of these statistical methods in order to determine: (1) the dominant patterns in the arcellaceans across the cores (arcellacean assemblages or groups); and (2) the relationship between the identified assemblages and the measured environmental variables (e.g. sand/silt%, C/N ratio and TOC).

We added a paragraph (last 6 lines in the Materials and Methods paragraph) and a figure on the statistical method used.

\* I am really concerned about the low subsampling resolution in this study. The majority of the cores were about 1 m in length. However, the number of samples representing each core was very low, ranging from 5 (core CF-F) to 16 (core CF-10) subsamples. Such low resolution does not provide an accurate representation of the general trends or changes in conditions in each core. I highly recommend enhancing the temporal resolution by analysing more subsamples from each core (at least 33 subsamples per core). The additional subsamples will surely enhance the resolution in particular and the quality of the paper in general.

As in the previous answer, according to the request of both reviewers, we increased the number of the samples especially in correspondence to the level of the previously seen changes.

### 4- Results

\* In some of the cores, the appearance of plant remains does not match the marked depths in figures 1 to 4. For example, layers of plants remains were reported at depths 8, 15, 18, 56, 67, 78 and 88. However, the depths marked on figure 1 were approximately 4, 14, 18, 57, 64 and 85.

OK, done (the depths in the text were all right while they were incomplete in the figure 1 and 4, now they're all right).

\* The first two sentences (P7 Line 7 to 12) in section 4.2 must be moved to the materials and methods section.

OK; moved.

\* In section 4.3, the term "Carbon contents" is used to describe the Total Organic Analysis (TOC) results. This is not stated clearly in this section.

The section has been re-written and now it is no more used the "carbon contents" notation but only TOC.

\* In core CF-3A, the reported change in the relative abundance of some arcellacean species does not match the results shown in figure 2. For instance, the author states that the abundance of *D. protaeiformis* "amphoralis" (DPA) abruptly decreases at the depth of 40 cm, but figure 2 shows that the decrease started approximately at 56 cm. Additionally, the change in the abundance of *D. oblonga* "oblonga" (DOO) in figure 2

also contradicts the reported results. Diffugi oblonga "oblonga" was reported to decrease from the bottom of the core up to 40 cm. However, the figure clearly shows a sudden increase in the numbers of DOO between approximately 56 and 40 cm.

We changed the diagram shape, no more with lines but with histograms.

#### 5- Discussion

\* The discussion section needs substantial improvement.

We changed the Paragraph according to new findings (statistical analyses) and the text has been improved.

\* There is little explanation to the relationships between the measured environmental variables and the changes in arcellaceans. These relationships must be explored with the help of statistical analysis (see 5. Materials and Methods) in order to better validate the results and conclusions in this study.

We performed the statistical analyses that confirms the data of the histograms.

\* A more in-depth discussion about the different pioneering arcellacean species is needed. The author may wish to clearly define these species, explain their importance and refer to some examples from the literature

We changed the discussion according to histograms and statistical analyses so that no more concepts about pioneering species are present.

\* The results are based on the assumption that all these interesting changes in the sand/silt%, C/N ratio and arcellacean abundances are occurring at the depth of 64 cm, which corresponds to the seismic events that modified the lake in 1949. However, the author does not confirm if that depth (64 cm) corresponds to the same year (1949) in all four cores.

We added a short paragraph (4.3) about age estimation in the cores. In the Discussion paragraph, the changes correspond to the occurrence of the 1949 earthquake. In some cases, it is not very clear the light/dark level couple due to some plant remains.

\* The results of this study are based on analyses that were performed on a small number of subsamples (low resolution). The results, therefore, provides a basic, yet incomplete, idea about the temporal changes of the investigated environmental variables and arcellacean abundance.

The increased number of samplings we performed in this second version may solve the problem.

#### 6- Conclusion

\* P12 Line 2. The author states that "arcellacean react quickly to environmental stress, either natural or anthropogenic". I have two comments on this conclusion. First, the source and nature of environmental stress needs to be discussed in more details in the discussion section. Secondly, the source of environmental stress in this study seems to be natural (the 1949 seismic event). Therefore, I suggest deleting the word "anthropogenic" from the conclusion since such source of environmental stress was not investigated in this study.

We changed the discussion paragraph so that relationships between seismic event and environmental changes should have been clarified. We deleted the word anthropogenic.

Please refer to the following address for further correspondence:

Dr. Mauro Caffau  
Istituto Nazionale di Oceanografia e di Geofisica Sperimentale (OGS) – Trieste (Italy)  
Borgo Grotta Gigante 42/C - 34016 Sgonico (Trieste)  
Fax +39-040327307; Tel +39-0402140350  
E-mail mcaffau@ogs.trieste.it

Trieste, June, 8, 2015

## \*Detailed Response to Reviewers

Ms. Ref. No.: SAMES-D-14-00186

Title: *First evidence of testate amoebae in Lago Fagnano (54° S), Tierra del Fuego (Argentina): Proxies to reconstruct environmental changes*

Dear Dr. Audemard,  
please find the revised version of the above-mentioned paper.

We have remarkably changed the text, according to the reviewers requests and comments. In the new text the changes are marked in red. Below you will find the description of how we modified the text.

The main modifications made on the text are:

- 1) Sampling frequency on samples has been improved (smaller sampling range)
- 2) Details on palaeo-earthquakes in the study area have been inserted
- 3) Cluster analyses on testate amoebae have been performed

Here below, detailed answers to the reviewers:

Reviewer #1:

This change should be clearly plotted on concerned core representation. Should it be possible to add a higher frequency sampling of the few cm bounding the change, in order to check the precise position of the change? If any textural characteristics of the sediment also display the abrupt change, please add it on the logs.

We increased the number of the studied samples (44 samples in the first version, actually 97 samples) and modified the way to display them in the diagrams.

Nevertheless, as the detected abrupt change is attributed to a major earthquake, a few references on this subject should be added to discuss this interpretation.

We have added some bibliographic citations about Holocene events related to earthquakes in the study area, as revealed from seismic data acquired in the Lago Fagnano.

Reviewer #2:

For the minor revisions, please refer to the comments in the attached edited manuscript file. I really hope that the author finds my comments helpful as he continues to revise the manuscript.

As far as concerns the minor edits in the text we accepted them and corrected accordingly.

Major Revisions

1- Title and Key Words

\* Tierra del Fuego is divided between Argentina and Chile, and so is Lago Fagnano. In the introduction, the author states that the samples were collected from the Argentine side, so I suggest adding the name of the country to the title and the key words.

OK, changed accordingly (see the new title).

2- Introduction

\*a general description of the test (shell) size and morphology is missing in this section. Therefore, the author may wish to add few sentence to describe the size and different types of test (e.g. autogenous vs. xenogenous tests; refer to Patterson and Kumar 2002).

OK, done.

\* I think the objectives for this study need to be made clearer. The second paragraph in the introduction (P2 Line 35 to 50) states the main finding of the study without mentioning the main objectives.

OK, in the Introduction paragraph we have made clearer the aim of the work.

\* Did the 1949 seismic event modify the morphology of Lago Fagnano in general, or was the impact exclusive to the eastern section of the lake? This needs to be made clearer.

OK, we have discussed a bit more the impact of this event in the morphology of the lake, and introduced more details on the palaeo-earthquake evidence (mass-wasting events recorded in the Lago Fagnano)

3- Materials and Methods

\* The author does not provide any rationale for the sampling design, the choice of sampling techniques and details about the sampling process. Therefore, I suggest adding a subsection (e.g. sampling design and field methods) under Materials and Methods to provide these important details.

In the Materials and Methods paragraph we have discussed all the methodology applied on the sampling material, and clearly indicated the adopted techniques.

\* There are some major inconsistencies between measurements or values reported in the methods, results and discussion sections and the tables provided with the paper. Below I list these in more details:

A. P4 Line 57. Subsamples for arcellacean analyses were collected from each core at approximately 4 cm intervals. However, this does not match what is shown in tables 1 to 4. Tables 1, 2 and 3 show that the interval was approximately 8 cm between most of the subsamples, while table 4 revealed that the interval between the analysed subsamples was approximately 16 cm.

We sampled with smaller ranges in correspondence of the critical points (every 2 cm). We have refined the sampling interval in general (every 4 cm).

B. The reported core length for cores CF-1B (76 cm) and CF-F (82.5 cm) does not match the depths shown in figures 3 (80 cm) and figure 5 (83.5 cm).

OK, corrected.

\* A major weakness in the methods section is the lack of any statistical analysis, such as cluster analysis, Detrended Correspondence Analysis (DCA), Principle Component Analysis (PCA), Redundancy Analysis (RDA) and/or Canonical Correspondence Analysis (CCA). The data presented in this study must be explored in more details with the help some of these statistical methods in order to determine: (1) the dominant patterns in the arcellaceans across the cores (arcellacean assemblages or groups); and (2) the relationship between the identified assemblages and the measured environmental variables (e.g. sand/silt%, C/N ratio and TOC).

We added a paragraph (last 6 lines in the Materials and Methods paragraph) and a figure on the statistical method used.

\* I am really concerned about the low subsampling resolution in this study. The majority of the cores were about 1 m in length. However, the number of samples representing each core was very low, ranging from 5 (core CF-F) to 16 (core CF-10) subsamples. Such low resolution does not provide an accurate representation of the general trends or changes in conditions in each core. I highly recommend enhancing the temporal resolution by analysing more subsamples from each core (at least 33 subsamples per core). The additional subsamples will surely enhance the resolution in particular and the quality of the paper in general.

As in the previous answer, according to the request of both reviewers, we increased the number of the samples especially in correspondence to the level of the previously seen changes.

#### 4- Results

\* In some of the cores, the appearance of plant remains does not match the marked depths in figures 1 to 4. For example, layers of plants remains were reported at depths 8, 15, 18, 56, 67, 78 and 88. However, the depths marked on figure 1 were approximately 4, 14, 18, 57, 64 and 85.

OK, done (the depths in the text were all right while they were incomplete in the figure 1 and 4, now they're all right).

\* The first two sentences (P7 Line 7 to 12) in section 4.2 must be moved to the materials and methods section.

OK; moved.

\* In section 4.3, the term "Carbon contents" is used to describe the Total Organic Analysis (TOC) results. This is not stated clearly in this section.

The section has been re-written and now it is no more used the "carbon contents" notation but only TOC.

\* In core CF-3A, the reported change in the relative abundance of some arcellacean species does not match the results shown in figure 2. For instance, the author states that the abundance of *D. protaeiformis* "amphoralis" (DPA) abruptly decreases at the depth of 40 cm, but figure 2 shows that the decrease started approximately at 56 cm. Additionally, the change in the abundance of *D. oblonga* "oblonga" (DOO) in figure 2

also contradicts the reported results. *Diffugi oblonga* "oblonga" was reported to decrease from the bottom of the core up to 40 cm. However, the figure clearly shows a sudden increase in the numbers of DOO between approximately 56 and 40 cm.

We changed the diagram shape, no more with lines but with histograms.

#### 5- Discussion

\* The discussion section needs substantial improvement.

We changed the Paragraph according to new findings (statistical analyses) and the text has been improved.

\* There is little explanation to the relationships between the measured environmental variables and the changes in arcellaceans. These relationships must be explored with the help of statistical analysis (see 5. Materials and Methods) in order to better validate the results and conclusions in this study.

We performed the statistical analyses that confirms the data of the histograms.

\* A more in-depth discussion about the different pioneering arcellacean species is needed. The author may wish to clearly define these species, explain their importance and refer to some examples from the literature

We changed the discussion according to histograms and statistical analyses so that no more concepts about pioneering species are present.

\* The results are based on the assumption that all these interesting changes in the sand/silt%, C/N ratio and arcellacean abundances are occurring at the depth of 64 cm, which corresponds to the seismic events that modified the lake in 1949. However, the author does not confirm if that depth (64 cm) corresponds to the same year (1949) in all four cores.

We added a short paragraph (4.3) about age estimation in the cores. In the Discussion paragraph, the changes correspond to the occurrence of the 1949 earthquake. In some cases, it is not very clear the light/dark level couple due to some plant remains.

\* The results of this study are based on analyses that were performed on a small number of subsamples (low resolution). The results, therefore, provides a basic, yet incomplete, idea about the temporal changes of the investigated environmental variables and arcellacean abundance.



The increased number of samplings we performed in this second version may solve the problem.

#### 6- Conclusion

\* P12 Line 2. The author states that "arcellacean react quickly to environmental stress, either natural or anthropogenic". I have two comments on this conclusion. First, the source and nature of environmental stress needs to be discussed in more details in the discussion section. Secondly, the source of environmental stress in this study seems to be natural (the 1949 seismic event). Therefore, I suggest deleting the word "anthropogenic" from the conclusion since such source of environmental stress was not investigated in this study.

We changed the discussion paragraph so that relationships between seismic event and environmental changes should have been clarified. We deleted the word anthropogenic.







The first findings of testate amoebae at high southern latitudes (Tierra del Fuego, 54° S) is reported. Twelve taxa have been recognized

Four gravity cores recovered in the Lago Fagnano (Tierra del Fuego, Argentina), have been analyzed.

A correlation between major textural/granulometrical changes occurred during a seismic event (1949, Moment magnitude 7.5 earthquake), and testate amoebae distribution was found.

Testate amoebae represent important indicators to detect changes occurring in the environment in which they live.

# First evidence of testate amoebae in Lago Fagnano (54° S), Tierra del Fuego (Argentina): Proxies to reconstruct environmental changes

Mauro Caffau<sup>a,\*</sup>, Davide Lenaz<sup>b</sup>, Emanuele Lodolo<sup>a</sup>, Massimo Zecchin<sup>a</sup>, Cinzia Comici<sup>a</sup>, Alejandro Tassone<sup>c</sup>

<sup>a</sup>Istituto Nazionale di Oceanografia e di Geofisica Sperimentale (OGS), Trieste - Italy

<sup>b</sup>Dipartimento di Matematica e Geoscienze, Università di Trieste - Italy

<sup>c</sup>Instituto de Geofísica “D. Valencio”, Universidad de Buenos Aires - Argentina

\*Corresponding author: [mcaffau@ogs.trieste.it](mailto:mcaffau@ogs.trieste.it)

---

## Abstract

We report here the first findings of **testate amoebae** at high southern latitudes (54° S) from four gravity cores recovered in the Lago Fagnano (Tierra del Fuego, Argentina), where twelve taxa have been recognized. Among them, *Centropyxis constricta* “**constricta**”, *Centropyxis elongata*, *Diffugia globulus*, *Diffugia oblonga* “**oblonga**”, and *Diffugia protaeiformis* “**amphoralis**” are always present, while other taxa are randomly distributed. According to the sand/silt ratio in the different cores, the Total Organic **Carbon** content and the **Carbon/Nitrogen** ratio, as well as the presence/disappearance and abundance of **testate amoebae from cluster analysis**, we infer a correlation between major textural/granulometrical changes found in the cores **and environmental changes**. A seismic event occurred on 1949, which substantially modified the morphology of the eastern Lago Fagnano shoreline and the supply pattern from two main eastern tributaries of the lake, **is recorded in the studied cores**. This event has in part modified the distribution of **testate amoebae** taxa within the studied cores. **Present results show that testate amoebae represent important indicators to detect changes occurring in the environment in which they live.**

**Keywords:** Tierra del Fuego, Lago Fagnano, **testate amoebae**, earthquake, environmental changes.

## 1. Introduction

**Testate amoebae (Protozoa: Rhizopoda)** are unicellular organism characterized by enclosed, morphologically distinct tests. There are several types of tests, referred to as xenogenous

1 agglutinated or autogenous tests. The former are exceedingly variable, composed of particles  
2 derived from the environment such lithic granules and sometimes organic detritus such as diatom  
3 frustules. Autogenous tests are composed of material secreted by the organism. (Charman, 2001;  
4 Scott et al., 2001; Patterson and Kumar, 2002). Testate amoebae are present in a wide range of  
5 terrestrial and aquatic environments, including freshwater lakes, brackish intertidal environment,  
6 wetlands, peatlands and, to a lesser extent, moist soil and forest litter (Charman, 2001; Scott et al.,  
7 2001, Booth, 2002, Charman et al., 2006). Testate amoebae have proven to be good environmental  
8 and paleoenvironmental indicators, both because their tests tend to be resistant to dissolution and  
9 well preserved in sediments, and because they are sensitive to a wide variety of environmental  
10 variables (Medioli and Scott, 1988; Warner and Charman, 1994; Charman 2001; Patterson and  
11 Kumar, 2002; Patterson et al., 2002) including metal and organic pollutant contamination, substrate  
12 type, salinity, levels of organics, oxygen concentration, water temperatures, water table fluctuations,  
13 humification and changes in intertidal flooding (e.g., Scott and Medioli, 1983; Patterson et al.,  
14 1996, 2002; Medioli et al., 1990; Collins et al., 1990; McCarthy et al., 1995; Asioli et al., 1996;  
15 Charman et al., 1998; Reinhardt et al., 1998; Patterson and Kumar, 2000, 2002). Most research on  
16 *testate amoebae* has focused on faunas found at higher latitudes (Scott and Medioli, 1983; Collins et  
17 al., 1990; Patterson et al., 1996, 2002; Tolonen et al., 1992; Kliza and Schröder, 1999). In South  
18 America *testate amoebae* have been reported, among others, in the wetlands of central Brazil  
19 (Green, 1975), in Lake Cocococha, Peru (Haman and Kohl, 1994), in a peat from the Puyehue  
20 National Park in Chile (Zapata et al., 2008), whereas the most southern occurrence was found in the  
21 Chilean Guamblin Island (Fernandez et al., 2012), located at about 45° S.

22  
23  
24  
25  
26  
27  
28  
29  
30  
31  
32  
33  
34  
35  
36  
37  
38  
39  
40  
41  
42  
43  
44  
45  
46  
47  
48  
49  
50  
51  
52  
53  
54  
55  
56  
57  
58  
59  
60  
61  
62  
63  
64  
65  
In this paper, we present the first evidence of the presence of *testate amoebae* at the latitude  
of 54° 30' S in Tierra del Fuego, which is the southernmost latitude at which these organisms have  
been found to date. In order to achieve a better understanding of sediment provenance, transport  
mechanisms, inputs of organic matter from terrestrial sources, and on the sensitivity of testate  
amoebae to environmental changes, we combined analyses of organic matter parameters (C and N)  
with testate amoebae assemblages, their distribution within four boreholes, and the physical  
properties of sediments analyzed from four boreholes taken in the eastern part of Lago Fagnano.

## 2. General morphology of Lago Fagnano and depositional setting

Lago Fagnano (54° 30' S, 68° W, 26 m above sea level), covering a total area of 596 km<sup>2</sup>, is located  
in the central part of the Tierra del Fuego, and represents the southernmost and largest ice-free lake  
of the Southern Hemisphere (Fig. 1A). The basin trends roughly E-W, is more than 105 km long,  
and its width varies from 2.8 to 9.7 km. It occupies a significant segment of the South America-

1 Scotia transform plate boundary, which in Tierra del Fuego is known as the Magallanes-Fagnano  
2 fault system (Lodolo et al., 2002, 2003, 2007; Tassone et al., 2005; Menichetti et al., 2008).  
3 Repeated bathymetric and seismic surveys carried out along the entire length of the basin  
4 (Waldmann et al., 2008, 2010a, 2011; Zanolla et al., 2011; Lodolo et al., 2012; Esteban et al., 2014)  
5 have delineated in detail its morphological setting and sedimentary architecture. Data show that the  
6 basin floor is divided into two principal parts, separated by a morphologically complex, shallow  
7 relief located in the central part of the lake. This suggests that the basin is composed of at least two  
8 sub-basins. The deepest depression, with a maximum water depth of 206 m, is found in the eastern  
9 sub-basin, whereas the western-half of the lake has a maximum water depth of 165 m (Zanolla et  
10 al., 2011). Combining geophysical data with field observations and regional maps, Waldmann et al.  
11 (2010b) asserted that the Lago Fagnano region has experienced several phases of glacier growth and  
12 retreat since the Last Glacial Maximum. The peculiar shape and morphology of the lake, and its  
13 location within the principal displacement zone of the Magallanes-Fagnano fault system, suggests  
14 that the tectonic activity along this lineament has controlled the formation and development of the  
15 basin at least in its early times. However, the Late Quaternary glacial activity played an important  
16 role in shaping the bottom morphology and the surrounding landscape, as testified by the geometry  
17 and sedimentary setting of the deposits filling the basin, and by the widespread glacial landforms  
18 surrounding the lake coastline (Coronato et al., 2009).

19  
20  
21  
22  
23  
24  
25  
26  
27  
28  
29  
30  
31  
32  
33  
34  
35  
36  
37  
38  
39  
40  
41  
42  
43  
44  
45  
46  
47  
48  
49  
50  
51  
52  
53  
54  
55  
56  
57  
58  
59  
60  
61  
62  
63  
64  
65  
66  
67  
68  
69  
70  
71  
72  
73  
74  
75  
76  
77  
78  
79  
80  
81  
82  
83  
84  
85  
86  
87  
88  
89  
90  
91  
92  
93  
94  
95  
96  
97  
98  
99  
100  
101  
102  
103  
104  
105  
106  
107  
108  
109  
110  
111  
112  
113  
114  
115  
116  
117  
118  
119  
120  
121  
122  
123  
124  
125  
126  
127  
128  
129  
130  
131  
132  
133  
134  
135  
136  
137  
138  
139  
140  
141  
142  
143  
144  
145  
146  
147  
148  
149  
150  
151  
152  
153  
154  
155  
156  
157  
158  
159  
160  
161  
162  
163  
164  
165  
166  
167  
168  
169  
170  
171  
172  
173  
174  
175  
176  
177  
178  
179  
180  
181  
182  
183  
184  
185  
186  
187  
188  
189  
190  
191  
192  
193  
194  
195  
196  
197  
198  
199  
200  
201  
202  
203  
204  
205  
206  
207  
208  
209  
210  
211  
212  
213  
214  
215  
216  
217  
218  
219  
220  
221  
222  
223  
224  
225  
226  
227  
228  
229  
230  
231  
232  
233  
234  
235  
236  
237  
238  
239  
240  
241  
242  
243  
244  
245  
246  
247  
248  
249  
250  
251  
252  
253  
254  
255  
256  
257  
258  
259  
260  
261  
262  
263  
264  
265  
266  
267  
268  
269  
270  
271  
272  
273  
274  
275  
276  
277  
278  
279  
280  
281  
282  
283  
284  
285  
286  
287  
288  
289  
290  
291  
292  
293  
294  
295  
296  
297  
298  
299  
300  
301  
302  
303  
304  
305  
306  
307  
308  
309  
310  
311  
312  
313  
314  
315  
316  
317  
318  
319  
320  
321  
322  
323  
324  
325  
326  
327  
328  
329  
330  
331  
332  
333  
334  
335  
336  
337  
338  
339  
340  
341  
342  
343  
344  
345  
346  
347  
348  
349  
350  
351  
352  
353  
354  
355  
356  
357  
358  
359  
360  
361  
362  
363  
364  
365  
366  
367  
368  
369  
370  
371  
372  
373  
374  
375  
376  
377  
378  
379  
380  
381  
382  
383  
384  
385  
386  
387  
388  
389  
390  
391  
392  
393  
394  
395  
396  
397  
398  
399  
400  
401  
402  
403  
404  
405  
406  
407  
408  
409  
410  
411  
412  
413  
414  
415  
416  
417  
418  
419  
420  
421  
422  
423  
424  
425  
426  
427  
428  
429  
430  
431  
432  
433  
434  
435  
436  
437  
438  
439  
440  
441  
442  
443  
444  
445  
446  
447  
448  
449  
450  
451  
452  
453  
454  
455  
456  
457  
458  
459  
460  
461  
462  
463  
464  
465  
466  
467  
468  
469  
470  
471  
472  
473  
474  
475  
476  
477  
478  
479  
480  
481  
482  
483  
484  
485  
486  
487  
488  
489  
490  
491  
492  
493  
494  
495  
496  
497  
498  
499  
500  
501  
502  
503  
504  
505  
506  
507  
508  
509  
510  
511  
512  
513  
514  
515  
516  
517  
518  
519  
520  
521  
522  
523  
524  
525  
526  
527  
528  
529  
530  
531  
532  
533  
534  
535  
536  
537  
538  
539  
540  
541  
542  
543  
544  
545  
546  
547  
548  
549  
550  
551  
552  
553  
554  
555  
556  
557  
558  
559  
560  
561  
562  
563  
564  
565  
566  
567  
568  
569  
570  
571  
572  
573  
574  
575  
576  
577  
578  
579  
580  
581  
582  
583  
584  
585  
586  
587  
588  
589  
590  
591  
592  
593  
594  
595  
596  
597  
598  
599  
600  
601  
602  
603  
604  
605  
606  
607  
608  
609  
610  
611  
612  
613  
614  
615  
616  
617  
618  
619  
620  
621  
622  
623  
624  
625  
626  
627  
628  
629  
630  
631  
632  
633  
634  
635  
636  
637  
638  
639  
640  
641  
642  
643  
644  
645  
646  
647  
648  
649  
650  
651  
652  
653  
654  
655  
656  
657  
658  
659  
660  
661  
662  
663  
664  
665  
666  
667  
668  
669  
670  
671  
672  
673  
674  
675  
676  
677  
678  
679  
680  
681  
682  
683  
684  
685  
686  
687  
688  
689  
690  
691  
692  
693  
694  
695  
696  
697  
698  
699  
700  
701  
702  
703  
704  
705  
706  
707  
708  
709  
710  
711  
712  
713  
714  
715  
716  
717  
718  
719  
720  
721  
722  
723  
724  
725  
726  
727  
728  
729  
730  
731  
732  
733  
734  
735  
736  
737  
738  
739  
740  
741  
742  
743  
744  
745  
746  
747  
748  
749  
750  
751  
752  
753  
754  
755  
756  
757  
758  
759  
760  
761  
762  
763  
764  
765  
766  
767  
768  
769  
770  
771  
772  
773  
774  
775  
776  
777  
778  
779  
780  
781  
782  
783  
784  
785  
786  
787  
788  
789  
790  
791  
792  
793  
794  
795  
796  
797  
798  
799  
800  
801  
802  
803  
804  
805  
806  
807  
808  
809  
810  
811  
812  
813  
814  
815  
816  
817  
818  
819  
820  
821  
822  
823  
824  
825  
826  
827  
828  
829  
830  
831  
832  
833  
834  
835  
836  
837  
838  
839  
840  
841  
842  
843  
844  
845  
846  
847  
848  
849  
850  
851  
852  
853  
854  
855  
856  
857  
858  
859  
860  
861  
862  
863  
864  
865  
866  
867  
868  
869  
870  
871  
872  
873  
874  
875  
876  
877  
878  
879  
880  
881  
882  
883  
884  
885  
886  
887  
888  
889  
890  
891  
892  
893  
894  
895  
896  
897  
898  
899  
900  
901  
902  
903  
904  
905  
906  
907  
908  
909  
910  
911  
912  
913  
914  
915  
916  
917  
918  
919  
920  
921  
922  
923  
924  
925  
926  
927  
928  
929  
930  
931  
932  
933  
934  
935  
936  
937  
938  
939  
940  
941  
942  
943  
944  
945  
946  
947  
948  
949  
950  
951  
952  
953  
954  
955  
956  
957  
958  
959  
960  
961  
962  
963  
964  
965  
966  
967  
968  
969  
970  
971  
972  
973  
974  
975  
976  
977  
978  
979  
980  
981  
982  
983  
984  
985  
986  
987  
988  
989  
990  
991  
992  
993  
994  
995  
996  
997  
998  
999  
1000

Currently, the main tributary of the eastern sector of the Lago Fagnano is the Rio Turbio. The climate of this region is Alpine, with a strong winter sub-polar Antarctic influence, and is under the south-westerly wind effect, bringing moisture and humidity to the region during austral summers (Rabassa et al., 2000). The average air temperature is about +5 °C (Tuhkanen, 1992). Precipitation is essentially of orographic nature (about 550 mm mean annual precipitation), and the drainage basin belongs to the vegetation zone of the subantarctic deciduous forest (Richter et al., 2010).

Petrophysical, sedimentological and geochemical studies of a Holocene lacustrine laminated succession, performed on selected cores (Waldmann et al., 2008, 2010a, 2011; Moy et al., 2011), revealed significant variations in major and trace elements, as well as in organic content, suggesting high variability in environmental conditions. Constituents of recent sediments are both biogenic and lithogenic, consisting of fresh water lacustrine biogenic material and variable amounts of reworked organic and clastic detritus delivered by rivers. In the area of Rio Turbio estuary, the sediment supply is characterized by silty suspended particles. **Since the stream provides old organic carbon from plants and grassland, soil or peat material eroded from the nearby land, Moy et al. (2011) pointed out that radiocarbon ages of bulk organic matter in core samples from the central area of the**

lake are unreliable. These datings demonstrate that the first 2.5 m of sediment has been deposited during Holocene time, but the ages of samples collected from the cores are irregularly distributed and range from ca. 0.5 to 11 kyr B.P. (Moy et al., 2011). Because the cores analyzed in this study are likely more directly under riverine influence than those of the central lake due to their vicinity to the Rio Turbio mouth, it is expected that the sedimentation rate, as well as the thickness of Holocene sediments and the amount of reworked organic matter are higher, which would make any <sup>14</sup>C date questionable. Therefore, tacking also into account that the thickness of present cores does not exceed 0.95 m, that is significantly lower than that of the cores of the central lake, we refer to the core material generally as Holocene sediment. In order to obtain an approximate age, Waldmann et al. (2008) counted the light/dark laminae couplets present in their core sediments of Lago Fagnano, considering that a couple of laminae deposited in one year.

Changes in the geochemical characteristics of lake sediments may indicate variations in the sedimentation pattern through time. These changes may indirectly occur following both strong climatic oscillations and seismic events. In the case of Lago Fagnano, a Moment magnitude = 7.5 earthquake occurred on December 17, 1949 (Lomnitz, 1970), which significantly modified the landscape, the hydrographic net, and the shoreline of the eastern Lago Fagnano (Figs. 1C; 1D). Following this event, a gravel barrier along the Lago Fagnano eastern shoreline was created, modifying the flow pattern of the two former principal tributaries represented by the Rio Turbio and Rio Tolhuin (Fig. 1B). A lagoon-like area (a sag pond) was thus created immediately following the earthquake (Fig. 1C,D), and the inundation caused a dead forest of *Nothofagus pumilio*, as presently observable (Costa et al., 2006). Today, the only output of the lagoon-like area is represented by the mouth of the Rio Turbio (Fig. 1B), as seen from satellite images which show the presence of currently buried stream outputs in correspondence of the northern sector of the sag pond. Fingerprints of palaeo-earthquakes enucleated along the Magallanes-Fagnano fault system have been also recognized in the study area by seismic records acquired in the Lago Fagnano, and represented by Holocene slope failures and megaturbidites (Waldmann et al., 2011).

### 3. Materials and methods

The meter-length gravity cores analyzed in this paper have been recovered in the vicinity of the eastern shore of Lago Fagnano on March 2010 (Lodolo et al., 2012). Sampling included smear slide analysis to characterize sediment type and the fossiliferous fraction. For the analysis of the lithological composition, 10 cm<sup>3</sup> sample volume of wet sediment for a total of 97 sediment samples were taken at approximate 4 cm intervals from four gravity cores. The percentage content of sand, silt and organic material, including macrophytes, terrestrial plant remains, wood remains,



1 Gyrogonites of Characeans, seeds and **testate amoebae**, was determined. Layers of macrophytes,  
2 terrestrial plant remains, and dark laminations are usually present and well distributed along the  
3 four cores, although at different concentrations in the analyzed samples. To calculate the percentage  
4 content of sand and silt of the samples, wet sediments were dried in oven at 40 °C and hence  
5 weighted. Samples were subsequently disaggregated with 10 V hydrogen peroxide, washed through  
6 0.063 mm sieve, dried and weighted. Each dried sediment was further analyzed at the optical  
7 microscope, to describe the biogenic and detrital inorganic material.

12 For organic content measurements, the sediment sample was freeze-dried and grounded in a  
13 mortar, then pestled and the fraction  $\geq 1$  mm was separated from the rest of the sample; a 1 mm  
14 mesh was used instead of 250  $\mu\text{m}$  because of the coarse grain size of the sediments. Two replicates  
15 of about 8-12 mg were weighted on a microultrabalance Perkin Elmer mod. AD-4 (0.1  $\mu\text{g}$   
16 accuracy). Total Nitrogen (TN) and Total Organic Carbon (TOC) were measured using an  
17 elemental analyzer CHNO-S Costech mod. ECS 4010. Known amounts of standard Acetanilide  
18 ( $\text{C}_8\text{H}_9\text{NO}$  – Carlo Erba; Assay  $\geq 99.5$  %) were used to calibrate the instrument. For TOC analysis,  
19 subsamples were treated with subsequent additions of hydrochloric acid at increasing concentration  
20 (0.1 N and 1 N) to remove the carbonates (Nieuwenhuize *et al.*, 1994). The detection limit (DL) of  
21 the instrument was 0.004 % dwt (dry weight) for TN and 0.14 % dwt for TOC.

31 For the analyses of the **testate amoebae**, 3  $\text{cm}^3$  sample volume of wet sediment were taken at  
32 approximate 4 cm intervals throughout the **cores** and prepared following the method described by  
33 Scott *et al.* (2001). Sediment samples were examined (Leica microscope) under x20 and x80  
34 magnification, and 200-300 **testate amoebae** were counted. Selected specimens were photographed  
35 on a SEM (Scanning Electron Microscope) Leica Stereoscan 430i. **Testate amoebae** were identified  
36 to the strain level using Burbige and Schöder-Adams (1998) and Reinhardt *et al.* (1998).

42 **X-ray diffraction patterns were obtained on bulk sediment samples spread out on aluminum  
43 plates using a STOE D 500 X-ray diffractometer at room temperature.  $\text{CuK}\alpha$  radiation was used  
44 through a flat graphite crystal monochromator. Samples were investigated in a range between 2 and  
45  $40^\circ$  of  $2\theta$  angle with a step scan of  $0.1^\circ$  and 2 s measure time.**

49 **R-mode cluster analysis was performed computing Bray and Curtis (1957) dissimilarity (i.e.,  
50 a statistic used to quantify the compositional dissimilarity between two different sites, based on  
51 counts at each site) between variables (i.e. species), and then using complete linkage as an  
52 agglomerative algorithm. Cluster analysis was carried out using a specific software ([http://www.r-  
53 project.org](http://www.r-project.org)). No other multivariate techniques were used since only one environmental parameter  
54 (sand/silt ratio) was available for a considerable number of samples.**

## 4. Results

### 4.1. Sedimentological analysis

In all the analyzed cores the sand percentage is always lower than the silt + clay fraction, with ratios ranging from 0.09 to 0.82, except for one sample (at 24 cm depth) in the core CF-1B. Notably, the amount of sand is usually higher in the upper 40 cm of all cores apart in the core CF-3A, where the amount of sand is higher in its lower part. It is also to be noted that in the two northern cores there is a concomitant increase of sand fraction (sand/silt ratio about 0.8) at 62-64 cm depth. Percentages of sand and silt + clay are shown in Figs. 2, 3, 4 and 5. Sands are mainly composed of quartz and subordinately of feldspars. Clay minerals are also present. No appreciable changes in the percentages of the recognized minerals are recorded among the cores.

Core CF-3A was taken at a water depth of 19.8 m and recovered 92 cm of lacustrine deposits (Fig. 2). The sediment is thoroughly laminated, consisting of an alternation between 1.5-2 mm thick dark-gray silty laminae and 1-2 mm thick slightly lighter-colored laminae of clayey silt. A peak in sand/silt ratio is found at 62-64 cm from the top. Several layers of plant remains are also found along the core; they are up to 2-3 mm thick and are present at depths of 88, 78, 67, 56, 18, 15 and 8 cm. Other patches of organic content, less than 1 mm thick, are frequent and distributed at about 5 cm intervals along the core. Gyrogonites of Characeans are found at depth of 56 cm. A wood remain, 1.5 cm thick, was recovered at depth of 18 cm. Inorganic components mainly consist of poorly selected, rounded quartz grains.

Core CF-1B was taken at a water depth of 20.4 m and recovered 80 cm of lacustrine deposits (Fig. 3). The examined sediments consist of interlaminated silt to sandy silt, silt and clay, forming planar lamination. In detail, from the bottom up to 38 cm, the recovered material consists of gray silt with dark-gray silty clay laminae 0.5-2 mm thick. These sediments contain relatively scarce organic content, in places found in small lenses 2 to 4 mm thick, whereas scattered patches of macrophytes remains, Gyrogonites of Characeans and terrestrial plant are found at depths of 48 and 40 cm. Between these depths, sand contents average is 27.4 %. A peak in sand/silt ratio is found at 62-64 cm depth. From 38 to 4 cm, the succession consists of sandy silt and silt laminae 1 to 2.5 mm thick. The abundance of lenses and layers of vegetal remains give a dark-gray color to these sediments. Abundant Gyrogonites of Characeans are found in two organic layer at depths of 24 and 8 cm.

Core CF-10 was retrieved at a water depth of 51 m and recovered 95 cm of lacustrine deposits (Fig. 4). The examined sediment consists of interlaminated clayey silt and sandy silt, forming planar lamination. An upward increase in sand/silt ratio is found starting from 64 cm depth.

1  
2  
3  
4  
5  
6  
7  
8  
9  
10  
11  
12  
13  
14  
15  
16  
17  
18  
19  
20  
21  
22  
23  
24  
25  
26  
27  
28  
29  
30  
31  
32  
33  
34  
35  
36  
37  
38  
39  
40  
41  
42  
43  
44  
45  
46  
47  
48  
49  
50  
51  
52  
53  
54  
55  
56  
57  
58  
59  
60  
61  
62  
63  
64  
65

In detail, from the bottom up to 64 cm, the sediment consists of dark-gray silt with black silty clay laminae 0.5-2 mm thick, apart from the lowermost part of the core (95-85 cm), where organic matter have been smeared during the core cutting, creating blackish waves. This is observed also in the 48-50 cm interval. Layers with abundant vegetal remains are present from depth of 64 cm to the top of the core. In particular, a 8 mm thick layer with abundant Characeans Gyrogonites is revealed at 30 cm depth. Other layers with abundant vegetal remains, containing some Characeans Gyrogonites, are up to 3-5 mm thick and are found at depths of 28, 20, 10 and 8 cm. Vegetal remains are present also in the upper 12 cm the core. Core CF-10 presents the lowest amount of sand, which progressively increases toward the top. The highest sand concentration, 32 wt. %, corresponds to an organic layer between 29 and 32 cm depth.

Core CF-F was retrieved at a water depth of 36 m and recovered 83 cm of lacustrine deposits (Fig. 5). The examined sediment consists of interlaminated clayey silt and sandy silt, forming planar lamination. An upward increase in sand/silt ratio is found starting from 64 cm depth. In detail, from the bottom up to 64 cm, the sediment is composed of dark-gray silt with black silty clay laminae 0.5-2 mm thick. This interval contains terrestrial plant remains that consist of wood fragments with sizes from 2 to 6 mm and seeds found in two thicker layers 14 and 5 mm thick, at depths of 75 cm and 66 cm, respectively. From 64 cm to the top of the core the sediment consists of gray sandy silt with dark-gray silt laminae, 1 to 2.5 mm thick. It contains terrestrial plant and macrophytes remains, 2-3 mm thick.

#### 4.2. Total Organic Carbon and Nitrogen analyses

The concentration of Total Organic Carbon (TOC) in a lake is fundamental for characterizing the abundance of organic matter in the sediments (Meyers and Teranes, 2001; Meyers, 2003), as its variation and distribution serve to trace its provenance and the mechanism of sedimentation. The Carbon/Nitrogen (C/N) ratio allows to differentiate between sources of organic matter in lake sediments, whether lacustrine or terrestrial (Meyers and Lallier-Vergès, 1999; Meyers and Teranes, 2001; Meyers, 2003). Lacustrine algae are cellulose-poor and protein-rich with low C/N values, usually between 4 and 10, while terrestrials plants are cellulose-rich and protein-poor, providing higher C/N values that could reach a range between 17 and 42 (Meyers, 2003). TOC contents (percentage of dry weight) and C/N ratio values obtained from the analysis of the four cores are presented in Figs. 2, 3, 4 and 5.

In core CF-3A, TOC content ranges between 0.9 and 1.3 %, with an increment at the top of the sample, where it reaches a value of 1.6 % dwt. Overall, TOC shows an increase upward (Fig. 2). The C/N values are more or less constant, ranging from 11 to 14. To note that the lowest values of

1 TOC (0.9 %) and C/N ratio (10) are found at 8 cm depth (Fig. 2). Core CF-1B shows TOC values  
2 ranging between 1.2 and 1.7 %, slightly higher than in the previous core. Overall, TOC increases  
3 upward up to 15 cm depth and then decreases (Fig. 3). The C/N ratio is between 14 and 16, and  
4 only at the top it decreases to 12 (Fig. 3). In core CF-10, TOC is usually lower than in cores CF-3A  
5 and CF-1B, and shows an overall upward increase (Fig. 4). Very low TOC values, 0.5 to 0.7 %,   
6 are reported in the lower part of the core, from 74 to 93 cm depth, while a value of 0.8 % is found at  
7 40 and 20 cm depth. Along the rest of the core, the value of TOC is in the range 1.1 - 2.1% dwt, the  
8 latter corresponding to a layer with a high concentration of vegetable remains at 29 cm depth. C/N  
9 ratios are low (8-11) from the bottom to 74 cm depth, while are higher (12-18) from 74 cm to the  
10 top, with the exception of a ratio equal to 10 at 20 cm depth. This is the only core where it is  
11 possible to find C/N values below 10, which is indicative of poor to absent terrestrial contribution  
12 (Fig. 4). The first peak in both C/N and TOC in core CF-10 is recorded at 64 cm depth (Fig. 4).  
13 TOC content in core CF-F does not show relevant variability, being comprised between 2 and 2.3 %  
14 dwt (Fig. 5). Slightly lower values are recorded towards the bottom (1.6 % dwt; 48 cm depth) and at  
15 the top (1.9 % dwt, 1 cm depth). Notably, these higher concentrations are determined by a higher  
16 volume of organic matter in the sediment. Similarly, C/N ratio variations are limited and comprised  
17 between 14 and 16 (Fig. 5).

### 31 4.3. Age estimation

32 The approximate time interval represented by the core sediments was estimated by counting the  
33 light/dark coupled laminae in the cores, according to the method applied by Waldmann et al. (2008)  
34 for cores recovered in the central part of the Lago Fagnano. The laminae counts suggest a mean  
35 sedimentation rate in the order of 1 cm/yr, suggesting that the studied core sediments accumulated  
36 during the last century.

### 45 4.4. *Testate amoebae* distribution in the cores

46 Species diversity of the *testate amoebae* is generally low, with 12 taxa identified from the four  
47 examined cores (Fig. 6 and Table 1, 2, 3 and 4). Preservation of tests was generally good in all the  
48 cores.

#### 54 4.4.1. Core CF-3A

55 Only four species, *Centropyxis elongata*, *Diffflugia oblonga* “oblonga”, *Diffflugia protaeiformis*  
56 “amphoralis” and *Diffflugia viscidula*, are present in all the collected samples (Table 1).

1 The relative abundance of *D. protaeiformis* “amphoralis” is present in a range of 25.0 to  
2 36.4 % from the bottom of the core up to 60 cm depth and then reaches a maximum value of 43.6 %  
3 at 56 cm depth. Subsequently it decreases at 40 cm with a minimum of 9.5 %, maintaining values  
4 below 16.1 % up to 28 cm. A trend of increasing values of its relative abundance is seen in the  
5 upper part of the core from 20.9 % at 24 cm up to 34.6 % at the top of the core. The abundance of  
6 *D. oblonga* “oblonga” is found below 17.2 % from the bottom of the core up to 44 cm depth,  
7 including a lower range of 3.7 to 6.8% from 80 to 56 cm, and subsequently increases to a maximum  
8 value of 28.6 % at 40 cm. At 28 cm there is another peak of 25%, followed by a range of 23.1 to  
9 13.5 % up to the top of the core. Similarly, the relative abundance of *Diffflugia oblonga* “linearis”  
10 presents a peak at 40 and 36 cm depth, although its overall abundance is very low, and the *Diffflugia*  
11 *oblonga* “glans” is mainly present between 40 and 20 cm depth. The relative abundance of *D.*  
12 *viscidula* decreases at 40 and 20 cm depth with values of 4.8 and 3.9 %, respectively. There is a  
13 trend of decreasing values from 16.7 to 7.4% of the relative abundance of *C. elongata* from the  
14 bottom to a depth of 72 cm. Above this depth its abundance is similar, showing slight fluctuations.  
15 *Lagenodiffflugia vas* appears at 64 cm depth and is always present in the upper part of the core,  
16 showing relative abundances between 12.9 and 30.8 %. Among the other taxa, *Diffflugia urceolata*  
17 “lageniformis” is mainly present below 40 cm depth and is very low between 32 and 28 cm depth.  
18 The relative abundance of *Cyclopyxis kahli* is low and it is present from 80 to 60 cm and from 16 to  
19 8 cm depth. Similarly, the abundance of *Diffflugia globulus* is very low and is mainly present in the  
20 lower part of the core and at 28 cm depth. *Centropyxis constricta* “constricta” shows a decreasing  
21 trend from the bottom towards the top of the core and disappears at 16 cm depth.  
22  
23  
24  
25  
26  
27  
28  
29  
30  
31  
32  
33  
34  
35  
36  
37  
38  
39

#### 40 4.4.2. Core CF-1B

41 As in the previous core, only four species *C. elongata*, *D. oblonga* “oblonga”, *D. protaeiformis*  
42 “amphoralis” and *D. viscidula* are present in all the collected samples (Table 2). The most common  
43 taxon is *D. protaeiformis* “amphoralis”, with slightly variations from the bottom to 24 cm depth,  
44 with a peak of 24.1% at a depth of 44 cm. Above 20 cm up to the top of the core there is an abrupt  
45 increase of its abundance ranging from 30.0 to 34.8 %. The second most abundant species is *D.*  
46 *oblonga* “oblonga”. Two intervals may be distinguished, a low abundance from the bottom of the  
47 core up to 48 cm with a range of 8.7 to 14.3 % and a higher abundance from 44 cm up to the top,  
48 ranging between 16.0 to 24.5 %. The distribution of *D. viscidula* is preserved with slightly  
49 variations between 11.1% and 18.2% along the core, with a peak of 20.3% at 50 cm depth and two  
50 minimal values of 7.7% and 8.2% at 32 and 20 cm respectively. *Centropyxis constricta* “constricta”  
51 and *C. elongata* are decreasing in abundance towards the top of the core, showing a reduction trend  
52  
53  
54  
55  
56  
57  
58  
59  
60  
61  
62  
63  
64  
65



1 from the bottom to the top. As in core CF-3A, *L. vas* appears at 64 cm depth with a low abundance  
2 that maintains also at 62 cm, and subsequently shows an increasing trend with slight variations up  
3 to the top of the core. *Cyclopyxis kahli*, *D. globulus*, *D. oblonga* “glans”, *D. oblonga* “linearis” and  
4 *D. urceolata* “lageniformis” occur discontinuously along the core. They are usually absent near the  
5 top of the core.  
6  
7

#### 8 9 10 11 4.4.3. Core CF-10

12 This core contains the highest number of taxa (Table 3). In fact, besides the species mentioned  
13 above for the previous cores, in this core occur the *D. oblonga* “tenuis”, although it is present only  
14 from 42 to 20 cm depth. The testate amoebae in this core are found only above 62 cm depth.  
15  
16

17 *Centropyxis constricta* “constricta” and *C. elongata* are the most abundant species. Two  
18 intervals may be distinguished, one of higher abundance: from 62 cm to 36 cm or 40 cm for both  
19 taxa respectively; and another with lower abundance up to the top of the core. *Cyclopyxis kahli*, *D.*  
20 *oblonga* “glans” and *D. urceolata* “lageniformis” are present in very low amounts (Table 3).  
21 *Diffflugia oblonga* “oblonga” and *L. vas* appear at 36 cm whereas *D. viscidula* occurs at 32 cm depth  
22 and the relative abundance of these three taxa show a discontinuous trend. *Diffflugia globulus*, *D.*  
23 *oblonga* “linearis” and *D. protaeiformis* “amphoralis” also show irregular or discontinuous trend,  
24 with relative abundance values well below 18.8% (registered at 8 cm depth for *D. protaeiformis*  
25 “amphoralis”), except for a peak of 22.4 % at 42 cm depth for *D. globulus*.  
26  
27  
28  
29  
30  
31  
32  
33  
34  
35

#### 36 4.4.4. Core CF-F

37 Five species, in particular *C. constricta* “constricta”, *C. elongata*, *D. globulus*, *D. oblonga*  
38 “oblonga”, and *D. protaeiformis* “amphoralis”, are present in all the collected samples (Table 4).  
39 The most abundant species is *D. oblonga* “oblonga”, the distribution of which shows a trend similar  
40 to that of TOC (Fig. 5). The second most abundant species is *D. protaeiformis* “amphoralis”,  
41 showing an increase from 48 cm depth to the top. *Centropyxis constricta* “constricta”, shows a  
42 variable abundance, being comprised between 13.6 and 16.7 %, with a marked decrease at 64 cm  
43 depth, where its abundance is 10.7 %. The abundance of *C. elongata* decreases from the bottom  
44 (about 10.5 %) to 48 cm depth (6.3 %), and then increases up to 11.1 %. *Diffflugia globulus* shows  
45 variable abundances and is less frequent at 48 cm depth. *Lagenodifflugia vas* appears at 64 cm (3.6  
46 %) and increases up to 10.4 % at 48 cm depth. *Diffflugia oblonga* “glans”, *D. oblonga* “linearis” *D.*  
47 *oblonga* “tenuis”, *D. urceolata* “lageniformis” and *D. viscidula* occur discontinuously along the  
48 core. Among these, *D. oblonga* “tenuis” is the only species present at the top of the core. As in core  
49  
50  
51  
52  
53  
54  
55  
56  
57  
58  
59  
60  
61  
62  
63  
64  
65

1  
2  
3  
4  
5  
6  
7  
8  
9  
10  
11  
12  
13  
14  
15  
16  
17  
18  
19  
20  
21  
22  
23  
24  
25  
26  
27  
28  
29  
30  
31  
32  
33  
34  
35  
36  
37  
38  
39  
40  
41  
42  
43  
44  
45  
46  
47  
48  
49  
50  
51  
52  
53  
54  
55  
56  
57  
58  
59  
60  
61  
62  
63  
64  
65

CF-3A, *L. vas* appears at 64 cm depth with a low abundance that maintains also at 62 cm, and subsequently shows an increasing trend with slight variations up to the top of the core.

## 5. Discussion

From the studied sedimentary cores collected in the eastern Lago Fagnano, it is possible to discriminate between the two cores that are closer to the coast (i.e., the northern cores CF-3A and CF-1B) and those that are more distant (i.e., the southern cores CF-F and CF-10) (Fig. 1B). The latter show a very similar trend, with an average upward sand content increase (Figs. 4 and 5). The northern cores show a similarity in their lowest part, with a peak of the sand/(silt + clay) ratio at 62-64 cm depth (Figs. 2 and 3). A minor sand content increment at this depth is found also in cores CF-F and CF-10 (Figs. 4 and 5). An average upward sand content increment above 50 cm depth characterizes also core CF-1B (Fig. 3), whereas this increment does not occur in core CF-3A (Fig. 2).

In general, TOC content shows an overall slight upward increment in cores CF-3A, CF-1B and CF 10, with a decrease in the uppermost part of core CF-1B, whereas a clear trend is not apparent in core CF-F (Figs. 2-5). The C/N ratio, instead, does not show a trend in the cores except in core CF-10, where it exhibits an overall upward increment, although discontinuous (Figs. 2-5). It is noteworthy that core CF-10 records the first input of terrestrial material at 64 cm depth, as the C/N ratio increases from about 8 to 15, and this is accompanied by an increment in TOC values at the same depth (Fig. 4).

A positive correlation is found between the percentage abundance of testate amoebae species indicative of abundant organic matter, such as *D. oblonga* “oblonga” and *D. protaeiformis* “amphoralis” (Collins et al., 1990; Lorencová, 2009), and TOC content. In fact, the overall upward TOC increment in cores CF-3A, CF-1B and CF-10 is accompanied by an overall upward increment of these species, although less evident in core CF-3A (Figs. 2-4). In contrast, a negative correlation is found between the percentage abundance of testate amoebae species that tolerate a low organic matter content, such as *C. constricta* “constricta” and *D. urceolata* “lageniformis” (Patterson et al., 1985, 1996), and TOC in the same cores, excluding the lower part of core CF-10 that does not contain testate amoebae (Figs. 2-4). The same trend is exhibited also by *C. elongata* (Fig. 2-4). *Centropyxis constricta* “constricta” and *D. urceolata* “lageniformis”, however, are also characteristic of cold climate conditions (Collins et al., 1990; Patterson et al., 1996). Tacking into account that the temperature of the study area rose during the last century (Rabassa, 2010), the observed upward decrease in percentage abundance of these two species, and their disappearance in

1 the upper part of almost all cores (Figs. 2-5), might be related to either overall warming or TOC  
2 content trend, or both. Also *D. globulus* is an indicator of cold climate (Schönborn, 1984; Collins et  
3 al., 1990; Asioli et al., 1996) and this species is more abundant in the lower part of all cores  
4 excepting for core CF-10 (Figs. 2-5).  
5

6  
7 The lack of a clear trend in TOC content in core CF-F is reflected by a relatively erratic  
8 abundance of *D. oblonga* “oblonga” and *C. constricta* “constricta”, whereas the percentage  
9 abundance of *D. protaeiformis* “amphoralis” overall increases upward, as observed in the other  
10 cores, and that of *D. urceolata* “lageniformis” decreases upward (Fig. 5). The reason for the upward  
11 increase of percentage abundance *D. protaeiformis* “amphoralis” in core CF-F is unclear, whereas  
12 the upward decrease in the same core of *D. urceolata* “lageniformis”, as well as of *D. globulus*,  
13 probably reflects the overall temperature increase during the last century. All these evidence suggest  
14 that, between the two species that record changes of both TOC content and temperature, *C.*  
15 *constricta* “constricta” is probably more sensitive to TOC, and less sensitive to temperature changes  
16 than *D. urceolata* “lageniformis”, as only the latter shows an upward decrease of percentage  
17 abundance in core CF-F (Fig. 5). In contrast to the species mentioned above, *C. kahli*, *D. oblonga*  
18 “glans”, *D. oblonga* “linearis”, *D. viscidula* and *L. vas* do not exhibit a clear relationship with TOC  
19 content (Figs. 2-5), possibly due to either a minor sensitivity to environmental changes or a more  
20 complex sensitivity to a combination of several variables.  
21

22  
23 The reason for the lack of testate amoebae in the lower part of core CF-10 is unclear. Some  
24 testate amoebae species (*C. constricta* “constricta”, *C. elongata*, *C. kahli*, *D. globulus*, *D. oblonga*  
25 “linearis”, *D. protaeiformis* “amphoralis”) appear at 62 cm from the top, and both *C. constricta*  
26 “constricta” and *C. elongata* dominate in the central part of this core (Fig. 4). In cores CF-1B and  
27 CF-3A (Figs. 2 and 3), a greater concentration of sand is recorded just at depth of 62-64 cm,  
28 whereas a minor increase in sand content is found in core CF-F (Fig. 5). Taking into account a mean  
29 sedimentation rate of 1 cm/yr for the studied sediments (see Section 4.3), the layer at 62-64 cm  
30 depth accumulated about 62-64 years ago. This marked change in the cores might be related to a  
31 large amount of both sand and suspended organic material made available just after the earthquake  
32 event occurred on 1949 (Lomnitz, 1970) and transported far from the shoreline by currents.  
33

34  
35 This textural variation in sediment cores, which occurred just after the earthquake as a  
36 consequence of a change in the river input (increment in the sand content at a 62-64 cm depth in the  
37 cores), has also had an influence on the distribution and abundance of testate amoebae. This is seen  
38 especially in cores CF-10 and CF-F, which are the most southern and more distant from the coast of  
39 Lago Fagnano, while in the two cores CF-3A and CF-1B this variation in the distribution of taxa  
40  
41  
42  
43  
44  
45  
46  
47  
48  
49  
50  
51  
52  
53  
54  
55  
56  
57  
58  
59  
60  
61  
62  
63  
64  
65

1  
2  
3  
4  
5  
6  
7  
8  
9  
10  
11  
12  
13  
14  
15  
16  
17  
18  
19  
20  
21  
22  
23  
24  
25  
26  
27  
28  
29  
30  
31  
32  
33  
34  
35  
36  
37  
38  
39  
40  
41  
42  
43  
44  
45  
46  
47  
48  
49  
50  
51  
52  
53  
54  
55  
56  
57  
58  
59  
60  
61  
62  
63  
64  
65

was milder (see Figs. 2 to 5). This might reflect a particular sensitivity of this species to the seismic event.

Cluster analyses performed on testate amoebae assemblages, has shown that taxa can be grouped into three clusters (Fig. 7). The first cluster is characterized by the presence of *D. oblonga* “tenuis”, *D. oblonga* “glans” and *D. oblonga* “linearis”. The second cluster is represented by *C. kahli*, *D. globulus* and *D. urceolata* “lageniformis”. The third one collects all the other species. The first and the second cluster are represented by species that are usually present in low amounts and where species are alternatively present, i.e. when *C. kahli*, *D. globulus* and *D. urceolata* are present *D. oblonga* “tenuis”, *D. oblonga* “glans” and *D. oblonga* “linearis” are missing and *vice versa*.

## 6. Conclusions

The conclusions of this study can be summarized as follows:

- (1) Here is reported the first findings of testate amoebae in the sediments of four gravity cores (CF-3A, CF-1B, CF-10, CF-F), acquired in the eastern part of the Lago Fagnano (Tierra del Fuego) at latitude of 54° S. This is the first evidence of these microorganism at those southern latitudes.
- (2) Twelve taxa have been recognized. Among them *C. constricta* “constricta”, *C. elongata*, *D. globulus*, *D. oblonga* “oblonga”, and *D. protaeiformis* “amphoralis” are always present. *D. oblonga* “tenuis” is present only in the southern cores, while *C. kahli* misses in the southernmost core.
- (3) An increase in sand/silt ratio at 62-64 cm depth is recognizable in all cores. Taking into account a mean sedimentation rate of 1 cm/yr for the studied sediments, the event found at 62-64 cm depth is inferred to be related to both sand and suspended organic material made available just after a 7.5 magnitude earthquake occurred on 1949, which severely modified the morphology of the eastern shoreline of the lake.
- (4) The content of Total Organic Carbon (TOC) shows an overall, although discontinuous, upward increase in some cores (CF-3A, CF-1B and CF 10); The Carbon/Nitrogen (C/N) ratio, instead, does not show a clear trend excepting in core CF-10, where it exhibits an overall upward increase with values lower than about 10 (i.e. lacustrine) below 64 cm and higher than 11 upwards (i.e. terrestrial).
- (5) Some testate amoebae show a relation with the variations of TOC. In particular, where TOC displays an overall increase, the percentage abundance of *D. oblonga* “oblonga” and *D. protaeiformis* “amphoralis” tend to increase, whereas that of *C. constricta* “constricta” and *D. urceolata* “lageniformis” dent to decrease. The latter two species, together with *D. globulus*, are also known to be characteristic of cold climate, and their percentage abundance exhibits a

1 marked upward decrease, very probably related to the overall increase of the mean temperature  
2 experienced by the study area in the last century.

3 (6) The appearance and disappearance of certain species as well as the change in sand/silt ratio  
4 found at 62-64 cm depth in the cores suggest that the earthquake event produced some  
5 environmental changes in all locations of the eastern Lago Fagnano.  
6  
7  
8  
9

10  
11  
12 **Acknowledgments.** Presented data have been acquired in the frame of an Italian-Argentinean  
13 scientific project funded by the Italian Ministry of Foreign Affairs, and jointly coordinated by the  
14 *Istituto Nazionale di Oceanografia e di Geofisica Sperimentale (OGS)* of Trieste and the *Instituto*  
15 *de Geofisica "D. Valencio"* of the *Universidad de Buenos Aires*. Horacio Lippai and Maurizio  
16 Grossi have contributed to data acquisition in the field. Francesca Vita kindly assisted at SEM. We  
17  
18  
19  
20  
21  
22  
23  
24  
25  
26  
27  
28  
29  
30  
31  
32  
33  
34  
35  
36  
37  
38  
39  
40  
41  
42  
43  
44  
45  
46  
47  
48  
49  
50  
51  
52  
53  
54  
55  
56  
57  
58  
59  
60  
61  
62  
63  
64  
65

deeply thanks Andrea Bussani for carrying out the statistical analysis used in this study.

## References

- Asioli, A., Medioli, F. S., Patterson, R. T., 1996. Thecamoebians as a tool for reconstruction of paleoenvironments in some Italian lakes in the foothills of the southern Alps (Orta, Varese and Candia). *Journal of Foraminiferal Research* **26**, 248-263.
- Booth, R. K., 2002. Testate amoebae as paleoindicators of surface-moisture changes on Michigan peat lands: modern ecology and hydrological calibration. *Journal of Paleolimnology* **28**, 329-348.
- Burbige, S. M., Schöder-Adams, C. J., 1998. Thecamoebians in Lake Winnipeg: a tool for Holocene paleolimnology. *Journal of Paleolimnology* **19**, 309-328.
- Bray, J. R. Curtis, J. T., 1957. An ordination of upland forest communities of southern Wisconsin. *Ecological Monographs* **27**, 325-349.
- Costa, C. H., Smalley, R., Schwartz, H. D., Ellis, M., Ahumada, E. A., Velasco, M. S., 2006. Paleoseismic observation of an onshore transform boundary: the Magallanes-Fagnano fault, Tierra del Fuego, Argentina. *Revista de la Asociación Geológica Argentina* **61**, 647-657
- Charman, D. J., Roe W.R., Gehrels, W. R., 1998. The use of testate amoebae in studies of sea-level change: a case study from the Taf Estuary, south Wales, UK. *Holocene* **8**, 209-218.
- Charman, D. J., Blundell, A., Accrotelm Members, 2006. A new European testate amoebae transfer function for palaeohydrological reconstruction on ombrotrophic peatlands. *Journal of Quaternary Science* **22**, 209-221.



- 1 Charman, D. J., 2001. Biostratigraphic and palaeoenvironmental applications of testate amoebae.  
2 *Quaternary Science review* **20**, 1753-1764.
- 3 Collins, E. S., McCarthy, F. M. G., Medioli, F. S., Scott, D. B., Honig, C. A., 1990. Biogeographic  
4 distribution of modern thecamoebians in a transect along the eastern North American coast.  
5 In: *Paleoecology, Biostratigraphy, Paleoceanography and Taxonomy of Agglutinated*  
6 *Foraminifera*. Hemleben C, Kaminski M. A, Kuhnt W, Scott D. B, (eds). NATO Advanced  
7 Study Institute Series. Serie C: Mathematical and Physical Science **327**, 783-791.
- 8 Coronato, A., Seppälä, M., Ponce, J. F., Rabassa, J., 2009. Glacial geomorphology of the  
9 Pleistocene Lake Fagnano ice lobe, Tierra del Fuego, southern South America.  
10 *Geomorphology* **112**, 67-81.
- 11 Esteban, F. D., Tassone, A., Lodolo, E., Menichetti, M., Lippai, H., Waldmann, N., Darbo, A.,  
12 Baradello, L., Vilas, J. F., 2014. Basement geometry and sediment thickness of Lago Fagnano  
13 (Tierra del Fuego). *Andean Geology* **41**, 293-313.
- 14 Fernandez, L. D., Zapata, J., Meisterfeld, R., 2012. First records and community pattern of  
15 Arcellinida inhabiting a pristine and remote island from Southeastern Pacific, Chile. *Acta*  
16 *Protozoologica* **51**, 139-154.
- 17 Green, J., 1975. Fresh water ecology in Mato Grosso, Central Brazil IV: associations of testate  
18 rhizopoda. *Journal of Natural History* **9**, 545-560.
- 19 Haman, D., Kohl, B., 1994. A thecamoebians assemblage from Lake Cocococha, Tambopata  
20 Reserve, Madre de Dios Province, Southeastern Peru. *Journal of Foraminiferal Research* **24**,  
21 226-232.
- 22 Kliza, D. A., Schöder-Adams, C. J., 1999. Holocene thecamoebians in freshwater lakes on Bylot  
23 Island, Northwest Territories *Journal of Foraminiferal Research* **29**, 26-39.
- 24 Lodolo, E., Menichetti, M., Tassone, A., Geletti, R., Sterzai, P., Lippai, H., Hormaechea, J.-L.  
25 2002. Researchers target a continental transform fault in Tierra del Fuego. *EOS, Trans.,*  
26 *AGU* **83**, 1-5.
- 27 Lodolo, E., Menichetti, M., Bartole, R., Ben-Avraham, Z., Tassone, A., Lippai, H., 2003.  
28 Magallanes-Fagnano continental transform fault (Tierra del Fuego, southernmost South  
29 America). *Tectonics* **22**, DOI: 10.1029/2003TC001500.
- 30 Lodolo, E., Lippai, H., Tassone, A., Zanolla, C., Menichetti, M., Hormaechea J.-L., 2007. Gravity  
31 map of the Isla Grande de Tierra del Fuego, and morphology of Lago Fagnano. *Geologica*  
32 *Acta* **4**, 307-314.

- 1  
2  
3  
4  
5  
6  
7  
8  
9  
10  
11  
12  
13  
14  
15  
16  
17  
18  
19  
20  
21  
22  
23  
24  
25  
26  
27  
28  
29  
30  
31  
32  
33  
34  
35  
36  
37  
38  
39  
40  
41  
42  
43  
44  
45  
46  
47  
48  
49  
50  
51  
52  
53  
54  
55  
56  
57  
58  
59  
60  
61  
62  
63  
64  
65
- Lodolo, E., Baradello, L., Darbo, A., Caffau, M., Tassone, A., Lippai, H., Lodolo, A., De Zorzi, G., Grossi, M., 2012. Occurrence of shallow gas in easternmost Lago Fagnano (Tierra del Fuego). *Near Surface Geophysics* **10**, 161-169.
- Lomnitz, C., 1970. Major earthquakes and tsunamis in Chile during the period 1535 to 1955. *Geologische Rundschau* **59**, 938-960.
- Lorencová, M., 2009. Thecamoebians from recent lake sediments from the Šumava Mts, Czech Republic. *Bulletin of Geosciences* **84**, 359-376.
- McCarthy, F. M. G., Collins, E. S., McAndrews, J. H., Kerr, H. A., Scott, D. B., Medioli F. S., 1995. A comparison of postglacial arcellacean (thecamoebians) and pollen succession in Atlantic Canada, illustrating the potential of arcellaceans for paleoclimatic reconstruction. *Journal of Paleontology* **69**, 980-993.
- Medioli, F. S., Scott, D. B., Collins, E. S., McCarthy F. M. G., 1990. Fossil thecamoebians: present status and prospects for the future. In: *Paleoecology, Biostratigraphy, Paleoceanography and Taxonomy of Agglutinated Foraminifera*. Hemleben, C, Kaminski. M. A., Kuhnt, W, Scott, D. B. (eds). NATO Advanced Study Institute Series. Serie C: Mathematical and Physical Science **327**, 813-840.
- Medioli, F. S., Scott, D. B., 1988. Lacustrine thecamoebians (mainly arcellaceans) as potential tools for palaeolimnological interpretations. *Palaeogeography, Palaeoclimatology, Palaeoecology* **62**, 361-386.
- Menichetti, M., Lodolo, E., Tassone, A., 2008. Structural geology of the Fuegian Andes and Magallanes fold-and-thrust belt - Tierra del Fuego Island. *Geologica Acta* **6**, 19-42.
- Meyers, P. A., 2003. Applications of organic geochemistry to paleolimnological reconstructions: a summary of example from the Laurentian Great Lake. *Organic Geochemistry* **34**, 261-289.
- Meyers, P. A., Lallier-Vergés, E., 1999. Lacustrine sedimentary organic matter records of Late Quaternary paleoclimates. *Journal of Paleolimnology* **21**, 345-372.
- Meyers, P.A., Teranes, J. L., 2001. Sediment organic matter. In: *Tracking environmental change using lake sediments: Physical and geochemical methods*. Last, W.M., Smol, J. P. (eds). Kluwer Academic Publisher. Netherlands **2**, 239-270.
- Moy, C. M., Dunbar, R. B., Guilderson, T.P., Waldmann, N., Mucciarone, D. A., Recasesn, C., Aritztegui, D., Austin, J. A., Anselmetti, F. S., 2011. A geochemical and sedimentary record of high southern latitude Holocene climate evolution from Lago Fagnano, Tierra del Fuego. *Earth and planetary Science Letters* **302**, 1-13.
- Nieuwenhuize, J., Maas, Y. E. M., Middelburg, J. J., 1994. Rapid analysis of organic carbon and nitrogen in particulate materials. *Marine Chemistry* **45**, 217-224.

- 1  
2  
3  
4  
5  
6  
7  
8  
9  
10  
11  
12  
13  
14  
15  
16  
17  
18  
19  
20  
21  
22  
23  
24  
25  
26  
27  
28  
29  
30  
31  
32  
33  
34  
35  
36  
37  
38  
39  
40  
41  
42  
43  
44  
45  
46  
47  
48  
49  
50  
51  
52  
53  
54  
55  
56  
57  
58  
59  
60  
61  
62  
63  
64  
65
- Patterson, R. T., MacKinnon, K. D., Scott, D. B., Medioli, F. S., 1985. Arcellaceans (thecamoebians) in small lake of New Brunswick and Nova Scotia: modern distribution and Holocene stratigraphic change. *Journal of Foraminiferal Research* **15**, 114-137.
- Patterson, R. T., Barker, T., Burbige, S. M., 1996. Arcellaceans (thecamoebians) as proxies of arsenic and mercury contamination in northeastern Ontario Lake. *Journal of Foraminiferal Research* **26**, 172-183.
- Patterson, R. T., Dalby, A. P., Kumar, A., Henderson, L. A., 2002. Arcellaceans (thecamoebians) as indicators of land-use change: settlement history of the Swan Lake area, Ontario as a case study. *Journal of Paleolimnology* **28**, 297-316.
- Patterson, R.T., Kumar, A., 2000. Assessment of arcellacea (thecamoebians) assemblages, species and strains as contaminated indicators in variability contaminated James Lake, northeastern Ontario. *Journal of Foraminiferal Research* **30**, 310-320.
- Patterson, R. T., Kumar, A., 2002. A review of current testate rhizopod (thecamoebians) research in Canada. *Palaeogeography, Palaeoclimatology, Palaeoecology* **180**, 225-251.
- Rabassa, J., 2010. El cambio climático global en la Patagonia desde el viaje de Charles Darwin hasta nuestro días. *Revista de la Asociación Geológica Argentina*. **67**, 139-156.
- Rabassa, J., Coronato, A., Bujaleski, G., Salemme, M., Roig, C., Maglioli, A., Heusser, C., Gordillo, S., Roig, F., Borrromei, A., Quattrocchio, M., 2000. Quaternary of Tierra del Fuego, Southernmost South America; an updated review. *Quaternary International* **68**, 217-240.
- Reinhardt, E. D., Dalby, A.P., Kumar, A., Patterson, R. T. 1998., Utility of arcellacean morphotypic variants as pollution indicators in mine tailing contaminated lakes near Cobalt, Ontario, Canada. *Micropaleontology* **44**, 1-18.
- Richter, A., Hormaechea, J.-L., Dietrich, R., Perdomo, R., Fritsche, M., Del Cogliano, D., Liebsch, G., Mendoza, L., 2010. Lake-level variation of Lago Fagnano, Tierra del Fuego: observations, modelling and interpretation. *Journal of Limnology* **69**, 29-41.
- Schönborn, W. E., 1984. Studies on remains of testacea in cores of the Great Woryty Lake (NE Poland). *Limnologica* **16**, 185-190.
- Scott, D. B., Medioli, F. S., 1983. Agglutinated rhizopod in Lake Erie: modern distribution and stratigraphic implication. *Journal of Paleontology* **57**, 809-820.
- Scott, D. B., Medioli, F. S., Schafer, C. T., 2001. Monitoring in coastal environments using foraminifera and thecamoebian indicators. Cambridge University Press, New York; 177 pp.
- Tassone, A., Lippai, H., Lodolo, E., Menichetti, M., Comba, A., Hormaechea, J. L., Vilas, J. F., 2005. A geological and geophysical crustal section across the Magallanes-Fagnano fault in Tierra del Fuego. *Journal of South American Earth Sciences* **19**, 99-109.

- 1 Tolonen, K., Warner, B. G., Vasander, H., 1992. Ecology of Testaceans in mires southern Finland:  
2 I. Autoecology. *Archiv für Protistenkunde* **142**, 119-138.
- 3 Tuhkanen, S., 1992. The climate of Tierra del Fuego from a vegetation geographical point of view  
4 and its ecoclimatic counterparts elsewhere. *Acta Botanica Fennica* **145**, 1-64.
- 5 Waldmann, N., Ariztegui, D., Anselmetti, F. S., Austin, J. A., Dunbar, R., Moy, C. M., Recasens,  
6 C., 2008. Seismic stratigraphy of Lago Fagnano sediments (Tierra del Fuego, Argentina): a  
7 potential archive of paleoclimatic change and tectonic activity since the Late Glacial.  
8 *Geologica Acta* **6**, 101-110.
- 9 Waldmann, N., Ariztegui, D., Anselmetti, F. S., Austin, J. A., Stern, C., Moy, C. M., Recasens, C.,  
10 Dunbar, R., 2010a. Holocene climatic fluctuations and positioning of the Southern  
11 Hemisphere westerlies in Tierra del Fuego (54° S), Patagonia. *Journal of Quaternary Science*  
12 **25**, 1063-1075.
- 13 Waldmann, N., Ariztegui, D., Anselmetti, F. S., Coronato, A., Austin, J. A., 2010b. Geophysical  
14 evidence of multiple glacier advances in Lago Fagnano (54° S), southernmost Patagonia.  
15 *Quaternary Science Reviews* **29**, 1188-1200.
- 16 Waldmann, N., Anselmetti, F. S., Ariztegui, D., Austin, J. A., Pirouz, M., Moy, C. M., Dunbar, R.,  
17 2011. Holocene mass-wasting events in Lago Fagnano, Tierra del Fuego (54° S): implications  
18 for paleoseismicity of the Magellanes-Fagnano transform fault. *Basin Research* **23**, 171-190.
- 19 Warner, B. G., Charman, D. J., 1994. Holocene change on a peat land in northwestern Ontario  
20 interpreted from testate amoeba (Protozoa) analysis. *Boreas* **23**, 270-279.
- 21 Zanolla, C., Lodolo, E., Lippai, H., Tassone, A., Menichetti, M., Baradello, L., Grossi, M.,  
22 Hormaechea, H. L., 2011. Bathymetric map of Lago Fagnano (Tierra del Fuego Island).  
23 *Bollettino di Geofisica Teorica ed Applicata* **52**, 1-8.
- 24 Zapata, J., Yáñez, M., Rudolph, E., 2008. Tecamebianos (Protozoa: Rhizopoda) de una turbera del  
25 Parque Nacional Puyehue (40°45'S; 72°19'W), Chile. *Gayana* **72**, 9-17.
- 26  
27  
28  
29  
30  
31  
32  
33  
34  
35  
36  
37  
38  
39  
40  
41  
42  
43  
44  
45  
46  
47  
48  
49  
50  
51  
52  
53  
54  
55  
56  
57  
58  
59

60 **FIGURE CAPTIONS**

61  
62  
63  
64  
65

1 **Figure 1.** (A) **Shaded-relief** map showing the general morphology of Tierra del Fuego, crossed by the Magallanes-  
2 Fagnano transform system (MFS). MS indicates the Magallanes Strait. White box indicates the easternmost sector of  
3 the Lago Fagnano where data have been acquired. (B) Google Earth® image of the eastern Lago Fagnano area, with the  
4 location of the gravity cores (black stars) used in this study. Bathymetric contours (in blue) every 10 m (Zanolla et al.  
5 2011). The blue area indicates the lagoon nearby the estuary of the Rio Turbio, and the yellow area indicates the buried  
6 streams of the Rio Tolhuin. (C) Photographs of the **lagoon** at the eastern shoreline of the Lago Fagnano, taken before  
7 the 1949 earthquake, compared with the present-day shoreline (D).

13 **Figure 2.** Lithology, sand/silt percentage, TOC percentage, C/N ratio and relative abundance of **testate amoebae** in core  
14 CF-3A.

18 **Figure 3.** Lithology, sand/silt percentage, TOC percentage, C/N ratio and relative abundance of **testate amoebae** in core  
19 CF-1B.

22 **Figure 4.** Lithology, sand/silt percentage, TOC percentage, C/N ratio and relative abundance of **testate amoebae** in core  
23 CF-10.

27 **Figure 5.** Lithology, sand/silt percentage, TOC percentage, C/N ratio and relative abundance of **testate amoebae** in core  
28 CF-F.

31 **Figure 6.** Scanning electron micrographs of selected **testate amoebae** from the eastern Lago Fagnano: (1) *Centropyxis*  
32 *constricta* “**constricta**” from core CF-3A; (2-3) *Centropyxis constricta* “constricta”, from core CF-10; (4) *Centropyxis*  
33 *elongata* from core CF-F; (5-6) *Centropyxis elongata* ventral and dorsal view from core CF-10, respectively; (7-8)  
34 *Cyclopyxis kahli* from core CF-10; (9-10) *Diffflugia globulus* from core CF-F; (11-12) *Diffflugia oblonga* “glans” from  
35 core CF-1B; (13-14) *Diffflugia oblonga* “oblonga” from core CF-3A and CF-F, respectively; (15-16) *Diffflugia oblonga*  
36 “linearis” from core CF-F and CF-10, respectively; (17-18); *Diffflugia oblonga* “tenuis” from core CF-F and CF-10,  
37 respectively; (19-20) *Diffflugia protaeiformis* “amphoralis” from core CF-3A and CF-F, respectively; (21-22) *Diffflugia*  
38 *urceolata* “lageniformis” from core CF-3A and CF-F, respectively; (23) *Diffflugia urceolata* “lageniformis” from core  
39 CF-10; (24-25) *Diffflugia viscidula* from core CF-3A and CF-1B, respectively; (26-27) *Lagenodiffflugia vas* from core  
40 CF-1B and CF-10, respectively.

47 **Figure 7.** Cluster analysis on testate amoebae. *ccc* = *Centropyxis constricta* “constricta”; *ce* = *Centropyxis elongate*; *ck*  
48 = *Cyclopyxis kahli*; *dg* = *Diffflugia globulus*; *dog* = *Diffflugia oblonga* “glans”; *doo* = *Diffflugia oblonga* “oblonga”; *dol* =  
49 *Diffflugia oblonga* “linearis”; *dot* = *Diffflugia oblonga* “tenuis”; *dpa* = *Diffflugia protaeiformis* “amphoralis”; *dul* =  
50 *Diffflugia urceolata* “lagenifomis”; *dv* = *Diffflugia viscidula*; *lv* = *Lagenodiffflugia vas*.

56 **Table 1.** Percentage abundance and total number of individuals counted of **testate amoebae** in core CF-3A.

58 **Table 2.** Percentage abundance and total number of individuals counted of **testate amoebae** in core CF-1B.



**Table 3.** Percentage abundance and total number of individuals counted of *testate amoebae* in core CF-10.

1  
2  
3  
4  
5  
6  
7  
8  
9  
10  
11  
12  
13  
14  
15  
16  
17  
18  
19  
20  
21  
22  
23  
24  
25  
26  
27  
28  
29  
30  
31  
32  
33  
34  
35  
36  
37  
38  
39  
40  
41  
42  
43  
44  
45  
46  
47  
48  
49  
50  
51  
52  
53  
54  
55  
56  
57  
58  
59  
60  
61  
62  
63  
64  
65

**Table 4.** Percentage abundance and total number of individuals counted of *testate amoebae* in core CF-F.

Figure 1  
[Click here to download high resolution image](#)






Figure 2  
[Click here to download high resolution image](#)



**Figure 3**  
[Click here to download high resolution image](#)






Figure 5  
[Click here to download high resolution image](#)






Figure 6  
[Click here to download high resolution image](#)





Figure 7  
[Click here to download high resolution image](#)

### Cluster Dendrogram



z4  
hclust ("complete")

CORE CF-3A water depth 19.8m												
Sample	CF-3A/1	CF-3A/4	CF-3A/8	CF-3A/12	CF-3A/16	CF-3A/20	CF-3A/24	CF-3A/28	CF-3A/32	CF-3A/36	CF-3A/40	CF-3A/44
Depth cm	1	4	8	12	16	20	24	28	32	36	40	44
<b>Species</b>												
<i>Centropyxis constricta</i> "constricta"	0,0	0,0	0,0	0,0	0,0	4,5	2,3	3,9	5,6	5,1	4,8	9,3
<i>Centropyxis elongata</i>	5,8	11,0	5,6	9,4	7,7	9,0	9,3	13,3	12,7	11,0	9,5	9,9
<i>Cyclopyxis kahli</i>	0,0	0,0	11,1	8,2	7,7	0,0	0,0	0,0	0,0	0,0	0,0	0,0
<i>Diffugia globulus</i>	0,0	0,0	0,0	0,0	0,0	0,0	0,0	1,6	0,0	0,0	0,0	0,0
<i>Diffugia oblonga</i> "glans"	0,0	0,0	0,0	1,2	0,0	14,2	11,6	7,0	11,1	9,3	9,5	2,5
<i>Diffugia oblonga</i> "oblonga"	13,5	17,8	16,7	20,0	23,1	18,7	16,3	25,0	16,7	18,6	28,6	13,0
<i>Diffugia oblonga</i> "linearis"	0,0	0,0	0,0	0,0	0,0	1,3	2,3	7,8	7,1	13,6	14,3	1,9
<i>Diffugia oblonga</i> "tenuis"	0,0	0,0	0,0	0,0	0,0	0,0	0,0	0,0	0,0	0,0	0,0	0,0
<i>Diffugia protaeiformis</i> "amphoralis"	34,6	35,6	27,8	25,9	23,1	27,7	20,9	13,3	14,3	16,1	9,5	24,2
<i>Diffugia urceolata</i> "lageniformis"	0,0	0,0	0,0	0,0	0,0	0,0	0,0	0,8	1,6	0,0	0,0	4,3
<i>Diffugia viscidula</i>	23,1	15,1	11,1	22,4	7,7	3,9	14,0	9,4	16,7	9,3	4,8	14,3
<i>Lagenodiffugia vas</i>	23,1	20,5	27,8	12,9	30,8	20,6	23,3	18,0	14,3	16,9	19,0	20,5
<b>Total number of individuals counted</b>	104	73	108	85	156	155	258	128	126	118	126	161

CORE CF-3A water depth 19.8m												
Sample	CF-3A/50	CF-3A/56	CF-3A/60	CF-3A/62	CF-3A/64	CF-3A/66	CF-3A/72	CF-3A/76	CF-3A/80	CF-3A/84	CF-3A/88	CF-3A/91
Depth cm	50	56	60	62	64	66	72	76	80	84	88	91
<b>Species</b>												
<i>Centropyxis constricta</i> "constricta"	9,2	5,1	7,0	8,0	7,8	9,8	11,1	14,8	9,8	14,1	16,7	17,7
<i>Centropyxis elongata</i>	7,9	7,7	7,4	8,7	7,8	10,2	7,4	10,2	11,8	14,1	16,7	16,1
<i>Cyclopyxis kahli</i>	0,0	0,0	2,8	4,0	3,9	7,3	11,1	6,8	3,9	0,0	0,0	0,0
<i>Diffugia globulus</i>	0,0	0,0	1,4	1,5	0,0	4,9	7,4	4,5	3,9	3,1	0,0	0,0
<i>Diffugia oblonga</i> "glans"	3,3	0,0	0,0	0,0	0,0	0,0	0,0	0,0	0,0	0,0	0,0	0,0
<i>Diffugia oblonga</i> "oblonga"	10,5	5,1	4,7	5,1	3,9	5,3	3,7	6,8	15,7	17,2	8,3	8,1
<i>Diffugia oblonga</i> "linearis"	0,0	0,0	0,0	1,5	0,0	0,0	0,0	0,0	2,0	0,0	0,0	0,0
<i>Diffugia oblonga</i> "tenuis"	0,0	0,0	0,0	0,0	0,0	0,0	0,0	0,0	0,0	0,0	0,0	0,0
<i>Diffugia protaeiformis</i> "amphoralis"	27,0	43,6	36,3	33,1	31,4	35,8	33,3	36,4	27,5	25,0	33,3	33,9
<i>Diffugia urceolata</i> "lageniformis"	3,9	7,7	9,8	8,7	7,8	10,6	7,4	6,8	7,8	9,4	8,3	8,1
<i>Diffugia viscidula</i>	11,2	12,8	13,5	15,3	13,7	16,3	18,5	13,6	17,6	17,2	16,7	16,1
<i>Lagenodiffugia vas</i>	27,0	17,9	17,2	14,2	23,5	0,0	0,0	0,0	0,0	0,0	0,0	0,0
<b>Total number of individuals counted</b>	152	258	215	275	156	246	162	88	51	64	72	62

Table 1. Percentage abundance and total number of individuals counted of testate amoebae in core CF-3A.

Table 2

1  
2  
3  
4  
5  
6  
7  
8  
9  
10  
11  
12  
13  
14  
15  
16  
17  
18  
19  
20  
21  
22  
23  
24  
25  
26  
27  
28  
29  
30  
31  
32  
33  
34  
35  
36  
37  
38  
39  
40  
41  
42  
43  
44  
45  
46  
47  
48  
49  
50  
51  
52  
53  
54  
55  
56  
57  
58  
59  
60  
61  
62  
63  
64  
65

CORE CF-1B water depth 20.4m											
Sample	CF-1B/1	CF-1B/4	CF-1B/8	CF-1B/12	CF-1B/16	CF-1B/20	CF-1B/24	CF-1B/28	CF-1B/32	CF-1B/36	CF-1B/40
Depth cm	1	4	8	12	16	20	24	28	32	36	40
<b>Species</b>											
<i>Centropyxis constricta</i> "constricta"	0,0	0,0	0,0	0,0	4,3	5,4	8,0	8,8	3,8	14,9	16,0
<i>Centropyxis elongata</i>	9,8	11	7,7	11,4	8,7	10,2	12,0	13,5	11,5	11,3	8,0
<i>Cyclopyxis kahli</i>	0,0	0,0	0	4	4,3	5,4	4,0	0,0	3,8	3,6	0,0
<i>Diffugia globulus</i>	0,0	0,0	0,0	0,0	0,0	0,0	0	1	3,8	3,6	12,0
<i>Diffugia oblonga</i> "glans"	0,0	0	0	0	0	0	0	0	0	0	4,0
<i>Diffugia oblonga</i> "oblonga"	19,5	22,2	23,1	23,6	17,4	24,5	24,0	18,8	19,2	19,6	16,0
<i>Diffugia oblonga</i> "linearis"	0,0	0,0	0,0	0,0	0	1,4	4,0	6,5	15,4	0,0	0
<i>Diffugia oblonga</i> "tenuis"	0,0	0	0	0	0	0	0	0	0	0	0
<i>Diffugia protaeiformis</i> "amphoralis"	34,1	33,3	30,8	30,0	34,8	32,7	20,0	21,2	23,1	19,0	16,0
<i>Diffugia urceolata</i> "lageniformis"	0,0	0,0	0,0	0,0	0,0	0,0	0,0	0,6	0,0	0,0	0,0
<i>Diffugia viscidula</i>	13,4	11,1	15,4	17,9	13,0	8,2	12,0	12,9	7,7	13,1	12,0
<i>Lagenodiffugia vas</i>	23,2	22,2	23,1	13,6	17,4	12,2	16,0	16,5	11,5	14,9	16,0
<b>Total number of individuals counted</b>	<b>82</b>	<b>81</b>	<b>117</b>	<b>140</b>	<b>207</b>	<b>147</b>	<b>225</b>	<b>170</b>	<b>234</b>	<b>168</b>	<b>225</b>

CORE CF-1B water depth 20.4m											
Sample	CF-1B/44	CF-1B/48	CF-1B/50	CF-1B/56	CF-1B/60	CF-1B/62	CF-1B/64	CF-1B/66	CF-1B/72	CF-1B/76	CF-1B/80
Depth cm	44	48	50	56	60	62	64	66	72	76	80
<b>Species</b>											
<i>Centropyxis constricta</i> "constricta"	16,0	14,3	17,0	14,0	11,3	16,3	17,4	15,0	14,3	12,5	18,2
<i>Centropyxis elongata</i>	13,2	14,3	15,8	17,5	15,0	11,0	13,0	10,0	11,3	18,8	18,2
<i>Cyclopyxis kahli</i>	0,0	0,0	0,0	0,0	3	7	8,7	8,2	9,1	6	9,1
<i>Diffugia globulus</i>	0,0	0	0	0	0	13	8,7	16,3	7,8	6,3	0,0
<i>Diffugia oblonga</i> "glans"	2,4	8,6	2,5	3,0	7,5	0,0	0	0	0	0,0	0,0
<i>Diffugia oblonga</i> "oblonga"	19,8	11,4	8,7	13,0	11,3	12,5	13,0	11,6	14,3	12,5	9,1
<i>Diffugia oblonga</i> "linearis"	0	2,9	1,7	0,0	0	0	0,0	1,3	0,9	0	0,0
<i>Diffugia oblonga</i> "tenuis"	0	0	0	0	0	0	0	0	0	0	0,0
<i>Diffugia protaeiformis</i> "amphoralis"	24,1	20,0	18,7	19,5	15,0	16,3	17,4	15,5	17,7	18,8	18,2
<i>Diffugia urceolata</i> "lageniformis"	1,4	0,0	2,5	5,5	11,3	7,7	4,3	7,6	9,1	12,5	9,1
<i>Diffugia viscidula</i>	13,7	14,3	20,3	15,5	11,3	13,9	13,0	14,5	15,6	12,5	18,2
<i>Lagenodiffugia vas</i>	9,0	14,3	12,9	12,0	15,0	2,7	4,3	0,0	0,0	0	0,0
<b>Total number of individuals counted</b>	<b>212</b>	<b>315</b>	<b>241</b>	<b>200</b>	<b>234</b>	<b>337</b>	<b>207</b>	<b>380</b>	<b>231</b>	<b>144</b>	<b>99</b>

Table 2. Percentage abundance and total number of individuals counted of testate amoebae in core CF-1B.

Table 3

1  
2  
3  
4  
5  
6  
7  
8  
9  
10  
11  
12  
13  
14  
15  
16  
17  
18  
19  
20  
21  
22  
23  
24  
25  
26  
27  
28  
29  
30  
31  
32  
33  
34  
35  
36  
37  
38  
39  
40  
41  
42  
43  
44  
45  
46  
47  
48  
49  
50  
51  
52  
53  
54  
55  
56  
57  
58  
59  
60  
61  
62  
63  
64  
65

CORE CF-10 water depth 51m													
Sample	CF-10/1	CF-10/4	CF-10/8	CF-10/12	CF-10/16	CF-10/20	CF-10/24	CF-10/29	CF-10/32	CF-10/36	CF-10/40	CF-10/42	CF-10/44
Depth cm	1	4	8	12	16	20	24	29	32	36	40	42	44
<b>Species</b>													
<i>Centropyxis constricta</i> "constricta"	13,0	12,7	15,6	15,0	12,8	16,7	6,3	15,4	14,3	37,5	33,3	28,6	43,2
<i>Centropyxis elongata</i>	8,7	8,0	3,1	6,9	10,3	5,6	12,5	7,7	14,3	16,7	41,7	22,4	32,4
<i>Cyclopyxis kahli</i>	4,3	7,3	6,3	0,0	0,0	5,6	9,4	3,8	7,1	4,2	0,0	0,0	0,0
<i>Diffugia globulus</i>	8,7	10,0	6,3	11,3	7,7	11,1	3,1	11,5	7,1	12,5	8,3	22,4	13,5
<i>Diffugia oblonga</i> "glans"	4,3	9,3	12,5	8,8	2,6	0,0	9,4	7,7	1,8	0,0	0,0	0,0	0,0
<i>Diffugia oblonga</i> "oblonga"	17,4	14,0	12,5	13,8	12,8	11,1	25,0	23,1	10,7	8,3	0,0	0,0	0,0
<i>Diffugia oblonga</i> "linearis"	8,7	10,0	9,4	5,6	0,0	0,0	0,0	0,0	3,6	0,0	8,3	8,2	10,8
<i>Diffugia oblonga</i> "tenuis"	0,0	0,0	0,0	0,0	0,0	11,1	12,5	7,7	5,4	4,2	8,3	18,4	0,0
<i>Diffugia protaeiformis</i> "amphoralis"	8,7	12,0	18,8	13,8	5,1	16,7	9,4	15,4	0,0	4,2	0,0	0,0	0,0
<i>Diffugia urceolata</i> "lageniformis"	0,0	0,0	0,0	1,9	5,1	0,0	6,3	0,0	0,0	0,0	0,0	0,0	0,0
<i>Diffugia viscidula</i>	8,7	7,3	12,5	10,0	23,1	11,1	0,0	0,0	14,3	0,0	0,0	0,0	0,0
<i>Lagenodiffugia vas</i>	17,4	9,3	3,1	13,1	20,5	11,1	6,3	7,7	21,4	12,5	0,0	0,0	0,0
<b>Total number of individuals counted</b>	<b>161</b>	<b>150</b>	<b>224</b>	<b>160</b>	<b>273</b>	<b>126</b>	<b>224</b>	<b>182</b>	<b>392</b>	<b>168</b>	<b>84</b>	<b>49</b>	<b>37</b>

CORE CF-10 water depth 51m													
Sample	CF-10/48	CF-10/52	CF-10/56	CF-10/58	CF-10/60	CF-10/61	CF-10/62	CF-10/63	CF-10/64	CF-10/74	CF-10/79	CF-10/84	CF-10/93
Depth cm	48	52	56	58	60	61	62	63	64	74	79	84	93
<b>Species</b>													
<i>Centropyxis constricta</i> "constricta"	44,4	48,4	40,0	43,4	41,2	40,9	40,0	0,0	0,0	0	0	0	0
<i>Centropyxis elongata</i>	33,3	20,3	33,3	20,8	11,8	29,5	23,3	0,0	0,0	0	0	0	0
<i>Cyclopyxis kahli</i>	0,0	0,0	6,7	7,5	11,8	4,5	8,3	0,0	0,0	0	0	0	0
<i>Diffugia globulus</i>	11,1	4,7	6,7	7,5	11,8	6,8	6,7	0,0	0,0	0	0	0	0
<i>Diffugia oblonga</i> "glans"	0,0	0,0	0,0	0,0	0,0	0,0	0,0	0,0	0,0	0	0	0	0
<i>Diffugia oblonga</i> "oblonga"	0,0	0,0	0,0	0,0	0,0	0,0	0,0	0,0	0,0	0	0	0	0
<i>Diffugia oblonga</i> "linearis"	11,1	17,2	6,7	11,3	17,6	11,4	13,3	0,0	0,0	0	0	0	0
<i>Diffugia oblonga</i> "tenuis"	0,0	0,0	0,0	0,0	0,0	0,0	0,0	0,0	0,0	0	0	0	0
<i>Diffugia protaeiformis</i> "amphoralis"	0,0	9,4	6,7	9,4	5,9	6,8	8,3	0,0	0,0	0	0	0	0
<i>Diffugia urceolata</i> "lageniformis"	0,0	0,0	0,0	0,0	0,0	0,0	0,0	0,0	0,0	0	0	0	0
<i>Diffugia viscidula</i>	0,0	0,0	0,0	0,0	0,0	0,0	0,0	0,0	0,0	0	0	0	0
<i>Lagenodiffugia vas</i>	0,0	0,0	0,0	0,0	0,0	0,0	0,0	0,0	0,0	0	0	0	0
<b>Total number of individuals counted</b>	<b>63</b>	<b>64</b>	<b>105</b>	<b>53</b>	<b>51</b>	<b>44</b>	<b>60</b>	<b>0</b>	<b>0</b>	<b>0</b>	<b>0</b>	<b>0</b>	<b>0</b>

Table 3. Percentage abundance and total number of individuals counted of testate amoebae in core CF-10.

Table 4

1  
2  
3  
4  
5  
6  
7  
8  
9  
10  
11  
12  
13  
14  
15  
16  
17  
18  
19  
20  
21  
22  
23  
24  
25  
26  
27  
28  
29  
30  
31  
32  
33  
34  
35  
36  
37  
38  
39  
40  
41  
42  
43  
44  
45  
46  
47  
48  
49  
50  
51  
52  
53  
54  
55  
56  
57  
58  
59  
60  
61  
62  
63  
64  
65

CORE CF-F water depth 36m													
Sample	CF-F1	CF-F4	CF-F8	CF-F12	CF-F16	CF-F20	CF-F24	CF-F28	CF-F30	CF-F32	CF-F36	CF-F40	CF-F44
Depth cm	1	4	8	12	16	20	24	28	30	32	36	40	44
<b>Species</b>													
<i>Centropyxis constricta</i> "constricta"	16,7	18,6	19,1	13,9	13,6	10,0	16,9	11,5	11,9	14,3	16,7	11,5	10,4
<i>Centropyxis elongata</i>	11,1	13,7	11,7	11,5	9,1	13,6	9,1	11,5	11,4	8,6	11,3	12,0	7,2
<i>Cyclopyxis kahli</i>	0,0	0,0	0,0	0,0	0,0	0,0	0,0	0,0	0,0	0,0	0,0	0,0	0,0
<i>Diffugia globulus</i>	5,6	6,9	6,4	5,7	9,1	10,7	8,4	7,7	8,6	8,6	9,1	5,7	5,0
<i>Diffugia oblonga</i> "glans"	0,0	0,0	0,0	0,0	0,0	0,0	0,0	0,0	0,0	0,0	0,0	0,0	0,0
<i>Diffugia oblonga</i> "oblonga"	27,8	27,5	24,5	33,6	31,8	24,3	27,9	35,5	35,7	28,6	18,3	27,1	19,5
<i>Diffugia oblonga</i> "linearis"	0,0	0,0	2,1	1,6	0,0	4,3	3,9	4,4	3,2	5,7	5,9	4,7	3,6
<i>Diffugia oblonga</i> "tenuis"	11,1	5,9	7,4	6,6	9,1	6,4	2,6	0,0	0,0	0,0	0,0	1,0	5,4
<i>Diffugia protaeiformis</i> "amphoralis"	22,2	23,5	20,2	18,0	18,2	15,0	11,7	10,9	11,4	14,3	17,2	15,6	18,6
<i>Diffugia urceolata</i> "lageniformis"	0,0	0,0	0,0	0,0	0,0	0,0	0,0	3,3	3,8	5,7	8,6	4,7	4,1
<i>Diffugia viscidula</i>	0,0	0,0	0,0	0,0	0,0	5,7	5,8	6,0	6,5	5,7	7,0	6,8	15,4
<i>Lagenodiffugia vas</i>	5,6	3,9	8,5	9,0	9,1	10,0	13,6	9,3	7,6	8,6	5,9	10,9	10,9
<b>Total number of individuals counted</b>	<b>144</b>	<b>102</b>	<b>94</b>	<b>122</b>	<b>176</b>	<b>140</b>	<b>154</b>	<b>183</b>	<b>185</b>	<b>280</b>	<b>186</b>	<b>192</b>	<b>221</b>

CORE CF-F water depth 36m												
Sample	CF-F48	CF-F52	CF-F56	CF-F60	CF-F62	CF-F64	CF-F66	CF-F68	CF-F72	CF-F76	CF-F80	CF-F82
Depth cm	48	52	56	60	62	64	66	68	72	76	80	82
<b>Species</b>												
<i>Centropyxis constricta</i> "constricta"	14,6	14,7	13,9	10,9	11,7	10,7	13,3	13,1	14,4	13,6	15,8	16,4
<i>Centropyxis elongata</i>	6,3	10,8	12,5	11,9	11,7	10,7	12,1	13,1	12,2	12,7	10,5	11,2
<i>Cyclopyxis kahli</i>	0,0	0,0	0,0	0,0	0,0	0,0	0,0	0,0	0,0	0,0	0,0	0,0
<i>Diffugia globulus</i>	4,2	2,5	3,4	3,0	11,7	7,1	15,6	16,2	10,0	10,0	5,3	5,2
<i>Diffugia oblonga</i> "glans"	0,0	0,0	4,3	5,5	3,3	3,6	1,2	0,0	0,0	0,0	0,0	0,0
<i>Diffugia oblonga</i> "oblonga"	22,9	15,7	23,1	27,4	23,9	28,6	27,7	27,3	23,3	30,0	26,3	23,9
<i>Diffugia oblonga</i> "linearis"	4,2	5,4	4,3	10,9	2,3	7,1	0,0	0,0	0,0	0,0	0,0	0,0
<i>Diffugia oblonga</i> "tenuis"	4,2	3,4	3,8	2,0	1,4	3,6	1,7	0,0	0,0	0,0	0,0	0,0
<i>Diffugia protaeiformis</i> "amphoralis"	14,6	16,7	11,1	8,0	9,9	7,1	6,9	9,1	12,2	8,2	10,5	11,2
<i>Diffugia urceolata</i> "lageniformis"	4,2	6,4	3,8	4,0	9,9	7,1	12,7	9,1	10,0	11,8	10,5	9,7
<i>Diffugia viscidula</i>	14,6	16,2	15,4	10,9	12,2	10,7	8,7	12,1	17,8	13,6	21,1	22,4
<i>Lagenodiffugia vas</i>	10,4	8,3	4,3	5,5	1,9	3,6	0,0	0,0	0,0	0,0	0,0	0,0
<b>Total number of individuals counted</b>	<b>384</b>	<b>204</b>	<b>208</b>	<b>201</b>	<b>213</b>	<b>224</b>	<b>173</b>	<b>99</b>	<b>90</b>	<b>110</b>	<b>152</b>	<b>134</b>

Table 4. Percentage abundance and total number of individuals counted of testate amoebae in core CF-F.