

RADIALLY SYMMETRIC SOLUTIONS OF AN ANISOTROPIC MEAN CURVATURE EQUATION MODELING THE CORNEAL SHAPE

CHIARA CORSATO

Università degli Studi di Trieste
Dipartimento di Matematica e Geoscienze
Via A. Valerio 12/1, 34127 Trieste, Italy

COLETTE DE COSTER

Université de Valenciennes et du Hainaut Cambrésis
LAMAV, FR CNRS 2956
Institut des Sciences et Techniques de Valenciennes
F-59313 Valenciennes Cedex 9, France

PIERPAOLO OMARI

Università degli Studi di Trieste
Dipartimento di Matematica e Geoscienze
Via A. Valerio 12/1, 34127 Trieste, Italy

ABSTRACT. We prove existence and uniqueness of classical solutions of the anisotropic prescribed mean curvature problem

$$-\operatorname{div} \left(\nabla u / \sqrt{1 + |\nabla u|^2} \right) = -au + b / \sqrt{1 + |\nabla u|^2}, \quad \text{in } B, \quad u = 0, \quad \text{on } \partial B,$$

where $a, b > 0$ are given parameters and B is a ball in \mathbb{R}^N . The solution we find is positive, radially symmetric, radially decreasing and concave. This equation has been proposed as a model of the corneal shape in the recent papers [13, 14, 15, 18, 17], where however a linearized version of the equation has been investigated.

1. Introduction. This note is devoted to the study of the existence, the uniqueness and the qualitative properties of classical solutions of the anisotropic prescribed mean curvature problem

$$\begin{cases} -\operatorname{div} \left(\frac{\nabla u}{\sqrt{1 + |\nabla u|^2}} \right) = -au + \frac{b}{\sqrt{1 + |\nabla u|^2}}, & \text{in } B, \\ u = 0, & \text{on } \partial B, \end{cases} \quad (1)$$

where $a > 0$ and $b > 0$ are given constants and $B = B(x_0, R)$ is the open ball in \mathbb{R}^N of center x_0 and radius R . This problem has been recently proposed in [13, 14, 15, 18, 17] as a mathematical model for the geometry of the human cornea: we refer to these articles for further references on the subject. However, in all these papers a simplified version of (1) has been investigated, where the curvature operator

$$\operatorname{div}(\nabla u / \sqrt{1 + |\nabla u|^2})$$

2010 *Mathematics Subject Classification*. Primary: 35J93, 35J25; Secondary: 35B07, 35B09, 35A09, 35A02, 35A24, 34A12.

Key words and phrases. Anisotropic prescribed mean curvature equation, Dirichlet boundary condition, radially symmetric solution, positive solution, existence, uniqueness, shooting method.

This paper was written under the auspices and with the support of INdAM-GNAMPA.

The first and the third named authors have been supported by the University of Trieste, in the frame of the 2013 FRA project “Nonlinear Ordinary Differential Equations: Qualitative Theory, Numerics and Applications”.

has been replaced by its linearization Δu around 0. In particular, it has been proved in [14] that, if

$$b \in \left] 0, \frac{3\sqrt{3}}{2} \frac{\sqrt{a}I_0(\sqrt{a})}{I_1(\sqrt{a})(2I_0(\sqrt{a}) - 1)} \right[,$$

with I_n ($n = 0, 1$) the n -order modified Bessel functions of the first kind, and B is a unit disk in \mathbb{R}^2 , then the (physically relevant) problem

$$\begin{cases} -\Delta u = -au + \frac{b}{\sqrt{1 + |\nabla u|^2}}, & \text{in } B, \\ u = 0, & \text{on } \partial B \end{cases}$$

has a unique radially symmetric solution which is the uniform limit of a sequence of successive approximations. We stress that, in the one-dimensional case, these limitations on the parameters have later been removed in [18], where it has also been pointed out the interest of studying the complete model (1).

Like in [6], dealing with the one-dimensional case, we tackle here the fully nonlinear problem (1) and we prove the existence of a unique solution for the whole range of positive parameters a, b and for any radius R . Precisely, we first prove a uniqueness result for a more general problem

$$\begin{cases} -\operatorname{div} \left(\frac{\nabla u}{\sqrt{1 + |\nabla u|^2}} \right) = -au + \frac{b}{\sqrt{1 + |\nabla u|^2}}, & \text{in } \Omega, \\ u = 0, & \text{on } \partial\Omega, \end{cases} \quad (2)$$

where the ball B is replaced by any bounded domain Ω in \mathbb{R}^N with a Lipschitz boundary $\partial\Omega$. Then we establish the existence of a classical radially symmetric solution of (1), by solving the problem

$$\begin{cases} - \left(\frac{r^{N-1} v'}{\sqrt{1 + v'^2}} \right)' = r^{N-1} \left(-av + \frac{b}{\sqrt{1 + v'^2}} \right), & \text{in }]0, R[, \\ v'(0) = 0, \quad v(R) = 0. \end{cases} \quad (3)$$

This radial solution is therefore the unique solution of (1). We also prove that it is positive, radially decreasing and concave. Our result is stated in the following theorem.

Theorem 1.1. *Let $a > 0$ and $b > 0$ be given and let $B = B(x_0, R)$ be the open ball in \mathbb{R}^N of center x_0 and radius R . Then there exists a unique solution $u \in C^2(\bar{B})$ of (1), which in addition satisfies:*

- *there exists a function $v \in C^2([0, R])$ such that $u(x) = v(|x - x_0|)$ for all $x \in \bar{B}$;*
- *$0 < v(r) < b/a$ for all $r \in [0, R[$;*
- *$v'(r) < 0$ for all $r \in]0, R[$;*
- *$v''(r) < 0$ for all $r \in [0, R]$.*

It is well-known that in general the study of mean curvature problems requires much care because of the possible occurrence of derivative blow-up phenomena, even in the one-dimensional case (see, e.g., [5, 12] and the references therein). However, in this case, we can show that an a priori bound in C^1 for a class of solutions of

$$- \left(\frac{r^{N-1} v'}{\sqrt{1 + v'^2}} \right)' = r^{N-1} \left(-av + \frac{b}{\sqrt{1 + v'^2}} \right) \quad (4)$$

can be obtained by an elementary argument which exploits the structure of the equation and the qualitative properties – positivity, monotonicity and concavity – of the solutions themselves. These estimates enable us to use a shooting method on a modification of equation (4) in order to prove the existence of a solution of (3) and hence of a radially symmetric solution of (1).

The proof of the uniqueness of solutions of (2) is instead based on converting, by a suitable change of variable, the original problem into a variational inequality, for which the uniqueness of solutions can be easily established by using a monotonicity argument.

The numerical solution of the quasilinear problem (3), but limited to the dimension $N = 1$, has been considered in [6] and [16], by using different approximation techniques.

We wish to mention that part of our results extends to the N -dimensional problem in a general domain: achieving this conclusion however requires a quite different approach, apparently even in the “simple” case of an annulus. This topic is discussed in [7].

We finally recall that anisotropic prescribed mean curvature equations have been recently considered, driven by different motivations, in [8, 9, 2, 10, 3, 1, 4, 11].

2. Existence, uniqueness and qualitative properties. The proof of Theorem 1.1 is based on Proposition 2.1 and Proposition 2.2 below.

Proposition 2.1. *Let $a > 0$ and $b > 0$ be given and let Ω be a bounded domain in \mathbb{R}^N with a Lipschitz boundary $\partial\Omega$. Then problem (2) has at most one solution $u \in C^2(\overline{\Omega})$.*

Proof. The proof consists of two steps.

Step 1. A variational inequality. We show that if $u \in C^2(\overline{\Omega})$ is a solution of (2), then $v = \exp(-bu)$ satisfies

$$\int_{\Omega} \sqrt{w^2 + b^{-2}|\nabla w|^2} dx - \int_{\Omega} \sqrt{v^2 + b^{-2}|\nabla v|^2} dx \geq - \int_{\Omega} ab^{-2} \ln v (w - v) dx \quad (5)$$

for all $w \in C^1(\overline{\Omega})$ with $\min_{\overline{\Omega}} w > 0$ and $w = 1$ on $\partial\Omega$. Indeed, it is easy to verify that, if $u \in C^2(\overline{\Omega})$ is a solution of (2), then $v = \exp(-bu)$ satisfies

$$\begin{cases} -\operatorname{div} \left(\frac{\nabla v}{\sqrt{v^2 + b^{-2}|\nabla v|^2}} \right) + \frac{b^2 v}{\sqrt{v^2 + b^{-2}|\nabla v|^2}} = -a \ln v, & \text{in } \Omega, \\ v = 1, & \text{on } \partial\Omega. \end{cases} \quad (6)$$

Pick any $w \in C^1(\overline{\Omega})$, with $\min_{\overline{\Omega}} w > 0$ and $w = 1$ on $\partial\Omega$, multiply the equation in (6) by $w - v$ and integrate by parts. The convexity and the differentiability in $\mathbb{R}_0^+ \times \mathbb{R}^N$ of the map $(s, \xi) \mapsto b^2 \sqrt{s^2 + b^{-2}|\xi|^2}$ then yields

$$\begin{aligned} - \int_{\Omega} a \ln v (w - v) dx &= \int_{\Omega} \frac{\nabla v \nabla (w - v)}{\sqrt{v^2 + b^{-2}|\nabla v|^2}} dx + \int_{\Omega} \frac{b^2 v (w - v)}{\sqrt{v^2 + b^{-2}|\nabla v|^2}} dx \\ &\leq \int_{\Omega} b^2 \sqrt{w^2 + b^{-2}|\nabla w|^2} dx - \int_{\Omega} b^2 \sqrt{v^2 + b^{-2}|\nabla v|^2} dx. \end{aligned}$$

Step 2. Uniqueness. Let us show that problem (2) has at most one solution $u \in C^2(\overline{\Omega})$. Suppose that $u_1, u_2 \in C^2(\overline{\Omega})$ are solutions of (2). Then, as $v_1 = \exp(-bu_1)$, $v_2 = \exp(-bu_2)$ satisfy (5), we have in particular

$$\int_{\Omega} \sqrt{v_2^2 + b^{-2}|\nabla v_2|^2} dx - \int_{\Omega} \sqrt{v_1^2 + b^{-2}|\nabla v_1|^2} dx \geq - \int_{\Omega} ab^{-2} \ln v_1 (v_2 - v_1) dx$$

and

$$\int_{\Omega} \sqrt{v_1^2 + b^{-2}|\nabla v_1|^2} dx - \int_{\Omega} \sqrt{v_2^2 + b^{-2}|\nabla v_2|^2} dx \geq - \int_{\Omega} ab^{-2} \ln v_2 (v_1 - v_2) dx.$$

Summing up and rearranging we get

$$0 \geq \int_{\Omega} ab^{-2} (\ln v_2 - \ln v_1) (v_2 - v_1) dx.$$

The strict monotonicity of the logarithm function yields $v_1 = v_2$ and hence $u_1 = u_2$. \square

Remark 1. We point out that, essentially by the same proof, we can obtain the following more general conclusion: if $u_1, u_2 \in W^{1,1}(\Omega) \cap L^\infty(\Omega)$ satisfy $u_1|_{\partial\Omega} = u_2|_{\partial\Omega}$ and, for $i = 1, 2$,

$$\int_{\Omega} \frac{\nabla u_i \nabla w}{\sqrt{1 + |\nabla u_i|^2}} dx = - \int_{\Omega} a u_i w dx + \int_{\Omega} \frac{b w}{\sqrt{1 + |\nabla u_i|^2}} dx$$

for all $w \in W_0^{1,1}(\Omega) \cap L^\infty(\Omega)$, then $u_1 = u_2$. We refer to [7] for further details.

Proposition 2.2. *For any given $a, b > 0$ and any $R > 0$, problem (3) has a solution $v \in C^2([0, R])$. In addition, the following conditions hold:*

- (i) $0 < v(r) < b/a$ for all $r \in [0, R]$;
- (ii) $v'(r) < 0$ for all $r \in]0, R]$;
- (iii) $v''(r) < 0$ for all $r \in [0, R]$.

Proof. The proof is divided into four steps.

Step 1. A modified problem. Set $c = \sqrt{\exp(2b^2/a) - 1}$ and define a function $\varphi : \mathbb{R} \rightarrow \mathbb{R}$ by

$$\varphi(s) = \begin{cases} \frac{s}{\sqrt{1 + s^2}}, & \text{if } |s| \leq c, \\ \frac{s + \operatorname{sgn}(s) c^3}{(1 + c^2)^{3/2}}, & \text{if } |s| > c. \end{cases}$$

Note that

$$\varphi'(s) = \begin{cases} \frac{1}{(1 + s^2)^{3/2}}, & \text{if } |s| \leq c, \\ \frac{1}{(1 + c^2)^{3/2}}, & \text{if } |s| > c \end{cases}$$

is bounded, bounded away from 0 and satisfies, for all $s \in \mathbb{R}$,

$$\varphi'(s) \geq \frac{1}{(1 + s^2)^{3/2}}. \quad (7)$$

Let us introduce the initial value problem

$$\begin{cases} -(r^{N-1} \varphi(v'))' = r^{N-1} \left(-av + \frac{b}{\sqrt{1 + v'^2}} \right), \\ v(0) = d, \quad v'(0) = 0, \end{cases} \quad (8)$$

with $d \in \mathbb{R}$. Clearly a function $v \in C^2([0, \omega[)$, for some $\omega \in \mathbb{R}_0^+ \cup \{+\infty\}$, is a solution of (8) if and only if it is a solution of

$$\begin{cases} v'' = \left(av - \frac{b}{\sqrt{1 + v'^2}} \right) \frac{1}{\varphi'(v')} - \frac{N-1}{r} \frac{\varphi(v')}{\varphi'(v')}, \\ v(0) = d, \quad v'(0) = 0. \end{cases} \quad (9)$$

In addition, as

$$\lim_{r \rightarrow 0} \frac{\varphi(v'(r))}{r \varphi'(v'(r))} = v''(0),$$

one has

$$v''(0) = \frac{ad - b}{N}. \quad (10)$$

Step 2. Global existence, uniqueness and continuous dependence. We are going to show that, for any given $d \in \mathbb{R}$, the initial value problem (8), or equivalently (9), has a unique solution $v \in C^2([0, R])$, which, moreover, depends continuously on the initial datum d and satisfies (10).

The local existence and uniqueness of solutions of (8) can be proved as follows. One first observes that, for any $\delta > 0$ small enough, the operator \mathcal{S} , defined by

$$(\mathcal{S}v)(r) = d + \int_0^r \varphi^{-1} \left(\int_0^s \left(\frac{t}{s} \right)^{N-1} \left(av(t) - \frac{b}{\sqrt{1+v'(t)^2}} \right) dt \right) ds,$$

is a contraction in the space $C^1([0, \delta])$, endowed with the usual norm; here the global Lipschitz continuity of φ^{-1} is in particular exploited. The C^2 -regularity up to 0 of the fixed point is verified directly; thus it gives rise to a local solution of (8). The continuous dependence of local solutions on the initial datum d is a consequence of the continuous dependence of the fixed points of \mathcal{S} on the parameter d .

Let us now denote by $g : \mathbb{R}_0^+ \times \mathbb{R} \times \mathbb{R} \rightarrow \mathbb{R}$, defined by

$$g(r, s, \xi) = \left(as - \frac{b}{\sqrt{1+\xi^2}} \right) \frac{1}{\varphi'(\xi)} - \frac{N-1}{r} \frac{\varphi(\xi)}{\varphi'(\xi)},$$

the function which appears at the right-hand side of the equation in (9). Since g is locally Lipschitz continuous in $[\delta, R] \times \mathbb{R} \times \mathbb{R}$ and grows linearly in $(s, \xi) \in \mathbb{R} \times \mathbb{R}$ uniformly in $r \in [\delta, R]$, any local solution of (9) can be uniquely continued to $[0, R]$.

Finally, the continuous dependence of these global solutions on the initial datum d is a standard consequence of the uniqueness.

Step 3. Qualitative properties. We shall show that, for any given $d < b/a$, the solution v of (9), defined on $[0, R]$, satisfies conditions (ii) and (iii).

Let us prove (ii). As $v'(0) = 0$ and $v''(0) < 0$, there exists $\delta_1 > 0$ such that $v'(r) < 0$ in $]0, \delta_1[$. Assume by contradiction that there exists $r_0 \in]0, R]$ such that $v'(r_0) \geq 0$. We can suppose that $v'(r_0) = 0$ and $v'(r) < 0$ in $]0, r_0[$. This yields in particular $v(r_0) < b/a$. As from the equation in (9) we have

$$v''(r_0) = av(r_0) - b < 0,$$

there exists $\delta_2 > 0$ such that $v'(r) > 0$ in $]r_0 - \delta_2, r_0[$, which is a contradiction.

Let us prove (iii). Assume by contradiction that there exists $\bar{r} \in]0, R]$ such that $v''(\bar{r}) \geq 0$. As $v''(0) < 0$, we can suppose that there exists $r_0 \in]0, \bar{r}[$ such that $v''(r_0) = 0$ and $v''(r) < 0$ in $[0, r_0[$. Define, for $r \in [0, R]$,

$$\psi(r) = v''(r)\varphi'(v'(r)) = av(r) - \frac{b}{\sqrt{1+v'(r)^2}} - \frac{N-1}{r}\varphi(v'(r)).$$

We have $\psi(r_0) = 0$ and, by (ii),

$$\begin{aligned} \psi'(r_0) &= av'(r_0) + b(1+v'(r_0)^2)^{-3/2}v'(r_0)v''(r_0) \\ &\quad + \frac{N-1}{r_0^2}\varphi(v'(r_0)) - \frac{N-1}{r_0}\varphi'(v'(r_0))v''(r_0) \\ &= av'(r_0) + \frac{N-1}{r_0^2}\varphi(v'(r_0)) < 0. \end{aligned}$$

This implies the existence of $\delta_3 > 0$ such that $\psi(r) > 0$ on $]r_0 - \delta_3, r_0[$ and in particular $v''(r) > 0$ in $]r_0 - \delta_3, r_0[$, which is a contradiction.

Step 4. Solvability. The map $\mathcal{T} : [0, b/a] \rightarrow \mathbb{R}$ defined by $\mathcal{T}(d) = v(R)$, where v is the solution of (9), is continuous and satisfies, according to condition (ii),

$$\mathcal{T}(0) < 0 < b/a = \mathcal{T}(b/a).$$

Then there exists $d \in]0, b/a[$ such that $\mathcal{T}(d) = 0$. The corresponding solution $v \in C^2([0, R])$ satisfies $v'(0) = 0 = v(R)$, as well as, by Step 3, conditions (i), (ii), (iii).

In order to show that v is the desired solution of (3), we prove that v also satisfies

$$\|v'\|_\infty \leq \sqrt{\exp(2b^2/a) - 1} = c,$$

or equivalently, by (iii),

$$v'(R) \geq -c.$$

From the equation in (9) we easily get

$$v''(r) \geq \frac{-b}{\varphi'(v'(r))\sqrt{1+v'(r)^2}} \geq -b(1+v'(r)^2),$$

and hence

$$\frac{v'(r)v''(r)}{1+v'(r)^2} \leq -bv'(r).$$

Integrating this inequality over $[0, R]$, we obtain

$$\frac{1}{2} \ln(1+v'(R)^2) \leq bv(0) < \frac{b^2}{a},$$

that is

$$|v'(R)| < \sqrt{\exp(2b^2/a) - 1} = c.$$

This concludes the proof. \square

Proof of Theorem 1.1. Proposition 2.1 guarantees that problem (1) has at most one solution $u \in C^2(\overline{B})$. Proposition 2.2 ensures that problem (3) has a solution $v \in C^2([0, R])$. Setting $u(x) = v(|x - x_0|)$ for all $x \in \overline{B}$, a simple calculation shows that $u \in C^2(\overline{B})$ is the unique solution of (1) which satisfies the cited conditions. Thus Theorem 1.1 follows. \square

REFERENCES

- [1] M. Athanassenas, J. Clutterbuck, A capillarity problem for compressible liquids, *Pacific J. Math.* **243** (2009), 213–232.
- [2] M. Athanassenas, R. Finn, Compressible fluids in a capillary tube, *Pacific J. Math.* **224** (2004), 201–229.
- [3] M. Bergner, The Dirichlet problem for graphs of prescribed anisotropic mean curvature in \mathbb{R}^{n+1} , *Analysis (Munich)* **28** (2008), 149–166.
- [4] M. Bergner, On the Dirichlet problem for the prescribed mean curvature equation over general domains, *Differential Geom. Appl.* **27** (2009), 335–343.
- [5] D. Bonheure, P. Habets, F. Obersnel, P. Omari, Classical and non-classical solutions of a prescribed curvature equation, *J. Differential Equations* **243** (2007), 208–237.
- [6] I. Coelho, C. Corsato, P. Omari, A one-dimensional prescribed curvature equation modeling the corneal shape, *Bound. Value Probl.* 2014, **2014**:127, 19 pp..
- [7] C. Corsato, C. De Coster, P. Omari, The Dirichlet problem for a prescribed anisotropic mean curvature equation: existence, uniqueness and regularity of solutions, preprint (2015), 39 pp., doi: 10.13140/2.1.3837.7766. Available from: <https://www.researchgate.net/publication/272826705>.
- [8] R. Finn, On the equations of capillarity, *J. Math. Fluid Mech.* **3** (2001), 139–151.
- [9] R. Finn, Capillarity problems for compressible fluids, *Mem. Differential Equations Math. Phys.* **33** (2004), 47–55.
- [10] R. Finn, G. Luli, On the capillary problem for compressible fluids, *J. Math. Fluid Mech.* **9** (2007), 87–103.
- [11] T. Marquardt, Remark on the anisotropic prescribed mean curvature equation on arbitrary domains, *Math. Z.* **264** (2010), 507–511.
- [12] F. Obersnel, P. Omari, Existence, regularity and boundary behaviour of bounded variation solutions of a one-dimensional capillarity equation, *Discrete Contin. Dyn. Syst.* **33** (2013), 305–320.
- [13] W. Okrański, L. Płociniczak, A nonlinear mathematical model of the corneal shape, *Nonlinear Anal. Real World Appl.* **13** (2012), 1498–1505.
- [14] W. Okrański, L. Płociniczak, Bessel function model of corneal topography, *Appl. Math. Comput.* **223** (2013), 436–443.
- [15] W. Okrański, L. Płociniczak, Regularization of an ill-posed problem in corneal topography, *Inverse Probl. Sci. Eng.* **21** (2013), 1090–1097.

- [16] L. Płociniczak, G.W.Griffits, W.E.Schiesser, ODE/PDE analysis of corneal curvature, *Comput. Biol. Med.* **53** (2014), 30–41.
- [17] L. Płociniczak, W. Okrasiński, Nonlinear parameter identification in a corneal geometry model, *Inverse Probl. Sci. Eng.* **23** (2015), 443–456.
- [18] L. Płociniczak, W. Okrasiński, J.J. Nieto, O. Domínguez, On a nonlinear boundary value problem modeling corneal shape, *J. Math. Anal. Appl.* **414** (2014), 461–471.

Received August 2014; revised January 2015.

E-mail address: chiara.corsato@phd.units.it, corsato.chiara@gmail.com

E-mail address: colette.decoster@univ-valenciennes.fr

E-mail address: omari@units.it