



Contrasting patterns of native and non-native plants in a network of protected areas across spatial scales

This is the peer reviewed (post-print) version of the following article:

Original

Contrasting patterns of native and non-native plants in a network of protected areas across spatial scales / Landini, S., Tordini, E., Amici, V., Bacaro, G., Carboni, M., Filibeck, G., Scoppola, A., Bagella, S.. - In: BIODIVERSITY AND CONSERVATION. - ISSN 0960-3115. - STAMPA. - 2020:(2020), pp. 1-51. [10.1007/s10531-020-01958-y]

Availability:

This version is available at: 11368/2959467 since: 2020-03-02T13:02:06Z

Publisher:

Published

DOI:10.1007/s10531-020-01958-y

Terms of use:

Some rights reserved. More details in the PostPrint rights section below.

License:

Copyright Editore

Publisher copyright

--

(Article begins on next page)

CONTRASTING PATTERNS OF NATIVE AND NON-NATIVE PLANTS IN A NETWORK OF PROTECTED AREAS ACROSS SPATIAL SCALES

Sara Landi^{1,2*}, Enrico Tordoni³, Valerio Amici², Giovanni Bacaro³, Marta Carboni⁴, Goffredo Filibeck⁵, Anna Scoppola⁵, Simonetta Bagella¹

¹ University of Sassari, Department of Chemistry and Pharmacy, Italy

² GREENARCO, Spin-Off Company of the Alma Mater Studiorum, University of Bologna, Italy.

³ University of Trieste, Department of Life Sciences, Italy

⁴ University of Roma Tre, Department of Sciences, Italy

⁵ University of Tuscia, Department of Agriculture and Forest Sciences (DAFNE), Italy

* Corresponding author e-mail address: slandi@uniss.it

Abstract: Networks of protected areas are fundamental for biodiversity conservation, but many factors determine their conservation efficiency. In particular, on top of other human-driven disturbances, invasions by non-native species can cause habitat and biodiversity loss. Jointly understanding what drives patterns of plant diversity and of non-native species in protected areas is therefore a priority. We tested whether the richness and composition of native and non-native plant species within a network of protected areas follow similar patterns across spatial scales. Specifically, we addressed three questions: *a*) what is the degree of congruence in species richness between native and non-native species? *b*) do changes in the composition of non-native species across ecological gradients reflect a similar turnover of native species along the same gradients? *c*) what are the main environmental and human disturbance drivers controlling species richness in these two groups of species?

Species richness and composition of native and non-native plant species were compared at two spatial scales: the plot scale (10 m x 10 m) and the Protected Area scale (PA). In addition, we fit Generalized Linear Models to identify the most important drivers of native and non-native species richness at each scale, focusing on environmental conditions (climate, topography) and on the main sources of human disturbance in the area (land use and roads). We found a significant positive correlation between the turnover of native and non-native species composition at both plot and PA scales, whereas their species richness was only correlated at the larger PA scale. The lack of congruence between the richness of native and non-native species at the plot scale was likely driven by differential responses to fine scale environmental factors, with non-natives favoring drier climates and milder slopes (climate and slope). In addition, more non-native species were found closer to road-ways in the reserve network. In contrast, the congruence in the richness of native and non-native species at the broader PA scale was mainly driven by the common influence of PA area, but also by similar responses of the two groups of species to climatic heterogeneity. Thus, our study highlights the strong spatial dependence of the relationship between native and non-native species richness and of their responses to environmental variation. Taken together, our results suggest that within the study region the introduction and establishment of non-native species would be more likely in warmer and dryer areas, with high native species richness at large spatial scale but intermediate levels of anthropogenic disturbances and mild slope inclinations and elevation at fine scale. Such an exhaustive understanding of the factors that influence

the spread of non-native species, especially in networks of protected areas is crucial to inform conservation managers on how to control or curb non-native species.

Keywords: plant species richness, Habitat Directive, habitat heterogeneity, Natura 2000, scale dependence, protected areas, human disturbance.

INTRODUCTION

The introduction and spread of non-native species (also known as alien or exotic species) is considered a major cause of habitat transformation and biodiversity erosion, especially in the Mediterranean biome (Sala et al. 2000, Lucy et al. 2016). Rather than only focusing on controlling species that are already established or naturalized in a territory, proactive management of biological invasions should aim at curbing the arrival or establishment of new non-native species (Spear et al. 2013) or controlling their populations during early-stages of the invasion *continuum* (*sensu* Richardson and Pyšek 2006). Identifying the features that make certain areas more likely to harbor many non-native species (which are also known to correlate with areas harboring many invasive species, Williamson & Fitter, 1996) is an important information for controlling non-native species spread. In addition, by understanding whether the same features also drive the distribution of native species it might be possible to highlight potential hotspots for biodiversity management. For example, if native species tend to be associated with similar conditions as non-native species, valuable areas of high biodiversity will inevitably overlap with hotspots of invasions (e.g. Stadler et al. 2000), potentially leading to greater risks of biodiversity loss as a consequence of non-native species impacts (Simberloff et al. 2013). These might thus represent joint priorities for conservation and non-native species control efforts.

An enormous body of literature has tried to identify general features or ecosystem properties facilitating the establishment of non-native species (*e.g.*, Ashton and Mitchell 1989; Faliński 1998; Sukopp 1998; Pyšek and Richardson 2006, Bjarnason et al. 2017), and to understand whether the same features also promote native species richness and high overall biodiversity (*e.g.*, Shea and Chesson 2002; Deutshevitz et al. 2003; Carboni et al. 2010; Tordoni et al. 2017). In general, environmental factors which influence patterns of species richness at regional scales include climate, landscape heterogeneity, and geomorphological processes, all of which typically affect native as well as non-native species (Davies et al. 2005; Moser et al. 2005; Carboni et al. 2010). But in addition, in the current Anthropocene era (Crutzen 2006), human disturbance and management practices are a major agent of change of species richness and diversity patterns across spatial scales (Maestre 2004; Gaston 2005). For example, human disturbance may generate environmental heterogeneity, which may increase extinction risk of native species, but also allow for resource partitioning by creating new niche opportunities (Shochat et al. 2006). These factors may thus facilitate the arrival of non-native species pre-adapted to such altered conditions (Callaway 2007). Indeed, while natural or near-natural ecosystems often display a certain ecological resistance against biological invasion (*e.g.*, Faliński 1998; Simberloff et al. 2013), densely populated areas or areas subject to strong human disturbance are typically found to be prone to higher levels of non-native species establishment and invasion success (*e.g.*, Pyšek et al. 1998; Sukopp 1998; McKinney 2002; Chytrý et al. 2008; Tordoni et al. 2017).

The scale dependence of biodiversity patterns is a well-known issue in ecology (*e.g.*, Huston 1999; Richardson and Pyšek 2006). In particular the relationship between native and non-native species richness seems to change across spatial scales, which led scientists to even coin the term “invasion paradox” (Shea and Chesson 2002; Frid-

ley et al. 2007). Specifically, a negative relationship is usually observed at small spatial scales, at which species typically interact (e.g., Cornell and Karlson 1997; Levine 2000; Tilman 1997), but this relationship tends to become positive when increasing the grain of the sampling units or the extent of the study area (Planty-Tabacchi et al. 1996; Lonsdale 1999; Levine 2000; Stadler et al. 2000; McKinney 2002; Sax 2002; Kumar et al. 2006; Stohlgren et al. 2006). Although there has been a heated debate to try to explain these contrasting patterns, this conundrum of invasion ecology is still far from being resolved. Further studies are thus needed in order to understand the relationships between native and non-native species patterns and the environmental and anthropogenic features which foster biological invasions across spatial scales.

In the Convention on Biological Diversity (art. 8 *In-situ Conservation*, <https://www.cbd.int/>), prevention measures, control, or eradication of problematic non-native species are called for. Identifying pathways for non-native species establishment and management priorities are possible actions in this direction. In particular, protected areas are a key component of the global response to environmental change and degradation (e.g., Hannah et al. 2007; Gaston et al. 2008; Foxcroft et al. 2017), and can be part of a framework to devise effective invasion control measures. Nevertheless, they face many challenges, such as the effectiveness of reserve design, governance (Pressey et al. 2015), and anthropogenic change (Foxcroft et al. 2017), with generally few restrictions currently in place for preventing the introduction of non-native species (Pyšek et al. 2003). Most protected areas in Europe are in a mosaic of land use types that can form a network of potential sources for non-native species introductions (e.g., Foxcroft et al. 2007; Meiners and Pickett 2013). In addition, recent evidence shows that there is almost no difference in the patterns of non-native and invasive species inside and outside protected areas, suggesting that currently habitat protection has little or no effect on non-native species richness (e.g. Moustakas et al. 2018). Studying which features are linked to higher invasion levels can thus help identify the main pathways that need regulation and which areas are most at risk, to guide future conservation planning within protected area networks.

In this study, we investigated the importance of biotic (represented by native species richness and composition), environmental (climate) and anthropogenic (road network and land uses) factors in driving non-native plant species spatial patterns within a network of protected areas in central Italy. We aimed at providing insights on the ecological mechanisms useful for the effective control of non-native species establishments, which is extremely important in the context of the management of reserve networks. Ideally, reserve networks strive to maximize the protection of biodiversity features, while non-native species clearly represent a potential threat for nature reserves and their management (Pyšek et al. 2003). If native and non-native species follow similar patterns within the network, an overlap of high native biodiversity and invasion hotspots is likely to emerge. Identifying such areas at high risk of biodiversity loss and the ecological features that may promote invasion or otherwise hamper biodiversity protection is therefore essential to improve control and management actions. We aim at testing, at different spatial scales, whether species richness levels of native and non-native vascular plants are correlated and whether similar factors control the turnover in species composition of native and non-native species along ecological gradients. Specifically, our research questions are: a) what is the degree of congruence in species richness between native and non-native plant species, and does this relationship vary across spatial scales? b) do changes in the composition of non-native species across environmental gradients reflect native species turnover along the same gradients? c) what are the main environmental and anthropogenic drivers controlling species richness in each of these two groups of species?

113 METHODS

114 Study area

115 This study was performed in the local network of protected areas (PAs) of the province of Siena (Italy), including
116 four Nature Reserves (designated under national or regional regulations) and 17 Special Areas of Conservation of
117 Natura 2000 network (SACs, designated under the EU Habitat Directive 93/43/EEC) (Figure 1). The size of the
118 single PAs within the network ranges from 268 ha to 13747 ha, while their elevation ranges from 122 m to 1660 m
119 a.s.l. (Chiarucci et al. 2012) with a cumulative area of 593 km² (15.6% of the Province).

120 The study area is characterized by a Mediterranean macro-climate, even though there is a strong variation across
121 sampling sites due to differences in morphology and local elevations (Castrignanò et al. 2006). Long term mean
122 annual precipitation ranges from 630 to 1275 mm (Barazzuoli et al. 1993). The highest precipitations (above 1000
123 mm on average) and lowest mean annual temperature values (lower than 12 °C) are found at higher elevations
124 (Monte Amiata). A relatively arid and warm zone (mean annual temperature of about 14 °C and average precipita-
125 tions of ca. 600 mm) is localized in the South-East of the province of Siena (Orcia river valley; Barazzuoli et al.
126 1993).

127 The geology is rather varied and complex (including, *inter alia*, limestone, clay, marl, metamorphic and volcanic
128 bedrock), resulting in highly heterogeneous morphology and a great variety of landscapes. The main land-cover
129 types include evergreen coppice woods (dominated by *Quercus ilex*) and deciduous coppice woods or forests
130 (thermophilous types dominated by *Quercus pubescens* or *Q. cerris* at lower elevations, and mesophilous types
131 dominated by *Castanea sativa* or *Fagus sylvatica* at higher elevations); evergreen Mediterranean shrublands
132 (characterized *e.g.* by *Erica arborea*, *E. scoparia*, *Phillyrea latifolia*, *Pistacia lentiscus*, *Arbutus unedo*, *Cistus*
133 *salvifolius*, *Juniperus communis* and *J. oxycedrus*); croplands (mainly wheat and horticultural crops), vineyards
134 and olive groves. Other relevant land cover types include pastures, meadows, garigues (on calcareous and ultra-
135 mafic substrates), conifer plantations and wetlands.

136 Sampling design

137 Plants were sampled by using an operational approach of plant communities (Chiarucci 2007), defined by a fixed
138 grain and uniform sample density within each protected area (PA). Sampling design was based on a grid of
139 1 km × 1 km cells, covering the whole study area, with a sampling point randomly selected within each cell (Chi-
140 arucci et al. 2008, 2012). A sampling unit represented by a square plot of 10 m x 10 m was centered at each sam-
141 pling point. This was further divided in 16 contiguous squared 2.5 m x 2.5 m subplots (Figure S1 of Supplemen-
142 tary material), on which the occurrence of all species of vascular plants was recorded.

143 The field data collection was performed from April to June, during the years 2005-2009. The total number of
144 sampled plots was 604.

145 Native and non-native species richness and composition

146 All the vascular plants recorded within each plot were identified at the species or subspecies level with standard
147 floras (Pignatti 1982; Tutin et al. 1964–1980, 1993) or monographs (Grunanger 2001; Weber 1995). Nomenclature
148 was standardized according to Conti et al. (2005). Plants were classified as native or non-native, depending on
149 their distributional status as given by Celesti-Gradow et al. (2011).

Composition and species richness for both native and non-native species were then separately assessed at plot and PA scale, i.e. two data matrices were prepared (species by plots and species by PAs). At plot scale, frequency of each species, calculated as the sum of occurrences within the 16 subplots, was used as a coarse measure of abundance. At PA scale, relative frequency of each species, measured as the ration between the number of occupied plots *versus* the number of recorded plots, was used as measure of abundance.

The PA-scale species lists were obtained by pooling the data from the plots included within each PA.

Environmental and human disturbance predictors

At plot scale, four groups of predictor variables for evaluating the environmental and human-mediated and spatial factors affecting species richness were considered: 1) climatic, 2) topographic, and 3) human disturbance (Table 1):

1) Climatic variables: we derived one synthetic variable named *climate*. We obtained this variable from 36 climatic variables related mainly to monthly temperatures and precipitations (e.g. maximum annual temperature, minimum annual temperature, annual rainfall) for each plot, from the LaMMA consortium (Laboratorio di Monitoraggio e Modellistica Ambientale per lo sviluppo sostenibile; <http://www.lamma.rete.toscana.it/en>). Specifically the LaMMA data consisted of local interpolated climatic grids with a resolution of 250 m x 250 m. Since these variables were highly inter-correlated, a Principal Component Analysis (PCA) was performed to reduce their multicollinearity (Taylor et al. 2002). Given that the first PCA factor explained more than 90% of variance (Figure S2 of Supplementary material), this was used as a single variable (hereafter called *climate*). This was found to be negatively correlated with all the variables related to total and summer rainfall (Pearson correlation coefficients from -1 to -0.85), positively correlated with the minimum temperatures (Pearson correlation coefficients from 0.77 to 0.83) and maximum ones (Pearson correlation coefficients from 0.84 to 0.90). This compound *climate* variable corresponds therefore to a gradient from wet and cold to warm and (summer-) dry conditions, that is from Temperate to Mediterranean meso-climate, moving from the negative to the positive extreme of the axis.

2) Topographic variables: *elevation*, *slope*, and potential *solar radiation* values (the latter was obtained from slope, aspect and latitude, following McCune and Keon 2002). The resolution used to develop these variables was 30 m.

3) Human disturbance variables: *distance to the nearest road* (paved or unpaved) and *land-use type* variables. We used the distance to the nearest road as a proxy of human disturbance as the road network is a well-known introduction pathway for non-native species (e.g. Pauchard and Alaback 2004; Arévalo et al. 2010). Road layer have been derived from GEOscopio Geoportal, Tuscany Region - Territorial and Environmental Information System (<http://www.Geografia.toscana.it/>)

In addition, *landscape diversity* (H_{plot}) and *landscape evenness* (E_{plot}) within a buffer area of 250 m radius around the plot were assessed based on the Shannon Index of diversity and the Pielou Index of evenness applied on the Corine Land Cover (CLC) map. At plot scale, the human disturbance variables linked to land use have not been considered due to the resolution of the CLC map which did not allow to detect artificial land use types in the plots or in related buffer areas. So we used artificial land use types as proxy of human disturbance only at protected area scale.

At PA scale, the same groups of predictor variables as at the plot scale were considered, with some unavoidable differences in the definition of the single predictors due to different spatial resolution (Table 2):

1) Climatic variables: the variability of climate across plots was accounted for by calculating two derived climatic variables at PA scale: *mean climate* (calculated as the centroid, along the first axis in the climate PCA, of the plots belonging to each PA) and *climate range* (calculated as the range of the same points along the PCA axis).

2) Topographic variables: *elevation range* and *mean elevation* within each PA.

3) Human disturbance variables: total *road density* and *land-use type* variables. We obtained the total road density as the linear extension of paved and unpaved roads per km² within each PA and was considered as a proxy of human disturbance and potential propagule pressure of non-native species.

Moreover, as human disturbance, we obtained 4 land-use type variables reflecting the different artificial land-use types found in each PA based on CLC map (I level) of the area extracted from the Copernicus database (<https://land.copernicus.eu/pan-european/corine-land-cover>). Land-use types variables, artificial, semi-natural and natural, were expressed as percentages of relative Corine land-use classes (human, agriculture, natural, wetland). The variables *landscape diversity* (H_{PA}) and *landscape evenness* (E_{PA}) were calculated using the Shannon Index of diversity and the Pielou Index of evenness on the III level CLC map. The calculation of the landscape diversity was done at the III level of CLC because this allowed for a finer classification of landscape units, that was not necessary for detecting the dominant land use type.

4) Geographical variable: *area* (expressed in km²) of each PA area.

STATISTICAL ANALYSES

Separate analyses were performed at the plot and PA scales. At each scale, we assessed: 1) the correlation between native and non-native species richness, 2) whether specific sets of non-native species tend to be associated with specific native species assemblages, and 3) which predictors best explained native and non-native species richness, respectively.

Relationships between native and non-native species

First, we assessed the correlation between native and non-native species richness by computing the Spearman correlation coefficient ρ at both spatial scales. This was done to assess whether native and non-native species richness within the reserve network followed 1) similar trends, suggesting that factors which favor high richness of native species, also increase opportunities for non-native species (Thuiller et al. 2010), leading to a positive native-non-native richness relationship (Shea and Chesson 2002), or 2) opposite trends suggesting higher biotic resistance of more diverse native communities, as interpreted by several authors (Cornell and Karlson 1997; Stohlgren et al. 1999; Levine 2000; Tilman 1997; Brown and Peet 2003; Davies et al. 2005; Souza et al. 2011).

Second, we verified if turnover in native species composition is associated also to a turnover in non-native species and thus if specific sets of non-native species tend to always be associated with the same natives. To do so, we first obtained plot-to-plot (and PA-to-PA) dissimilarities in species composition, separately for non-native and for native species, using Bray Curtis pairwise dissimilarities on $\log(x+1)$ transformed species frequencies. Then, we tested whether the pairwise dissimilarity matrices (or distance matrices) of native and non-native species were

linearly independent by performing a Mantel test (Spearman correlation, 999 permutations, McCune and Keon 2002). For this we considered only the plots or PAs that had at least one non-native species.

Determinants of species richness across spatial scales

To assess the relationship between vascular plant richness (separately for native and non-native species) and the environmental and human disturbance predictors, we used two different approaches depending on the scale of investigation (plot or PA scale). In both cases, we first evaluated the presence of spatial autocorrelation in the response variables by calculating Moran's I coefficient using the R package 'ape' (Paradis and Schliep 2018), based on the geographical coordinates of the plots and of the centroid of the PA, respectively. Also, multicollinearity among explanatory variables at each scale was tested by computing Spearman's (correlation matrix available in Table S1 of the Supplementary material) and ensuring that there were no pairs of variables in the models where $|\rho| > 0.7$ (Dormann et al. 2013).

At plot scale, we fit Generalized Linear Mixed Models (GLMM) using R package "lme4" (Bates et al. 2015) and assuming Poisson family errors. PA was considered as random effect to control for the spatial dependence of plots within PA. At PA scale, Generalized Linear Models (GLM) were used using Poisson family errors; in case of overdispersion in the data a quasipoisson family was used instead. A Minimum Adequate Model (MAM) and a set of models with good support were thus obtained by performing a stepwise variable selection procedure through AICc minimization using R package "MuMIn" (Barton 2019). Only the models with $\Delta AICc \leq 2$ compared to the best model were considered to have good support (Burnham and Anderson 2002). In case of more models with similar AICc values, the one which retained a lower number of predictors was considered as minimum adequate model (MAM) according to Occam's razor. Note that, since results were qualitatively similar across the set of models with good support (Table S2 of Supplementary material), we present and discuss only the MAMs in the main text. In case of overdispersion in the data, it was not possible to use an AICc-based selection approach, then the amount of deviance accounted for by the GLM adjusted by the number of predictors was used instead (D^2_{adjusted} ; Guisan and Zimmermann 2000; Barbosa et al. 2014). In addition, R^2 statistics (marginal effect) were derived for GLMMs using the "r2glmm" R package (Jaeger 2017). Prior to analysis, quantitative variables were standardized in order to have mean zero and unit variance. Any residual spatial autocorrelation after modeling was assessed by means of spline-correlograms using the R package "necf" (Bjørnstad 2019). Specifically, 95% pointwise bootstrap confidence intervals were calculated from 5000 bootstrap samples of Pearson residuals after accounting for the level of spatial autocorrelation explained by the explanatory variables in each model.

All the statistical analyses were conducted using R 3.6.1 (R Core Team 2019).

RESULTS

Species patterns at plot scale

In total, 993 native species and 48 non-native species were recorded in 604 plots. Plant species richness ranged from 0 to 117 (mean 31.0) for native species and from 0 to 9 (mean 0.6) for non-native species. Native and non-native species richness exhibited a slight positive correlation, though not significant (Spearman $\rho = 0.06$, $p = 0.11$). In contrast, distance matrices based on plot-to-plot compositional dissimilarity showed a significant positive correlation (Mantel test, Spearman $\rho = 0.24$, $p = 0.001$).

Table S2 reports the full list of candidate models for species richness predictors derived from the model selection procedure. Table 3 describes the best set of predictors for species richness in the GLMM MAM. Specifically, minimum adequate models showed, on the one hand, that native species richness was positively associated with *landscape diversity* (H_{plot}) and negatively with the first axis of PCA derived from climatic variables (*climate*) and slope. On the other hand, non-native species richness was negatively related to *road distance* and *slope*, and positively associated with *climate*. However both models, in particular the native-species model, are characterized by a weak goodness of fit ($R^2 = 0.02$ and $R^2 = 0.11$ for native and non-native species, respectively).

Species patterns at protected area scale

Plant species richness at PA scale ranged between 22 and 547 (mean 219.8 species) for native species and 0-24 (mean 6.6 species) for non-native species. Species richness values of natives and non-native species (Figure 2) were positively and significantly correlated (Spearman $\rho = 0.75$, $p < 0.001$). The relative proportion of non-native species (ratio of non-native to native richness) in PAs was positively correlated with the richness of native species ($R^2 = 0.32$, $p < 0.01$), increasing up to a limit of about 5% (Figure 2). Similarly, the Mantel correlation between native and non-native species dissimilarity matrices was significant (Spearman $\rho = 0.33$ with $p = 0.003$). Further, according to the MAMs, species richness at PA scale was positively correlated with *area* and *climate range* for both species groups (Table 3). In addition, native species richness showed a positive relationship with *mean climate* and a negative, but not significant, relationship with *% wetland*. In contrast, non-native species richness was negatively related with *mean elevation*. In both species groups, the deviance accounted for by the model was relatively high ($D^2_{adjusted} = 0.76$ and $D^2_{adjusted} = 0.71$ for native and non-native species model, respectively).

All four spline correlograms (Figure S3 of Supplementary material) failed to reveal any evidence of spatial autocorrelation in the residuals, thereby allowing us to exclude its influence on model parameter estimates.

DISCUSSION

The Italian flora is currently estimated to include 7634 *taxa* (species and subspecies), 13.4% of which (1023 *taxa*) are considered to be non-native (Celesti-Grapow et al. 2011). In our dataset, collected using a probabilistic sampling strategy within the protected areas of the Siena province only, we recorded a rich flora (1041 species were recorded by this survey) and a relatively low proportion of non-native species (4.9% of the whole sample). However, even if the province of Siena is characterized, overall, by a well preserved landscape of traditional land uses (Geri et al 2010), only protected areas were included in this study and thus most of the sampled sites are on average less disturbed than the remainder of the landscape at province (or country) scale. Even though the proportion of non-native species in the investigated system is generally low, we found evidence of scale dependency of the relationship between native and non-native species richness. While there was a strong positive relationship at the PA scale, native and non-native species richness were not correlated at the plot scale. This was the consequence of different responses of the two groups of species to environmental and human factors at the fine scale, while similar factors (chiefly reserve area) explained both native and non-native species richness at large scale.

Species patterns at the plot scale

At the plot scale, we did not observe a relation between native and non-native species richness but we observed a relationship between the compositional gradients of native and non-native composition species. This suggests that non-native species tend to be associated with specific sets of native species (*i.e.*, native plant communities), either because they have similar environmental requirements or because of more direct biotic interactions (*e.g.*, facilitation or competitive exclusion). However, the lack of significant correlation between native and non-native species richness instead suggests a limited role for biotic interactions and for biotic resistance through competitive exclusion in these plant communities.

Indeed, our results suggested that different factors affected the species richness of the two groups of species at plot scale. Specifically, native species richness was only very weakly related to the variables we measured at this scale, and our best model only explained a very small proportion of variability ($R^2 = 0.02$). Nevertheless, we found evidence that native richness was positively associated with *landscape diversity* around the plot (250 m radius) and negatively associated with the *climatic* gradient ranging from Temperate to Mediterranean meso-climatic conditions. The first result is in agreement with previous studies suggesting that greater *landscape diversity* may be related to a higher number of available niches, potentially hosting species with different ecological requirements (Deutschewitz et al. 2003; Kumar et al. 2006). In our study area, higher values of small-scale *landscape diversity* might specifically indicate fragmentation and the survival of more natural ecosystems within an otherwise homogeneous agricultural landscape. The negative relationship between native species and the temperate-mediterranean gradient indicates that more native species were found in the cooler and wetter, rather than in the drier, meso-climatic conditions. This is likely to be connected to the regional context, in which higher temperatures at plot scale might represent a significant ecological constrain in summer, while lower temperatures are not likely to be a limiting factor in winter.

In contrast, more variability could be explained for non-native species richness. Non-native species were negatively related with *road distance*, and *slope* inclination and positively related to the temperate-mediterranean climatic gradient.

As expected, non-native species richness was higher in plots in close proximity to roadways. This is in accordance with the well-documented notion that roads, and roadside habitats, are a major source from which non-native species colonize natural areas (Parendes and Jones 2000; Pauchard and Alaback 2004; Bacaro et al. 2015; Ullmann and Heindl 1989; Ullmann et al. 1995; Arévalo et al. 2010). Roads may facilitate the dispersal of non-native species inducing habitat fragmentation and altering (micro-) environmental conditions, facilitating the human-mediated dispersal of propagules (via air movement associated with vehicle traffic, and via the seeds attached to the vehicles themselves) and facilitating the colonisation by non-native species by suppressing the growth or removing stands of native species (Trombulak and Frissel 2000; Bacaro et al. 2015). Therefore, limitation to the construction of new roads within or close to protected areas is important for preserving local biodiversity both directly, but also indirectly via the reduction of non-native introductions.

In terms of environmental factors, non-native species richness at plot scale was positively associated with warmer and drier Mediterranean meso-climatic conditions and negatively correlated with slope inclination. This can partly be a consequence of the well-known conservative role of steep slopes, because of soil- and microclimatic- limiting

factors: for instance, Bennie et al. (2006) found that in Britain, grassland swards on steep slopes were more resistant to invasion by competitive grass species than those on flatter sites, due to phosphorus limitation in shallow minerogenic soils, and to increased drought events. Filibeck et al. (2016) showed that, in grasslands in a protected area at the periphery of Rome, the removal or leveling of some steep debris heaps led to an increase in non-native and cosmopolite species. However, our dataset is not restricted to grasslands and involves a huge variety of land-cover types, so it is also possible that the observed effect is mediated by disturbance levels and land-use: in the study region, a high slope steepness usually allows only land-use types that feature an inherently low human disturbance (*e.g.*, woods or extensive rangelands), while flat morphologies are usually exploited with intensive crops or urban land-use, typically favouring non-native taxa invasions.

Species patterns at the protected area scale

We found that the turnover in non-native species composition was correlated with native species composition also at the scale of protected area. At this spatial grain, however, we also detected a strong congruence in the richness of the two groups of species. These findings, highlight that the protected areas that are most biodiverse are also the ones that are most easily colonized by alien species (and thus, potentially, most at risk of invasion). More generally, these findings support the “biotic acceptance hypothesis” or “the rich get richer” hypothesis, according to which sites with high native species richness are the most readily invaded by non-native species (*e.g.*, Stohlgren et al. 1999, 2006; Fridley et al. 2007; Pyšek and Richardson 2006; Sandel and Corbin 2010, Bartomeus et al. 2012). This would be explained because habitats that are generally ‘good’ for native species would also be ‘good’ for non-native ones (*e.g.*, McKinney 2002; Souza et al. 2011). This hypothesis can also be linked to Grime’s theory (Grime 1973), according to which all species (meaning both native and non-native species) respond, to some degree, in a similar way to stress, competition and disturbance (Tomasetto et al. 2013).

We therefore tested whether the similar compositional and richness patterns emerged as a consequence of similar responses to environmental factors and human disturbances. Indeed, we found that two main variables were retained in both the models for native and non-native species richness at the PA scale: *area* of the protected area, and *climate range* (*i.e.* spatial heterogeneity of climate within the PA). Both these variables showed positive effects, even if with different weights for native and non-native species richness. Thus, our research shows a positive native-non-native relationship related to the same response of native and non-native species to the available area in the PA and to the main gradients at large scale. The positive effect of *area* on both native and non-native species richness is likely dependent on the higher availability of niches in larger protected areas for both groups of species. The congruence of native and non-native species richness is thus largely the net result of concordant well-known species-area relationships, that show similar patterns for both groups of species. Indeed a congruence in species-area relationships for native and non-native species had already been reported in this network of protected areas (Chiarucci et al. 2012), as well as in other reserve networks (Pyšek et al. 2002a) or insular systems (*e.g.*, the islands of the Tuscan archipelago, Chiarucci et al. 2017). In addition, the positive effect on richness of the climatic variables at the scale of the protected area (*climate range*) indicates that more climatically heterogeneous areas enhance regional native richness but also promote non-native species establishment. Indeed, environments with greater spatial heterogeneity (*e.g.*, higher habitat diversity or spatial variability in resources or conditions; Davies et al. 2005) have been shown to support higher numbers of both native and non-native species at broad spatial scales in numerous other systems. Thus, overall we conclude that in our reserve network “rich protected areas get

richer” because of more available area (as expected) but also because of more favorable conditions in terms of available niches and of climates.

In addition to the drivers that were common to both natives and non-natives, the model for native species at PA scale also included a significant positive effect of mean climate, which suggests that reserves with warmer and drier average climates tend to support greater species richness. The model for non-native species included a significant effect of mean elevation, that was negatively related to non-native species richness indicating that fewer non-native species were found at higher altitudes. This finding is in agreement with previous studies that showed a negative relationship between non-native plant species richness and elevation in various systems (*e.g.*, Pyšek et al. 2002b; Stevens 1992; Pausas 1994; Rey-Benayas 1995; Marini et al. 2009; Siniscalco et al. 2011; Barni et al. 2012; Bacaro et al. 2015). Nevertheless it is important to caution that upward movements of non native species are increasingly being detected in many montane ecosystems (Kalwij et al. 2015), and are also predicted to accelerate in the future (Carboni et al. 2018). Hence, even if our results show that currently higher elevations are of lesser concern for non-native plants, potential future invasion risks should not be discarded.

Conclusions and cross-scale comparisons

Overall, our results support the idea that the relationship between the number of native and non-native species is altered when changing the scale of analysis. Indeed, it is widely acknowledged that the strength and form of this relationship (in many studies known as NERR – native exotic richness relationship – *e.g.*, Souza et al. 2011; Symonds and Pither 2012) is scale-dependent. Here, we found evidence that the lack of congruence at plot scale was likely driven by differential responses to fine scale environmental factors (*climate* and *landscape diversity*) and human disturbances (*road distance*), while at the larger PA scale native and non-native species largely responded in a similar way to available *area* and to *climate* variability (climate range). While at coarse scales the species richness of native species (and non-native species) was generally higher in warmer Mediterranean climates, at the fine plot scale the native species richness was even mildly associated with cooler meso-climatic conditions. Overall richness of non-native species in the protected area network increased under moderate anthropogenic disturbances at at fine scale, coupled with high levels of habitat and climatic heterogeneity at large scale. Thus, our data suggest that within the study region the introduction and establishment of non-native species would be more likely in warmer and dryer areas, with high native species richness at large spatial scale but intermediate levels of anthropogenic disturbances and mild slope inclinations and elevation. We also found potential conservation hotspots, especially because the very biodiverse protected areas potentially feature a higher risk of invasion, due to higher establishment success of non-native species in general. These areas should thus be prioritized for invasion monitoring. Finally, the results also highlight that both 1) the measure of the proportion of non-native species and 2) the relationship with potential predictors should be studied at the appropriate spatial scale in order to be comparable among different regions and informative for conservation purposes.

405 **Acknowledgment**

406 We would like to thank Prof. Alessandro Chiarucci for the opportunity to conduct this research on the data of the
407 project MOBISIC he coordinated when he was at the University of Siena. We would also thank Francesco Geri for
408 his assistance with geographical information. G.F. and A.S. carried out this research partially in the frame of the
409 MIUR (Italian Ministry for Education, University and Research) initiative “Department of excellence” (Law
410 232/2016).

411

412 **Author Contributions:**

413 All the authors contributed to the interpretation of results and writing.

REFERENCES

- Amici V, Landi S, Frascaroli F, Rocchini D, Santi E, Chiarucci A. 2015. Anthropogenic drivers of plant diversity: perspective on land use change in a dynamic cultural landscape. *Biodiversity and Conservation* 24: 3185-3199.
- Arévalo JR, Delgado JD, Otto R, Naranjo A, Salas M, Fernández-Palacio JM. 2005. Distribution of alien vs. native plant species in roadside communities along an altitudinal gradient in Tenerife and Gran Canaria (Canary Islands). *Perspectives in Plant Ecology, Evolution and Systematics* 7: 185–202.
- Ashton PJ, Mitchell DS. 1989. Aquatic plants: patterns and modes of invasion, attributes of invading species and assessment of control programmes. In: Drake JA, Mooney HA, di Castri F, Groves RH, Kruger FJ, Rejmánek M, Williamson M (eds) *Biological invasions: a global perspective*. Wiley, Chichester, pp 111-154.
- Bacaro G, Maccherini S, Chiarucci A, Jentsch A, Rocchini D, Torri D, Gioria M, Otto R, Escudero CG, Fernandez-Lugo S, Fernandez-Palacios JM, Arévalo JR. 2015. Distributional patterns of endemic, native and alien species along a roadside elevational gradient in Tenerife, Canary Islands. *Community Ecology* 16: 223-234. doi: 10.1556/168.2015.16.2.10
- Barazzuoli P, Guasparri G, Salleolini M. 1993. Il clima. In: Giusti F (ed) *La storia naturale della Toscana meridionale*. Monte dei Paschi di Siena, Amilcare Pizzi Editore, pp 141-171.
- Barbosa AM, Brown JA, Real R. 2014. modEvA – an R package for model evaluation and analysis. R package, version 0.1.
- Barni E, Bacaro G, Rocchini D, Falzoi S, Spanna F, Siniscalco C. 2012. Establishing climatic constraints shaping the distribution of alien plant species along the elevation gradient in the Alps. *Plant Ecology* 213: 757-767.
- Bartomeus I, Sol D, Pino J, Vicente P, Font X. 2012. Deconstructing the native-exotic richness relationship in plants. *Global Ecology and Biogeography* 21:524–533.
- Barton K. 2019. MuMIn: Multi-Model Inference. R package version 1.43.6. <https://CRAN.R-project.org/package=MuumIn>
- Bates D, Maechler M, Bolker B, Walker S. 2015. Fitting Linear Mixed-Effects Models Using lme4. *Journal of Statistical Software*, 67(1): 1-48. doi:10.18637/jss.v067.i01.
- Bazzichetto M, Malavasi M, Barták V, Acosta A, Rocchini D, Carranza M. 2018. Plant invasion risk: A quest for invasive species distribution modelling in managing protected areas. *Ecological Indicators* 95: 311-319. doi:10.1016/j.ecolind.2018.07.046
- Bennie J, Hill MO, Baxter R, Huntley B. 2006. Influence of slope and aspect on long-term vegetation change in British chalk grasslands. *Journal of Ecology* 94: 355–368.
- Brown RL, Peet RK. 2003. Diversity and invasibility of southern Appalachian plant communities. *Ecology* 84: 32–39.

446 Bjarnason A., Katsanevakis S., Galanidis A., Vogiatzakis I.N., Moustakas A. 2017. Evaluating Hypotheses
 447 of Plant Species Invasions on Mediterranean Islands: Inverse Patterns between Alien and Endemic species.
 448 *Frontiers Ecol. Evolut.*, 5 (91). Mediterranean.

449 Bjornstad ON. 2019. ncf: Spatial Covariance Functions. R package version 1.2-8. [https://CRAN.R-](https://CRAN.R-project.org/package=ncf)
 450 [project.org/package=ncf](https://CRAN.R-project.org/package=ncf)

451 Burnham KP, Anderson DR. 2002. Model selection and multimodel inference: a practical information-
 452 theoretic approach. Springer-Verlag, New York, New York, USA.

453 Calcagno V. 2013. glmulti: Model selection and multimodel inference made easy. R package version 1.0.7.

454 Callaway RM. 2007. Positive interactions and interdependence in plant communities. New York, Springer-Verlag.

455 Carboni M, Thuiller W, Izzi F, Acosta A. 2010. Disentangling the relative effects of environmental versus human
 456 factors on the abundance of native and alien plant species in Mediterranean sandy shores. *Diversity and Distribu-*
 457 *tions* 16: 537–546. doi: 10.1111/j.1472-4642.2010.00677.x

458 Carboni M, Guéguen M, Barros C, Georges D, Boulangeat I, Douzet R, Dullinger S, Klöner G, van Kleunen M,
 459 Essl F, Bossdorf O, Haeuser E, Talluto MV, Moser D, Block S, Conti L, Dullinger I, Münkemüller T, Thuiller W.
 460 2018. Simulating plant invasion dynamics in mountain ecosystems under global change scenarios. *Global Change*
 461 *Biology* 24: 289-302.

462 Castrignanò A, Costantini EAC, Barbetti R, Lopez N. 2006. Assessing and describing multi-scale spatial variabil-
 463 ity using factorial kriging. *Geophysical Research Abstracts* 8:10245

464 Celesti-Grapow L, Alessandrini A, Arrigoni PV, Banfi E, Bernardo L, Bovio M, Brundu G, Cagiotti MR, Camarda
 465 I, Carli E, Conti F, Fascetti S, Galasso G, Gubellini L, La Valva V, Lucchese F, Marchiori S, Mazzola P,
 466 Peccenini S, Poldini L, Pretto F, Prosser F, Siniscalco C, Villani MC, Viegi L, Wilhalm T, Blasi C. 2009. Invento-
 467 ry of the non-native flora of Italy . *Plant Biosystems* 143: 386–430.

468 Chiarucci A. 2007. To sample or not to sample. That is the question. *Folia Geobotanica* 42: 209-216.

469 Chiarucci A, Bacaro G, Rocchini D. 2008. Quantifying plant species diversity in a Natura 2000 network: Old ideas
 470 and new proposals. *Biological Conservation* 141: 2608-2618.

471 Chiarucci A, Bacaro G, Filibeck G, Landi S, Maccherini S, Scoppola A. 2012. Scale dependence of plant species
 472 richness in a network of protected areas. *Biodiversity and Conservation* 21: 503-516.

473 Chiarucci A, Fattorini S, Foggi B, Landi S, Lazzaro L, Podani J, Simberloff D. 2017. Plant recording across two
 474 centuries reveals dramatic changes in species diversity of a Mediterranean archipelago. *Scientific Reports* 7: 5415.
 475 <https://doi.org/10.1038/s41598-017-05114-5>

476 Chytrý M, Jarošík V, Pyšek P, Hájek O, Knollová I, Tichý L, Danihelka J. 2008. Separating habitat invasibility by
 477 alien plants from the actual level of invasion. *Ecology* 89:1541–53.

478 Chytrý M, Pyšek P, Wild J, Pino J, Maskell LC, Vilà M. 2009. European map of alien plant invasions based on the
479 quantitative assessment across habitats. *Diversity and Distributions* 15: 98–107.

480 Conti F, Abbate G, Alessandrini A, Blasi C. 2005. An Annotated Checklist of the Italian Vascular Flora. Palombi,
481 Roma.

482 Cornell HV, Karlson RH. 1997. Local and regional processes as controls of species richness. In: Tilman D, Ka-
483 reiva P (eds) *Spatial ecology: the role of space in population dynamics and interspecific interactions*. Princeton
484 University Press, Princeton, pp 250–268.

485 Crutzen PJ. 2006. The “anthropocene”. In *Earth system science in the anthropocene*. Springer, Berlin, Heidelberg,
486 pp 13-18.

487 Davies KF, Chesson P, Harrison S, Inouye BD, Melbourne BA, Rice KJ. 2005. Spatial heterogeneity explains the
488 scale dependence of the native-exotic diversity relationship. *Ecology* 86: 1602-1610.

489 Deutschewitz K, Lausch A, Kühn I, Klotz S. 2003. Native and alien plant species richness in relation to spatial
490 heterogeneity on a regional scale in Germany. *Global Ecology and Biogeography* 12: 299-311.

491 Dimitrakopoulos PG, Koukoulas S, Galanidis A, Delipetrou P, Gounaridis D, Touloumi K, Arianoutsou M. 2017.
492 Factors shaping alien plant species richness spatial patterns across Natura 2000 Special Areas of Conservation of
493 Greece. *Science of the Total Environment* 601: 461–468.

494 Dormann CF, Elith J, Bacher S, Buchmann C, Carl G, Carré G, Garcia Marquéz JR, Gruber B, Lafourcade B,
495 Leitão PJ, Münkemüller T, McClean C, Osborne PE, Reineking B, Schröder B, Skidmore Ak, Zurell D, Lauten-
496 bach S. 2013. Collinearity: A review of methods to deal with it and a simulation study evaluating their perfor-
497 mance. *Ecography* 36: 27–46.

498 Faliński JB. 1998. Invasive alien plants and vegetation dynamics. In: Starfinger U, Edwards KR, Kowarik I,
499 Williamson M (eds) *Plant invasions: ecological mechanisms and human responses*. Backhuys Publishers, Leiden,
500 pp 3–21.

501 Filibeck G, Petrella P, Cornellini P. 2016. All ecosystems look messy, but some more so than others: a case-study
502 on the management and acceptance of Mediterranean urban grasslands. *Urban Forestry and Urban Greening*
503 15:32–39.

504 Foxcroft LC, Pyšek P, Richardson DM, Genovesi P, MacFadyen S. 2017. Plant invasion science in protected areas:
505 progress and priorities. *Biological Invasions* 19:1353–1378.

506 Fridley JD, Stachowicz JJ, Naeem S, Sax DF, Seabloom EW, Smith MD, Stohlgren TJ, Tilman D, Von Holle B.
507 2007. The invasion paradox: reconciling pattern and process in species invasions. *Ecology* 88:3–17.
508 [http://dx.doi.org/10.1890/0012-9658\(2007\)88\[3:TIPRPA\]2.0.CO;2](http://dx.doi.org/10.1890/0012-9658(2007)88[3:TIPRPA]2.0.CO;2)

509 Gaston KJ. 2005. Biodiversity and extinction: species and people. *Progress in Physical Geography* 29: 239–247.

510 Gaston KJ, Jackson SF, Cantu-Salazar L, Cruz-Pinon G. 2008. The ecological performance of protected areas.
511 Annual Review of Ecology, Evolution, and Systematics 39: 93-113.
512 doi:10.1146/annurev.ecolsys.39.110707.173529

513 Geri F., Amici V., Rocchini D. 2010. Human activity impact on the heterogeneity of Mediterranean landscape.
514 Applied Geography 30: 370-379. doi:10.1016/j.apgeog.2009.10.006

515 Grime JP. 1973. Competitive exclusion in herbaceous vegetation. Nature 242: 344–347.

516 Grunanger P. 2001. Orchidacee d'Italia. Quaderni di Botanica Ambientale e Applicata 11: 3–80.

517 Guisan A, Zimmermann NE. 2000. Predictive habitat distribution models in ecology. Ecological Modelling
518 135:147-186.

519 Hannah L, Midgley G, Anelman S, Araujo M, Hughes G, Martinez-Meyer E, Pearson R, Williams P. 2007. Pro
520 tected area needs in a changing climate. Frontiers in Ecology and the Environment 5:131–138. doi:10.1890/1540

521 Huston MA. 1999. Local processes and regional patterns: appropriate scales for understanding variation in the
522 diversity of plants and animals. Oikos 86: 393–401.

523 Jaeger B. 2017. r2glmm: Computes R Squared for Mixed (Multilevel) Models. R package version 0.1.2

524 Kalwij JM, Robertson MP, van Rensburg BJ. 2015. Annual monitoring reveals rapid upward movement of exotic
525 plants in a montane ecosystem. Biological Invasions 17: 3517-3529.

526 Kumar S, Stohlgren TJ, Chong GW. 2006. Spatial heterogeneity influences native and nonnative plant species
527 richness. Ecology 87: 3186-3199.

528 Levine J, D'Antonio C. 1999. Elton revisited: a review of evidence linking diversity and invasibility. Oikos 87:
529 15–26.

530 Levine JM. 2000. Species diversity and biological invasions: relating local process to community pattern. Science
531 288: 852–854.

532 Lonsdale WM. 1999. Global patterns of plant invasions and the concept of invasibility. Ecology 80: 1522–1536.

533 Lucy FE, Roy H, Simpson A, Carlton JT, Hanson JM, Magellan K, Campbell ML, Costello MJ, Pagad S, Hewitt
534 CL, McDonald J, Cassey P, Thomaz SM, Katsanevakis S, Zenetos A, Tricarico E, Boggero E, Groom QJ, Adri-
535 aens T, Vanderhoeven S, Torchin M, Hufbauer R, Fuller P, Carman MR, Conn DB, Vitule JRS, Canning-Clode J,
536 Galil BS, Ojaveer H, Bailey SA, Theriault TW, Claudi R, Gazda A, Dick JTA, Caffrey J, Witt A, Kenis M,
537 Lehtiniemi M, Helmsaari H, Panov VE. 2016. INVASIVESNET towards an international association for open
538 knowledge on invasive alien species. Management of Biological Invasions 7:131–139.

539 Maestre FT. 2004. On the importance of patch attributes, environmental factors and past human impacts as deter-
540 minants of perennial plant species richness and diversity in Mediterranean semiarid steppes. Diversity and Distri-
541 butions 10: 21–29.

542 Marini L, Gaston KJ, Prosser F, Hulme PE. 2009. Contrasting response of native and alien plant species richness
543 to environmental energy and human impact along alpine elevation gradients. *Global Ecology and Biogeography*
544 18:652–661.

545 McCune B, Keon D. 2002. Equations for potential annual direct incident radiation and heat load. *Journal of Vege-*
546 *tation Science* 13: 603–606.

547 McKinney ML. 2002. Do human activities raise species richness? Contrasting patterns in United States plants and
548 fishes. *Global Ecology and Biogeography* 11: 343–348.

549 Meiners SJ, Pickett STA. 2013. Plant invasion in protected landscape: exception or expectation? In: Foxcroft LC,
550 Richardson DM, Pyšek P, Genovesi P (eds) *Plant invasions in Protected Areas*. Springer, New York

551 Moser D, Dullinger S, Englisch T, Niklfeld H, Plutzer C, Sauberer N, Zechmeister HG, Grabherr G. 2005. Envi-
552 ronmental determinants of vascular plant species richness in the Austrian Alps. *Journal of Biogeography*
553 32: 1117–1127. doi: 10.1111/j.1365-2699.2005.01265.x

554 Moustakas A., Voutsela A., Katsanevakis S. 2018. Sampling alien species inside and outside protected areas: Does
555 it matter? *Science of the Total Environment* 625: 194-198. doi: 10.1016/j.scitotenv.2017.12.198

556 Paradis E, Schliep K. 2018. ape 5.0: an environment for modern phylogenetics and evolutionary analyses in R.

557 Parendes LA, Jones JA. 2000. Role of light availability and dispersal in exotic plant invasion along roads and
558 streams in the H.J. Andres Experimental Forest, Oregon. *Conservation Biology* 14: 64–75.

559 Pauchard A, Alaback PB. 2004. Influence of elevation, land use, and landscape context on patterns of alien plant
560 invasions along roadsides in protected areas of south central Chile. *Conservation Biology* 18: 238–248.

561 Pausas JG. 1994. Species richness patterns in the understorey of Pyrenean *Pinus sylvestris* forest. *Journal of*
562 *Vegetation Science* 5: 517- 524.

563 Pignatti S. 1982. *Flora d'Italia* (3 vols). Edagricole, Bologna.

564 Planty-Tabacchi AM, Tabacchi E, Naiman RJ, Deferrari C, Decamps H. 1996. Invasibility of species-rich com-
565 munities in riparian zones. *Conservation Biology* 10: 598– 607.

566 Pressey RL, Visconti P, Ferraro PJ. 2015. Making parks make a difference: poor alignment of policy, planning and
567 management with protected-area impact, and ways forward. *Philosophical Transactions of the Royal Society B*
568 370: 20140280. doi:10.1098/rstb.2014.0280

569 Pyšek P, Prach K, Mandák B. 1998. Invasions of alien plants into habitats of Central European Landscape: an his-
570 torical pattern. In Starfinger U, Edwards KR, Kowarik I, Williamson M (eds) *Plant invasions: ecological mecha-*
571 *nisms and human responses*. Backhuys Publishers, Leiden, pp 23-32.

572 Pyšek P, Jarosík V, Kucera T. 2002a. Patterns of invasion in temperate nature reserves. *Biological Conservation*
573 104: 13–24.

574 Pyšek P, Sadlo J, Mandak B. 2002b. Catalogue of alien s plants of the Czech Republic. *Preslia* 74:97–186.

Pyšek P, Jarošík V, Kučera T. 2003. Inclusion of native and alien species in temperate nature reserves: an historical study from Central Europe. *Conservation Biology* 17: 1414-1424.

Pyšek P, Richardson DM. 2006. The biogeography of naturalization in alien plants. *Journal of Biogeography* 33: 2040-2050.

Quantum GIS Development Team 2016, version 2.14. QGIS Geographic Information System. Open Source Geospatial Foundation Project. <http://qgis.osgeo.org>

R Core Team 2019. R: A language and environment for statistical computing. R Foundation for Statistical Computing, Vienna, Austria. ISBN 3-900051-07-0, URL <http://www.R-project.org/>

Rey-Benayas JM. 1995. Patterns of diversity in the strata of forest in British Columbia. *Journal of Vegetation Science* 6: 95-98.

Richardson DM, Pyšek P. 2006. Plant Invasions: Merging the Concepts of Species Invasiveness and Community Invasibility. *Progress in Physical Geography*, 30, 409-431. <http://dx.doi.org/10.1191/0309133306pp490pr>

Sala OE, Chapin IFS, Armesto JJ, Berlow E, Bloomfield J, Dirzo R, Huber Sanwald E, Huenneke LF, Jackson RB, Kinzig A, Leemans R, Lodge DH, Mooney HA, Oesterheld M, Leroy Poff N, Sykes MT, Walker BH, Walker M, Wall DH. 2000. Global biodiversity scenarios for the year 2100. *Science* 287: 770-1774.

Sandel B, Corbin JD. 2010. Scale, disturbance and productivity control the native-exotic richness relationship. *Oikos* 119: 1281-1290

Sax DF. 2002. Native and naturalized plant diversity are positively correlated in scrub communities of California and Chile. *Diversity and Distributions* 8: 193 -210.

Shea K, Chesson P. 2002. Community ecology theory as a framework for biological invasions. *Trends in Ecology & Evolution* 17: 170-176.

Shochat E, Warren PS, Feath SH, McIntyre NE, Hope D. 2006. From patterns to emerging processes in mechanistic urban ecology. *Trends in Ecology & Evolution* 21: 186-191.

Simberloff D, Martin JL, Genovesi P, Maris V, Wardle DA, Aronson J, Courchamp F, Galil B, García-Berthou E, Pascal M, Pyšek P, Sousa R, Tabacchi E, Vilà M. 2013. Impacts of biological invasions: what's what and the way forward. *Trends in Ecology & Evolution* 28: 58-66.

Siniscalco C, Barni E, Bacaro G. 2011. Non-native species distribution along the elevation gradient in the western Italian Alps. *Plant Biosystems* 145: 150-158.

Souza L, Bunn WA, Simberloff D, Lawton RM, Sanders NJ. 2011. Biotic and abiotic influences on native and exotic richness relationship across spatial scales: favourable environments for native species are highly invasible. *Functional Ecology* 25: 1106-1112.

Spear D, Foxcroft LC, Bezuidenhout H, McGeoch MA. 2013. Human population density explains alien species richness in protected areas. *Biological Conservation* 159: 137-147. doi:10.1016/j.biocon.2012.11.022

608 Stadler J, Trefflich A, Klotz S, Brandl R. 2000. Exotic plant species invade diversity hot spots: the alien flora of
609 northwestern Kenya. *Ecography* 23: 169–176.

610 Stevens GC. 1992. The elevational gradient in altitudinal range: an extension of Rapoport's latitudinal rule to alti-
611 tude. *American Naturalist* 140: 893-911.

612 Stohlgren TJ, Binkley D, Chong GW, Kalkhan MA, Schell LD, Bull KA, Otsuki Y, Newman G, Bashkin M, Son
613 Y. 1999. Exotic plant species invade hot spots of native plant diversity. *Ecological Monographs* 69: 25–46.

614 Stohlgren TJ, Jarnevich C, Chong GW, Evangelista PH. 2006. Scale and plant invasions: a theory of biotic ac-
615 ceptance. *Preslia* 78: 405–426.

616 Sukopp H. 1998. On the study of anthropogenic plant migrations in central Europe. In: Starfinger U, Edwards KR,
617 Kowarik I, Williamson M (eds) *Plant invasions: ecological mechanisms and human responses*. Backhuys Publish-
618 ers, Leiden, pp 43–56.

619 Symonds JE, Pither J. 2012. Multi-scale analyses of exotic and native plant species diversity within Canada's en-
620 dangered antelope-brush ecosystem. *Plant Ecology* 213:1263–1275.

621 Taylor AH, Allen JI, Clark PA. 2002. Extraction of a weak climatic signal by an ecosystem. *Nature* 416: 629–632.

622 Thuiller W, Gallien L, Boulangeat I, de Bello F, Munkemuller T, Roquet-Ruiz C, Lavergne S. 2010. Resolving
623 Darwin's naturalization conundrum: a quest for evidence. *Diversity and Distributions* 16: 461–475.

624 Tilman D. 1997. Community invasibility, recruitment limitation, and grassland biodiversity. *Ecology* 78:81–92.

625 Tomasetto F, Duncan RP, Hulme PE. 2013. Environmental gradients shift the direction of the relationship between
626 native and alien plant species richness. *Diversity and Distributions* 19:49-59.

627 Tordoni E, Napolitano R, Nimis P, Castello M, Altobelli A, Da Re D, Zago S, Chines A, Martellos S, Maccherini
628 S, Bacaro G. 2017. Diversity patterns of alien and native plant species in Trieste port area: exploring the role of
629 urban habitats in biodiversity conservation. *Urban Ecosystems* 20: 1151-1160.

630 Trombulak SC, Frissell CA. 2000. Review of ecological effects of roads on terrestrial and aquatic communities.
631 *Conservation Biology* 14: 18-30.

632 Tutin TG, Burges NA, Chater AO, Edmondson JR, Heywood VH, Moore DM, Valentine DH, Walters SM, Webb
633 DA. (eds) 1993. *Flora Europaea*, Vol. 1, 2nd edn. Cambridge University Press, Cambridge. Thomas CD, Palmer G
634 2015. Non-native plants add to the British flora without negative consequences for native diversity. *PNAS* 112:
635 4387-4392.

636 Tutin TG, Heywood VH, Burges NA, Moore DM, Valentine DH, Walters SM, Webb DA. 1964–1980. *Flora Eu-*
637 *ropaea* (5 vols.). Cambridge University Press, Cambridge .

638 Ullmann I, Heindl B. 1989. Geographical and Ecological Differentiation of roadside vegetation in temperate Eu-
639 rope. *Botanica Acta* 102: 261–269.

- 640 Ullmann I, Bannister P, Wilson JB. 1995. The vegetation of roadside verges with respect to environmental gradi-
641 ents in southern New Zealand. *Journal of Vegetation Science* 6: 131–142.
- 642 Venables WN, Ripley BD. 2002. *Modern Applied Statistics with S*. Fourth Edition. Springer, New York. ISBN 0 -
643 387-95457-0
- 644 Weber HE. 1995. *Rubus* L. In: Hegi G (ed) *Illustrierte Flora von Mitteleuropa*, Band IV/2A, 3 edn. Blackwell
645 Wissenschafts-Verlag, Berlin, pp 284–595.
- 646 Williamson M, Fitter A. 1996. The varying success of invaders. *Ecology* 77(6):1661–1666

Click here to view linked References

1 CONTRASTING PATTERNS OF NATIVE AND NON-NATIVE PLANTS IN A NETWORK OF PRO- 2 TECTED AREAS ACROSS SPATIAL SCALES

3 Sara Landi^{1,2*}, Enrico Tordoni³, Valerio Amici², Giovanni Bacaro³, Marta Carboni⁴, Goffredo Filibeck⁵, Anna
4 Scoppola⁵, Simonetta Bagella¹

5 1 University of Sassari, Department of Chemistry and Pharmacy, Italy

6 2 GREENARCO, Spin-Off Company of the Alma Mater Studiorum, University of Bologna, Italy.

7 3 University of Trieste, Department of Life Sciences, Italy

8 4 University of Roma Tre, Department of Sciences, Italy

9 5 University of Tuscia, Department of Agriculture and Forest Sciences (DAFNE), Italy

10 * Corresponding author e-mail address: slandi@uniss.it

11

12

13

14 **Abstract:** Networks of protected areas are fundamental for biodiversity conservation, but many factors determine
15 their conservation efficiency. In particular, on top of other human-driven disturbances, invasions by non-native
16 species can cause habitat and biodiversity loss. Jointly understanding what drives patterns of plant diversity and of
17 potential invasions non-native species in protected areas is therefore a priority. We tested whether the richness and
18 composition of native and non-native plant species (not necessarily invasive species) within a network of protected
19 areas follow similar patterns across spatial scales. Specifically, we addressed two-three questions: a) what is the
20 degree of congruence in species richness between species richness and composition of native and non-
21 native plant species? b) what is the degree of congruence between changes in the composition of native and
22 non-native species across ecological gradients reflect a similar turnover of native species along the same gradients,
23 in floristic assemblage variation environmental gradients? c) what are the main environmental and human disturb-
24 ance drivers controlling for species richness in these two groups of species?

25 Species richness and composition of native and non-native plant species were compared at two spatial scales,
26 namely at: the plot scale (10 m x 10 m) and the Protected Area scale (PA). In addition, we fit Generalized Linear
27 Mixed-Models (GLMMs) and Generalized Linear Models (GLMs) were applied on native and non-native species
28 richness to identify the most important drivers of species native and non-native species richness at plot and
29 PA each scale, focusing on environmental conditions (climate, topography) and on the main sources of human dis-
30 turbance in the area (land use and roads). We found a significant positive correlation was observed between the
31 response turnover of native and non-native species composition composition at both plot and PA scales, whereas
32 their species richness was only correlated at the larger PA scale. The lack of congruence between the richness of
33 native and non-native species at fine the plot scale was likely driven by differential responses to fine scale envi-

Commented [A1]:

Rev4: The only larger issue
the terms invasive and non-
interchangeably.
The distinction between the
as only a small proportion o
become invasive and so the
species is not necessarily c
invaders. Are
you able to distinguish any
non-natives? I have indicate
text
needs to be clarified within

Commented [A2]:

Rev2: overall, the abstract
clear. Many things are not e
meant by "moderate anthro
this paper? What types of d
why? Natura 2000 sites are
very end of the abstract, wh
didn't see reference to them
paper. The structure could b
questions and help the read
design and findings.
Sometimes you're comparin
plants, and sometimes you'
general. Throughout, you co
For example, question a in
sounds like two questions -
natives, and richness vs cor
you're grouping native and
no need to split those out. T
difficult to read, because the
when a comparison is being
groups of plants, and when
being compared to a third fa
species richness of native a
correlated at the PA scale?
with a third factor?).
Abstract is also a bit vague
respond to different drivers
be more specific?

Commented [A3]:

Rev 4: What kinds of drivers

Commented [A4]:

Rev 4: Between what? Nat
richness?

ronmental factors, with non-natives favoring drier climates and milder slopes and human disturbances (climate and slope). In addition, more non-native species were found closer to road-ways in the reserve network. In contrast, the congruence in the richness of native, while at the larger PA and non-native species at the broader PA scale native and non-native species largely was mainly driven by the common influence of PA area, but also by similar responses of the two groups of species to responded in a similar way to resource climatic availability and variability heterogeneity. Across spatial scales, richness of non-native species increased under moderate anthropogenic disturbances. Thus, our study highlights the strong spatial dependence of the relationship between native and non-native species richness and of their responses to environmental variation. Taken together, our results suggest that within the study region the introduction and establishment of non-native species would be more likely in warmer and dryer areas, with high native species richness at large spatial scale but intermediate levels of anthropogenic disturbances and mild slope inclinations and elevation at fine scale. In particular, non-native species seem to respond to different drivers at fine scale, highlighting the primary role of local abiotic conditions (such as climatic and slope) and habitat heterogeneity. At PA scale, native and non-native species richness was more affected by regional-scale factors such as climatic variables and anthropogenic disturbances. Such an exhaustive understanding of the factors that influence the spread of non-native species (some of which are invasive species such as *Robinia pseudoacacia*, or *Amaranthus retroflexus*) drivers of invasion, especially in networks of protected areas such as Natura 2000 sites is may be crucial to inform conservation managers on how to control or curb of those that could be or become problems of biological invasions non-native species especially in the light of ongoing global changes.

Keywords: plant species richness, Habitat Directive, habitat heterogeneity, Natura 2000, scale dependence, protected areas, human disturbance.

INTRODUCTION

The introduction and spread of non-native species (also known as alien or exotic species) is considered a major cause of habitat transformation and biodiversity erosion, especially in the Mediterranean biome (Sala et al. 2000, Lucy et al. 2016). Rather than only focusing on controlling species that are already established or naturalized in a territory, proactive management of biological invasions should aim at curbing the arrival or establishment of new non-native species (Spear et al. 2013) or to controlling their populations during early-stages of the invasion continuum (*sensu* Richardson and Pyšek 2006). Thus, identifying the features that make certain areas more likely to harbor many non-native species (which are also known to correlate with areas harboring many invasive species, Williamson & Fitter, 1996) along with the driving factors of previously successful invasions is an important baseline information to be used for a proper management of controlling non-native species spread and conservation goal strategies. In addition, understanding by understanding whether the same features also drive the distribution of native species can then help it might be possible to highlight potential conflicts or hotspots for biodiversity management. For example, if native species tend to be associated with similar conditions as non-native species, valuable areas of high biodiversity will inevitably overlap with hotspots of invasions (e.g. Stadler et al. 2000), potentially leading to greater risks of biodiversity loss as a consequence of non-native species impacts (Simberloff et al. 2013). These might thus represent joint priorities for conservation and non-native species control efforts, especially within reserve networks or specific areas with conservation interest protected area networks, such as Natura 2000 sites.

Commented [A5]:
Rev 4: Different than what?

73 An ~~enormous-large~~ body of literature has tried to identify general features or ecosystem properties facilitating
 74 the establishment of non-native species (e.g., Ashton and Mitchell 1989; Faliński 1998; Sukopp 1998; Pyšek and
 75 Richardson 2006; [Bjarnason et al. 2017](#)), and to understand whether the same features also promote native species
 76 richness and high overall biodiversity (e.g., Shea and Chesson 2002; Deutschevitz et al. 2003; Carboni et al. 2010;
 77 Tordoni et al. 2017). In general, environmental factors which influence patterns of species richness at regional
 78 scales include climate, landscape heterogeneity, ~~spatial patterns, and~~ geomorphological processes ~~and level of pro-~~
 79 ~~tection~~, all of which typically affect native as well as non-native species (Davies et al. 2005; Moser et al. 2005;
 80 Carboni et al. 2010). But in addition, in the current Anthropocene era (Crutzen 2006), human disturbance ~~and~~
 81 ~~management practices are is~~ a major agent of change of species richness and diversity patterns across spatial scales
 82 (Maestre 2004; Gaston 2005). For example, human disturbance may generate environmental heterogeneity, which
 83 may increase extinction risk of native species, but also allow for resource partitioning by creating new niche op-
 84 portunities (Shochat et al. 2006). These factors may thus facilitate the arrival of non-native species pre-adapted to
 85 such altered conditions (Callaway 2007). Indeed, while natural or near-natural ecosystems often display a certain
 86 ecological resistance against biological invasion (e.g., Faliński 1998; Simberloff et al. 2013), densely populated
 87 areas or areas subject to strong human disturbance are typically found to be prone to higher levels of [non-native](#)
 88 [species establishment and invasion success](#) (e.g., Pyšek et al. 1998; Sukopp 1998; McKinney 2002; Chytrý et al.
 89 2008; Tordoni et al. 2017).

90 The scale dependence of biodiversity patterns is a well-known issue in ecology (e.g., Huston 1999; Richardson
 91 and Pyšek 2006). In particular the relationship between native and non-native species richness seems to change
 92 across spatial scales, which led scientists to even coin the term “invasion paradox” (Shea and Chesson 2002; Frid-
 93 ley et al. 2007). Specifically, a negative relationship is usually observed at ~~a small—finer~~ small spatial scales, ~~that~~
 94 ~~are those in at~~ which species typically interact (e.g., Cornell and Karlson 1997; Levine 2000; Tilman 1997), but
 95 this relationship tends to become positive when increasing the grain of the sampling units or the extent of the study
 96 area (Planty-Tabacchi et al. 1996; Lonsdale 1999; Levine 2000; Stadler et al. 2000; McKinney 2002; Sax 2002;
 97 Kumar et al. 2006; Stohlgren et al. 2006). [Although](#) there has been a heated debate to try to explain these con-
 98 trasting patterns, this conundrum of invasion ecology is still far from being resolved. Further studies are thus
 99 needed in order to understand the relationships between native and non-native species [patterns](#) and the environ-
 100 mental and anthropogenic features which foster biological invasions across spatial scales.

101 In the Convention on Biological Diversity (art. 8 *In-situ Conservation*, <https://www.cbd.int/>), prevention measures,
 102 control, or eradication of problematic non-native species are called for. Identifying pathways for non-native spe-
 103 cies establishment and management priorities are possible actions in this direction. In particular, protected areas
 104 are a key component of the global response to environmental change and degradation (e.g., Hannah et al. 2007;
 105 Gaston et al. 2008; Foxcroft et al. 2017), and can ~~offer be part of~~ a framework to devise effective invasion control
 106 measures. Nevertheless, they face many challenges, such as the effectiveness of reserve design, governance (Pres-
 107 sey et al. 2015), and anthropogenic change (Foxcroft et al. 2017), with generally ~~little few~~ restrictions currently in
 108 place for preventing the introduction of non-native species (Pyšek et al. 2003). Most protected areas in Europe are
 109 in a mosaic of land use types that can form a network of potential sources for non-native species introductions
 110 (e.g., Foxcroft et al. 2007; Meiners and Pickett 2013). [In addition, there is no recent evidence shows that there is](#)
 111 [almost no difference in the patterns of non-native and invasive species inside and outside protected areas.](#)

Formatted: English (United Kingdom)

Commented [A6]:
 Rev4: Maybe this shouldn't
 environmental factors if you
 from humans, but in the nex
 factors.

112 ~~i.e. suggesting that currently habitat protection has little or no effect on non-native species richness (e.g. Moustakas~~
113 ~~et al. 2018).~~ ~~SStudyingStudying~~ which features are linked to higher invasion levels can thus help identify the main
114 pathways that need regulation and which areas are most at risk, to guide future conservation planning within pro-
115 tected area networks.

116 In this study, we investigated the importance of biotic (represented by native species richness and composition),
117 environmental (climate) ,~~landscape diversity~~ and anthropogenic (road network ~~and land uses~~) factors in driving
118 non-native plant species spatial patterns within a network of protected areas in central Italy. We aimed at provid-
119 ing insights on the ecological mechanisms useful for the effective control of ~~biological invasions~~~~non-native spe-~~
120 ~~cies establishments~~, which is extremely important in the context of the management of reserve networks. Ideally,
121 reserve networks strive to maximize the protection of biodiversity features, while non-native species clearly repre-
122 sent a potential threat for nature reserves and their management (Pyšek et al. 2003). If native and non-native spe-
123 cies follow similar patterns within the network, ~~local conflicts between an overlap of~~ high native biodiversity and
124 invasion hotspots ~~can be likely to potentially~~ emerge. Identifying such ~~potential conflicts between natives and non-~~
125 ~~native species~~~~areas at high risk of biodiversity loss~~ and the ecological features that may promote invasion or oth-
126 erwise hamper ~~high~~-biodiversity protection is therefore essential to improve control and management actions.

127 Specifically, ~~w~~We aim at testing, ~~at different spatial scales~~, whether ~~composition and~~ species richness ~~patterns-~~
128 ~~levels of native and non-native plants- vascular plants~~ ~~species follow the same patterns~~~~are correlated of native-~~
129 ~~species at different spatial scales and whether similar factors control the turnover in species composition of native~~
130 ~~and non-native species along ecological gradients. Specifically, by answering the following specific~~~~our~~ research
131 questions are: ~~a) what is the degree of congruence in species richness between native and non-native plant spe-~~
132 ~~cies, and does this relationship vary across spatial scales? b) do changes in the composition of non-native species~~
133 ~~across environmental gradients reflect native species turnover along the same gradients? what is the degree of-~~
134 ~~congruence, between native and non-native species, in floristic assemblage variation across environmental gradi-~~
135 ~~ents? c) what are the main biotic environmental and abiotic anthropogenic drivers controlling species richness in~~
136 ~~each of these two groups of species?~~

137

138 ~~a) is there a relationship between composition and richness of native and non-native species and does this relation-~~
139 ~~vary across spatial scales? b) is native species richness affected by the same abiotic and human-mediated factors-~~
140 ~~affecting non-native species across spatial scales?~~

141 METHODS

142 Study area

143 This study was performed in the local network of protected areas (PAs) of the province of Siena (Italy), including
144 ~~four~~ Nature Reserves (designated under national or regional regulations) and 17 Special Areas of Conservation
145 (SACs, designated under the EU Habitat Directive 93/43/EEC) (Figure 1). The size of the single PAs within the
146 network ranges from 268 ha to 13747 ha, while their elevation ranges from 122 m to 1660 m a.s.l. (Chiarucci et al.
147 2012) with a cumulative area of 593 km² (15.6% of the Province).

Commented [GF7]:

vedi commento mio più sopra
questa faccenda del conflitto
sostenere, oppure va spiegato
perdere l'idea del conflitto co-
studiare la congruenza fra le
come motivazione dello studio
genericamente l'interesse di
concentreranno più naturali
richness delle native può es-
predittore delle zone di possi-
aliene). Un altro motivo di i-
(forse era questo che intenc-
proprio laddove ci sono più
state istituite le riserve, c'è
naturalizzazioni di aliene e
particolarmente in campana
cosa veramente teorica....

Marta: ah ecco, non mi ricordo
qualcosa qui... va bene, sono
può impostare anche come
forse cambiato qua e là in ta-
ho lasciato un po' com'era to-
"conflicts" che creava confu-
discorso di overlap delle ale-
vista conservazionistico. Per
articoli di conservazione su
diversità funzionale e tassono-
congruence, ecc, ma con il
sarebbe più auspicabile il co-
solo accennato velocemente
bene... Però il fatto che se le
native e rispondono agli ste-
overlap delle aree più biodi-
sembra solo teorico: è per f

148 The study area is characterized by a Mediterranean macro-climate, even though there is a strong variation across
149 sampling sites due to differences in morphology and local elevations (Castrignanò et al. 2006). Long term mean
150 annual precipitation ranges from 630 to 1275 mm (Barazzuoli et al. 1993). The highest precipitations (above 1000
151 mm on average) and lowest mean annual temperature values (lower than 12 °C) are found at higher elevations
152 (Monte Amiata). A relatively arid and warm zone (mean annual temperature of about 14 °C and average precipita-
153 tions of ca. 600 mm) is localized in the South-East of the province of Siena (Orcia river valley; Barazzuoli et al.
154 1993).

155 The geology is rather varied and complex (including, *inter alia*, limestone, clay, marl, metamorphic and volcanic
156 bedrock), resulting in highly heterogeneous morphology and a great variety of landscapes. The main land-cover
157 types include evergreen coppice woods (dominated by *Quercus ilex*) and deciduous coppice woods or forests
158 (thermophilous types dominated by *Quercus pubescens* or *Q. cerris* at lower elevations, and mesophilous types
159 dominated by *Castanea sativa* or *Fagus sylvatica* at higher elevations); evergreen Mediterranean shrublands
160 (characterized e.g. by *Erica arborea*, *E. scoparia*, *Phillyrea latifolia*, *Pistacia lentiscus*, *Arbutus unedo*, *Cistus*
161 *salvifolius*, *Juniperus communis* and *J. oxycedrus*); croplands (mainly wheat and horticultural crops), vineyards
162 and olive groves. Other relevant land cover types include pastures, meadows, garigues (on calcareous and ultra-
163 mafic substrates), conifer plantations and wetlands.

164 **Sampling design**

165 Plants were sampled by using an operational approach of plant communities (Chiarucci 2007), defined by a fixed
166 grain and uniform sample density within each protected area (PA). Sampling design was based on a grid of
167 1 km × 1 km cells, covering the whole study area, with a sampling point randomly selected within each cell (Chi-
168 arucci et al. 2008, 2012). A sampling unit represented by a square plot of 10 m x 10 m (plot) was centered at each
169 sampling point. This was further divided in 16 contiguous squared 2.5 m x 2.5 m subplots (Figure S1 of Supple-
170 mentary material), on which the presence/occurrence of all species (or subspecies) of vascular plants was recorded.
171 At plot scale, frequency of each species, calculated as the sum of occurrences within the 16 subplots, was
172 used as a coarse measure of abundance. The field data collection was performed from April to June, during the
173 years 2005-2009. The total number of sampled plots was 604.

174 **Native and non-native species richness and composition**

175 All the vascular plants recorded within each plot were identified at the species or subspecies level with standard
176 floras (Pignatti 1982; Tutin et al. 1964–1980, 1993) or monographs (Grunanger 2001; Weber 1995). Nomenclature
177 was standardized according to Conti et al. (2005). Plants were classified as native or non-native, depending on
178 their distributional status as given by Celesti-Grapow et al. (2011).

179 Composition and species richness for both native and non-native species were then separately assessed at plot and
180 PA scale, i.e. two data matrices were prepared as separate community matrices (species by plots and species by
181 PAs). At plot scale, frequency of each species, calculated as the sum of occurrences within the 16 subplots, was
182 used as a coarse measure of abundance. At PA scale, relative frequency of each species, measured as the ratio
183 between the number of occupied plots versus the number of recorded plots, was used as measure of abundance.

184 Species composition and richness values at the PA-scale [species lists](#) were obtained by pooling the data from
185 the plots included within each PA.

186 Environmental and human disturbance predictors

187 At plot scale, four [categories-groups](#) of predictor variables for evaluating the environmental ([first two](#)), and human-
188 mediated ([last two](#)) [and spatial](#) factors affecting [native and non-native](#) species richness were considered: 1) climat-
189 ic, 2) topographic, [and](#) 3) [land use, and human](#) disturbance [variables](#), and 4) [geographical variables](#) ([Quantum](#)
190 [GIS Development Team 2016, version 2014](#); Table 1):

191 1) Climatic variables: [we derived one synthetic variable named climate. We obtained this variable was derived](#)
192 [based on from](#) 36 climatic variables [related mainly to annual and monthly temperatures and precipitations, in-](#)
193 [cluding \(e.g. mean annual temperature, maximum annual temperature, minimum annual temperature, absolute-](#)
194 [maximum temperature, absolute minimum temperature, and total annual rainfall\) mean monthly data for total](#)
195 [rainfall, minima and maxima temperatures obtained](#) for each plot, [the Data source was from the](#) LaMMa consortium
196 (Laboratorio di Monitoraggio e Modellistica Ambientale per lo sviluppo sostenibile;
197 [http://www.lamma.rete.toscana.it/en](#)), [which Specifically the LaMMa data consisted of has available provides](#) local
198 [interpolated](#) climatic [data for on a](#) grids with a resolution of 250 m x 250 m [for the area](#). Since these variables were
199 highly [inter](#)-correlated, a Principal Component Analysis (PCA) was performed to reduce their multicollinearity
200 (Taylor et al. 2002). Given that the first PCA factor explained more than 90% of variance ([Figure S2 of Supple-](#)
201 [mentary material](#)), this was used as a single variable (hereafter called *climate*). This was found to be negatively
202 correlated with [all the variables related to](#) total and summer rainfall (Pearson correlation coefficients between -1
203 and -0.85), positively correlated with the minimum temperatures (Pearson correlation coefficients between 0.77
204 and 0.83) and maximum ones (Pearson correlation coefficients between 0.84 and 0.90). This compound *climate*
205 variable corresponds therefore to a gradient from wet and cold to warm and (summer-) dry conditions, that is from
206 Temperate to Mediterranean [conditions](#) [meso-climate](#), moving from the negative to the positive extreme of the axis.

207 [ables: latitude and longitude of each center](#) |

209 2) Topographic variables: *elevation*, *slope*, and potential *solar radiation* values (the latter was obtained from slope,
210 aspect and latitude, following McCune and Keon 2002). [The resolution used to develop these variables was 30 m.](#)

211 3) [Human disturbance variables: distance to the nearest road \(paved or unpaved\) and land-use type variables. We](#)
212 [used the distance to the nearest road as a proxy of human disturbance as the road network is a well-known intro-](#)
213 [duction pathway for non-native species \(e.g. Pauchard and Alaback 2004; Arévalo et al. 2010\). Road layer have](#)
214 [been derived from GEOScopio Geoportal, Tuscany Region - Territorial and Environmental Information System](#)
215 [\(http://www.Geografia.toscana.it/\)](#)
216 [We obtained 9 land-use type variables we obtained 9 dummy variables reflecting the different land-use types found](#)
217 [in the area by recording the II level of the CORINE Land Cover \(CLC\) of each plot based on a Corine Land Cover](#)
218 [map \(II level\) \(III level\) of the area from the Copernicus database \(https://land.copernicus.eu/pan-european/corine-](#)
219 [land-cover\).](#) In addition, [we calculated landscape diversity \(\$H_{plot}\$ \) and landscape evenness \(\$E_{plot}\$ \) within a buffer](#)
220 [area of 250 m radius around the plot were assessed. For this we used the based on the](#) Shannon Index of diversity
221 and the Pielou Index of evenness applied on the CLC map. [At plot scale, the human disturbance variables linked to](#)

Commented [MC8]:
notare che qui land use era
human/antropogenic variab

come decidiamo di proced
avere senso... ma facciamo
e altre parti a trattarle così

Commented [A9]:
Rev2: Line 158: what was th
topographic model used to

Formatted: Line spacing: 1

land use have not been considered due to the resolution of the CLC map which did not allow to detect artificial land use types in the plots or in related buffer areas. So we used artificial land use types as proxy of human disturbance only at protected area scale.

–(III level).

|

Human disturbance variable: the distance to the nearest road (paved or unpaved) was used to calculate the disturbance-related variable *road distance* as a proxy of human disturbance as the road network is a well-known introduction pathway for non-native species (e.g. Pauchard and Alaback 2004; Arévalo et al. 2010). At PA scale, we considered the same groups of predictor variables as at the plot scale were considered, with some unavoidable differences in the definition of the single predictors due to different spatial resolution (Quantum GIS Development Team 2016, version 2.14; Table 2):

1) Climatic variables: the variables per PA were based on estimated values on the two of climatic variables calculated per each plot, as described above. In particular, data for each plot as previously reported. Then, since each PA contains more than one plot, the variability of climate across plots was accounted for by calculating two derived climatic variables at PA scale: two climatic variables per PA were obtained as the follow: from the PCA described above: mean climate (calculated as the centroid, along the first axis in the climate PCA, axis of the cloud of points-plots belonging to each PA) and climate range (calculated as the range of the same cloud of points along the PCA axis). The gradient remains the same described above for the climate variable at plot scale.

2) Planar surface Geographical variables: area of each PA expressed in km².

3) Topographic variables: elevation range and mean elevation within each PA. |

3) Human disturbance variables: total road density and land use type variables. We obtained the total road density as the linear extension of paved and unpaved roads per km² within each PA and was considered as a proxy of human disturbance and potential propagule pressure of non-native species.

Moreover, as human disturbance, we obtained 4 land-use type variables reflecting the different artificial land use types found in each PA based on Corine Land Cover map (I level) of the area extracted from the Copernicus database (<https://land.copernicus.eu/pan-european/corine-land-cover>). Land use types variables, artificial, semi-natural and natural, were expressed as percentages of relative Corine land use classes (human, agriculture, natural, wetland). The variables *landscape diversity* (H_{PA}) and *landscape evenness* (E_{PA}) were calculated using the Shannon Index of diversity and the Pielou Index of evenness on the III level CLC map. The calculation of the landscape diversity was done at the III level of CLC because this allowed for a finer classification of landscape units, that was not necessary for detecting the dominant land use type.

Moreover, as human disturbance, we obtained 4 land-use type variables reflecting the different land use types found in each PA based on a Corine Land Cover map (I level) of the area from the Copernicus database (<https://land.copernicus.eu/pan-european/corine-land-cover>). Land use cover type variables were as percentages of land use classes (human, agriculture, natural, wetland, and water body) were calculated from the CLC map (I

Commented [A10]:

Rev2: Line 166: what level were used for roads? At least road density can be very sensitive. Or were you using aerial images?

Commented [A11]:

Rev4: Could you explain what variables here but only one

Commented [A12]:

Valerio potresti controllare l

Commented [A13]:

Rev3: lines 176-181 and 182 are using the level 3 (most detailed) classification of land use types: artificial and agricultural land. The authors have chosen to use anthropogenic pressures or are part of a CLC classification of artificial land use types in the used.

See for example the methodology <https://doi.org/10.3389/feart.2019.00011>. In my view the authors should use habitat richness or cover in the disturbance variable.

level) to obtain 5 variables for the land cover type. Then, the variables *landscape diversity* (H_{PA}) and *landscape evenness* (E_{PA}) were calculated using the Shannon Index of diversity and the Pielou Index of evenness on the III level CLC map. The calculation of the landscape diversity was done at the III level of CLC because this allowed for a finer classification of landscape units, that was not necessary for detecting the dominant land-use type.

5) Human disturbance variable: total road density was measured as the linear extension of for paved and unpaved roads per km² within each PA and was considered as a proxy of human disturbance and potential propagule pressure of non-native species. 5) Geographical variables: *latitude and longitude of the centroid, and area* (expressed in km²) of each PA area.

STATISTICAL ANALYSES

Separate analyses were performed at the plot and PA scales. At each scale, we assessed: 1) the correlation between native and non-native species richness, 2) whether specific sets of non-native species tend to be associated with specific native species assemblages, 2) the correlation between native and non-native species richness, and 3) which predictors best explained native and non-native species richness, respectively by using Minimum Adequate Model selection procedures on Generalized Linear Mixed-effects Models (GLMMs, plot scale) and Generalized Linear Models (GLMs, PA scale). Multicollinearity among explanatory variables at each scale was tested computing Spearman's ρ (correlation matrix available in Table S1 of the Supplementary material) and ensuring that in the minimum adequate models there were not variables where $|\rho| > 0.7$ (Dormann et al. 2013).

Relationships between native and non-native species

First, we assessed the correlation between native and non-native species richness by computing the Spearman correlation coefficient ρ at both spatial scales. This was done to assess whether native and non-native species richness within the reserve network followed 1) similar trends, suggesting that factors which favor high richness of native species, also increase opportunities for non-native species (Thuiller et al. 2010), leading to a positive native-non-native richness relationship (Shea and Chesson 2002), or 2) opposite trends suggesting higher biotic resistance of more diverse native communities, as interpreted by several authors (Cornell and Karlson 1997; Stohlgren et al. 1999; Levine 2000; Tilman 1997; Brown and Peet 2003; Davies et al. 2005; Souza et al. 2011).

Second, in order to we verified whether the same environmental and anthropogenic turnover in native species composition was associated also to turnover in non-native species and thus if specific sets of non-native species tend to always be associated with the same set of natives. To do so, a comparison we first obtained plot-to-plot (and PA-to-PA) compositional dissimilarities in species composition, separately for non-native and for native species, using Bray Curtis pairwise dissimilarities on $\log(x+1)$ transformed species frequencies. was

performed. To do so, we tested whether the pairwise dissimilarity matrices (or distance matrices) of distances among plots in native and non-native sub-communities were linearly independent by performing a Mantel test (Spearman correlation, 999 permutations, McCune and Keon 2002). For this we considered only the plots or PAs that had at least one non-native species. Then, we assessed the correlation between native and non-native species richness by computing the Spearman correlation coefficient ρ at both spatial scales available. This was done to assess whether native and non-native species richness within the reserve network followed 1) similar trends, suggesting that factors which favor high richness of native species, also increase opportunities for non-native species (Thuiller et al. 2010), leading to a positive native-non-native richness relationship (Shea and Chesson 2002), or 2) opposite trends suggesting higher biotic resistance of more diverse native communities, as interpreted by several authors (Cornell and Karlson 1997; Stohlgren et al. 1999; Levine 2000; Tilman 1997; Brown and Peet 2003; Davies et al. 2005; Souza et al. 2011). Also this relationship was tested at the two spatial scales.

Determinants of species richness across spatial scales

To assess the relationship between vascular plant richness (separately for native and non-native species) vs. response variables and the environmental and human disturbance predictors, we used two different approaches depending on the scale of investigation (plot or PA scale). In both cases, we first evaluated the presence of spatial autocorrelation in the response variables by calculating Moran's I coefficient using the R package 'ape' (Paradis and Schliep 2018), based on the geographical coordinates of the plots and of the centroid of the PA, respectively. Also, multicollinearity among explanatory variables at each scale was tested by computing Spearman's ρ (correlation matrix available in Table S1 of the Supplementary material) and ensuring that there were no pairs of variables in the models where $|\rho| > 0.7$ (Dormann et al. 2013).

At plot scale, we fit Generalized Linear Mixed Models (GLMM) using R package "lme4" (Bates et al. 2015) and assuming Poisson family errors. Furthermore, PA was considered as random effect to control for the spatial dependence of plots within PA. At PA scale, Generalized Linear Models (GLM) approach was used using Poisson family errors; in case of overdispersion in the data a quasipoisson family was used instead. A Minimum Adequate Model (MAM) and a set of models with good support were thus obtained by performing a stepwise variable selection procedure through AICc minimization using R package "MuMIn" (Barton 2019). Only the models with $\Delta AICc \leq 2$ compared to the best model were considered to have good support (Burnham and Anderson 2002). In case of more models with similar AICc values, the one which retained a lower number of predictors was considered as minimum adequate model (MAM) according to Occam's razor. Note that, since results were qualitatively similar across the set of models with good support (Table S2 of Supplementary material), we present and discuss only the MAMs in the main text. In case of overdispersion in the data, it was not possible to use an AICc-based selection approach, then the amount of deviance accounted for by the GLM adjusted by the number of predictors was considered instead ($D^2_{adjusted}$; Guisan and Zimmermann 2000; Barbosa et al. 2014). In addition, R^2 statistics (marginal effect) were derived for GLMMs using the "r2glmm" R package (Jaeger 2017). Prior to analysis, quantitative variables were standardized in order to have mean zero and unit variance. Any residual spatial autocorrelation after modeling was assessed by means of spline-correlograms using the R package "ncl" (Bjørnstad 2019). Specifically, 95% pointwise bootstrap confidence intervals were calculated from 5000 bootstrap samples of Pearson residuals after accounting for the level of spatial autocorrelation explained by the explanatory variables in each model.

Commented [A15]:
Rev2: Line 207: the relation parentheses isn't clear

Formatted: Font: Times Ne

Formatted: HTML Preform

Formatted: Font: Times Ne

Formatted: Font: Times Ne

334 sampling units were strongly spatially autocorrelated (natives: Moran's $I = 0.05$; $p < 0.001$; non-natives: Moran's I
 335 $= 0.03$, $p < 0.001$). For this reason, for further analyses we used GLMMs via Penalized Quasi-Likelihood
 336 ("GLMMPL"), using R package "MASS" (Venables and Ripley 2002) and assuming a Poisson family error.
 337 Spatial autocorrelation was taken into account by adding a matrix describing the correlation structure of the data
 338 (Gaussian correlation). Furthermore, PA was considered as random effect to control for the spatial dependence of
 339 plots within PA. Prior to analysis, quantitative variables were standardized in order to have mean zero and unit
 340 variance. A Minimum Adequate Model (MAM) was thus obtained performing a manual backward selection from
 341 the full dataset and considering potential unimodal relationships taking care to treat independently the two most
 342 correlated variables (*landscape diversity*, H_{plot} and *landscape evenness*, E_{plot}). Goodness of fit was evaluated com-
 343 paring R^2 statistics derived using "r2glmm" R package (Jaeger 2017).
 344 Unlike at the plot scale, no signals of spatial autocorrelation were detected at the PA scale for either native or non-
 345 native species richness (natives: Moran's $I = 0.05$; $p = 0.95$; non-natives: Moran's $I = -0.04$, $p = 0.80$). For this
 346 reason, classical Generalized Linear Models (GLMs) with a Poisson family error were used, and MAMs were ob-
 347 tained using a stepwise procedure aiming at AICc minimization by means of package "GLMULTI" R package
 348 (Calcagno 2013) plus a backward selection. As a measure of goodness of fit, the amount of deviance accounted for
 349 by each GLM (adjusted by the number of predictors) was computed ($D^2_{adjusted}$; Guisan and Zimmermann 2000;
 350 Barbosa et al. 2014).
 351 All the statistical analyses were conducted using R 3.6.5.1 (R Core Team 2019).

352 RESULTS

353 Species patterns at plot scale

354 In total, 993 native species and 48 non-native species were recorded in 604 plots. At the plot scale, plant species
 355 richness ranged from 0 to 117 (mean 31.0) for native species (one plot was found to harbor no species at all) and
 356 from 0 to 9 (mean 0.6) for non-native species. Native and non-native species richness exhibited a slight positive
 357 correlation, though not significant (Spearman $\rho = 0.06$, $p = 0.11$). In contrast, distance matrices based on plot-to-
 358 plot compositional dissimilarity showed a significant positive correlation (Mantel test, Spearman $\rho = 0.24$, $p =$
 359 0.001).

360 Table S23 reports the full list of candidate models for species richness predictors derived from the model selection
 361 procedure. Table 34 describes the best set of predictors for species richness in the GLMM MAM.
 362 Specifically, minimum adequate models showed, on the one hand, that native species richness were positively
 363 associated with landscape diversity (H_{plot}) and negatively with the first axis of PCA derived from climatic varia-
 364 bles (climate) and slope. On the other hand, non-native species richness were negatively related to elevation,
 365 road distance and slope, and positively associated with climate. However both models, however and in particular
 366 the native-species model, are characterized by a weak goodness of fit ($R^2 = 0.024$ and $R^2 = 0.116$ for native and
 367 non-native species, respectively).

368 Species patterns at protected area scale

369 Plant species richness within at PA scale ranged between 22 and 547 (mean 219.8 species) for native species and
 370 0-24 (mean 6.6 species) for non-native species. Species richness values of natives and non-native species (Figure 2)

Commented [A16]:
Rev4: Could you explain ho
matrix?

Formatted: Font: Times Ne
Formatted: HTML Preform

371 were positively and significantly correlated (Spearman $\rho = 0.75$, $p < 0.001$). Similarly, the Mantel correlation be-
 372 tween native and non-native species dissimilarity matrices was significant (Spearman $\rho = 0.33$ with $p = 0.003$).
 373 The relationship between the ratio of non-native species richness to native species richness and the native species
 374 richness proportion of non-native species per PA significantly increased with the size of the native flora. The rela-
 375 tive proportion of non-native species (ratio of non-native to native richness) in PAs was positively correlated with
 376 the richness of native species ($R^2 = 0.32$, $p < 0.01$), increasing up to a limit of about 5% (Figure 2). Similarly, the
 377 Mantel correlation between native and non-native species dissimilarity matrices was significant (Spearman ρ
 378 $= 0.33$ with $p = 0.003$). Further, according to the MAMs, species richness at PA scale was positively correlated
 379 with area and climate range for both species groups (Table 34). Table 4xx2 shows a positive relationship with
 380 area, mean climate and climate range was observed in both species' groups. In addition, native species richness
 381 showed a positive relationship with mean climate and a negative and not, although but not significant,
 382 relationship with % wetland. In contrast, non-native species richness was negatively related with mean elevation.
 383 showed a unimodal relationship with road density and a negative relationship with % agricultural area (Table 3).
 384 In both cases species groups, the deviance accounted for by the model was relatively high ($D^2_{\text{adjusted}} = 0.6976$ and
 385 $D^2_{\text{adjusted}} = 0.781$ for native and non-native species model, respectively).
 386 All four spline correlograms (Figure S32 of Supplementary material) failed to reveal any evidence of spatial auto-
 387 correlation in the residuals, thereby allowing us to exclude its influence on model parameter estimates.

388

389 DISCUSSION

390 The Italian flora is currently estimated to include 7634 taxa (species and subspecies), 13.4% of which (1023 taxa)
 391 are considered to be non-native (Celesti-Grapow et al. 2011). In our dataset, collected using a probabilistic sam-
 392 pling strategy within the protected areas of the Siena province only, the protected areas of the Siena province host
 393 a relatively we recorded a rich flora (1041 species were recorded by this survey) and a relatively low proportion of
 394 non-native species (4.9% of the whole sample and on average of 3.9% at the plot scale and 4.9% at the whole
 395 sample scale, respectively). However, even if the province of Siena is characterized, overall, by well a well pre-
 396 served landscape which is experiencing a process of abandonment of traditional land uses (Geri et al 2010), we
 397 should consider the fact that only protected areas have been were included into then this study and thus only better
 398 preserved most of the habitat are likely to have been sampled sites are on average less disturbed than the remain-
 399 der of the landscape at province (or country) scale. Even though the proportion of non-native species in the inves-
 400 tigated system is generally low, we found evidence of scale dependent effects of the relationship between native
 401 and non-native species richness. proportion of — relationship between native and non-native species rich-
 402 ness that increase from the plot to the PA scale and the non-native richness from not being correlated at the plot
 403 scale to haigh — and a strong While there was a strong positive relationship at the PA scale, native and non-native
 404 species richness were not correlated at the plot scale. This was the consequence of different responses of non-
 405 native species the two groups of species to environmental and anthropogenic factors at the fine scale, while similar
 406 factors (chiefly reserve area) explained both native and non-native species richness at large scale, in primis a simi-
 407 lar scale dependence (Chiarucci et al. 2012).

Commented [A17]:

Rev2: Line 248: unclear wh
 non-native species per PA
 the
 size of the native flora". Doe
 as native richness increase
 non-native species increase

Commented [A18]:

Rev4: I didn't see any discu
 shown in Figure 2 in the dis

Commented [A19]:

Rev4: These sentences are
 what point you are trying to

408 Species patterns at the plot scale¹

409 At the plot scale, we ~~did not observe a relation between native and non-native species richness but we~~ observed a
410 relationship between ~~the compositional gradients of native species composition~~ and non-native composition~~species~~.
411 This suggests that non-native species tend to be associated with specific sets of native species (*i.e.*, native plant
412 communities), either because they have similar environmental requirements or because of more direct biotic inter-
413 actions (*e.g.*, facilitation or competitive exclusion). However, ~~we also observed no~~ the lack of significant correla-
414 tion between native and non-native species richness~~which~~ instead suggests a limited role for biotic interactions
415 and for biotic resistance through competitive exclusion in these plant communities.

416 ~~Further~~Indeed, our results suggested that different factors affected the species richness of the two groups of spe-
417 cies ~~at plot scale~~. ~~Specifically, native species richness was only very weakly related to the variables we measured~~
418 ~~at this scale, and our best model only explained a very small proportion of variability ($R^2 = 0.02$). Nevertheless, we~~
419 ~~found evidence that native richness was~~ positively associated with *landscape diversity* around the plot (250 m ra-
420 dius) and negatively associated with the *climatic* gradient ranging from Temperate to Mediterranean ~~micromacro-~~
421 ~~meso-~~climatic conditions. The first result is in agreement with previous studies suggesting that greater *landscape*
422 *diversity* may be related to a higher number of available niches, potentially hosting species with different ecologi-
423 cal requirements (Deutschewitz et al. 2003; Kumar et al. 2006). In our study area, higher values of ~~fine~~small-scale
424 *landscape diversity* might specifically indicate fragmentation and the survival of more natural ecosystems within
425 an otherwise homogeneous agricultural landscape. The negative relationship between native species and the tem-
426 perate-mediterranean gradient indicates that more native species were found in the cooler and wetter, rather than in
427 the drier, ~~micromeso-~~climatic conditions. This is likely to be connected to the regional context, in which higher
428 temperatures at plot scale might represent a significant ecological constrain in summer, while lower temperatures
429 are not likely to be a limiting factor in winter.

430 In contrast, ~~more variability could be explained for non-native species richness.~~ Non-native species were nega-
431 tively related with ~~elevation~~, road distance, and slope ~~activity~~inclination and positively related to the temperate-
432 ~~mediterranean climatic gradient~~. ~~Elevation was important in this model, in agreement with previous studies that~~
433 ~~showed a negative relationship between non-native plant species richness and elevation in various systems (*e.g.*,~~
434 ~~Pyšek et al. 2002b; Stevens 1992; Pausas 1994; Rey-Benayas 1995; Marini et al. 2009; Siniscalco et al. 2011;~~
435 ~~Barni et al. 2012; Bacaro et al. 2015). However, upward movements of non-native species are increasingly being~~
436 ~~detected in montane ecosystems (Kalwij et al. 2015), probably due to the short residence time of the species or to~~
437 ~~niche unfilling, and are also predicted to accelerate under future global change scenarios (Carboni et al. 2018).~~
438 This suggests that, even though higher elevations are ~~currently less invaded in this protected area network, caution~~
439 ~~is needed and potential future invasion risks should not be discarded.~~

440 ~~Further~~, ~~a~~s expected, non-native species richness was higher in plots in close proximity to roadways. This is in
441 accordance with ~~many studies reporting that the well-documented notion that~~ roads, and roadside habitats, are a
442 major source from which non-native species ~~invade~~colonize natural areas (Parendes and Jones 2000; Pauchard
443 and Alaback 2004; Bacaro et al. 2015; Ullmann and Heindl 1989; Ullmann et al. 1995; Arévalo et al. 2010). ~~Roads~~
444 ~~are one of the main anthropogenic features that affect the distribution of non-native species and represent a path-~~
445 ~~way for their spread (*e.g.*, Pauchard and Alaback 2004).~~ Roads may facilitate the dispersal of non-native species
446 inducing habitat fragmentation and altering (micro-) environmental conditions, facilitating the human-mediated

Commented [A20]:
Rev2: No acknowledgment
low R2 values for the plot le
is not an informative model.

Commented [A21]:
Rev4: I think this would fit b
next paragraph

Commented [A22]:
Rev2: Line 286: this is not a

Commented [A23]:
Rev4: You shouldn't use the
you actually looked at invas
species.

447 dispersal of propagules (via air movement associated with vehicle traffic, and via the seeds attached to the vehicles
448 themselves) and facilitating the colonisation by non-native species by suppressing the growth or removing stands
449 of native species (Trombulak and Frissel 2000; Bacaro et al. 2015). Therefore, limitation to the construction of
450 new roads within or close to protected areas is important for preserving local biodiversity both directly, but also
451 indirectly via the reduction of non-native introductions.

452 ~~Finally~~In terms of environmental factors, non-native species richness at plot scale was positively associated with
453 warmer and drier Mediterranean meso-climatic conditions and ~~also~~ negatively correlated with slope inclination.
454 This can partly be a consequence of the well-known conservative role of steep slopes, because of soil- and micro-
455 climatic- limiting factors: for instance, Bennie et al. (2006) found that in Britain, grassland swards on steep slopes
456 were more resistant to invasion by competitive grass species than those on flatter sites, due to phosphorus limita-
457 tion in shallow minerogenic soils, and to increased drought events. Filibeck et al. (2016) showed that, in grass-
458 lands in a protected area at the periphery of Rome, the removal or leveling ~~by the management~~ of some steep de-
459 bris heaps led to an increase in non-native and cosmopolite species. However, our dataset is not restricted to grass-
460 lands and involves a huge variety of land-cover types, so it is also possible that the observed effect is mediated by
461 disturbance levels and land-use: in the study region, a high slope steepness usually allows only land-use types that
462 feature an inherently low human disturbance (*e.g.*, woods or extensive rangelands), while flat morphologies are
463 usually exploited with intensive crops or urban land-use, typically favouring non-native taxa invasions.

464 Species patterns at the protected area scale

465 ~~Based on the Mantel test,~~We found that the turnover in non-native species composition was correlated with native
466 species composition also at the scale of protected area. ~~In this~~At this spatial grain, ~~ease~~ however, we also detected
467 a strong congruence in the richness of the two groups of species ~~across protected areas~~. These findings, ~~first of all,~~
468 highlight ~~potential conservation hotspots, since that~~ the protected areas that are most biodiverse are also the ones
469 that are most ~~at risk of easily colonized by alien species (and thus, potentially, most at risk of invasion)~~. ~~Second-~~
470 ~~ly~~More generally, these findings support the ~~hypothesis that has been sometimes termed~~ “biotic acceptance hy-
471 pothesis” or “the rich get richer” hypothesis, according to which sites with high native species richness are the
472 most readily invaded by non-native species (*e.g.*, Stohlgren et al. 1999, 2006; Fridley et al. 2007; Pyšek and Rich-
473 ardson 2006; Sandel and Corbin 2010, Bartomeus et al. 2012). This would ~~result be explained~~ because habitats
474 that are generally ‘good’ for native species would also be ‘good’ for non-native ones (*e.g.*, McKinney 2002; Souza
475 et al. 2011). This hypothesis can also be linked to Grime’s theory (Grime 1973), according to which all species
476 (meaning both native and non-native species) respond, to some degree, in a similar way to stress, competition and
477 disturbance (Tomasetto et al. 2013).

478 We therefore tested whether the similar compositional and richness patterns emerged as a consequence of similar
479 responses to environmental factors and human disturbances. Indeed, we found that ~~three~~two main variables were
480 retained in both the models for native and non-native species richness at the PA scale: *area* of the protected area,
481 ~~mean climate~~ and *climate range* (*i.e.* spatial heterogeneity of climate within the PA). ~~All~~Both these variables
482 showed positive effects, even if with different weights for native and non-native species richness. Thus, our re-
483 search shows a positive native-non-native relationship related to the same response of native and non-native spe-
484 cies to the available area in the PA and to the main gradients (~~area and mean climate~~ at large scale. The positive
485 effect of *area* on both native and non-native species richness is likely dependent on the higher availability of nich-

es in larger protected areas for both groups of species. The congruence of native and non-native species richness is thus largely the net result of concordant well-known species-area relationships, that show similar patterns for both groups of species. Indeed similar-a congruence in species-area relationships for native and non-native species had already been reported in this network of protected areas (Chiarucci et al. 2012), as well as in other reserve networks (Pyšek et al. 2002a) or insular systems (e.g., the islands of the Tuscan archipelago, Chiarucci et al. 2017). In addition, the positive effect on richness of the climatic variables at the scale of the protected area (mean climate-and-climate range) indicates, on the one hand, that warmer and drier average climates support greater species richness for both groups of native and non-native species (Barni et al. 2012) and, on the other hand, that more climatically heterogeneous areas enhance regional native richness but also promote non-native species establishment. Indeed, environments with favorable (mean) abiotic conditions (e.g., higher soil fertility or optimal climate conditions; Levine and D'Antonio 1999; Stohlgren et al. 1999) and greater spatial heterogeneity (e.g., higher habitat diversity or spatial variability in resources or conditions; Davies et al. 2005) have been shown to support higher numbers of both native and non-native species at broad spatial scales in numerous other systems. Thus, overall we conclude that in our reserve network "rich protected areas get richer" because of more available area (as expected) but also because of more favorable conditions in terms of available area-and-niches and of climates.

In addition to the drivers that were common to both natives and non-natives, to the drivers that were in-common with native species, the model for native species at PA scale also included a significant positive effect of mean climate, which suggests that reserves with warmer and drier average climates tend to support greater species richness confirms what expressed above in particular for native species. Instead, non-native species also included a non-significant effect of road density, which was similar to the effect of roads found at the plot scale, and the percentage of agricultural area within the protected area. A positive effect of road density is a typical finding in studies explaining the number of non-native species at broad scales, because roads serve as introduction pathways as outlined above. Here we found instead a unimodal relationship with non-native species richness, suggesting that non-native species were most abundant at intermediate levels of anthropogenic disturbance (while their spread was potentially hampered by very high levels of fragmentation at higher road densities). Perhaps, the high degree of naturalness of the protected areas studied here, and the small number of intensively used roads could have driven this unimodal relationship. In contrast, the percentage of agricultural land-wetland was negatively related to non-native richness, which counters what has been observed in many other agricultural landscapes. Chytrý et al. (2009), for example, showed that the highest levels of non-native invasion among the CORINE land-cover classes in Europe are predicted not only in urban and industrial areas, but also on arable lands. In our case the negative correlation between non-native species and agricultural land-wetlands use might be due to the specific local agricultural practices. They are represented by the lakes of Chiusi (CHU) and Montepulciano (MPU), where the anthropic disturbance is extremely high. In fact these lakes are surrounded by agricultural fields and vineyards and their water is used for irrigation.... Relatedly, the model for non-native species, included a significant effect of mean elevation, that was negatively related to non-native species richness indicating that fewer non-native species were found at higher altitudes. Mean Elevation and in general elevation, was important in this model. This finding is in agreement with previous studies that showed a negative relationship between non-native plant species richness and elevation in various systems (e.g., Pyšek et al. 2002b; Stevens 1992; Pausas 1994; Rey-Benayas 1995; Marini et al. 2009; Siniscalco et al. 2011; Barni et al. 2012; Bacaro et al. 2015). Nevertheless it is important to caution that upward movements of non native species are increasingly being detected in many montane ecosystems (Kalwij et al.

526 2015), probably due to the short residence time of the species or to niche-unfilling, and are also predicted to
527 accelerate under the future global change scenarios (Carboni et al. 2018). Hence, even if our results show that
528 currently higher elevations are of lesser concern for non-native plants, it suggests that caution is needed and po-
529 tential future invasion risks should not be discarded.

Commented [A24]:
Rev2: Line 286: this is not a

530 , that are still largely done according to traditional techniques, on small-sized fields interspersed with many hedges,
531 remnants of natural forests (Amici et al. 2015) and grasslands. The traditional agriculture present in many parts of
532 the province of Siena might thus promote resistance to non-native species invasion, rather than facilitating inva-
533 sions. Alternatively, this result might reflect the lack of negative correlation of largely agricultural protected areas
534 with their percentage of urban land cover, w PA, that result which might indicate less potential introduction
535 sources of non-native species.

Commented [A25]:
Rev4: The wording here is

Commented [A26]:
changed

536 Conclusions and cross-scale comparisons

537 In conclusion, ~~o~~Our overall ~~Overall~~ our results suggest that native and non-native species richness are only weakly-
538 related at a fine scale, but share similar patterns at the PA scale. This supports the idea that the relationship between
539 the number of native and non-native species is altered when changing the scale of analysis. Indeed, it is widely
540 acknowledged that the strength and form of this relationship (in many studies known as NERR – native exotic
541 richness relationship – e.g., Souza et al. 2011; Symonds and Pither 2012) is scale-dependent. Here, we found evi-
542 dence that the lack of congruence at ~~fine~~small plot scale was likely driven by differential responses to fine scale
543 environmental factors (climate and landscape diversity) and human disturbances (road distance), while at the larg-
544 er PA scale native and non-native species largely responded in a similar way to ~~variability of~~ available area and
545 to climate variability (climate range). Interestingly, this pattern also resulted from the fact that the effect of climat-
546 ic variables (climate range and mean climate) on native species changed when moving from the coarse to the fine
547 scale. While at coarse scales the species richness of native species (and non-native species) was generally more
548 abundant higher in warmer Mediterranean climates, at the fine plot scale the native species richness were was even
549 mildly more associated with cooler micromeso-climatic conditions. Overall richness of non-native species in the
550 protected area network increased under moderate anthropogenic disturbances at at fine both scale, coupled with
551 high levels of habitat and structural-climatic heterogeneity at large scale. Thus, our data suggest that within the
552 study region the introduction and establishment of non-native species would be more likely in warmer and dryer
553 areas, with high native species richness at large spatial scale but intermediate levels of anthropogenic disturbances;
554 but with limited fine-scale heterogeneity and mild slope inclinations and elevation. We also found potential con-
555 servation conflicts and hotspots, especially because the very biodiverse protected areas are potentially most at fea-
556 ture a higher risk of invasion, due to the presence of invasive higher establishment success of non-native species in
557 general. These areas should thus be prioritized for invasion monitoring. Finally, the results also highlight that both
558 1) the measure of the proportion of non-native species and 2) the relationship with potential predictors should be
559 studied at the appropriate spatial scale in order to be comparable among different regions and informative for con-
560 servation purposes.

Commented [MC27]:
toglierei perchè climate range

Commented [G28]:
però a plot scale il modello
fatto questo effetto del clima

Commented [A29]:
Rev4: I think you mean native
higher, which is not the same
more abundant.

561 Although biological invasions represent a potential threat for the biodiversity in Europe, few studies have attempt-
562 ed to propose model-based methodologies for preventing the expansion of invasive species in Natura 2000 sites
563 (Dimitrakopoulos et al. 2017; Bazzichetto et al. 2018). ~~job work we do not consider invasive species, but rather~~
564 ~~focus on non-native species but only the alien ones, we think that these results could increase the knowledge about~~

565 the presence of non-native species and help to prevent spread of biological invasions. In fact, biological inva-
566 sions represent a potential threat for the biodiversity in Europe and few studies have attempted to propose model-
567 based methodologies for preventing the expansion of invasive species in Natura 2000 sites (Dimitrakopoulos et al.
568 2017; Bazzichetto et al. 2018).

569 **Acknowledgment**

570 We would like to thank Prof. Alessandro Chiarucci for the opportunity to conduct this research on the data of the
571 project MOBISIC he coordinated when he was at the University of Siena. We would also thank Francesco Geri for
572 his ~~assistence~~ assistance with ~~geografical~~ geographical informations. G.F. and A.S. carried out this research partial-
573 ly in the frame of the MIUR (Italian Ministry for Education, University and Research) initiative “Department of
574 excellence” (Law 232/2016).

575

576 **Author Contributions:**

577 All the authors contributed to the interpretation of results and writing.

578 **REFERENCES**

- 579 Amici V, Landi S, Frascaroli F, Rocchini D, Santi E, Chiarucci A. 2015. Anthropogenic drivers of plant diversity:
580 perspective on land use change in a dynamic cultural landscape. *Biodiversity and Conservation* 24: 3185-3199.
- 581 Arévalo JR, Delgado JD, Otto R, Naranjo A, Salas M, Fernández-Palacio JM. 2005. Distribution of alien vs. na-
582 tive plant species in roadside communities along an altitudinal gradient in Tenerife and Gran Canaria (Canary Is-
583 lands). *Perspectives in Plant Ecology, Evolution and Systematics* 7: 185–202.
- 584 Ashton PJ, Mitchell DS. 1989. Aquatic plants: patterns and modes of invasion, attributes of invading species and
585 assessment of control programmes. In: Drake JA, Mooney HA, di Castri F, Groves RH, Kruger FJ, Rejmánek M,
586 Williamson M (eds) *Biological invasions: a global perspective*. Wiley, Chichester, pp 111-154.
- 587 Bacaro G, Maccherini S, Chiarucci A, Jentsch A, Rocchini D, Torri D, Gioria M, Otto R, Escudero CG,
588 Fernandez-Lugo S, Fernandez-Palacios JM, Arévalo JR. 2015. Distributional patterns of endemic, native and alien
589 species along a roadside elevational gradient in Tenerife, Canary Islands. *Community Ecology* 16: 223-234. doi:
590 10.1556/168.2015.16.2.10
- 591 Barazzuoli P, Guasparri G, Salleolini M. 1993. Il clima. In: Giusti F (ed) *La storia naturale della Toscana*
592 *meridionale*. Monte dei Paschi di Siena, Amilcare Pizzi Editore, pp 141-171.
- 593 Barbosa AM, Brown JA, Real R. 2014. modEvA – an R package for model evaluation and analysis. R package,
594 version 0.1.
- 595 Barni E, Bacaro G, Rocchini D, Falzoi S, Spanna F, Siniscalco C. 2012. Establishing climatic constraints shaping
596 the distribution of alien plant species along the elevation gradient in the Alps. *Plant Ecology* 213: 757-767.
- 597 Bartomeus I, Sol D, Pino J, Vicente P, Font X. 2012. Deconstructing the native-exotic richness relationship in
598 plants. *Global Ecology and Biogeography* 21:524–533.
- 599 [Barton K. 2019. MuMIn: Multi-Model Inference. R package version 1.43.6. https://CRAN.R-](https://CRAN.R-project.org/package=MumIn)
600 [project.org/package=MumIn](https://CRAN.R-project.org/package=MumIn)
- 601 [Bates D, Maechler M, Bolker B, Walker S. 2015. Fitting Linear Mixed-Effects Models Using lme4. Journal of](https://doi.org/10.18637/jss.v067.i01)
602 [Statistical Software, 67\(1\): 1-48. doi:10.18637/jss.v067.i01.](https://doi.org/10.18637/jss.v067.i01)
- 603 Bazzichetto M, Malavasi M, Barták V, Acosta A, Rocchini D, Carranza M. 2018. Plant invasion risk: A quest for
604 invasive species distribution modelling in managing protected areas. *Ecological Indicators* 95: 311-319.
605 10.1016/j.ecolind.2018.07.046
- 606 Bennie J, Hill MO, Baxter R, Huntley B. 2006. Influence of slope and aspect on long-term vegetation change in
607 British chalk grasslands. *Journal of Ecology* 94: 355–368.
- 608 Brown RL, Peet RK. 2003. Diversity and invasibility of southern Appalachian plant communities. *Ecology* 84:
609 32–39.

610 [Bjarnason A., Katsanevakis S., Galanidis A., Vogiatzakis I.N., Moustakas A. 2017. Evaluating Hypotheses](#)
611 [of Plant Species Invasions on Mediterranean Islands: Inverse Patterns between Alien and Endemic species.](#)
612 [Frontiers Ecol. Evolut., 5 \(91\). Mediterranean.](#)

613 [Bjornstad ON. 2019. ncf: Spatial Covariance Functions. R package version 1.2-8. \[https://CRAN.R-\]\(https://CRAN.R-project.org/package=ncf\)](#)
614 [project.org/package=ncf](#)

615 [Burnham KP, Anderson DR. 2002. Model selection and multimodel inference: a practical information-](#)
616 [theoretic approach. Springer-Verlag, New York, New York, USA.](#)

617 Calcagno V. 2013. glmulti: Model selection and multimodel inference made easy. R package version 1.0.7.

618 Callaway RM. 2007. Positive interactions and interdependence in plant communities. New York, Springer-Verlag.

619 Carboni M, Thuiller W, Izzi F, Acosta A. 2010. Disentangling the relative effects of environmental versus human
620 factors on the abundance of native and alien plant species in Mediterranean sandy shores. Diversity and Distribu-
621 tions 16: 537–546. doi: 10.1111/j.1472-4642.2010.00677.x

622 Carboni M, Guéguen M, Barros C, Georges D, Boulangeat I, Douzet R, Dullinger S, Klonner G, van Kleunen M,
623 Essl F, Bossdorf O, Haeuser E, Talluto MV, Moser D, Block S, Conti L, Dullinger I, Münkemüller T, Thuiller W.
624 2018. Simulating plant invasion dynamics in mountain ecosystems under global change scenarios. Global Change
625 Biology 24: 289-302.

626 Castrignanò A, Costantini EAC, Barbetti R, Lopez N. 2006. Assessing and describing multi-scale spatial variabil-
627 ity using factorial kriging. Geophysical Research Abstracts 8:10245

628 Celesti-Grapow L, Alessandrini A, Arrigoni PV, Banfi E, Bernardo L, Bovio M, Brundu G, Cagiotti MR, Camarda
629 I, Carli E, Conti F, Fascetti S, Galasso G, Gubellini L, La Valva V, Lucchese F, Marchiori S, Mazzola P,
630 Peccenini S, Poldini L, Pretto F, Prosser F, Siniscalco C, Villani MC, Viegi L, Wilhalm T, Blasi C. 2009. Invento-
631 ry of the non-native flora of Italy . Plant Biosystems 143: 386–430.

632 Chiarucci A. 2007. To sample or not to sample. That is the question. Folia Geobotanica 42: 209-216.

633 Chiarucci A, Bacaro G, Rocchini D. 2008. Quantifying plant species diversity in a Natura 2000 network: Old ideas
634 and new proposals. Biological Conservation 141: 2608-2618.

635 Chiarucci A, Bacaro G, Filibeck G, Landi S, Maccherini S, Scoppola A. 2012. Scale dependence of plant species
636 richness in a network of protected areas. Biodiversity and Conservation 21: 503-516.

637 Chiarucci A, Fattorini S, Foggi B, Landi S, Lazzaro L, Podani J, Simberloff D. 2017. Plant recording across two
638 centuries reveals dramatic changes in species diversity of a Mediterranean archipelago. Scientific Reports 7: 5415.
639 <https://doi.org/10.1038/s41598-017-05114-5>

640 Chytrý M, Jarošík V, Pyšek P, Hájek O, Knollová I, Tichý L, Danihelka J. 2008. Separating habitat invasibility by
641 alien plants from the actual level of invasion. Ecology 89:1541–53.

- 642 Chytrý M, Pyšek P, Wild J, Pino J, Maskell LC, Vilà M. 2009. European map of alien plant invasions based on the
643 quantitative assessment across habitats. *Diversity and Distributions* 15: 98–107.
- 644 Conti F, Abbate G, Alessandrini A, Blasi C. 2005. An Annotated Checklist of the Italian Vascular Flora. Palombi,
645 Roma.
- 646 Cornell HV, Karlson RH. 1997. Local and regional processes as controls of species richness. In: Tilman D, Ka-
647 reiva P (eds) *Spatial ecology: the role of space in population dynamics and interspecific interactions*. Princeton
648 University Press, Princeton, pp 250–268.
- 649 Crutzen PJ. 2006. The “anthropocene”. In *Earth system science in the anthropocene*. Springer, Berlin, Heidelberg,
650 pp 13–18.
- 651 Davies KF, Chesson P, Harrison S, Inouye BD, Melbourne BA, Rice KJ. 2005. Spatial heterogeneity explains the
652 scale dependence of the native-exotic diversity relationship. *Ecology* 86: 1602–1610.
- 653 Deutschewitz K, Lausch A, Kühn I, Klotz S. 2003. Native and alien plant species richness in relation to spatial
654 heterogeneity on a regional scale in Germany. *Global Ecology and Biogeography* 12: 299–311.
- 655 Dimitrakopoulos PG, Koukoulas S, Galanidis A, Delipetrou P, Gounaridis D, Touloumi K, Arianoutsou M. 2017.
656 Factors shaping alien plant species richness spatial patterns across Natura 2000 Special Areas of Conservation of
657 Greece. *Science of the Total Environment* 601: 461–468.
- 658 [Dormann CF, Elith J, Bacher S, Buchmann C, Carl G, Carré G, García Márquez JR, Gruber B, Lafourcade B,](#)
659 [Leitão PJ, Münkemüller T, McClean C, Osborne PE, Reineking B, Schröder B, Skidmore Ak, Zurell D, Lauten-](#)
660 [bach S. 2013. Collinearity: A review of methods to deal with it and a simulation study evaluating their perfor-](#)
661 [mance. *Ecography* 36: 27–46.](#)
- 662 Faliński JB. 1998. Invasive alien plants and vegetation dynamics. In: Starfinger U, Edwards KR, Kowarik I,
663 Williamson M (eds) *Plant invasions: ecological mechanisms and human responses*. Backhuys Publishers, Leiden,
664 pp 3–21.
- 665 Filibeck G, Petrella P, Cornelini P. 2016. All ecosystems look messy, but some more so than others: a case-study
666 on the management and acceptance of Mediterranean urban grasslands. *Urban Forestry and Urban Greening*
667 15:32–39.
- 668 Foxcroft LC, Pyšek P, Richardson DM, Genovesi P, MacFadyen S. 2017. Plant invasion science in protected areas:
669 progress and priorities. *Biological Invasions* 19:1353–1378.
- 670 Fridley JD, Stachowicz JJ, Naeem S, Sax DF, Seabloom EW, Smith MD, Stohlgren TJ, Tilman D, Von Holle B.
671 2007. The invasion paradox: reconciling pattern and process in species invasions. *Ecology* 88:3–17.
672 [http://dx.doi.org/10.1890/0012-9658\(2007\)88\[3:TIPRPA\]2.0.CO;2](http://dx.doi.org/10.1890/0012-9658(2007)88[3:TIPRPA]2.0.CO;2)
- 673 Gaston KJ. 2005. Biodiversity and extinction: species and people. *Progress in Physical Geography* 29: 239–247.

674 Gaston KJ, Jackson SF, Cantu-Salazar L, Cruz-Pinon G. 2008. The ecological performance of protected areas.
675 Annual Review of Ecology, Evolution, and Systematics 39: 93-113.
676 doi:10.1146/annurev.ecolsys.39.110707.173529

677 [Geri F., Amici V., Rocchini D. 2010. Human activity impact on the heterogeneity of Mediterranean landscape.](#)
678 [Applied Geography 30: 370-379. doi:10.1016/j.apgeog.2009.10.006](#)

679 Grime JP. 1973. Competitive exclusion in herbaceous vegetation. Nature 242: 344–347.

680 Grunanger P. 2001. Orchidacee d'Italia. Quaderni di Botanica Ambientale e Applicata 11: 3–80.

681 Guisan A, Zimmermann NE. 2000. Predictive habitat distribution models in ecology. Ecological Modelling
682 135:147-186.

683 Hannah L, Midgley G, Anelman S, Araujo M, Hughes G, Martinez-Meyer E, Pearson R, Williams P. 2007. Pro
684 tected area needs in a changing climate. Frontiers in Ecology and the Environment 5:131–138. doi:10.1890/1540

685 Huston MA. 1999. Local processes and regional patterns: appropriate scales for understanding variation in the
686 diversity of plants and animals. Oikos 86: 393–401.

687 Jaeger B. 2017. r2glmm: Computes R Squared for Mixed (Multilevel) Models. R package version 0.1.2

688 Kalwij JM, Robertson MP, van Rensburg BJ. 2015. Annual monitoring reveals rapid upward movement of exotic
689 plants in a montane ecosystem. Biological Invasions 17: 3517-3529.

690 Kumar S, Stohlgren TJ, Chong GW. 2006. Spatial heterogeneity influences native and nonnative plant species
691 richness. Ecology 87: 3186-3199.

692 Levine J, D'Antonio C. 1999. Elton revisited: a review of evidence linking diversity and invasibility. Oikos 87:
693 15–26.

694 Levine JM. 2000. Species diversity and biological invasions: relating local process to community pattern. Science
695 288: 852–854.

696 Lonsdale WM. 1999. Global patterns of plant invasions and the concept of invasibility. Ecology 80: 1522–1536.

697 Lucy FE, Roy H, Simpson A, Carlton JT, Hanson JM, Magellan K, Campbell ML, Costello MJ, Pagad S, Hewitt
698 CL, McDonald J, Cassey P, Thomaz SM, Katsanevakis S, Zenetos A, Tricarico E, Boggero E, Groom QJ, Adri-
699 aens T, Vanderhoeven S, Torchin M, Hufbauer R, Fuller P, Carman MR, Conn DB, Vitule JRS, Canning-Clode J,
700 Galil BS, Ojaveer H, Bailey SA, Theriault TW, Claudi R, Gazda A, Dick JTA, Caffrey J, Witt A, Kenis M,
701 Lehtiniemi M, Helmsaari H, Panov VE. 2016. INVASIVESNET towards an international association for open
702 knowledge on invasive alien species. Management of Biological Invasions 7:131–139.

703 Maestre FT. 2004. On the importance of patch attributes, environmental factors and past human impacts as deter-
704 minants of perennial plant species richness and diversity in Mediterranean semiarid steppes. Diversity and Distri-
705 butions 10: 21–29.

- 706 Marini L, Gaston KJ, Prosser F, Hulme PE. 2009. Contrasting response of native and alien plant species richness
707 to environmental energy and human impact along alpine elevation gradients. *Global Ecology and Biogeography*
708 18:652–661.
- 709 McCune B, Keon D. 2002. Equations for potential annual direct incident radiation and heat load. *Journal of Vege-*
710 *tation Science* 13: 603–606.
- 711 McKinney ML. 2002. Do human activities raise species richness? Contrasting patterns in United States plants and
712 fishes. *Global Ecology and Biogeography* 11: 343–348.
- 713 Meiners SJ, Pickett STA. 2013. Plant invasion in protected landscape: exception or expectation? In: Foxcroft LC,
714 Richardson DM, Pyšek P, Genovesi P (eds) *Plant invasions in Protected Areas*. Springer, New York
- 715 Moser D, Dullinger S, Englisch T, Niklfeld H, Plutzer C, Sauberer N, Zechmeister HG, Grabherr G. 2005. Envi-
716 ronmental determinants of vascular plant species richness in the Austrian Alps. *Journal of Biogeography*
717 32: 1117–1127. doi: 10.1111/j.1365-2699.2005.01265.x
- 718 [Moustakas A., Voutsela A., Katsanevakis S. 2018. Sampling alien species inside and outside protected areas: Does
719 it matter? *Science of the Total Environment* 625: 194-198. doi: 10.1016/j.scitotenv.2017.12.198](#)
- 720 Paradis E, Schliep K. 2018. ape 5.0: an environment for modern phylogenetics and evolutionary analyses in R.
- 721 Parendes LA, Jones JA. 2000. Role of light availability and dispersal in exotic plant invasion along roads and
722 streams in the H.J. Andres Experimental Forest, Oregon. *Conservation Biology* 14: 64–75.
- 723 Pauchard A, Alaback PB. 2004. Influence of elevation, land use, and landscape context on patterns of alien plant
724 invasions along roadsides in protected areas of south central Chile. *Conservation Biology* 18: 238–248.
- 725 Pausas JG. 1994. Species richness patterns in the understorey of Pyrenean *Pinus sylvestris* forest. *Journal of*
726 *Vegetation Science* 5: 517– 524.
- 727 Pignatti S. 1982. *Flora d'Italia* (3 vols). Edagricole, Bologna.
- 728 Planty-Tabacchi AM, Tabacchi E, Naiman RJ, Deferrari C, Decamps H. 1996. Invasibility of species-rich com-
729 munities in riparian zones. *Conservation Biology* 10: 598– 607.
- 730 Pressey RL, Visconti P, Ferraro PJ. 2015. Making parks make a difference: poor alignment of policy, planning and
731 management with protected-area impact, and ways forward. *Philosophical Transactions of the Royal Society B*
732 370: 20140280. doi:10.1098/rstb.2014.0280
- 733 Pyšek P, Prach K, Mandák B. 1998. Invasions of alien plants into habitats of Central European Landscape: an his-
734 torical pattern. In Starfinger U, Edwards KR, Kowarik I, Williamson M (eds) *Plant invasions: ecological mecha-*
735 *nisms and human responses*. Backhuys Publishers, Leiden, pp 23-32.
- 736 Pyšek P, Jarosík V, Kucera T. 2002a. Patterns of invasion in temperate nature reserves. *Biological Conservation*
737 104: 13–24.
- 738 Pyšek P, Sadlo J, Mandak B. 2002b. Catalogue of alien s plants of the Czech Republic. *Preslia* 74:97–186.

- 739 Pyšek P, Jarošík V, Kučera T. 2003. Inclusion of native and alien species in temperate nature reserves: an historical study from Central Europe. *Conservation Biology* 17: 1414-1424.
- 740
- 741 Pyšek P, Richardson DM. 2006. The biogeography of naturalization in alien plants. *Journal of Biogeography* 33: 2040-2050.
- 742
- 743 Quantum GIS Development Team 2016, version 2.14. QGIS Geographic Information System. Open Source Geospatial Foundation Project. <http://qgis.osgeo.org>
- 744
- 745 R Core Team 2019. R: A language and environment for statistical computing. R Foundation for Statistical Computing, Vienna, Austria. ISBN 3-900051-07-0, URL <http://www.R-project.org/>
- 746
- 747 Rey-Benayas JM. 1995. Patterns of diversity in the strata of forest in British Columbia. *Journal of Vegetation Science* 6: 95-98.
- 748
- 749 Richardson DM, Pyšek P. 2006. Plant Invasions: Merging the Concepts of Species Invasiveness and Community Invasibility. *Progress in Physical Geography*, 30, 409-431. <http://dx.doi.org/10.1191/0309133306pp490pr>
- 750
- 751 Sala OE, Chapin IFS, Armesto JJ, Berlow E, Bloomfield J, Dirzo R, Huber Sanwald E, Huenneke LF, Jackson RB, Kinzig A, Leemans R, Lodge DH, Mooney HA, Oesterheld M, Leroy Poff N, Sykes MT, Walker BH, Walker M, Wall DH. 2000. Global biodiversity scenarios for the year 2100. *Science* 287: 770-1774.
- 752
- 753
- 754 Sandel B, Corbin JD. 2010. Scale, disturbance and productivity control the native-exotic richness relationship. *Oikos* 119: 1281-1290
- 755
- 756 Sax DF. 2002. Native and naturalized plant diversity are positively correlated in scrub communities of California and Chile. *Diversity and Distributions* 8: 193-210.
- 757
- 758 Shea K, Chesson P. 2002. Community ecology theory as a framework for biological invasions. *Trends in Ecology & Evolution* 17: 170-176.
- 759
- 760 Shochat E, Warren PS, Feath SH, McIntyre NE, Hope D. 2006. From patterns to emerging processes in mechanistic urban ecology. *Trends in Ecology & Evolution* 21: 186-191.
- 761
- 762 Simberloff D, Martin JL, Genovesi P, Maris V, Wardle DA, Aronson J, Courchamp F, Galil B, García-Berthou E, Pascal M, Pyšek P, Sousa R, Tabacchi E, Vilà M. 2013. Impacts of biological invasions: what's what and the way forward. *Trends in Ecology & Evolution* 28: 58-66.
- 763
- 764
- 765 Siniscalco C, Barni E, Bacaro G. 2011. Non-native species distribution along the elevation gradient in the western Italian Alps. *Plant Biosystems* 145: 150-158.
- 766
- 767 Souza L, Bunn WA, Simberloff D, Lawton RM, Sanders NJ. 2011. Biotic and abiotic influences on native and exotic richness relationship across spatial scales: favourable environments for native species are highly invisable. *Functional Ecology* 25: 1106-1112.
- 768
- 769
- 770 Spear D, Foxcroft LC, Bezuidenhout H, McGeoch MA. 2013. Human population density explains alien species richness in protected areas. *Biological Conservation* 159: 137-147. doi:10.1016/j.biocon.2012.11.022
- 771

- 772 Stadler J, Trefflich A, Klotz S, Brandl R. 2000. Exotic plant species invade diversity hot spots: the alien flora of
773 northwestern Kenya. *Ecography* 23: 169–176.
- 774 Stevens GC. 1992. The elevational gradient in altitudinal range: an extension of Rapoport's latitudinal rule to alti-
775 tude. *American Naturalist* 140: 893–911.
- 776 Stohlgren TJ, Binkley D, Chong GW, Kalkhan MA, Schell LD, Bull KA, Otsuki Y, Newman G, Bashkin M, Son
777 Y. 1999. Exotic plant species invade hot spots of native plant diversity. *Ecological Monographs* 69: 25–46.
- 778 Stohlgren TJ, Jarnevich C, Chong GW, Evangelista PH. 2006. Scale and plant invasions: a theory of biotic ac-
779 ceptance. *Preslia* 78: 405–426.
- 780 Sukopp H. 1998. On the study of anthropogenic plant migrations in central Europe. In: Starfinger U, Edwards KR,
781 Kowarik I, Williamson M (eds) *Plant invasions: ecological mechanisms and human responses*. Backhuys Publish-
782 ers, Leiden, pp 43–56.
- 783 Symonds JE, Pither J. 2012. Multi-scale analyses of exotic and native plant species diversity within Canada's en-
784 dangered antelope-brush ecosystem. *Plant Ecology* 213:1263–1275.
- 785 Taylor AH, Allen JI, Clark PA. 2002. Extraction of a weak climatic signal by an ecosystem. *Nature* 416: 629–632.
- 786 Thuiller W, Gallien L, Boulangeat I, de Bello F, Munkemuller T, Roquet-Ruiz C, Lavergne S. 2010. Resolving
787 Darwin's naturalization conundrum: a quest for evidence. *Diversity and Distributions* 16: 461–475.
- 788 Tilman D. 1997. Community invasibility, recruitment limitation, and grassland biodiversity. *Ecology* 78:81–92.
- 789 Tomasetto F, Duncan RP, Hulme PE. 2013. Environmental gradients shift the direction of the relationship between
790 native and alien plant species richness. *Diversity and Distributions* 19:49–59.
- 791 Tordoni E, Napolitano R, Nimis P, Castello M, Altobelli A, Da Re D, Zago S, Chines A, Martellos S, Maccherini
792 S, Bacaro G. 2017. Diversity patterns of alien and native plant species in Trieste port area: exploring the role of
793 urban habitats in biodiversity conservation. *Urban Ecosystems* 20: 1151–1160.
- 794 Trombulak SC, Frissell CA. 2000. Review of ecological effects of roads on terrestrial and aquatic communities.
795 *Conservation Biology* 14: 18–30.
- 796 Tutin TG, Burges NA, Chater AO, Edmondson JR, Heywood VH, Moore DM, Valentine DH, Walters SM, Webb
797 DA. (eds) 1993. *Flora Europaea*, Vol. 1, 2nd edn. Cambridge University Press, Cambridge. Thomas CD, Palmer G
798 2015. Non-native plants add to the British flora without negative consequences for native diversity. *PNAS* 112:
799 4387–4392.
- 800 Tutin TG, Heywood VH, Burges NA, Moore DM, Valentine DH, Walters SM, Webb DA. 1964–1980. *Flora Eu-*
801 *ropaea* (5 vols.). Cambridge University Press, Cambridge .
- 802 Ullmann I, Heindl B. 1989. Geographical and Ecological Differentiation of roadside vegetation in temperate Eu-
803 rope. *Botanica Acta* 102: 261–269.

- 804 Ullmann I, Bannister P, Wilson JB. 1995. The vegetation of roadside verges with respect to environmental gradi-
805 ents in southern New Zealand. *Journal of Vegetation Science* 6: 131–142.
- 806 Venables WN, Ripley BD. 2002. *Modern Applied Statistics with S*. Fourth Edition. Springer, New York. ISBN 0 -
807 387-95457-0
- 808 Weber HE. 1995. *Rubus L.* In: Hegi G (ed) *Illustrierte Flora von Mitteleuropa*, Band IV/2A, 3 edn. Blackwell
809 Wissenschafts-Verlag, Berlin, pp 284–595.
- 810 Williamson M, Fitter A. 1996. The varying success of invaders. *Ecology* 77(6):1661–1666

TABLES

Table 1.

Measurement units and variability (Min=minimum, Max=maximum and Mean values) of the predictor variables used to model native and non-native species richness at the plot scale.

Predictors	Measurement Unit	Min	Max	Mean
<i>Climatic variable</i>				
Climate (Scores on 1 st PCA axis)		-7.64	2.61	0.00
<i>Topographic variables</i>				
Elevation	m	122.00	1660.00	406.63
Slope	radiants	0.00	26.82	7.67
Solar radiation	MJ*cm ⁻² *year ⁻¹	4555.86	6975.20	5639.80
<i>Human disturbance variables</i>				
Road distance	m	0.00	3492.85	775.57
Landscape diversity (Shannon's H) - H _{Plot}		0.00	1.22	0.42
Landscape Eveness (Pielou's E) - E _{Plot}		0.00	0.10	0.03

Table 2. Measurement units and variability (Min=minimum, Max=maximum and Mean values) of the predictor variables used to model native and non-native species richness at the PA scale.

Predictors	Measurement Unit	Min	Max	Mean
<i>Climatic variables</i>				
Mean climate (scores on 1 st PCA axis)		-5.23	2.03	-0.26
Climate range (scores on 1 st PCA axis)		-1.47	4.04	1.66
<i>Topographic variable</i>				
Elevation range	m	22.00	989.00	390.76
Mean elevation	m	215.70	1242.06	454.87
<i>Human disturbance variables</i>				
Road density	m/km ²	0.00	1.23	0.58
% anthropogenic area	%	0.00	0.28	0.04
% agricultural area	%	0.00	92.90	25.19
% natural area	%	0.00	99.72	67.97
% wetlands	%	0.00	50.94	4.16
Landscape diversity (Shannon's H) - H _{PA}		0.00	1.55	0.96
Landscape evenness (Pielou's E) - E _{PA}		0.00	0.09	0.06
<i>Geographical variables</i>				
Area	km ²	2.68	137.47	28.24

Table 3

Summary output of the minimum adequate model for native and non-native species across spatial scales. Please note that for GLMM, Wald confidence (CI) intervals were computed.

	<i>Native species</i>		<i>Non-native species</i>	
	<i>Estimate</i>	<i>CI (2.5%, 97.5%)</i>	<i>Estimate</i>	<i>CI (2.5%, 97.5%)</i>
<i>Plot scale (GLMMs)</i>				
(Intercept)	2.77***	2.45, 3.09	3.10***	1.40, 4.80
Climate	-18.64***	-23.82, -13.45	107.59***	63.73, 151.45
H _{plot}	6.48***	3.81, 9.15	-	-
Slope	-0.06***	-0.08, -0.05	-0.27**	-0.44, -0.09
Road distance	-	-	-0.22**	-0.38, -0.06
<i>PA scale (GLMs)</i>				
(Intercept)	4.97***	4.70, 5.22	0.68***	0.26, 1.03
Area	6.53E-05*	1.33E-05, 1.14E-04	1.01E-04**	3.91E-05, 1.60E-04
% Wetland	-0.37	-0.82, -0.05	-	-
Mean climate	0.36*	0.12, 0.61	-	-
Climate range	0.43*	0.14, 0.73	0.62**	0.23, 1.03
Mean elevation	-	-	-0.90***	-1.44, -0.45

*** $p < 0.001$; ** $p < 0.01$; * $p < 0.05$.

FIGURES




Figure 1. Survey area and the II level of CORINE Land Cover (CLC) of plots on each PA (Protected Areas, including 17 SACs and 4 Nature Reserves) in Siena province. The sampling plots within each PA are shown in black. AMI: Cono Vulcanico del Monte Amiata; AVM: Alta Val di Merse; BOG: Bogatto; BVM: Basso Merse; CAS: Castelvechio; CET: Monte Cetona; CHN: Monti del Chianti; CHU: Lago di Chiusi; COR: Cornate e Fosini; FOR: Crete dell'Orcia e del Formone; LEO: Crete di Camposodo e Crete di Leonina; LUC: Lucciolabella; MNS: Montagnola Senese; MOL: Monte Oliveto Maggiore e Crete di Asciano; MPU: Lago di Montepulciano; FIG: Foreste del Siele e del Pigelletto di Piancastagnaio; PPO: Pietraporciana; RIP: Ripa d'Orcia, SAG: Bosco di Sant'Agnese; TOC: Tocchi; VFA: Val di Farma.




Figure 2. Relationship between the ratio of non-native species richness to native species richness and the native species richness.

[Click here to view linked References](#)



Click here to access/download
Attachment to Manuscript
Supplementary material.doc

