



Ultrasound evaluation of varicoceles: guidelines and recommendations of the European Society of Urogenital Radiology Scrotal and Penile Imaging Working Group (ESUR-SPIWG) for detection, classification, and grading

This is the peer reviewed (post-print) version of the following article:

Original

Ultrasound evaluation of varicoceles: guidelines and recommendations of the European Society of Urogenital Radiology Scrotal and Penile Imaging Working Group (ESUR-SPIWG) for detection, classification, and grading / Freeman, S., Bertolotto, M., Richenberg, J., Belfield, J., Dogra, V., Huang, D.Y., Lotti, F., Markiet, K., Nikolic, O., Ramanathan, S., Ramchandani, P., Rocher, L., Secil, M., Sidhu, P.S., Skrobisz, K., Studniarek, M., Tsili, A., Tuncay Turgut, A., Pavlica, P., Derchi, L.E.. - In: EUROPEAN RADIOLOGY. - ISSN 0938-7994. - STAMPA. - 30:1(2020), pp. 11-25. [10.1007/s00330-019-06280-y]

Availability:

This version is available at: 11368/2959535 since: 2020-03-02T17:42:35Z

Publisher:

Published

DOI:10.1007/s00330-019-06280-y

Terms of use:

Some rights reserved. More details in the PostPrint rights section below.

License:

Attribution-NonCommercial-NoDerivatives 4.0 International

Publisher copyright

--

(Article begins on next page)

Ultrasound evaluation of varicoceles: guidelines and recommendations of the European Society of Urogenital Radiology Scrotal and Penile Imaging Working Group (ESUR-SPIWG) for detection, classification, and grading

Simon Freeman¹  • Michele Bertolotto² • Jonathan Richenberg³ • Jane Belfield⁴ • Vikram Dogra⁵ • Dean Y. Huang⁶ • Francesco Lotti⁷ • Karolina Markiet⁸ • Olivera Nikolic⁹ • Subramaniyan Ramanathan¹⁰ • Parvati Ramchandani¹¹ • Laurence Rocher^{12,13} • Mustafa Secil¹⁴ • Paul S. Sidhu⁶ • Katarzyna Skrobisz⁸ • Michal Studniarek⁸ • Athina Tsili¹⁵ • Ahmet Tuncay Turgut¹⁶ • Pietro Pavlica¹⁷ • Lorenzo E. Derchi¹⁸ • members of the ESUR-SPIWG WG

Abstract

Varicoceles are relatively common particularly in asymptomatic men and are even more prevalent in subfertile men, representing the most common potentially correctable cause of male infertility. Ultrasound (US) is the imaging modality of choice for varicocele evaluation, but there is no widely accepted consensus on examination technique, diagnostic criteria, or classification. In view of this uncertainty, the guideline writing group (WG) of the European Society of Urogenital Radiology (ESUR) Scrotal and Penile Imaging Working Group (ESUR-SPIWG) undertook a literature review and assessment of the quality of relevant evidence. The group then produced evidence-based recommendations for varicocele US examination, interpretation, and classification by consensus agreement. The results are presented in the form of 15 clinical questions with a brief summary of the relevant evidence and the authorised recommendations from the SPIWG. This paper provides a short summary of the evidence evaluation and the complete recommendations.

Key Points

- *Varicocele is a common clinical problem; it is highly prevalent amongst subfertile men and the most common potentially correctable cause of male infertility.*
- *Ultrasound is the imaging modality of choice for varicocele assessment, but there is no generally agreed consensus on the US examination technique or the criteria that should be used for diagnosis, grading, and classification.*
- *This paper summarises the recommendations of the ESUR-SPIWG for standardising the US assessment of varicoceles. This includes examination technique, image interpretation, classification, and reporting.*

Keywords Varicocele · Scrotum · Infertility, male · Ultrasonography · Ultrasonography, Doppler

Abbreviations

AGREE	Appraisal of Guidelines for Research and Evaluation
AVMs	Arteriovenous malformations
ESUR	European Society of Urogenital Radiology

Simon Freeman and Michele Bertolotto are joint first authors (making an equal contribution to this manuscript).

✉ Simon Freeman
simonfreeman@nhs.net

GRADE	Grading of Recommendations Assessment, Development and Evaluation
H	Height
L	Length
OCEBM	Oxford Centre for Evidence Based Medicine
PICO	Patient Problem (or Population), Intervention, Comparison or Control, Outcome
Sec	Seconds
SPIWG	Scrotal and Penile Imaging Working Group
V	Volume
W	Width
WG	Guideline writing group

Extended author information available on the last page of the article

Introduction

Varicoceles are a common urological problem. Although often detected incidentally, they are more prevalent in patients with infertility or chronic scrotal discomfort [1] and represent the commonest potentially correctable cause of male infertility.

Clinical assessment is unreliable and highly dependent on the clinician’s expertise [2]. US is the imaging modality of choice, although the need for US is debated [3, 4] and there is no consensus on the technique, diagnostic criteria or which of many different classification systems should be used [2]. Despite this, there is evidence that accurate imaging of varicoceles is helpful in treatment decisions [5–7].

Given these uncertainties, the guideline writing group (WG) of the ESUR-SPIWG undertook a literature review with the aim of producing evidence-based recommendations for standardising the technique and interpretation of the US examination. This paper summarises the evidence and provides the authorised SPIWG guidelines.

Materials and Methods

Guidelines were developed in accordance with the Appraisal of Guidelines for Research and Evaluation (AGREE) II document [8]. The WG formulated 15 questions after deliberating difficulties in their clinical practice, and these were transferred into PICO format [9, 10]. Four biomedical literature databases (PubMed, Web of Science, Embase, and the Cochrane Library) were searched for all relevant articles published before 2018 in English, German, French, or Italian. The quality of evidence was rated according to the Oxford Centre for Evidence Based Medicine (OCEBM) 2011 levels of evidence (Table 1) [11] and recommendations graded using the Grading of Recommendations Assessment, Development and Evaluation (GRADE) system (Table 2) [12]. The recommendations were circulated to all members of the SPIWG and discussed in detail at the SPIWG meeting in September 2018 and modifications agreed. Consensus was obtained by vote according to the modified Delphi technique [13] with

members able to approve, reject, or abstain from each recommendation.

Results

Question 1: What is the evidence for correlation between varicocele, spermatogenesis damage, and infertility?

There is a male contribution to infertility in approximately half the cases where a cause can be identified [6]. Varicocele is the most common treatable cause [14] and has an estimated prevalence of 15% in the general population [15], 40% in primary, and 75–81% in secondary male infertility [16–18]. Seventy-five percent of men with a varicocele are however fertile [19] and 80% have normal semen parameters [20].

Many different mechanisms have been proposed to explain the link between varicoceles and impaired spermatogenesis. A multifactorial etiology is likely involving heat, oxidative stresses, adrenal catecholamine reflux, and androgen deprivation [21–25].

Although sperm quality frequently improves following varicocele treatment, there is conflicting evidence regarding the value in improving male fertility [7, 26–32]; many studies have concluded that the current level of evidence is insufficient for a definitive opinion. Guidelines for varicocele repair in subfertility are also inconsistent [3, 6, 33].

Question 2: How are varicoceles classified by ultrasound?

Several different US sonographic classifications have been proposed for varicocele assessment [34–44] (Table 3). Unfortunately, there is no standardisation; US technique, parameters assessed, and diagnostic criteria are highly variable, making direct comparison or meta-analysis of pooled data problematic. A clear consensus has not been reached and there is no universally recognized system. Potentially conflicting results may be obtained in the same patient depending on

Table 1 Oxford Centre for Evidence-Based Medicine 2011 Levels of Evidence (diagnosis)

Level	
1	Systematic review of cross sectional studies with consistently applied reference standard and blinding
2	Individual cross sectional studies with consistently applied reference standard and blinding
3	Non-consecutive studies, or studies without consistently applied reference standards
4	Case-control studies, or “poor or non-independent reference standard”*
5	Mechanism-based reasoning

Source: OCEBM Levels of Evidence Working Group. “The Oxford 2011 Levels of Evidence.” Oxford Centre for Evidence-Based Medicine. <http://www.cebm.net/index.aspx?o=5653>. Accessed: 9 May 2018

Table 2 Grading of Recommendations Assessment, Development and Evaluation (GRADE)

Code	Quality of Evidence	Definition
A	High	Several high-quality studies with consistent results In special cases: one large, high-quality multicentre trial
B	Moderate	One high-quality study Several studies with some limitations
C	Low	One or more studies with severe limitations
D	Very low	Expert opinion No direct research evidence One or more studies with very severe limitations

Source: https://www.essentialevidenceplus.com/product/ebm_loe.cfm?show=grade. Accessed 9 May 2018

the classification system used, and there is a low predictive value for treatment outcomes.

Recommendation 1

Grey-scale and Doppler US modes are used to assess the parameters required for varicocele classification. There is no universally recognized classification system (LoE 3, GoR C). Strong consensus (21 approve/0 reject /0 abstain: 100%)

Question 3: Should the size of the dilated veins be measured? Should measurements be performed standing or supine, at rest or during the Valsalva maneuver? Which size threshold should be used for the dilated veins?

There is a wide variation in the threshold venous diameter recommended for diagnosis of varicoceles, the measurement site, patient position, and need for Valsalva [38, 45–54]. A venous diameter of 3 mm is commonly considered to be the threshold for diagnosis of a varicocele although higher and lower values have been advocated [39, 49, 50, 53, 55, 56] (Fig. 1).

It is critically important to document the patient’s position and the sampling site. The majority of investigators examine the patient in both supine and erect positions and during Valsalva, measuring the largest vein irrespective of location.

Recommendation 2

Given the widespread methodological variability that exists in measurements of venous diameter in varicocele assessment, it is critically important to document the patient’s position, whether measurement was made at rest or during the Valsalva manoeuvre, and the location of the measured veins relative to the spermatic cord or testis (LoE 1, GoR A). Strong consensus (21/0/0, 100%)

Recommendation 3

Measurement of the largest vein, irrespective of location, with the patient in the upright position and during the Valsalva manoeuvre is recommended (LoE 5, GoR D) Strong consensus (21/0/0, 100%)

Recommendation 4

A maximum venous diameter of 3mm or more can be considered diagnostic for a varicocele when measured with the patient in the upright position and during the Valsalva manoeuvre, (LoE 2, GoR B). Strong consensus (20/1/0, 95%)

Question 4: When to measure testicular size at ultrasound, and how?

Venous reflux due to a varicocele is associated with reduced testicular volume which may be reversed by varicocele repair [57–59]. Normal testicular volume, measured by US, varies widely between different studies and in different populations. A total testicular volume greater than 20–24 ml is associated with normal testicular function in Caucasian and African men [60–63]. Lower testicular volumes are reported in infertile patients [63–66].

Three different formulae are used to calculate testicular volume from US measurements of length, width, and height. The most accurate has been shown to be Lambert’s formula ($V = L \times W \times H \times 0.71$) [67–69].

Recommendation 5


Testicular volume should be measured in all cases as it correlates with testicular function in both infertile patients and patients with a varicocele (LoE 1, GoR A). Strong consensus (21/0/0, 100%)

Table 3 Classifications of varicocele according to ultrasonographic criteria

Study, year	Patient position	Grades			
Hirsh et al (1980) [35]	Standing	[1] No spontaneous reflux, inducible with Valsalva [1] Reflux < 2 sec	[2] Intermittent spontaneous reflux [2] Reflux > 2 sec	[3] Continuous spontaneous reflux [3] Spontaneous reflux increasing with Valsalva	[5] Enlarged vessels in supine and standing positions, with caliber not increasing with Valsalva. Reflux at rest, not increasing during Valsalva. Testicular hypotrophy. Intratesticular varices may be present
Dhabuwala et al (1989) [36]	Supine			[3] Peritesticular reflux only during Valsalva in enlarged vessels. Visible but not dilated vessels when supine. Enlarged when standing	
Sarteschi et al (1993) [37]	Standing and supine	[1] Inguinal reflux only during Valsalva in not enlarged vessels	[2] Supra-testicular varicosities with reflux only during Valsalva		
Hoekstra and Witt (1995) [38]	Standing	[1] Dilated veins < 2.5 mm without flow reversal after Valsalva Maximum vein diameter (mm) 0: < 2.5 mm 1: 2.5–2.9 mm 2: 3.0–3.9 mm 3: ≥ 4 mm	[2] Dilated veins 2.5–3.5 mm and flow reversal after Valsalva Plexus/sum of diameter of veins 0: no plexus identified 1: plexus with sum diameter > 3 mm 2: plexus with sum diameter 3–5.9 mm 3: plexus with sum diameter ≥ 6 mm	[3] Dilated veins > 3.5 mm and flow reversal after Valsalva Change of flow velocity on Valsalva maneuver 0: < 2 cm/sec or duration < 1 sec 1: 2–4.9 cm/sec 2: 5–9.9 cm/sec 3: ≥ 10 cm/sec	[4] Enlarged vessels in supine and standing positions, with increasing caliber with Valsalva. Reflux at rest, increasing during Valsalva. Possible testicular hypotrophy Total score 0–9 ≥ 4: presence of varicocele
Chiou et al (1997) [39]	Supine				
Comud et al (1999) [40]	Not specified	[1] Brief reflux < 1 sec	[2] Intermediate reflux < 2 sec decreasing during and stopping prior to the end of Valsalva	[3] Permanent reflux > 2 sec and with a plateau aspect throughout the abdominal strain	
Oyen (2002) [41]	Supine	[1] Slight reflux (< 2 sec) during Valsalva	[2] Reflux (> 2 sec) during Valsalva, not continuous	[3] Reflux at rest or continuous during the entire Valsalva maneuver	[4] Obvious vessel enlargement with basal reflux that does not increase during Valsalva
Pauroso et al (2011) [42]	Supine	[1] No varicosities seen. Reflux in the vessels of the inguinal canal that is observed only during Valsalva	[2] Small varicosities with reflux seen only during Valsalva		
Iosa and Lazzarini (2013) [43]	Standing and supine	[1] Reflux > 1 sec only during Valsalva	[2] Spontaneous, discontinuous reflux not increasing by Valsalva	[3] Spontaneous, discontinuous reflux increased by Valsalva	
Patil et al (2016) [44]	Standing	[0] Reflux time < 1 sec	[1] Reflux time 1–2.5 sec	[2] Reflux time 2.5–4 sec	[3] Reflux time > 4 sec

Scores are reported in square brackets

Fig. 1 Grey-scale image of a varicocele. There are multiple serpiginous dilated veins with a venous diameter of more than 3 mm



Recommendation 6

Accurate measurement of the three diameters of the testis is required to obtain testicular volume estimation. Use of Lambert's formula ($V=LxWxHx0.71$) is recommended. The mathematical formula used to calculate the volume should be reported (LoE 2, GoR B). Broad consensus (19/1/1, 95%)

Question 5: How should US be performed in patients with varicoceles?

The patient should be examined at rest and during Valsalva in the supine and erect positions; the standing position is often more informative. Bilateral grey-scale, colour, and spectral Doppler analysis are recommended.

Testicular volume should be calculated and the size of the largest vein measured, irrespective of venous location (within the inguinal canal or scrotum), patient position (supine or erect), or whether measurement was obtained with or without Valsalva. Colour and spectral Doppler should be used to detect and characterise venous reflux at the level of the inguinal canal, suprastesticular region, and level of the testis (Fig. 2).

Recommendation 7

A standardised protocol is required for varicocele ultrasound examination. A grey-scale and colour Doppler examination, with spectral Doppler analysis, should be performed bilaterally with the patient supine and standing, during spontaneous breathing and during the Valsalva manoeuvre. (LoE 2, GoR B). Strong consensus (21/0/0, 100%)

Question 6: Is Doppler evaluation of venous reflux needed and which parameters should be measured?

Venous reflux is believed to be the primary cause of testicular damage [70] which may be reversed if reflux is eliminated [71]. Evaluation of reflux is critical for the diagnosis of varicocele and prediction of treatment outcomes; continuous flow reversal or reflux triggered by Valsalva is strongly associated with post-operative improvement in semen quality [52, 70, 72].

Assessment of flow direction, changes with patient positioning, and Valsalva are made with colour Doppler and verified with spectral Doppler which should also measure the duration of reflux [44].

Recommendation 8

Demonstrating and evaluating reflux flow in patients being assessed for varicoceles is the most important part of the Doppler ultrasound study (LoE 3, GoR C.) Strong consensus (21/0/0, 100%)

Recommendation 9

Colour Doppler interrogation should be supplemented with spectral Doppler analysis. Reflux duration is the essential parameter to be measured (LoE 3, GoR C). Measurement of the reflux peak velocity is optional (LoE 5, GoR D). Strong consensus (21/0/0, 100%)

Question 7: How long should reflux last to make the diagnosis of varicocele?

The duration of venous reflux recommended for the diagnosis and grading of varicoceles varies significantly between different studies [36, 40, 41, 43, 44, 55, 73, 74]. Brief reflux

(< 1 sec) during Valsalva is often considered to be physiological [40, 43, 44]. Intermediate reflux has been defined as lasting < 2 sec with decreasing velocity during, and finishing before the end of, Valsalva and is considered to be an indeterminate finding. Permanent reflux lasts > 2 sec and shows a plateau throughout Valsalva; studies suggest that this degree of reflux is always associated with abnormality on venography [40, 60]. Reflux lasting > 2 sec is therefore considered to represent the most useful parameter for varicocele diagnosis (Fig. 3).

Recommendation 10

Reflux in the testicular veins lasting more than 2 seconds with the patient standing and during the Valsalva manoeuvre should be considered to be abnormal (LoE 4, GoR C). Strong consensus (21/0/0, 100%)

Question 8: Is reflux velocity clinically important, and how should it be measured?

Several investigations suggest that measurement of peak reflux venous velocity might be useful to predict the need for varicocele repair [55, 75–82]. There is a lack of consistency between the studies regarding the US technique. Mostly, angle correction was not performed and therefore velocity measurements cannot be regarded as accurate. Measurement of this parameter cannot be recommended in routine clinical practice.

Recommendation 11

There is insufficient data to recommend using reflux peak velocity measurements as a factor in determining the need for varicocele repair (LoE 5, GoR C). Strong consensus (21/0/0, 100%)

Question 9: How should US examinations be reported in patients with varicoceles?

The SPWIG believes that the following information should be included in the US report:

- Testicular volume, echogenicity, and echotexture.
- Testicular or extra-testicular abnormalities.
- Presence of varices at grey-scale and colour Doppler US, and relationships to the testis (inguinal canal, suprastesticular, around the testis, intratesticular).
- Diameter of the largest vein measured while standing and during the Valsalva maneuver.
- Changes of flow at colour and spectral Doppler according to the patient's position, before and during Valsalva. Evaluation of reflux peak velocity is optional.

A standardised report may be useful for this purpose (Table 4). It is not considered mandatory to grade varicoceles according to a classification system but, if utilised, the Sarteschi classification is preferred [20, 37].

Recommendation 12

When issuing reports on patients with varicoceles the examination technique should be described (LoE 1, GoR A). Strong consensus (21/0/0, 100%)

Recommendation 13

Grading varicoceles according to the Sarteschi classification may be helpful in clinical practice. For standardisation purposes it is recommended that all the US parameters used to evaluate the patient are also reported. (LoE 5, GoR D). Strong consensus (21/0/0, 100%)

Question 10: Is evaluation of intratesticular Doppler waveforms worthwhile in imaging varicoceles?

It has been proposed that increased intratesticular venous pressure secondary to varicocele may decrease testicular arterial blood supply resulting in impaired spermatogenesis [83, 84].

Studies evaluating intratesticular arterial blood flow have shown conflicting results, some showing no changes [85–87] but other showing alterations in volume flow [88], peak systolic velocity, and resistive or pulsatility indices [89, 90]. These studies are of interest in further understanding the pathophysiology of varicoceles but are not currently useful for routine clinical practice.

Recommendation 14

Evaluation of intra-testicular blood flow in patients with varicoceles is an active research field which might provide a valuable insight into the mechanisms that create testicular parenchymal damage. At present however, this evaluation cannot be recommended for clinical use. (LoE 3, GoR C) Strong consensus (21/0/0, 100%)

Question 11: Can evidence-based recommendations be provided for imaging right-sided varicoceles?

Clinical varicoceles are usually left-sided and unilateral; bilateral varicoceles are diagnosed in approximately 10% of patients [91]. Subclinical right-sided varicocele is however much more frequent [92–96] and bilateral disease underdiagnosed by clinical examination. Patients with left-sided varicoceles should therefore be carefully examined for right-sided disease by US [93–96]. Isolated right-sided

Fig. 2 Small varicocele lying above the testis (Sarteschi grade II). Colour Doppler images obtained at rest (a) and during Valsalva (b) showing that reflux is detected only during Valsalva




Fig. 3 Spectral Doppler analysis during Valsalva. Flow reversal demonstrated on the implementation of Valsalva (arrowhead) and persisting for more than 2 sec




Table 4 Example of a standardised US report for varicocele

	R	L
Testicular diameters (L, W, H) in mm		
Testicular volume (Lambert's formula: LxWxHx0.71) in mls.		
Testicular echogenicity abnormality Yes/No		
Intratesticular varicocele Yes/No		
SUPINE position		
Diameter of the largest vein (mm)		
Location of the largest vein (Spermatic cord /peritesticular)		
Spontaneous reflux at colour Doppler (Yes/No)		
Reflux during Valsalva manoeuvre Yes/No		
Level where reflux is identified (inguinal canal, supratesticular, peritesticular)		
Reflux velocity (optional)		
STANDING position		
Diameter of the largest vein (mm)		
Location of the largest vein (Spermatic cord /peritesticular)		
Spontaneous reflux at colour Doppler (Yes/No)		
Reflux during Valsalva manoeuvre Yes/No		
Duration of reflux (s)		
Level where reflux is identified (inguinal canal, supratesticular, peritesticular)		
Reflux velocity (optional)		

CONCLUSION:

Testicular atrophy: Yes/No

Varicocele: Yes/No

Varicocele Grade (Sarteschi’s classification):

Any other relevant findings:

varicocele is found in < 1% of patients [91], is usually subclinical and identified on US only during Valsalva [94, 96].

Due to the infrequency of clinical right-sided varicoceles, there is a belief that this should prompt the search for a secondary retroperitoneal cause. The literature does not support this approach in the majority of cases although US may rarely identify congenital venous anatomical variants [93, 97] and venous obstruction in high-risk cases.

Recommendation 15

Bilateral colour Doppler US should be performed in patients with left-sided varicoceles as it will frequently reveal subclinical right-sided varicoceles. (LoE 3, GoR B) Strong consensus (21/0/0, 100%)

Recommendation 16

In patients with an isolated clinical right-sided varicocele, US can be extended to the abdomen to look for abdominal and retroperitoneal pathology, as well as congenital vascular anomalies. (LoE 5, GoR D). Strong consensus (21/0/0, 100%)

Question 12: Is imaging follow-up necessary for subclinical varicoceles?

The need for treatment of subclinical varicoceles is controversial [6], especially in pre-pubertal boys and adolescents where the significant risk of over-treatment [98] must be weighed against the risk of progression to clinically evident disease and reduced testicular function [99, 100]. Practice varies widely regarding follow-up for non-operated patients but is specifically recommended in two groups who are at increased risk of progressive testicular dysfunction: adolescents with testicular hypotrophy and adolescents and young adults with normal testicular volume and semen analysis. Annual follow-up should include physical examination, US, and semen analysis [101].

Recommendation 17

In patients with subclinical varicoceles imaging follow-up is recommended in all adolescents who have not undergone surgical repair and in young adults with normal semen analysis and normal testicular volume (LoE 3, GoR C). Strong consensus (20/0/1 100%)

Question 13: Should patients be followed up after varicocele treatment?

Following treatment, US can be used for the assessment of early complications and later for the detection of persistent or recurrent disease [102–115]. Accurate clinical examination

following surgery is highly dependent on the experience of the clinician and will frequently detect enlarged veins, particularly for treated high-grade disease, even following successful treatment. Only US is able to determine whether there is persisting venous reflux that may require additional treatment. There are currently no generally accepted recommendations for imaging following varicocele repair.

Practice varies regarding the need for routine US following treatment [3, 103, 107]. Where subfertility is the indication for surgery, semen analysis should form the basis for follow-up with US reserved for cases where this remains unsatisfactory.

Recommendation 18

After varicocele repair US can be used to identify early post-operative complications (LoE 3, GoR C). Strong consensus (21/0/0, 100%)

Recommendation 19

Sperm analysis forms the basis of follow-up following varicocele repair. The data available does not support the routine use of US (LoE 1, GoR A). Broad consensus (18/1/2, 95%)

Recommendation 20

Colour Doppler US can be used after varicocele repair if semen analysis remains unsatisfactory to evaluate testicular volume and identify signs of persistent or recurrent disease (LoE 2, GoR B). Strong consensus (21/0/0, 100%).

Question 14: Is it always necessary to examine the abdomen for tumours in patients with a newly discovered varicocele?

Case reports and small series suggest that patients with retroperitoneal tumours may present with a varicocele [116–120]; however, varicoceles are common and retroperitoneal tumours rare. There is no compelling evidence to suggest that a retroperitoneal or renal tumour in a patient with a varicocele is more likely than in a man without a varicocele [42, 117, 119, 121]. Varicoceles are almost never the only clinical feature of a retroperitoneal tumour and usually a feature of advanced cancer stage which will frequently be detected by clinical examination [122–126]. The incidence of secondary right-sided varicocele is probably lower than for the left-sided varicocele [127].

Evidence does not support mandatory abdominal US in all patients with varicoceles. An exception is in children aged below 9 years of age; varicoceles are rare in this age group [128, 129] and the US examination should include the abdomen [130].

Although the incidence of an occult retroperitoneal mass is very low, the sonologist is justified in extending the scan to the abdomen whenever the varicocele is of acute onset, large or fails to decompress in the supine position (Fig. 4).


Recommendation 21

Extended US examination to the abdomen is recommended in children less than 9 years of age presenting with acute varicocele (LoE 2, GoR B). Strong consensus (21/0/0, 100%)

Recommendation 22

There is insufficient evidence to conclude that an extension of the ultrasound examination to the abdomen is mandatory in all adult patients with a varicocele. The ultrasound

Fig. 4 A patient presenting with an acute-onset large right-sided varicocele. On scrotal ultrasound, multiple dilated veins are seen around the testis, remaining distended with the patient in the supine position (a). Extending the examination to the abdomen (b) demonstrates a retroperitoneal mass (arrow) anterior to the aorta (Ao) and IVC, subsequently proven to represent an asymptomatic lymphoma mass



practitioner should use their clinical judgement to decide whether to proceed to an abdominal examination, particularly if the varicocele is large, of recent onset and persists with the patient in the supine position (LoE 5, GoR D). Strong consensus (21/0/0, 100%)

Question 15: What are the pitfalls in US when imaging varicoceles?

Spermatoceles, cysts, tubular ectasia, post-vasectomy changes, and Zinner syndrome [131] can mimic varicoceles but can usually be differentiated with colour Doppler.

Uncommon extra-testicular masses may present with a clinical suspicion of a varicocele; these include arteriovenous malformations (AVMs), cavernous haemangiomas, and lymphangiomas [132–134]. The correct diagnosis can usually

be made with US, although MRI is the modality of choice for detailed evaluation. Intratesticular varicoceles can be mimicked by a number of conditions [135] including tubular ectasia of the rete testis, intratesticular AVMs, and haemangiomas.

Recommendation 23

In patients being investigated for a clinically detected varicocele, the possibility of rare varicocele mimics should be considered. The correct diagnosis can usually be made by combining the grey-scale and Doppler US features (LoE 5, GoR D.) Strong consensus (21/0/0, 100%)

Funding The authors state that this work has not received any funding.

Compliance with ethical standards

Guarantor The scientific guarantor of this publication is Dr. Simon Freeman on behalf of the Scrotal and Penile Imaging Working Group of the European Society of Urogenital Radiology (ESUR-SPIWG).

Conflict of interest The authors of this manuscript declare no relationships with any companies, whose products or services may be related to the subject matter of the article.

Statistics and biometry No complex statistical methods were necessary for this paper.

Informed consent Written informed consent was not required for this study because this paper provides a summary of available evidence and expert opinion; no new studies on patients were involved.

Ethical approval Institutional Review Board approval was not required because this is an evidence summary and expert recommendation paper. No new studies on human patients were undertaken specifically for this paper.

Study subjects or cohorts overlap All previous studies cited as evidence in this paper are acknowledged in the references.

Methodology

• Multicentre authored paper on behalf of the Scrotal and Penile Imaging Working Group of the European Society of Urogenital Radiology (ESUR-SPIWG). Literature review, evidence summary, and expert recommendations.

References

1. Paick S, Choi WS (2019) Varicocele and testicular pain: a review. *World J Mens Health* 37(1):4–11
2. Dubin L, Amelar RD (1970) Varicocele size and results of varicocelectomy in selected subfertile men with varicocele. *Fertil Steril* 21(8):606–609
3. Practice Committee of the American Society for Reproductive Medicine; Society for Male Reproduction and Urology (2014) Report on varicocele and infertility: a committee opinion. *Fertil Steril* 102(6):1556–1560

4. Khera M, Lipshultz LI (2008) Evolving approach to the varicocele. *Urol Clin North Am* 35(2):183–189
5. Zini A, Dohle G (2011) Are varicoceles associated with increased deoxyribonucleic acid fragmentation? *Fertil Steril* 96(6):1283–1287
6. Jungwirth A, Giwercman A, Tournaye H et al (2012) European Association of Urology guidelines on male infertility: the 2012 update. *Eur Urol* 62(2):324–332
7. Agarwal A, Deepinder F, Cocuzzza M et al (2007) Efficacy of varicocelectomy in improving semen parameters: new meta-analytical approach. *Urology*. 70(3):532–538
8. Brouwers MC, Kho ME, Browman GP et al (2010) AGREE II: advancing guideline development, reporting and evaluation in health care. *CMAJ*. 182(18):E839–E842
9. van der Molen AJ, Reimer P, Dekkers IA et al (2018) Post-contrast acute kidney injury - Part 1: definition, clinical features, incidence, role of contrast medium and risk factors : recommendations for updated ESUR Contrast Medium Safety Committee guidelines. *Eur Radiol* 28(7):2845–2855
10. Guyatt GH, Oxman AD, Kunz R et al (2011) GRADE guidelines: 2. Framing the question and deciding on important outcomes. *J Clin Epidemiol* 64(4):395–400
11. Medicine OCfE-B. The Oxford 2011 Levels of Evidence 2011 Available from: <https://www.cebm.net/2016/05/ocbm-levels-of-evidence/>. Accessed April 2019
12. Plus EE. GRADE: Grading of Recommendations Assessment, Development and Evaluation (A, B, C, D): John Wiley & Sons, Inc; 2018. Available from: http://www.essential-evidence-plus.com/product/concept_infomastery.cfm. Accessed April 2019
13. Dalkey N, Helmer O (1963) An experimental application of the DELPHI method to the use of experts. *Manag Sci* 9(3):458–467
14. Jungwirth A, Diemer T, Kopa Z, Krausz C, Minhas S, Tournaye H (2018) Guidelines on male infertility, European Association of Urology. EAU Guidelines Office, Arnhem. Available from: <https://uroweb.org/wp-content/uploads/EAU-Guidelines-on-Male-Infertility-2018-large-text.pdf>. Accessed April 2019
15. Saypol DC (1981) Varicocele. *J Androl* 2(2):61–71
16. Alsaikhan B, Alrabeeh K, Delouya G, Zini A (2016) Epidemiology of varicocele. *Asian J Androl* 18(2):179–181
17. Dubin L, Amelar RD (1971) Etiologic factors in 1294 consecutive cases of male infertility. *Fertil Steril* 22(8):469–474
18. Kibar Y, Seckin B, Erduran D (2002) The effects of subinguinal varicocelectomy on Kruger morphology and semen parameters. *J Urol* 168(3):1071–1074
19. Zini A, Boman JM (2009) Varicocele: red flag or red herring? *Semin Reprod Med* 27(2):171–178
20. Liguori G, Trombetta C, Garaffa G et al (2004) Color Doppler ultrasound investigation of varicocele. *World J Urol* 22(5):378–381
21. Marmar JL (2001) Varicocele and male infertility: Part II: the pathophysiology of varicoceles in the light of current molecular and genetic information. *Hum Reprod Update* 7(5):461–472
22. Will MA, Swain J, Fode M, Sonksen J, Christman GM, Ohl D (2011) The great debate: varicocele treatment and impact on fertility. *Fertil Steril* 95(3):841–852
23. Naughton CK, Nangia AK, Agarwal A (2001) Varicocele and male infertility: Part II: pathophysiology of varicoceles in male infertility. *Hum Reprod Update* 7(5):473–481
24. Papanikolaou F, Chow V, Jarvi K, Fong B, Ho M, Zini A (2000) Effect of adult microsurgical varicocelectomy on testicular volume. *Urology*. 56(1):136–139
25. Jung A, Schuppe HC (2007) Influence of genital heat stress on semen quality in humans. *Andrologia*. 39(6):203–215
26. Evers JH, Collins J, Clarke J (2008) Surgery or embolisation for varicoceles in subfertile men. *Cochrane Database Syst Rev* 2008(3): CD000479. <https://doi.org/10.1002/14651858.CD000479.pub3>

27. Kroese AC, De Lange NM, Collins JA, Evers JL (2013) Varicocele surgery, new evidence. *Hum Reprod Update* 19(4):317
28. Kroese ACJ, de Lange NM, Collins J, Evers JL (2012) Surgery or embolization for varicoceles in subfertile men. *Cochrane Database Syst Rev* (10):CD000479. <https://doi.org/10.1002/14651858.CD000479.pub5>
29. Smit M, Romijn JC, Wildhagen MF, Veldhoven JL, Weber RF, Dohle GR (2010) Decreased sperm DNA fragmentation after surgical varicocelectomy is associated with increased pregnancy rate. *J Urol* 183(1):270–274
30. Baazeem A, Belzile E, Ciampi A et al (2011) Varicocele and male factor infertility treatment: a new meta-analysis and review of the role of varicocele repair. *Eur Urol* 60(4):796–808
31. Macleod R, Biyani CS, Cartledge J, Eardley I (2015) Varicocele. *BMJ Clin Evid* 2015
32. Kim KH, Lee JY, Kang DH, Lee H, Seo JT, Cho KS (2013) Impact of surgical varicocele repair on pregnancy rate in subfertile men with clinical varicocele and impaired semen quality: a meta-analysis of randomized clinical trials. *Korean J Urol* 54(10):703–709
33. Fertility: assessment and treatment for people with fertility problems. National Institute for Health and Clinical Excellence: Guidance [CG11]. London (UK) 2004. www.nice.org.uk/guidance/cg11. Accessed April 2019
34. Lotti F, Maggi M (2014) Ultrasound of the male genital tract in relation to male reproductive health. *Hum Reprod Update* 21(1): 56–83
35. Hirsh AV, Cameron KM, Tyler JP, Simpson J, Pryor JP (1980) The Doppler assessment of varicoceles and internal spermatic vein reflux in infertile men. *Br J Urol* 52(1):50–56
36. Dhabuwala CB, Kumar AB, Kerkar PD, Bhutawala A, Pierce J (1989) Patterns of Doppler recordings and its relationship to varicocele in infertile men. *Int J Androl* 12(6):430–438
37. Sarteschi LM, Paoli R, Bianchini M, Menchini Fabris GF (1993) Lo studio del varicocele con eco-color-Doppler. *G Ital Ultrasonologia* 4(2):43–49
38. Hoekstra T, Witt MA (1995) The correlation of internal spermatic vein palpability with ultrasonographic diameter and reversal of venous flow. *J Urol* 153(1):82–84
39. Chiou RK, Anderson JC, Wobig RK et al (1997) Color Doppler ultrasound criteria to diagnose varicoceles: correlation of a new scoring system with physical examination. *Urology*. 50(6):953–956
40. Cornud F, Belin X, Amar E, Delafontaine D, Helenon O, Moreau JF (1999) Varicocele: strategies in diagnosis and treatment. *Eur Radiol* 9(3):536–545
41. Oyen RH (2002) Scrotal ultrasound. *Eur Radiol* 12(1):19–34
42. Pauroso S, Di Leo N, Fulle I, Di Segni M, Alessi S, Maggini E (2011) Varicocele: ultrasonographic assessment in daily clinical practice. *J Ultrasound* 14(4):199–204
43. Iosa G, Lazzarini D (2013) Hemodynamic classification of varicoceles in men: our experience. *J Ultrasound* 16(2):57–63
44. Patil V, Shetty SC, Das SK (2016) Redefining the criteria for grading varicoceles based on reflux times: a clinicoradiological correlation. *Ultrasound Q* 32(1):82–85
45. Wolverson MK, Houttuin E, Heiberg E, Sundaram M, Gregory J (1983) High-resolution real-time sonography of scrotal varicocele. *AJR Am J Roentgenol* 141(4):775–779
46. Gonda RL Jr, Karo JJ, Forte RA, O'Donnell KT (1987) Diagnosis of subclinical varicocele in infertility. *AJR Am J Roentgenol* 148(1):71–75
47. Orda R, Sayfan J, Manor H, Witz E, Sofer Y (1987) Diagnosis of varicocele and postoperative evaluation using inguinal ultrasonography. *Ann Surg* 206(1):99–101
48. McClure RD, Khoo D, Jarvi K, Hricak H (1991) Subclinical varicocele: the effectiveness of varicocelectomy. *J Urol* 145(4):789–791
49. Eskew LA, Watson NE, Wolfman N, Bechtold R, Scharling E, Jarow JP (1993) Ultrasonographic diagnosis of varicoceles. *Fertil Steril* 60(4):693–697
50. Pilatz A, Altinkilic B, Kohler E, Marconi M, Weidner W (2011) Color Doppler ultrasound imaging in varicoceles: is the venous diameter sufficient for predicting clinical and subclinical varicocele? *World J Urol* 29(5):645–650
51. Jarow JP, Ogle SR, Eskew LA (1996) Seminal improvement following repair of ultrasound detected subclinical varicoceles. *J Urol* 155(4):1287–1290
52. Schiff JD, Li PS, Goldstein M (2006) Correlation of ultrasound-measured venous size and reversal of flow with Valsalva with improvement in semen-analysis parameters after varicocelectomy. *Fertil Steril* 86(1):250–252
53. Karami M, Mazdak H, Khanbabapour S, Adibi A, Nasr N (2014) Determination of the best position and site for color Doppler ultrasonographic evaluation of the testicular vein to define the clinical grades of varicocele ultrasonographically. *Adv Biomed Res* 3: 17
54. Kim YS, Kim SK, Cho IC, Min SK (2015) Efficacy of scrotal Doppler ultrasonography with the Valsalva maneuver, standing position, and resting-Valsalva ratio for varicocele diagnosis. *Korean J Urol* 56(2):144–149
55. Cina A, Minnetti M, Pirroni T et al (2006) Sonographic quantitative evaluation of scrotal veins in healthy subjects: normative values and implications for the diagnosis of varicocele. *Eur Urol* 50(2):345–350
56. Metin A, Bulut O, Temizkan M (1991) Relationship between the left spermatic vein diameter measured by ultrasound and palpated varicocele and Doppler ultrasound findings. *Int Urol Nephrol* 23(1):65–68
57. Zampieri N, Cervellione RM (2008) Varicocele in adolescents: a 6-year longitudinal and followup observational study. *J Urol* 180(4):1653–1656
58. Zhou T, Zhang W, Chen Q et al (2015) Effect of varicocelectomy on testis volume and semen parameters in adolescents: a meta-analysis. *Asian J Androl* 17(6):1012–1016
59. Sakamoto H, Saito K, Ogawa Y, Yoshida H (2008) Effects of varicocele repair in adults on ultrasonographically determined testicular volume and on semen profile. *Urology*. 71(3):485–489
60. Cornud F, Amar E, Hamida K, Thiounn N, Helenon O, Moreau JF (2000) Imaging in male hypofertility and impotence. *BJU Int* 86: 153–163
61. Goede J, Hack WW, Sijtermans K et al (2011) Normative values for testicular volume measured by ultrasonography in a normal population from infancy to adolescence. *Horm Res Paediatr* 76(1):56–64
62. Kim ED, Lipshultz LI (1996) Role of ultrasound in the assessment of male infertility. *J Clin Ultrasound* 24(8):437–453
63. Condorelli R, Calogero AE, La Vignera S (2013) Relationship between testicular volume and conventional or nonconventional sperm parameters. *Int J Endocrinol* 2013:6
64. Tijani KH, Oyende BO, Awosanya GO, Ojewola RW, Yusuf AO (2014) Assessment of testicular volume: a comparison of fertile and sub-fertile West African men. *Afr J Urol* 20(3):136–140
65. Sakamoto H, Ogawa Y, Yoshida H (2008) Relationship between testicular volume and testicular function: comparison of the Prader orchidometric and ultrasonographic measurements in patients with infertility. *Asian J Androl* 10(2):319–324
66. Sakamoto H, Yajima T, Nagata M, Okumura T, Suzuki K, Ogawa Y (2008) Relationship between testicular size by ultrasonography and testicular function: measurement of testicular length, width, and depth in patients with infertility. *Int J Urol* 15(6):529–533

67. Lambert B (1951) The frequency of mumps and of mumps orchitis and the consequences for sexuality and fertility. *Acta Genet Stat Med* 2(Suppl. 1):1–166
68. Mbaeri TU, Orakwe JC, Nwofor AME, Oranusi CK, Mbonu OO (2013) Ultrasound measurements of testicular volume: comparing the three common formulas with the true testicular volume determined by water displacement. *Afr J Urol* 19(2):69–73
69. Sakamoto H, Saito K, Oohta M, Inoue K, Ogawa Y, Yoshida H (2007) Testicular volume measurement: comparison of ultrasonography, orchidometry, and water displacement. *Urology*. 69(1):152–157
70. Liguori G, Trombetta C, Ollandini G et al (2009) Predictive factors of better improvement in semen quality after sclerotization of varicocele: preliminary report. *J Androl Sci* 16:47–53
71. Abdel-Meguid TA, Al-Sayyad A, Tayib A, Farsi HM (2011) Does varicocele repair improve male infertility? An evidence-based perspective from a randomized, controlled trial. *Eur Urol* 59(3):455–461
72. Liguori G, Ollandini G, Pomara G et al (2010) Role of renoepididymal basal reflow and age on semen quality improvement after sclerotization of varicocele. *Urology*. 75(5):1074–1078
73. Cornud F, Amar E, Hamida K, Helenon O, Moreau JF (2001) Ultrasound findings in male hypofertility and impotence. *Eur Radiol* 11(11):2126–2136
74. Kocakoc E, Serhatlioglu S, Kiris A, Bozgeyik Z, Ozdemir H, Bodakci MN (2003) Color Doppler sonographic evaluation of inter-relations between diameter, reflux and flow volume of testicular veins in varicocele. *Eur Radiol* 47(3):251–256
75. Kozakowski KA, Gjertson CK, Decastro GJ, Poon S, Gasalberti A, Glassberg KI (2009) Peak retrograde flow: a novel predictor of persistent, progressive and new onset asymmetry in adolescent varicocele. *J Urol* 181(6):2717–2722
76. Tasci AI, Resim S, Caskurlu T, Dincel C, Bayraktar Z, Gurbuz G (2001) Color Doppler ultrasonography and spectral analysis of venous flow in diagnosis of varicocele. *Eur Urol* 39(3):316–321
77. Mohseni MJ, Nazari H, Amini E et al (2011) Shunt-type and stop-type varicocele in adolescents: prognostic value of these two different hemodynamic patterns. *Fertil Steril* 96(5):1091–1096
78. Glassberg KI (2014) My indications for treatment of the adolescent varicocele (and why?). *Trans Androl Urol* 3(4):402–412
79. Paduch DA, Niedzielski J (1997) Repair versus observation in adolescent varicocele: a prospective study. *J Urol* 158(3):1128–1132
80. Poon SA, Gjertson CK, Mercado MA, Raimondi PM, Kozakowski KA, Glassberg KI (2010) Testicular asymmetry and adolescent varicoceles managed expectantly. *J Urol* 183(2): 731–734
81. Morteza T, Ehsan A (2016) Quantitative Doppler assessment of flow characteristics in pampiniform venous plexus, in patients with varicocele: is there any relationship with semen analysis? *PJR*. 20(3):121–124
82. Van Batavia JP, Badalato G, Fast A, Glassberg KI (2013) Adolescent varicocele-is the 20/38 harbinger a durable predictor of testicular asymmetry? *J Urol* 189(5):1897–1902
83. Hsu HS, Chang LS, Cheng MT, Wei YH (1994) Decreased blood flow and defective energy metabolism in the varicocele-bearing testicles of rats. *Eur Urol* 25:71–75
84. Zhang M, Du L, Liu Z, Qi H, Chu Q (2014) The effects of varicolectomy on testicular arterial blood flow: laparoscopic surgery versus microsurgery. *Urol J* 11(5):1900–1906
85. Grasso FL, Pepe P, Panella P, Pepe F (1997) Volocimetric evaluation of spermatic vessels with echo color Doppler in patients with idiopathic varicocele. *Minerva Urol Nefrol* 49(4):179–182
86. Ross JA, Watson JN Jr, Jarow JP (1994) The effect of varicoceles on testicular blood flow in man. *Urology*. 44(4):535–539
87. Akcar N, Turgut M, Adapinar B, Ozkan IR (2004) Intratesticular arterial resistance and testicular volume in infertile men with sub-clinical varicocele. *J Clin Ultrasound* 32(8):389–393
88. Tarhan S, Gumus B, Gunduz I, Ayyildiz V, Goktan C (2003) Effect of varicocele on testicular artery blood flow in men color Doppler investigation. *Scand J Urol Nephrol* 37(1):38–42
89. Biagiotti G, Cavallini G, Modenini F, Vitali G, Gianaroli L (2002) Spermatogenesis and spectral echo-colour Doppler traces from the main testicular artery. *BJU Int* 90(9):903–908
90. Unsal A, Turgut AT, Taskin F, Kosar U, Karaman CZ (2007) Resistance and pulsatility index increase in capsular branches of testicular artery: indicator of impaired testicular microcirculation in varicocele? *J Clin Ultrasound* 35(4):191–195
91. Baigorri BF, Dixon RG (2016) Varicocele: a review. *Semin Intervent Radiol* 33(3):170–176
92. (1985) Comparison among different methods for the diagnosis of varicocele. *World Health Organisation.. Fertil Steril* 43(4):575–582
93. Cariati M, Pieri S, Agresti P et al (2012) Diagnosis of right-sided varicocele: a retrospective comparative study between clinical examination, Doppler findings, US imaging and vascular anatomy at phlebography. *Eur J Radiol* 81(9):1998–2006
94. DeCastro GJ, Shabsigh A, Poon SA, Laor L, Glassberg KI (2009) Adolescent varicolectomy - is the potential for catch-up growth related to age and/or Tanner stage? *J Urol* 181(1):322–327
95. Gat Y, Bachar GN, Zukerman Z, Belenky A, Gorenish M (2004) Physical examination may miss the diagnosis of bilateral varicocele: a comparative study of 4 diagnostic modalities. *J Urol* 172(4):1414–1417
96. Pfeiffer D, Berger J, Schoop C, Tauber R (2006) A Doppler-based study on the prevalence of varicocele in German children and adolescents. *Andrologia*. 38(1):13–19
97. Preziosi P, Miano R, Bitelli M, Ciolfi MG, Micali S, Micali F (2001) Right varicocele associated with inferior vena cava malformation in situs inversus: percutaneous treatment with retrograde sclerotherapy. *J Endourol* 15(10):1001–1003
98. Ding H, Tian J, Du W, Zhang L, Wang H, Wang Z (2012) Open non-microsurgical, laparoscopic or open microsurgical varicolectomy for male infertility: a meta-analysis of randomized controlled trials. *BJU Int* 110(10):1536–1542
99. Cervellione RM, Corroppo M, Bianchi A (2008) Subclinical varicocele in the pediatric age group. *J Urol* 179(2):717–719
100. Zampieri N, Dall'Agnola A (2011) Subclinical varicocele and sports: a longitudinal study. *Urology*. 77(5):1199–1202
101. Diamond DA (2007) Adolescent varicocele. *Curr Opin Urol* 17(4):263–267
102. Cvitanic OA, Cronan JJ, Sigman M, Landau ST (1993) Varicoceles: postoperative prevalence-a prospective study with color Doppler US. *Radiology*. 187(3):711–714
103. Rotker K, Sigman M (2016) Recurrent varicocele. *Asian J Androl* 18(2):229–233
104. Glassberg KI, Badalato GM, Poon SA, Mercado MA, Raimondi PM, Gasalberti A (2011) Evaluation and management of the persistent/recurrent varicocele. *Urology*. 77(5):1194–1198
105. Kwak N, Siegel D (2014) Imaging and interventional therapy for varicoceles. *Curr Urol Rep* 15(4):399
106. El-Haggag S, Nassef S, Gadalla A, Latif A, Mostafa T (2012) Ultrasonographic parameters of the spermatic veins at the inguinal and scrotal levels in varicocele diagnosis and post-operative repair. *Andrologia*. 44(3):210–213
107. Winkelbauer FW, Ammann ME, Karnel F, Lammer J (1994) Doppler sonography of varicocele: long-term follow-up after venography and transcatheter sclerotherapy. *J Ultrasound Med* 13(12):953–958
108. Palazzo S, Boezio F, Mancini V et al (2002) Significance of scrotal Doppler color ultrasonography in the postoperative control of

- patients treated with varicocelectomy. *Arch Ital Urol Androl* 74(4):247–249
109. Hauser R, Yorgev L, Greif M et al (1997) Sperm binding and ultrasound changes after operative repair of varicocele: correlation with fecundity. *Andrologia*. 29(3):145–147
 110. Zucchi A, Mearini L, Mearini E, Fioretti F, Bini V, Porena M (2006) Varicocele and fertility: relationship between testicular volume and seminal parameters before and after treatment. *J Androl* 27(4):548–551
 111. Trombetta C, Salisci E, Deriu M et al (1993) Echo-flowmetric control 6 years after percutaneous treatment of varicocele. *Arch Ital Urol Androl* 65(4):363–367
 112. Haller J, Gritzmann N, Czemberek H et al (1986) Pulsed duplex sonographic follow-up of varicoceles following sclerotherapy. *Rofo*. 145(5):547–550
 113. Hamm B, Sorensen R, Berger T (1986) Non-invasive imaging procedures in the diagnosis and control of therapy of varicoceles. 2. Importance of thermography and sonography for therapy control following sclerotherapy of varicoceles. *Rofo*. 144(6):665–669
 114. Mastrojeni C, Pagano G, Scalisi M, Leonello G, Manfrè A, Versaci A (2001) Idiopathic varicocele. Diagnosis and follow-up with CW-Doppler and echo-color-Doppler: our experience. *G Chir*. 22(1-2):41–44
 115. Cil AS, Bozkurt M, Bozkurt DK, Gok M (2015) Investigating the relationship between persistent reflux flow on the first postoperative day and recurrent varicocele in varicocelectomy patients. *J Clin Med Res* 7(1):29–32
 116. El-Saeity NS, Sidhu PS (2006) “Scrotal varicocele, exclude a renal tumour”. Is this evidence based? *Clin Radiol* 61(7):593–599
 117. Spittel JA Jr, DeWeerd JH, Shick RM (1959) Acute varicocele: a vascular clue to renal tumor. *Proc Staff Meet Mayo Clin* 34(5): 134–137
 118. Gibbons RP, Monte JE, Correa RJ Jr, Mason JT (1976) Manifestations of renal cell carcinoma. *Urology*. 8(3):201–206
 119. Skinner DG, Colvin RB, Vermillion CD, Pfister RC, Leadbetter WF (1971) Diagnosis and management of renal cell carcinoma. A clinical and pathologic study of 309 cases. *Cancer*. 28(5):1165–1177
 120. El Abiad Y, Qarro A (2016) Acute varicocele revealing renal cancer. *N Engl J Med* 374(21):2075
 121. Bate J (1927) Symptomatic varicocele. *J Urol* 18(6):649–662
 122. Ding GX, Song NH, Feng CC et al (2012) Is there an association between advanced stage of renal cell carcinoma and paraneoplastic syndrome? *Med Princ Pract* 21(4):370–374
 123. Ates N, Yuksel M, Yilmaz S, Habibi M, Ipekci T (2016) Retroperitoneal paraganglioma presenting as right-sided varicocele: case report. *Ann Saudi Med* 36(2):148–151
 124. Arrabal-Polo MA, Merino-Salas S (2013) Acute left varicocele in an adult. *CMAJ*. 185(4):323
 125. Chandramohan S, Chakravertry S (2007) Re: “Scrotal varicocele, exclude a renal tumour”. Is this evidence based? *Clin Radiol* 62(2):192–193
 126. Espinosa Bravo R, Lemourt Oliva M, Pérez Monzón AF et al (2003) Carcinoma de células renales y varicocele izquierdo de forma simultánea. *Arch Esp Urol* 56(5):533–535
 127. Robson J, Wolstenhulme S, Knapp P (2012) Is there a co-association between renal or retroperitoneal tumours and scrotal varicoceles? A systematic review. *Ultrasound*. 20(4):182–191
 128. Oster J (1971) Varicocele in children and adolescents: an investigation of the incidence among Danish school children. *Scand J Urol Nephrol* 5(1):27–32
 129. Akbay E, Cayan S, Doruk E, Duce MN, Bozlu M (2000) The prevalence of varicocele and varicocele-related testicular atrophy in Turkish children and adolescents. *BJU Int* 86(4):490–493
 130. Erginel B, Vural S, Akin M et al (2014) Wilms’ tumor: a 24-year retrospective study from a single center. *Pediatr Hematol Oncol* 31(5):409–414
 131. Pavan N, Bucci S, Mazzon G, Bertolotto M, Trombetta C, Liguori G (2015) It’s not always varicocele: a strange case of Zinner syndrome. *Can Urol Assoc J* 9(7-8):E535–E538
 132. Conzi R, Damasio MB, Bertolotto M et al (2017) Sonography of scrotal wall lesions and correlation with other modalities. *J Ultrasound Med* 36(10):2149–2163
 133. Secil M, Bertolotto M, Rocher L et al (2017) Imaging features of paratesticular masses. *J Ultrasound Med* 36(7):1487–1509
 134. Yilmaz C, Arslan M, Arslan M (2009) Intrascrotal arteriovenous malformation simulating varicocele. *AJR Am J Roentgenol* 192(6):W351
 135. Bhatt S, Jafri SZ, Wasserman N, Dogra VS (2011) Imaging of non-neoplastic intratesticular masses. *Diagn Interv Radiol* 17(1): 52–63

Publisher’s note Springer Nature remains neutral with regard to jurisdictional claims in published maps and institutional affiliations.

Affiliations

Simon Freeman¹  • Michele Bertolotto² • Jonathan Richenberg³ • Jane Belfield⁴ • Vikram Dogra⁵ • Dean Y. Huang⁶ • Francesco Lotti⁷ • Karolina Markiet⁸ • Olivera Nikolic⁹ • Subramanayan Ramanathan¹⁰ • Parvati Ramchandani¹¹ • Laurence Rocher^{12,13} • Mustafa Secil¹⁴ • Paul S. Sidhu⁶ • Katarzyna Skrobisz⁸ • Michal Studniarek⁸ • Athina Tsili¹⁵ • Ahmet Tuncay Turgut¹⁶ • Pietro Pavlica¹⁷ • Lorenzo E. Derchi¹⁸ • members of the ESUR-SPIWG WG

¹ University Hospitals Plymouth NHS Trust, Derriford Hospital, Derriford Road, Crownhill, Plymouth, Devon PL6 8DH, UK

² Department of Radiology, Ospedale di Cattinara, University of Trieste, Strada di Fiume 447, 34149 Trieste, Italy

³ Brighton and Sussex University Hospitals NHS Trust, Royal Sussex County Hospital, Eastern Road, Brighton BN2 5BE, UK

⁴ Royal Liverpool University Hospital, Prescot street, Liverpool L7 8XP, UK

⁵ Department of Radiology, University of Rochester, 601 Elmwood ave, Rochester, NY 14642, USA

⁶ King’s College Hospital NHS Foundation Trust, Department of Radiology, King’s College Hospital, Denmark Hill, London SE5 9RS, UK

- ⁷ Andrology, Female Endocrinology and Gender Incongruence Unit, Department of Experimental and Clinical Biomedical Sciences, University of Florence, Viale Pieraccini 6, I-50139 Florence, Italy
- ⁸ Department of Radiology, Medical University of Gdansk, Mariana Smoluchowskiego 17, 80-214 Gdansk, Poland
- ⁹ Faculty of Medicine, Center of Radiology, Clinical Center of Vojvodina, University of Novi Sad, Hajduk Veljkova 1-9, Novi Sad 21000, Serbia
- ¹⁰ Department of Clinical Imaging, Weill Cornell Medical College, Al-Wakra Hospital, Hamad Medical Corporation, PO Box 82228, Doha, Qatar
- ¹¹ Perelman School of Medicine, University of Pennsylvania, 3400 Spruce Street, Philadelphia, PA 19104, USA
- ¹² Department of Adult Diagnostic and Interventional Radiology, Bicetre University Hospital, 78 avenue du Général Leclerc, 94270 Le Kremlin Bicêtre, France
- ¹³ Imagerie par Résonance Magnétique Médicale et Multi-Modalités, CNRS UMR8081, Service Hospitalier Frédéric Joliot, 4, Place du Gal Leclerc, 91401 Orsay Cedex, France
- ¹⁴ Faculty of Medicine, Department of Radiology, Dokuz Eylul University, Izmir, Turkey
- ¹⁵ Department of Clinical Radiology, Medical School, University of Ioannina, University Campus, 45110 Ioannina, Greece
- ¹⁶ Department of Radiology, Faculty of Medicine, Istinye University, Istanbul, Turkey
- ¹⁷ Private Hospital S.M. Maddalena, Occhiobello, Rovigo, Italy
- ¹⁸ Department of Health Sciences (DISSAL), Emergency Radiology, University of Genoa, Policlinico San Martino, Largo R. Benzi, I16122 Genoa, Italy