



Dermoscopic features of a primary scalp melanoma and its cutaneous metastases

This is the peer reviewed (post-print) version of the following article:

Original

Dermoscopic features of a primary scalp melanoma and its cutaneous metastases / Pizzichetta, M.A., Canzonieri, V., Militello, L., Borsatti, E., Lo Re, G., Bolzonello, S., Olivieri, M., Pinzani, C., Magaton-Rizzi, G., Puglisi, F., Zalaudek, I.. - In: ITALIAN JOURNAL OF DERMATOLOGY AND VENEREOLOGY. - ISSN 2784-8450. - 156:4(2021), pp. 499-501. [10.23736/S2784-8671.20.06742-5]

Availability:

This version is available at: 11368/2976149 since: 2020-12-09T18:17:50Z

Publisher:

Published

DOI:10.23736/S2784-8671.20.06742-5

Terms of use:

Some rights reserved. More details in the PostPrint rights section below.

License:

Copyright Editore

Publisher copyright

--

(Article begins on next page)

ARTICLE ONLINE FIRST

This provisional PDF corresponds to the article as it appeared upon acceptance.

A copyedited and fully formatted version will be made available soon.

The final version may contain major or minor changes.

Dermoscopic features of a primary scalp melanoma and its cutaneous metastases

Maria Antonietta PIZZICHETTA, Vincenzo CANZONIERI, Loredana MILITELLO ,
Eugenio BORSATTI, Giovanni LO RE, Silvia BOLZONELLO, Matteo OLIVIERI,
Caterina PINZANI, Giovanni MAGATON RIZZI, Fabio PUGLISI, Iris ZALAUDEK

Giornale Italiano di Dermatologia e Venereologia 2020 Nov 23

DOI: 10.23736/S0392-0488.20.06742-5

Article type: Letter to the Editor

© 2020 EDIZIONI MINERVA MEDICA

Article first published online: November 23, 2020

Manuscript accepted: August 5, 2020

Manuscript revised: July 25, 2020

Manuscript received: May 26, 2020

Subscription: Information about subscribing to Minerva Medica journals is online at:

<http://www.minervamedica.it/en/how-to-order-journals.php>

Reprints and permissions: For information about reprints and permissions send an email to:

journals.dept@minervamedica.it - journals2.dept@minervamedica.it - journals6.dept@minervamedica.it

Dermoscopic features of a primary scalp melanoma and its cutaneous metastases

Running title: Scalp melanoma with simultaneous cutaneous metastases

Maria A. Pizzichetta^{1,2*}, Vincenzo Canzonieri^{3,4}, Loredana Militello², Eugenio Borsatti⁵, Giovanni Lo Re², Silvia Bolzonello², Matteo Olivieri⁶, Caterina Pinzani², Giovanni Magaton-Rizzi¹, Fabio Puglisi^{7,2}, Iris Zalaudek¹

¹ Dermatologic Clinic, University of Trieste, Trieste, Italy;

² Department of Medical Oncology, Centro di Riferimento Oncologico di Aviano (CRO) IRCCS, Aviano, Italy

³ Department of Medical, Surgical and Health Sciences, University of Trieste, Trieste, Italy

⁴ Division of Pathology, Centro di Riferimento Oncologico di Aviano (CRO) IRCCS, Aviano, Italy

⁵ Division of Nuclear Medicine, Centro di Riferimento Oncologico di Aviano (CRO) IRCCS, Aviano, Italy

⁶ Division of Surgery, Centro di Riferimento Oncologico di Aviano (CRO) IRCCS, Aviano, Italy

⁷ Department of Medicine (DAME), University of Udine, Italy

Correspondence to: Maria Antonietta Pizzichetta, MD, Division of Medical Oncology and

Preventive Oncology, Centro di Riferimento Oncologico, Via Franco Gallini 2, I-33081 Aviano

(PN), Italy. Phone: +39-0434-659480; Fax: +39-0434-659453; E-mail: pizzichetta@cro.it

Cutaneous melanoma metastases (CMM) occur frequently in the natural history of melanoma, but the simultaneous occurrence of CMM with its primary lesion has so far been reported once, in a case of primary choroidal and cutaneous melanomas.¹ We describe an unusual case of the simultaneous occurrence of cutaneous metastases from a primary scalp amelanotic melanoma.

A 44-year-old man presented with multiple growing, pigmented papules and plaques on his scalp and trunk, first noticed approximately 3 months earlier. Dermoscopically, these lesions showed globular, multicomponent and pseudo-lacunar pattern suggestive of cutaneous metastases.

A careful inspection of the scalp revealed a 1.5 cm firm, skin-coloured, ulcerated nodule on the vertex, partially covered by hair and yellowish-brown crust (Fig. 1a). Dermoscopically, the lesion had a structureless pattern with white shiny structures (Fig. 1b). There also was a polymorphous vascular pattern of dotted, linear (irregular, hairpin, coiled, and helical), and arborizing vessels with milky-red globules. In addition, the right part of the lesion had a reddish-orange haemorrhagic crust from ulceration (Fig. 1b).

Considering the patient's medical history and the lesions' dermoscopic features, we made a diagnosis of primary scalp amelanotic melanoma with simultaneous CMM. The lesion on the vertex (Fig. 1a) and one on the left upper trunk (Fig. 1e) were excised. The histopathological examination of the scalp lesion revealed an ulcerated, polypoid cutaneous melanoma composed of atypical pigmented epithelioid cells, with a dense lymphoid infiltrate at the lesion base (Figs. 1c, d). It was a Clark level IV tumour, with a Breslow thickness of 4.7 mm and mitotic index $>5/\text{mm}^2$, without evident neoplastic vascular invasion. Instead, the left upper trunk lesion had sharp delimitation and relatively preserved symmetry. Atypical melanocytes were rare at the dermo-epidermal junction and tumour cells filled the papillary dermis (Figs. 1g, h). These features were consistent with the trunk lesion being a metastasis.

Immunohistochemical analysis showed that the neoplastic cells expressed HMB45, MITF1, BCL2, CD117/c-kit, p53, and Ki67, in the primary and putative secondary lesions to the same extent (not shown). Mutational analysis revealed a V600E mutation in the *BRAF* gene (c.1799T>A) in both lesions. PET/CT did not reveal suspicious hypermetabolic lesions in internal organs. The patient received the combined treatment with the B-Raf inhibitor dabrafenib and MEK inhibitor trametinib. After 18 months, there was complete remission of all CMM

In this case, clinical-dermoscopic whole body skin examination, including a scalp inspection, allowed us to diagnose a difficult amelanotic nodular melanoma of the scalp. Clinically, the nodule on the vertex could have gone unnoticed because it was a skin-coloured lesion that was covered by hair and simulated a dermal naevus. However, its dermoscopic pattern was completely different from that of scalp dermal naevi; these latter have a globular pattern frequently associated with central hypopigmentation (“eclipse naevi”) or perifollicular hypopigmentation.²

Dermoscopy helped identify not only CMM³ but also the primary scalp amelanotic nodular melanoma thanks to visualization of polymorphous vessels and milky-red globules associated with melanoma diagnosis.⁴ However, the analysis of these features does not lead to a specific diagnosis.

The use of reflectance confocal microscopy (RCM) could increase the accuracy of dermoscopic diagnosis of scalp melanoma because RCM features of this latter are similar to those of melanomas on other sites.⁵ The limitations for the RCM examination on the scalp are the presence of hair and the scalp convexity require the use of handheld confocal microscope which allows to scan the lesion but only in small fields of view preventing the visualization of the entire lesion.⁵

Total body examination, including the scalp, may facilitate the early diagnosis of scalp melanoma, possibly improving its prognosis.

Acknowledgments

The authors wish to thank Valerie Matarese for scientific editing and Nancy Michilin for technical support.

References

1. Eide N, Foerster A. Simultaneous occurrence of primary choroidal and cutaneous malignant melanoma and skin metastasis. *Acta Ophthalmol.* 1993; 71:496-499.
2. Stanganelli I, Argenziano G, Sera F et al. Dermoscopy of scalp tumours: a multi-centre study conducted by the International Dermoscopy Society. *J Eur Acad Dermatol Venereol* 2012; 26:953-963.
3. Costa J, Ortiz-Ibanez K, Salerni G et al. Dermoscopic patterns of melanoma metastases: interobserver consistency and accuracy for metastasis recognition. *Br J Dermatol.* 2013; 169:91-99.
4. Pizzichetta MA, Kittler H, Stanganelli I, et al. Dermoscopic diagnosis of amelanotic/hypomelanotic melanoma. *Br J Dermatol* 2017; 177(2):538-540.
5. Benati E, Longo C, Piana S, Moscarella E. Preliminary evaluation of reflectance confocal microscopy features of scalp melanoma. *Australasian Journal of Dermatology* 2017; 58:312-316.

Conflict of interest: The authors certify that there is no conflict of interest with any financial organization regarding the material discussed in the manuscript.

Authors' contributions: Maria A. Pizzichetta wrote the manuscript and provided images. All authors contributed to manuscript revision, read and approved the final version of the manuscript

Figure legend

Figure 1. Clinical, dermoscopic e histological images of primary amelanotic nodular melanoma of the scalp, and a cutaneous melanoma metastasis.

(a-d) Lesion on the vertex of the scalp. (a) The clinical image shows a skin-coloured, symmetrical, nodular lesion, covered by hair and a yellowish-brown crust from ulceration. (b) The dermoscopic image of the same lesion reveals a structureless pattern with white shiny lines and areas, and a polymorphous vascular pattern with dotted, linear (irregular, hairpin, coiled, and helical) and arborizing vessels, and milky-red areas and globules. The globules are poorly demarcated polygonal structures, often containing atypical central vessels, separated from each other by blurred shiny white lines. A haemorrhagic crust from ulceration is seen in the right part of the lesion.

Histopathological analysis (c, d) shows atypical pigmented epithelioid cells with obvious pagetoid intraepidermal spread. No evidence of neoplastic vascular invasion. A dense lymphoid infiltrate is present within the lesion. H&E, Original magnification 100x (b) and 200x (c).

(e-h) Lesion on the left upper trunk. (e) Clinically, the lesion is a firm dark brown papule with darker central pigmentation and regular borders. (f) The dermoscopic image of same lesion shows a pseudo-lacunar pattern with irregular, unfocused reddish-blue clods separated from each other by blurred whitish- blue lines. In addition, there is a bluish-white veil, some irregular brown globules and peripheral polymorphous vessels combined with milky-red globules. Histopathological analysis (g, h) reveals sharp delimitation and relatively preserved symmetry. Atypical melanocytes are rare at the dermo-epidermal junction and tumor cells fill the papillary dermis. A slight lymphoid infiltrate is present at the base of the lesion. These features are consistent with a metastatic lesion. H&E, Original magnifications, 50x (e) and 200x (f).

