

Journal Pre-proof

The evolutionary puzzle solution for the origins of the partial loss of the *Cr2* exon in notothenioid fishes

Alessia Ametrano, Marco Gerdol, Maria Vitale, Samuele Greco, Umberto Oreste, Maria Rosaria Coscia



PII: S1050-4648(21)00143-1

DOI: <https://doi.org/10.1016/j.fsi.2021.05.015>

Reference: YFSIM 7504

To appear in: *Fish and Shellfish Immunology*

Received Date: 18 December 2020


Revised Date: 29 April 2021

Accepted Date: 20 May 2021

Please cite this article as: Ametrano A, Gerdol M, Vitale M, Greco S, Oreste U, Coscia MR, The evolutionary puzzle solution for the origins of the partial loss of the *Cr2* exon in notothenioid fishes, *Fish and Shellfish Immunology*, <https://doi.org/10.1016/j.fsi.2021.05.015>.

This is a PDF file of an article that has undergone enhancements after acceptance, such as the addition of a cover page and metadata, and formatting for readability, but it is not yet the definitive version of record. This version will undergo additional copyediting, typesetting and review before it is published in its final form, but we are providing this version to give early visibility of the article. Please note that, during the production process, errors may be discovered which could affect the content, and all legal disclaimers that apply to the journal pertain.

© 2021 Published by Elsevier Ltd.



Journal Pre-proof

1 **The evolutionary puzzle solution for the origins of the partial loss of the *C τ 2* exon in**
 2 **notothenioid fishes**

3

4 Alessia Ametrano^{a,b}, Marco Gerdol^c, Maria Vitale^{a,d}, Samuele Greco^c, Umberto Oreste^a, Maria Rosaria
 5 Coscia^{a,*}

6

7 ^a Institute of Biochemistry and Cell Biology - National Research Council of Italy, Naples, Italy

8 ^b Department of Environmental, Biological and Pharmaceutical Sciences and Technologies, University of
 9 Campania Luigi Vanvitelli, Caserta, Italy

10 ^c Department of Life Sciences, University of Trieste, Trieste, Italy

11 ^d Department of Molecular Medicine and Medical biotechnology, University of Naples Federico II, Naples,
 12 Italy (Present address)

13 *Corresponding author:

14 E-mail address: mariarosaria.coscia@ibbc.cnr.it (Maria Rosaria Coscia)

15

16 **Abstract:** Cryonotothenioidea is the main group of fishes that thrive in the extremely cold Antarctic
 17 environment, thanks to the acquisition of peculiar morphological, physiological and molecular adaptations. We
 18 have previously disclosed that IgM, the main immunoglobulin isotype in teleosts, display typical cold-adapted
 19 features. Recently, we have analyzed the gene encoding the heavy chain constant region (CH) of the IgT isotype
 20 from the Antarctic teleost *Trematomus bernacchii* (family Nototheniidae), characterized by the near-complete
 21 deletion of the CH2 domain. Here, we aimed to track the loss of the CH2 domain along notothenioid phylogeny
 22 and to identify its ancestral origins. To this end, we obtained the *IgT* gene sequences from several species
 23 belonging to the Antarctic families Nototheniidae, Bathydraconidae and Artedidraconidae. All species display a
 24 CH2 remnant of variable size, encoded by a short *C τ 2* exon, which retains functional splicing sites and therefore
 25 is included in the mature transcript. We also considered representative species from the three non-Antarctic
 26 families: Elegendinopsioidea (*Eleginops maclovinus*), Pseudaphritioidea (*Pseudaphritis urvillii*) and Bovichtidae
 27 (*Bovichtus diacanthus* and *Cottoperca gobio*). Even though only *E. maclovinus*, the sister taxa of
 28 Cryonotothenioidea, shared the partial loss of *C τ 2*, the other non-Antarctic notothenioid species displayed early
 29 molecular signatures of this event. These results shed light on the evolutionary path that underlies the origins of
 30 this remarkable gene structural modification.

31

32 **Keywords:** teleost fish; Notothenioidei; genome modifications; IgT; exonic remnant; immunoglobulin domain;
 33 Antarctic marine environment; molecular evolution

34

35 **Abbreviations**

Ig	Immunoglobulin
Ig VH	Immunoglobulin heavy chain Variable region
Ig CH	Immunoglobulin heavy chain Constant domain
<i>Cτ</i>	IgT heavy chain constant exon
TbeL	<i>Trematomus bernacchii</i> IgT heavy chain Long variant
TbeS	<i>Trematomus bernacchii</i> IgT heavy chain Short variant
TbeSts	<i>Trematomus bernacchii</i> IgT heavy chain Shortest variant

36 1. Introduction

37 The peculiar architecture of the Immunoglobulin molecule (Ig) comprises two types of
38 domains with similar secondary and tertiary structures, but different amino acid sequences:
39 the variable domains (V), which provide immunoglobulins the ability to recognize and bind
40 foreign antigens, and the constant domains (C), which perform effector functions. Another
41 important difference between C and V domains derives from their underlying genomic
42 configuration. While C domains are always encoded by individual exons, the V domains of
43 both Ig heavy (VH) and light chains (VL) are generated by a recombination activating gene
44 -mediated rearrangement of multiple sets of *Variable (V)*, *Diversity (D)* and *Joining (J)* gene
45 segments. Each Ig gene encodes both a membrane-bound heavy chain form, which is present
46 on the B-cell surface as B cell receptor complex, and a secreted heavy chain, which can be
47 mainly found in blood. Both forms are generated by an alternative splicing process that
48 involves the 3' exons but does not alter the rearrangement of *VDJ* gene segments.

49 Different Ig isotypes can be distinguished based on the properties of their heavy chain
50 constant (CH) domains. Unlike mammals, which possess five Ig isotypes, teleosts only
51 possess three different Ig isotypes, i.e., IgM, IgD and IgT. IgM, mainly found in the blood but
52 also present in mucosal compartments, are the main players in systemic immune responses
53 against a broad range of pathogens [1]. Although the IgD isotype emerged early in vertebrate
54 evolution, its functional role in teleosts has remained unclear for a long time. Previous data
55 suggested that IgD participate in immune tolerance and in basophil-mediated immunity [2].
56 Most recently, Perdiguero et al. [3] have shown that secreted IgD interact with the gut
57 microbiota, playing a relevant role in maintaining mucosal homeostasis in rainbow trout. In
58 2005 a new Ig isotype, named "IgT" (for Teleost) or IgZ (for Zebrafish), was discovered
59 simultaneously in two teleost species, i.e., rainbow trout and zebrafish [4, 5]. Early studies
60 had initially linked IgT to a specialized mucosal role in the gut [6]. However, further
61 evidence has expanded this function to other tissues, which include the skin [7], the
62 nasopharyngeal tract [8, 9], the gills [9] and the buccal cavity [10]. In a recent work carried
63 out in rainbow trout, Yu et al. [11] have demonstrated that, after a parasitic challenge, IgT are
64 locally produced at the nasal mucosa, providing protection as the main Ig isotype in the nasal
65 adaptive immunity.

66 The growing body of molecular data obtained from other teleost species has
67 progressively revealed that IgT is a particularly heterogeneous Ig isotype. First, the number
68 of *IgT* genes per species can largely vary, from zero (e.g., channel catfish and medaka) to
69 several paralogous copies (e.g., rainbow trout and seabass) [12, 13]. The teleost *Ig heavy*
70 *chain (IgH) gene* locus organization had been originally described as similar to the
71 mammalian "translocon" type, consisting of multiple sets of *V* gene segments located
72 upstream of multiple *D* and *J* segments, followed by *C μ* and *C δ* exons, coding for the CH
73 regions of the IgM and IgD isotypes, respectively. Later, a comparative analysis of the *IgH*
74 locus in several teleost species overturned this concept, uncovering a novel and incredibly
75 diversified genomic organization [14]. In most cases, another set of *D τ -J τ -C τ* elements
76 encoding the IgT heavy chain (τ) was found upstream of previously described *D-J-C μ -C δ*
77 elements for IgM and IgD heavy chains. In this configuration, upstream *V* segments can be
78 rearranged either to *D τ J τ C τ* for the IgT VH region, or to *DJC μ* for the same region of IgM.

79 The canonical structure of IgT, exemplified by zebrafish, comprises a CH region
80 encoded by four *C τ* exons, hereinafter referred to as *C τ 1*, *C τ 2*, *C τ 3* and *C τ 4*. However,
81 remarkable variations have been observed in some teleosts. The pufferfish IgT heavy chain
82 gene only comprises two *C τ* exons, homologous to the zebrafish *C τ 1* and *C τ 4* exons,
83 respectively [15]. The Nile tilapia and the common carp also display unusual IgT domain
84 architectures: the former only possesses the first two CH domains [16], and the latter is
85 characterized by the presence of at least three chimeric IgM/IgT molecules, encoded by
86 different gene copies [17, 18]. The first report of an IgT CH region gene in a teleost species
87 living under extreme conditions dates back to 2015, when our group reported the
88 unprecedented case of a nearly complete truncation of the CH2 domain in the Antarctic fish
89 *Trematomus bernacchii* [19]. Two out of the three variants identified in this species, termed
90 Long (TbeL) and Short (TbeS), respectively) were encoded by alleles characterized by a
91 large deletion and only displayed a short remnant of the *C τ 2* exon, which was entirely
92 skipped by alternative splicing in the third isoform, termed Shortest (TbeSts).

93 Most Antarctic fishes belong to the Perciform suborder Notothenioidei
94 (Cryonotothenioidea), which comprises five families (Nototheniidae, Harpagiferidae,
95 Artedidraconidae, Bathydraconidae, Channichthyidae) and about 130 species of marine
96 fishes found in the Southern Ocean, with a circum-Antarctic distribution, but also found in
97 the more temperate coastal waters of the southern hemisphere [20]. Notothenioidei are
98 among the most intensively studied lineages of marine fishes since they are a rare example
99 of massive adaptive radiation driven by the same selective pressures (e.g., isolation and
100 cooling) that may have led to the dramatic extinction of most fish fauna in the Southern
101 Ocean [21]. The evolutionary success of Notothenioidei is marked by the acquisition of key
102 adaptive features that enabled cold adaptation, such as the expression of antifreeze
103 glycoproteins [22]. At the same time, Notothenioidei lost other traits universally shared by
104 non-Antarctic metazoans, such as the inducible heat shock response [23] and, in the family
105 Channichthyidae, hemoglobin [24].

106 Apart from the Antarctic Clade [25], three non-Antarctic lineages, distributed in
107 proximity of the Southern Ocean, i.e., the southern regions of South America, around the
108 Falkland Islands, Tristan da Cunha, New Zealand and south-eastern Australia, are currently
109 recognized [21, 26]. While the first one, Bovichtidae, includes six species, the two other
110 families are monotypic and therefore include a single species, i.e., *Pseudaphritis urvillii*
111 (family Pseudaphritiidae) and *Eleginops maclovinus* (family Eleginopsioidea). More
112 recently, much attention has been dedicated to these notothenioid taxa, due to their early
113 divergence from the polar lineage, which occurred before the climatic and geographic
114 isolation of Antarctica, the drastic reduction in water temperature, and prior to the
115 morpho-physiological diversification of cryonotothenioid species [27]. The study of the
116 evolutionary history of Bovichtidae has been an essential factor for clarifying the process
117 that drove the diversification between Cryonotothenioidea and their non-Antarctic relatives
118 [27]. Moreover, the revised positioning of *E. maclovinus* as the sister lineage of
119 Cryonotothenioidea [28], provides another key information for the study of the evolution of
120 these fishes.

121 In light of recent molecular and morphological studies, which have allowed a revision of
122 the notothenioid phylogeny [28-32], the present work aims to extend the molecular analysis
123 of IgT to the other notothenioid families and to solve the key question as to whether the
124 features of the *T. bernacchii* IgT are unique to this species or they are shared by other
125 Antarctic species. In order to track the evolutionary history of the *C τ 2* exon and to pinpoint
126 the timing of its partial loss, we obtained and comparatively investigated the *IgT* sequences
127 of representatives from each of the five Antarctic notothenioid families and the three
128 non-Antarctic lineages of Bovichtidae (*Bovichtus diachantus* and *Cottoperca gobio*),
129 Eleginopsioidea (*E. maclovinus*) and Pseudaphritioidae (*P. urvillii*). The findings reported
130 here bring further insights into the molecular evolution of the *IgT* gene in Antarctic fishes,
131 marking the loss of the *C τ 2* exon before the split between the Eleginopsioidea and
132 Cryonotothenioidea lineages, and revealing early signatures of this event in the other
133 early-branching non-Antarctic Notothenioidei.

134

135 2. Materials and methods

136

137 2.1 Biological samples

138

139 The biological material was collected from a set of species representing the Antarctic
140 families Nototheniidae (*Gobionotothen gibberifrons*); Artedidraconidae (*Histiodraco*
141 *velifer*); Bathydraconidae (*Gymnodraco acuticeps*); Channichthyidae (*Chionodraco*
142 *hamatus*, *Chionodraco rastrospinosus*, *Chaenocephalus aceratus*, and *Pageotopsis*
143 *macropterus*); and the three non-Antarctic lineages Bovichtidae (*Bovichtus diachantus* and
144 *Cottoperca gobio*), Eleginopsioidea (*Eleginops maclovinus*) and Pseudaphritioidae
145 (*Pseudaphritis urvillii*). *H. velifer* specimens, as well as the Channichthyidae *C. hamatus* and
146 *P. macropterus* specimens, were caught in the Ross Sea, in the proximity of the Italian
147 "Mario Zucchelli" Station at 74°42' S, 164°07' E, during the XXV Italian Antarctic
148 Expedition (2009-2010). Specimens of the other two species of the family Channichthyidae,
149 *C. rastrospinosus* and *C. aceratus*, were collected by bottom trawling from the research
150 vessel L.M. Gould near Low and Braband Islands in the Palmer Archipelago during the XIX
151 Italian Antarctic expedition (2003-2004). The activity permit, released by the Italian
152 National Program for Antarctic Research (PNRA), was in agreement with the "Protocol on
153 environmental protection to the Antarctic Treaty" Annex V. *P. urvillii*, *E. maclovinus*, *G.*
154 *acuticeps*, and *G. gibberifrons* specimens were collected during the ICEFISH cruise 2004
155 (International Collaborative Expedition to collect and study Fish Indigenous to
156 Sub-Antarctic Habitats). Spleen and head kidney samples were collected and immediately
157 frozen in liquid nitrogen. All samples, including the blood from *P. urvillii* and *G.*
158 *gibberifrons*, were kept at -80 °C until use. The tissue sample of each species and
159 classification of the non-Antarctic and Antarctic sampled specimens are summarized in
160 Table S1.

161

162 2.2 RNA extraction, PCR amplification, and cloning of cDNA encoding the notothenioid IgT 163 CH region

164

165 Considering the variable number of specimens and tissues available for each species, we
166 selected tissue samples of two individuals per species for RNA extraction. The one exception
167 was *E. maclovinus*, as only a single specimen was available for this species. Following the
168 instructions of the SV Total RNA Isolation System kit (Promega), RNA extractions were
169 carried out from 150-200 mg tissue samples, homogenized by Potter-Elvehjem glass-Teflon
170 in TriPure Isolation reagent (Roche). Initial quality checks revealed the successful isolation
171 of RNA suitable for downstream analyses in all samples, with the only exception of *P.*
172 *urvillii*, where heavy degradation of RNA (obtained from blood cells), incompatible with
173 reverse transcription and PCR applications, was observed.

174 RNA quality and concentration were assessed by the presence of sharp rRNA bands on 1%
175 agarose gel and by calculating the ratio of absorbance at 260 nm and 280 nm measured by a
176 NanoDrop 1000 Spectrophotometer (Thermo Scientific). cDNA was obtained from 5 µg of
177 total RNA using Maxima H Minus Reverse Transcriptase (Thermo Scientific). The target
178 sequence was amplified in a final volume of 25 µl using 2 µl cDNA (20 ng), 1,25 µM of
179 specific primers (1,0 µM), 0,5 µl dNTP Mix (0,2 µM), 2,5 µl 10X DreamTaq Buffer, 0,5 µl
180 (1 U) of DreamTaq DNA polymerase (Thermo Scientific), up to volume with H₂O as
181 follows: 95 °C for 3 min, 35 cycles of 95 °C (30 s), 60 °C (30 s), and 72 °C (1 min) with a
182 final extension at 72 °C for 10 min. In the case of *E. maclovinus*, a second amplification was
183 carried out following the same conditions as the first PCR in order to increase the amount of
184 the specific product. Primers used in all the PCR experiments are reported in Table S2, which
185 also shows the targeted constant domain of the IgT heavy chain for each primer. PCR
186 products were analyzed on 1% agarose gel, purified by NucleoSpin[®] Gel and PCR Clean-up
187 (Macherey-Nagel), and cloned into pGEM[®]-T Easy Vector (Promega). Positive clones were
188 identified by the blue/white screening method and sequenced on both strands on an ABI
189 PRISM 3100 automated sequencer at Eurofins Genomics Europe Sequencing GmbH
190 (Jakob-Stadler-Platz 7, 78467 Konstanz, Germany). The 5' end of nucleotide sequences of
191 the positive cDNA clones matched the forward primer except for *E. maclovinus* (Fig. S1). In
192 this case, the obtained sequences were missing several nucleotides at the 5' end due to a high
193 background noise in the chromatogram. The number of positive cDNA clones encoding the
194 Ig C τ 2 variants identified both in non-Antarctic and Antarctic species are shown in Table S3.

195

196 2.3 DNA extraction, PCR amplification, and cloning of the notothenioid IgT CH region gene

197

198 DNA was isolated from 100 mg of head kidney sample of the single *E. maclovinus*
199 specimen available, using the PureLink[®] Genomic DNA Mini Kit (Thermo Scientific),
200 following the manufacturer's instructions. The same experimental procedure was applied for
201 *G. gibberifrons*, *H. velifer* and *G. acuticeps* spleen and for *N. coriiceps* and *P. urvillii* blood.
202 PCR amplification of genomic DNA was performed using 150 ng of template gDNA and the
203 same Taq polymerase described in the previous section. Primers used are reported in
204 supplementary table 1. PCR products were analysed on 1% agarose gel, purified by
205 NucleoSpin[®] Gel and PCR Clean-up (Macherey-Nagel), and cloned into pGEM[®]-T Easy
206 Vector (Promega). Positive clones were identified by the blue/white screening method and

207 sequenced on both strands on ABI PRISM 3100 automated sequencer Eurofins Genomics
208 Europe Sequencing GmbH (Jakob-Stadler-Platz 7, 78467 Konstanz, Germany). Not all 5'
209 ends of nucleotide sequences of the positive genomic clones matched the forward primer
210 (Fig. S2 and S3) due to the high background noise in the chromatogram. The number of
211 positive genomic clones and the corresponding encoded Ig $C\tau 2$ variants identified both in
212 non-Antarctic and Antarctic species are summarized in Table S4.

213

214 2.4 Additional sequences obtained from public repositories

215

216 Sequence data concerning the *IgT* gene and cDNA sequences from *T. bernacchii* (family
217 Nototheniidae) and from *B. diacanthus* (family Bovichtidae) were retrieved from a previous
218 study carried out by our research team [19]. The genomic DNA and predicted cDNA
219 sequences of the *IgT* genes of *Notothenia coriiceps* (family Nototheniidae) and *C. gobio* were
220 retrieved from publicly available annotated genome assemblies. In detail, the *N. coriiceps*
221 genome refers to the study by Shin et al. [29] whereas the genome of *C. gobio* (v.
222 fCotCob3.1) has been recently released within the frame of the Vertebrate Genome Project
223 [33].

224

225 2.5 Computational analysis

226

227 Sequencing chromatograms were visualized using the program FinchTV (version 1.3.0).
228 The nucleotide sequences obtained were verified by sequence similarity searches against the
229 GenBank database, using the BLAST program [34]. Amino acid sequences were deduced
230 from nucleotide sequences using the ExPASy Translate Tool tool. The amino acid
231 composition was analysed using the ExPASy ProtParam and Pep-Calc (www.pepcalc.com)
232 tools. The GC-content of introns was calculated with the GC Content Calculator (Biologics
233 International Corp, Indianapolis, USA). Multiple sequence alignments were performed with
234 ClustalW [35]. Sequons and putative *N*-Glycosylation sites were identified using the
235 NetNGlyc 4.0 Server [36] (at <https://www.expasy.org/proteomics>). A 3D molecular model
236 was built for the *IgT* CH domains of *E. maclovinus* (sequence Ema2) using the Phyre2 tool
237 [37] available at <http://www.sbg.bio.ic.ac.uk/phyre2>. The structure of the human secretory
238 IgA1 was used as template (PDB entry: 3CHN).

239

240 2.6 Association between *IgT* genes and repeated elements

241

242 The reference genomes assembly of *C. gobio* [33] was analyzed with RepeatScout v.1.05
243 [38] to generate a species-specific repeat library. The *IgT* genomic DNA sequences obtained
244 in this study, as well as those identified in the genome of *C. gobio*, were analyzed with
245 RepeatMasker v.4.09 [39], with particular attention to the $C\tau 1$ - $C\tau 2$ and $C\tau 2$ - $C\tau 3$ intronic
246 regions. All sequences were screened for the presence of repeats against the Dfam v.3.1 [40]
247 library of known repeats found in the genomes of Actinopterygii. The *IgT* gene of *C. gobio*
248 was subjected to an additional round of screening against the custom species-specific repeat
249 libraries generated as described above.

250

251 *2.7 Molecular evolution of the C τ 1-C τ 2 and C τ 2-C τ 3 introns*

252

253 We investigated the evolutionary history of the *IgT* genes identified in Notothenioidei,
254 with particular focus on the genomic region subjected to the highest molecular diversity, i.e.,
255 the region spanning the *C τ 1-C τ 2* and *C τ 2-C τ 3* introns. Due to the truncation of the *C τ 2* exon
256 in Cryonotothenioidea (see the results section), this region was disregarded. The nucleotide
257 sequences of the two introns were separately aligned with MUSCLE v.3.8.31 [41]. The
258 multiple sequence alignments were manually refined and processed with GBLOCKS v.0.91b
259 [42] to remove phylogenetically uninformative positions. The two sequence blocks were
260 concatenated and tested with ModelTest-NG v.0.1.3 [43] to identify the best-fitting model of
261 molecular evolution. This was found to be the GTR+I model (Generalized time-reversible,
262 with a proportion of invariable sites) [44], based on the corrected Akaike information
263 criterion [45]. The multiple sequence alignment file was used as an input for Bayesian
264 phylogenetic inference, carried out with MrBayes v.3.2.7a [46], run for one million
265 generations and two parallel Markov Chain Monte Carlo (MCMC) analyses. The
266 convergence of the two independent analyses was checked with Tracer [47], by evaluating
267 that all the estimated parameters reached an ESS value higher than 200.

268

269 *2.8 Selection analyses*

270

271 The *Ig C τ* cDNA sequences of the available species from Cryonotothenioidea, either
272 determined by cloning or inferred from genomic DNA, were aligned with a strategy aimed at
273 preserving the integrity of codon triplets. This was achieved by aligning the translated amino
274 acid sequences with MUSCLE v.3.8.31 [41] within the MEGAX environment [48] and
275 back-translating the aligned sequences to the original gapped coding nucleotide sequence.
276 The multiple sequence analysis was subjected to tests aimed at detecting signatures of
277 selection with the DataMonkey adaptive evolution platform [49]. Sites evolving under
278 pervasive positive and negative selection were detected with FEL (Fixed Effect Likelihood)
279 [50] and those with evidence of episodic positive selection were identified with MEME
280 (Mixed Effects Model of Evolution) [51] using default p-value thresholds.

281

282 *2.9 Identification of short conserved sequence motifs in Antarctic species*

283

284 The unaligned *IgT* protein sequences of Cryonotothenioidea were analyzed to detect the
285 presence of short ungapped conserved amino acid motifs of 3-8 aa length with MEME
286 (Multiple Em for Motif Elicitation) [52] within the MEME suite v.5.0.1 environment [53].
287 The specific association of the detected motifs with Cryonotothenioidea was subsequently
288 tested by assessing their absence in the *IgT* sequences of non-Antarctic Notothenioidei and
289 temperate Perciformes. Finally, the relevance of the identified motifs in the context of
290 evolution in the Antarctic environment was evaluated by inspecting their overlap with
291 negatively selected sites, identified as explained in the previous section.

292 The *C τ 1-C τ 2* and *C τ 2-C τ 3* introns of all the available Notothenioidei sequences were
293 similarly screened, looking for conserved nucleotide motifs with size comprised between 6
294 and 50 base pairs, allowing any number of motif repeats for each sequence.

295

296 2.10 *In silico* gene expression analysis

297

298 We evaluated the expression levels of the *IgT* and *IgM* genes in two species
299 representative of Antarctic (*T. bernacchii*) and non-Antarctic (*E. maclovinus*)
300 notothenioids, based on the analysis of publicly available RNA-seq datasets. Namely, the
301 raw sequencing data were recovered from the Bioprojects PRJNA471228 and
302 PRJNA401363 for *T. bernacchii* and *E. maclovinus*, respectively. In both cases, available
303 data concerned a single individual. The selected tissues, for which RNA-seq data were
304 available in both species, were: intestine, gills, head kidney, spleen, skin. In detail, reads
305 were imported in the CLC Genomics Workbench v.20 (Qiagen, Hilden, Germany)
306 environment, trimmed to remove low quality nucleotides and mapped to the reference *IgT*
307 and *IgM* sequences of the two species. This procedure was carried out with the following
308 parameters: *length fraction*: 0.5; *similarity fraction*: 0.9. To ensure a reliable comparison
309 between the gene expression levels between the two species and avoid possible
310 quantification issues linked with the presence of alternatively spliced forms encoding
311 secretory and membrane-bound Igs, the mapping was carried out on the *C τ 3/C τ 4* and
312 *C μ 3/C μ 4* exons only. Gene expression levels were reported as Counts Per Million (CPM).

313

314 2.11 Data availability

315

316 Genomic DNA and/or cDNA sequences from *P. urvillii*, *E. maclovinus*, *G. gibberifrons*,
317 *N. coriiceps*, *H. velifer*, and *G. acuticeps* have been deposited in the GenBank database
318 (<http://www.ncbi.nlm.nih.gov/genbank/>) are available with the following accession
319 numbers: MN583561 (gDNA clone gPur1); MN583559 (gDNA clone gEma1); MN583560
320 (gDNA clone gEma3); MN583563 (gDNA clone gGgi2); MN583562 (gDNA clone gNco1);
321 MN602803 (gDNA clone gGac1); MN602804 (clone gGgi3, pseudogene); MN602807
322 (gDNA clone gHve2); MN602808 (gDNA clone gHve5); MN583555 (cDNA clone Ema1);
323 MN583556 (cDNA clone Ema2); MN583557 (cDNA clone Ggi1); MN602805 (cDNA clone
324 Hve1); MN602806 (cDNA clone Hve2); MN583558 (cDNA clone Gac1).

325

326 3. Results

327

328 3.1 The *IgT* cDNA sequence of *Eleginops maclovinus* provides new evidence about the origin 329 of the loss of *C τ 2* exon in Antarctic species

330

331 Our investigations on the *IgT* cDNA sequences targeted several Antarctic species
332 belonging to the Nototheniidae, Artetidraconidae, Bathydraconidae and Channichthyidae
333 families (see the Materials and methods section for details), but also included the

334 non-Antarctic species *E. maclovinus* due to its phylogenetic placement as a sister lineage of
335 Cryonotothenioidea.

336 We obtained partial cDNA sequences, coding for the CH region of the IgT secreted form
337 in all species analyzed, with the single exception of *E. maclovinus*, where the IgT
338 membrane-bound form was obtained. The multiple sequence alignment highlighted the high
339 conservation of the *C τ 1* and *C τ 4* exons in all species, as expected from previous publications
340 (Fig. 1 and Fig. S1). Mirroring the previously reported case of *T. bernacchii*, all Antarctic
341 teleosts displayed a truncated *C τ 2* exon, whose size ranged from 24 to 51 nt (Fig. 1, Table 1).

342 The two variants cloned in *E. maclovinus* displayed a partially deleted *C τ 2* exon (30 or
343 36 nt long in Ema1 and Ema2, respectively; see Table 1), revealing a *C τ 2* exon structure
344 similar to Antarctic Notothenioidei. On the other hand, we have previously shown that *B.*
345 *diacanthus*, a non-Antarctic species more distantly related with Cryonotothenioidea, retains
346 the entire *C τ 2* exon (285 nt) [27] (Table 1).

347 Although Antarctic species and *E. maclovinus* shared the peculiar structure of *C τ 2*, their
348 sequences differed due to several characteristic codon indels. In detail, the *C τ 3* exon of all
349 Antarctic species lacked one codon at positions 502, 598 and 617 (the latter was not missing
350 in *N. coriiceps*), and two consecutive codons at position 562. On the other hand, a single
351 codon insertion was found at position 526 in the Antarctic lineage (Fig. S1). Interestingly, all
352 Cryonotothenioidea also presented four additional codons in the highly conserved *C τ 4* exon
353 at positions 715, 718, 778 and 784 (Fig. S1).

354

355 *3.2 Genomic analysis of the IgT gene in Antarctic species and Eleginops maclovinus: putting*
356 *together the first pieces of the puzzle*

357

358 The next step in the exploration of the molecular mechanisms behind the partial loss of
359 the *C τ 2* exon in Antarctic IgT gene was the extension of our analyses to the neighboring
360 genomic regions (i.e., the *C τ 1-C τ 2* and the *C τ 2-C τ 3* introns). Based on the data reported in
361 the previous paragraph, we used *E. maclovinus* as a reference for comparative analyses (Fig.
362 2 and Fig. S2).

363 The two IgT genomic variants found in this sub-Antarctic species (Ema1 and Ema3)
364 were characterized by a *C τ 2* exon of variable size (30/36 nt) and matched those identified at
365 the cDNA level. The *C τ 1-C τ 2* intron of the isoform with the shortest *C τ 2* exon (Ema1)
366 displayed the insertion of a CAAACA sequence immediately before the splicing acceptor site
367 of the *C τ 2* exon (Fig. 2). On the other hand, the *C τ 2-C τ 3* intron had identical length in both
368 isoforms.

369 The two introns showed a significant size variation among Antarctic species. The
370 *C τ 1-C τ 2* intron ranged from 212 nt in *N. coriiceps* to 318 nt in *T. bernacchii*, which contained
371 two 46-nt long repeated regions (Table 2, Fig. 2). *N. coriiceps* and *G. gibberifrons* displayed
372 an insertion of additional 15 nt at the 3' end of the intron, which was paired with the presence
373 of the shortest *C τ 2* exon out of all the species analyzed (24 nt, 8 aa). The size and structure of
374 *C τ 2-C τ 3* intron was much more conserved across Antarctic species, ranging from 115 nt in *G.*
375 *gibberifrons* to 128 nt in *H. velifer*. The short length of the *G. gibberifrons* intron was due to

376 a 9-nt long deletion, which matched the position of a 7-nt indel evidenced by the multiple
377 sequence alignment in *E. maclovinus* (Fig. 2).

378

379 *3.3 The genomic analysis of the IgT gene from non-Antarctic species sheds light on the*
380 *stepwise process that led to C τ 2 exon loss*

381

382 To elucidate the processes by which the *C τ 2* exon was progressively lost along the
383 evolution of Notothenioidei, we extended our analyses to the *IgT*-encoding genomic
384 sequences of three additional key non-Antarctic notothenioid species (i.e., *B. diacanthus*, *P.*
385 *urvillii*, *C. gobio*, see the Materials and Methods section).

386 The alignment of the partial genomic sequences (Fig. 3 and Fig. S3) hinted that the
387 evolutionary process that led to the partial loss of the *C τ 2* exon might have already started
388 before the split between the Eleginopsioidea and Cryonotothenioidea lineages, as early as in
389 the late Cretaceous. Indeed, the gene of *P. urvillii* (Pseudaphritioidea) was characterized by
390 early molecular signatures of erosion shared with the Antarctic species or in their sister taxa
391 *E. maclovinus*. In particular, a few informative small deletions, matching the position of
392 similar gaps in the sequence of *E. maclovinus*, were found in the *C τ 1-C τ 2* intron, but not in
393 the *C τ 2-C τ 3* intron, which only included a few indels shared by all species, regardless of their
394 position in the phylogeny of Notothenioidei (Fig. 3). Unlike *E. maclovinus*, *P. urvillii*
395 retained a complete *C τ 2* exon, which only lacked the first three nt at its 5' end.

396 The sequences of the two species belonging to Bovichtidae, the most early-branch of the
397 Notothenioidei lineage, also retained a complete *C τ 2* exon (e.g., 282 nt – 94 aa in *C. gobio*
398 and 285 nt – 95 aa in *B. diacanthus*, see Table 1) and displayed much larger introns than *E.*
399 *maclovinus* (Table 2).

400 However, the *C τ 1-C τ 2* intron of *C. gobio* was also characterized by the presence of a few
401 indels in similar positions to those observed in *P. urvillii* and *E. maclovinus* (Fig. 3).

402 The multiple sequence alignment of the *C τ 2* exon of non-Antarctic Notothenioidei
403 interestingly revealed that, despite the nearly complete deletion of the *C τ 2* exon, *E.*
404 *maclovinus* retained the six nucleotides found at the 3' end of its remnant nearly intact (with
405 the single synonymous substitution TCG/TCT, see position 537 in Fig. 3). This observation
406 is in line with the high conservation of the canonical donor and acceptor splicing sites of the
407 remnant *C τ 2* exon observed in all species (Fig. 3).

408

409 *3.4 The absence of repeats in the introns of the IgT gene rules out the involvement of*
410 *transposable elements in the C τ 2 exon loss*

411

412 As a further step in the investigation of the molecular mechanisms that might have led
413 to the shortening of the *C τ 2* exon, we investigated whether any repeated element could be
414 identified in the *C τ 1-C τ 2* and *C τ 2-C τ 3* introns of the *IgT* gene from Antarctic notothenioids
415 and in their sister lineage Eleginopsioidea. The activity of transposable elements (TEs) is
416 well known to be associated with accelerated mutation, disrupting exons [54], altering
417 splicing patterns [55], and shuffling the position of entire exons or of their parts by moving
418 them to different genomic locations [56]. Hence, the presence of repeats could be indicative

419 of the presence of active TEs, which may have possibly accelerated the evolution of the $C\tau 2$
420 region. Our analyses revealed that neither the $C\tau 1$ - $C\tau 2$ nor the $C\tau 2$ - $C\tau 3$ introns contained
421 traces of repeats in any of the species analyzed in this study.

422

423 3.5 The IgT gene sequence phylogeny is consistent with the phylogenetic relationships 424 among Notothenioidei

425

426 In line with the observations provided above, we found that the molecular evolution of
427 the $C\tau 1$ - $C\tau 2$ and $C\tau 2$ - $C\tau 3$ introns closely followed the currently accepted phylogenetic
428 relationship among Notothenioidei. In detail, all the sequences from Cryonotothenioidea
429 were placed in a highly supported monophyletic clade (100% posterior probability, Fig. 4).
430 As the sequence divergence between the different Antarctic species, which have been subject
431 to a fast evolutionary radiation [57], was minimal, the topology of the Cryonotothenioidea
432 subtree was characterized by very short branches. However, the two variants found in *G.*
433 *gibberifrons* and *H. velifer* were closely related (100% posterior probability), suggesting that
434 both have been originated by species-specific gene duplications (the most likely hypothesis
435 in *G. gibberifrons*, since one of the two variants is pseudogenic) or that they represent allelic
436 variants.

437 As expected, the sequences from the Bovichtidae *C. gobio* and *B. diacanthus* were
438 placed as outgroups in a monophyletic clade at the base of the notothenioid IgT tree, whereas
439 the sequences from *P. urvillii* and *E. maclovinus* occupied, with high statistical support
440 (100% posterior probability in both cases), intermediate positions. These were well
441 consistent with the recently proposed position of Pseudaphritioidea and Eleginopsioidea
442 [27]. Curiously, the two variants from *E. maclovinus* shared closer homology with the
443 Antarctic species than with *P. urvillii* and Bovichtidae, confirming the high relevance of this
444 key species for the investigation of IgT evolution in Notothenioidei and further supporting
445 the observation of shared indels, which may indicate a process of progressive loss of the
446 $C\tau 1$ - $C\tau 2$ intron.

447 Overall, the structure of this intron can be summarized by the presence of 5 distinct
448 conserved sequence motifs (Fig. 5). Starting from the 5' end, the first 15 nt-long motif
449 (named motif 5 in Fig. 5) was most likely involved in the recognition of the $C\tau 1$ donor
450 splicing site. This was followed by the 42 nt-long motif 2, which was the least conserved
451 among those identified and ancestrally duplicated at the 3' end of the intron, where it may be
452 involved in the $C\tau 2$ splicing acceptor site. The subsequent 29 nt-long motif 3 was found in all
453 species, except *B. diacanthus* and *N. coriiceps*, whereas the highly conserved motif 1 (50
454 nt-long) was found in all species and corresponds to the previously mentioned duplicated
455 region found in *T. bernacchii* (Fig. 5). The last motif identified was the highly conserved 29
456 nt-long motif 4, which preceded the second repeat of motif 2. Curiously, the long $C\tau 1$ - $C\tau 2$
457 intron of *T. bernacchii* was characterized by the presence of an additional copy of motif 1 and
458 motif 2 at its 3' end.

459 The analysis of the nucleotide composition of the $C\tau 1$ - $C\tau 2$ intron did not reveal any
460 significant bias in Antarctic species (Fig. 6 and Fig. S6), but at the same time it revealed an
461 interesting trend in the $C\tau 2$ - $C\tau 3$ intron. Indeed, in line with the placement of *E. maclovinus* as

462 a sister group Cryonotothenioidea [30], the intron of this species had an AT content similar to
463 the Antarctic species, which all showed a similar AT content (>70%) regardless of their
464 length, and significantly higher than the other three non-Antarctic species (see Fig. 6).

465

466 3.6 Amino acid multiple sequence alignment sums up the main features of the notothenioid 467 IgT CH domains

468

469 Consistent with the previously reported analysis of the IgT cDNA and genomic
470 sequences (Fig. 1 and 2, Table 1), the multiple sequence alignment of the deduced amino acid
471 sequences highlighted the high conservation of the first, third and fourth CH domains (Fig.
472 7). On the contrary, the size of the CH2 domain varied significantly from species to species,
473 ranging from 8 to 95 aa, depending on its partial deletion (in all Antarctic species plus *E.*
474 *maclovinus*) or full retention (in the most basal notothenioid lineages) (Table 1).

475 All the complete constant domains were characterized by the presence of two
476 constitutive cysteines and one tryptophan, labeled as Cys 23, Cys 104, and Trp 41,
477 according to the IMGT unique numbering (<http://www.imgt.org>). These residues are
478 required to allow the correct folding of immunoglobulin domains, together with the
479 hydrophobic residue (in this case a valine) typically found at the conserved position 89. The
480 remnant portion of the CH2 domain just maintained the second of the two aforementioned
481 canonical cysteines, whereas an additional conserved cysteine residue, known to be involved
482 in the formation of an interchain disulfide bridge with the IgL chain, was found in CH1.
483 While the presence of other cysteine residues in addition to those reported above is quite
484 uncommon in temperate species, we observed several such cases in both Antarctic and
485 non-Antarctic Notothenioidei (Fig. 7). In detail, one extra cysteine was found in the
486 N-terminal end of the CH3 domain of one of the two isoforms of *H. velifer*, precisely
487 between the two APV repeats, and a second one was found at the C-terminal end of CH3 of
488 *G. acuticeps*. The two non-Antarctic species *B. diacanthus* and *E. maclovinus* also showed
489 extra cysteines, in CH2 and in the extracellular membrane proximal region, respectively.

490 We investigated whether the IgT sequences of Antarctic species were associated with the
491 presence of conserved motifs that could be considered as possible “cold hallmarks”, due to
492 their absence in non-Antarctic Notothenioidei and temperate Perciformes, combined with
493 signatures of purifying selection. Although a few short motifs conserved in Antarctic species
494 and progressively fading out in their sister lineages were found (e.g., VV[NK]V in CH1,
495 [IF]DCVI[SR] and TSTMXL[ST] in CH3) (Fig. 7), none of these contained sites subject to
496 purifying selection (Fig. 8).

497 We could detect, however, a few sites subject to significant purifying selection in the
498 two exons flanking the *Ct2* exonic remnant. In detail, four and one negatively selected sites
499 were detected in the *Ct1* and *Ct3* exons, respectively. The former exon included codons
500 encoding highly conserved Leu, Asn and Val residues (plus a Glu/Val residue found at the
501 N-terminus of the domain, but just observed in three sequences), whereas the latter included
502 a single codon encoding a Phe/Ile residue (Fig. 8). Interestingly, in spite of its short length,
503 two of the five positively selected sites detected by our analysis were located within CH2.
504 The first of such sites was placed in a region subject to indel in a few species (e.g., *G.*

505 *gibberifrons* and *N. coriiceps*), close to the N-terminal end of the CH2 remnant. The second
506 hypervariable site was found buried at the center of the domain remnant, adjacent to the
507 aforementioned conserved cysteine. The three other sites evolving under positive selection
508 were found at >20 residues of distance from the boundaries of the CH2 remnant. In detail,
509 the single positively selected codon found in CH1 (encoding either Pro, Thr or Lys) was
510 found near the negatively selected Val-encoding codon mentioned above (Fig. 8). The two
511 hypervariable sites of CH3 were close to each other and located approximately at the center
512 of the domain (Fig. 8). We evaluated the amino acid composition of the CH2 domain in both
513 Antarctic species and non-Antarctic species, comparing it with temperate teleosts. The most
514 represented amino acid residues were Ala and Lys in all species, with Val, Thr, Ser and Leu
515 being more abundant in temperate species and Ala and Pro being more frequently found in
516 Antarctic species.

517 Since the attached glycans can influence solubility, structural stability, and biological
518 function of Ig molecules, we assessed whether Asn-X-Ser/Thr sequons, i.e. possible
519 glycosylation sites, were present in IgT sequences. Antarctic species showed a high number
520 of sequons (Fig. 7 and Fig. S4), potentially subject to a higher degree of glycosylation (65%
521 in all Cryonotothenioidea, 100% in the family Nototheniidae) compared with both
522 non-Antarctic (40%) and temperate species (47%), suggesting a significant role of
523 glycosylation for cold-adapted proteins. Unlike temperate and non-Antarctic species, where
524 sequons are evenly distributed along CH1 and CH3, in Cryonotothenioidea most
525 glycosylation sites were found in CH3 (Fig. S5).

526 Although *N. coriiceps* and *G. gibberifrons* presented an Asn-Pro-Ser sequon in the CH2
527 remnant (also found in the CH4 domain of Artedidraconidae and Bathydraconidae), this is
528 unlikely to be a real glycosylation site due to the proximity between a Pro and an Asn residue,
529 which is expected to make the Asn residue inaccessible [59].

530

531 3.7 *In silico* evaluation of IgT gene expression in multiple tissues, compared with IgM

532

533 To provide some indications concerning the functional role of IgT in notothenioids, we
534 evaluated their gene expression levels in multiple tissues of the Antarctic species *T.*
535 *bernacchii* and the non-Antarctic species *E. maclovinus*, in comparison with a better
536 characterized Ig isotype, IgM (Fig. 9). Although the *in silico* approach we used has some
537 limitations (i.e., the available data regarded a single specimen and could therefore be affected
538 by inter-individual variation), the results presented here can be considered as a preliminary
539 indication of the tissue specificity of the *IgT* gene and its biological relevance compared with
540 IgM.

541 In general, IgT and IgM displayed a similar tissue specificity in both species,
542 highlighting the head kidney, gills and spleen as the most relevant sites of expression,
543 followed by the intestine. In *E. maclovinus*, the expression of IgM was generally much higher
544 than IgT (e.g., >100 folds higher in gills and head kidney, ~70 folds higher in spleen), but this
545 difference was much smaller in *T. bernacchii* (e.g., ~2.5 folds higher in gills, ~5 folds higher
546 in the head kidney and ~20 folds higher in spleen). Hence, the direct comparison between *T.*
547 *bernacchii* and *E. maclovinus* suggests that the *IgT* gene is expressed at higher levels in the

548 Antarctic species.

549

550 4. Discussion

551

552 The immunoglobulins of Antarctic fishes have been fascinating us since the early
553 discovery of unforeseen features of IgM from Cryonotothenioidea [60-62]. For several years,
554 our studies have been mostly focused on IgM, an ancient Ig isotype that first appeared in
555 jawed fish along with the emergence of an adaptive immune system [63]. However, we
556 recently moved our attention to the study of the heavy chain gene of IgT, a fish-specific Ig
557 isotype, whose discovery revealed the origins of the most ancient Ig specialized in mucosal
558 immune response. We disclosed that the gene of Antarctic species *T. bernacchii*, unlike the
559 early-branching non-Antarctic notothenioid species *B. diacanthus* and most species living in
560 temperate environments, displayed an unusual truncated $C\tau 2$ exon, which only encoded a
561 short remnant of the CH2 domain [19]. This finding was the starting point of the present
562 work, which extends our molecular investigations to other Antarctic species and to the
563 early-branching non-Antarctic notothenioid lineages, in the attempt to pinpoint the origins of
564 this partial exon loss event.

565 The groundwork for placing down the first piece of this evolutionary puzzle was
566 provided by the observation that the $C\tau 2$ exon was nearly completely missing also in *E.*
567 *maclovinus*, a non-Antarctic species which belongs to Eleginopsioidea, the sister group of
568 Cryonotothenioidea. This allowed us to move backwards through the phylogeny of
569 Notothenioidei, characterizing the IgT sequence of *C. gobio*, belonging to the most basal
570 group of non-Antarctic Notothenioidei, i.e., Bovichtidae. This species showed a nearly
571 complete $C\tau 2$ exon, except for a small deletion located at its 3' end of the exon, which
572 matched a similar indel carried by *E. maclovinus*. Moreover, *C. gobio* also displayed a
573 shorter $C\tau 1$ - $C\tau 2$ intron than its close relative *B. diacanthus*, with a few deletions shared with
574 *E. maclovinus*.

575 The cornerstone of the puzzle was provided by the analysis of the IgT sequence of *P.*
576 *urvillii*, belonging to the monotypic family Pseudaphritioidea, which covers an intermediate
577 position in notothenioid evolution, between Bovichtidae and the Eleginopsioidea +
578 Cryonotothenioidea lineage. The IgT heavy chain gene of this species showed evident
579 signatures of the evolutionary process that led to the $C\tau 2$ exon loss in Cryonotothenioidea,
580 which included a small deletion of three nucleotides at the 5' end of the $C\tau 2$ exon and several
581 short deletions within the $C\tau 1$ - $C\tau 2$ intron, shared with *E. maclovinus*.

582 Based on these observations, we propose a timeline for the loss of the $C\tau 2$ exon, which
583 might have occurred after the divergence of Pseudaphritioidea from the main notothenioid
584 lineage, but before the split between Eleginopsioidea and Cryonotothenioidea. This temporal
585 placement would be consistent with the recently revised phylogeny of Notothenioidei [27].
586 Several observations suggest that this process may have been initiated by the modifications
587 of the two introns flanking the $C\tau 2$ exon. First, intron length followed a clear trend towards
588 reduction in the Antarctic species, consistent in particular with the progressive loss of the
589 upstream $C\tau 1$ - $C\tau 2$ intron. Second, the phylogenetic analysis, the organization of the motifs
590 and shared indels found in each of the two introns neighboring $C\tau 2$ were fully consistent with

591 notothenioid phylogeny and marked a close similarity between the introns of *E. maclovinus*
592 and those of Antarctic species, with a progressive disappearance of the Antarctic features in
593 the most early-branching notothenioid species. Third, a significant bias towards a rich AT
594 content in the *C τ 2-C τ 3* intron was observed in Antarctic species and their sister taxa *E.*
595 *maclovinus*, but not in the early-branching non-Antarctic lineages. The latter observation
596 raises the interesting question of whether the high AT content may be interpreted as an
597 adaptive feature to improve replication, transcription or other molecular processes in a
598 cooling environment. AT-rich introns are a well-known peculiar feature of most teleost
599 genomes, which may be viewed as an ancestral characteristic of ectothermic vertebrates [64].
600 An integrated revision of the genomes readily available at present for many notothenioid
601 species may provide a useful framework for assessing this issue. These observations are in
602 line with key role that introns cover in the dynamic process of genome evolution [65]. The
603 role of introns in the adaptation to varying environmental pressure may have been
604 particularly relevant in teleosts, where these elements are found in higher number and
605 usually have a shorter length than other vertebrates, as a consequence of the teleost-specific
606 whole-genome duplication event [66]. We have previously revealed, as a key example of
607 this potential for adaptation, the peculiar rearrangement of the exon/intron architecture of
608 the region encoding the C-terminal Extracellular Membrane-Proximal Domain in the IgM
609 heavy chain gene locus of Antarctic fish [67].

610 Although Transposable Elements (TEs), found in 20–60% of introns in vertebrate
611 genomes, can also provide a significant contribution to the modification of gene
612 architecture [68], we could not find any active TEs in the two introns flanking the *C τ 2* exon,
613 neither in Antarctic, nor in non-Antarctic species. However, we cannot exclude the
614 possibility that TEs that were lost along evolution and that are not detectable anymore have
615 played a role in the process that led to the loss of the *C τ 2* exon.

616 Most certainly, the reconstruction of this scenario suffers from missing data and
617 significant “evolutionary gaps” due to the high phylogenetic distance between the genera we
618 took into account [27]. In any case, the *IgT* sequence features outlined above for
619 non-Antarctic species are “molecular fossils” that can provide useful information to infer the
620 stepwise *C τ 2* exon loss that occurred in the Antarctic lineage.

621 The evolutionary dynamics of exon–intron architecture is another significant aspect of
622 genome adaptation [69], as they accommodate a high variation of intron and exon size but
623 require an extreme conservation of the splicing site motifs located at their boundaries. We
624 show that the *C τ 2* exon remnant of all Antarctic species, despite its short length, retains both
625 functional acceptor (5') and donor (3') splice sites. While the conservation of the donor splice
626 site can be explained by the preservation of the six nucleotides at the 3' end of this exon, the
627 entire 5' end of *C τ 2* was deleted in Antarctic species, along with the acceptor splicing site. We
628 speculate that the correct splicing might still occur in this region thanks to the presence of an
629 AGGCAA motif, which nearly matches one of the five predicted splicing regulatory
630 hexamers of mammals [69].

631 The recurrent reorganization of *Ig* gene loci during vertebrate evolution has often led to
632 the generation of multiple functional gene copies and pseudogenes [70]. A similar
633 evolutionary process might have targeted Antarctic *IgT* genes, thereby generating novel

634 sequence variants that could have granted the acquisition of new structural arrangements and
635 functions, favorable for the adaptation to the polar environment, as in the previously reported
636 case of *T. bernacchii*.

637 Most of the variants reported in this work had a length similar to the variant S of *T.*
638 *bernacchii* (24-39 nt - 8-13 aa, vs 33 nt - 11aa). However, due to the limited number of
639 specimens available for each species, we cannot exclude that other Notothenioid species
640 share a similar repertoire of transcript variants. Interestingly, we could not identify any
641 variant characterized, like the Sts isoform of *T. bernacchii*, by the skipping of the *C τ 2* exon
642 remnant. Therefore, Sts may be considered a species-specific isoform, possibly linked with
643 the significant genomic reorganization of Trematominae [71]. Most likely, the availability of
644 additional fully sequenced Notothenioid genomes will clarify the orthology and paralogy
645 relationships of the different sequence variants identified in this work.

646 A question that remains to be solved is why Antarctic species retained a very short
647 remnant of the IgT CH2 domain, instead of completely losing it. A possible explanation
648 might reside in the structural utility of this region as a linker, with the function to keep the
649 first and second CH domains apart from each other, and to provide the Ig molecule with
650 greater flexibility of the Fab arms, as suggested by (i) its high solvent exposure predicted by
651 3D molecular modeling (Fig. 10); (ii) the abundance of amino acid residues typically found
652 in hinge regions (i.e. proline, glycine and cysteine), recalling in particular the human IgA1
653 hinge region, which is also of similar length. The finding that the *C τ 2* exon remnant, despite
654 its short length, contained two sites subject to diversifying selection was intriguing. While
655 the first one was found in a region characterized by indels in some species, the second one,
656 encoding Pro/Ser/Thr, was adjacent to a highly conserved cysteine residue. This
657 observation will undoubtedly lead to further investigations aimed at clarifying the
658 functional role of these residues in the context of cold adaptation, which presently remains
659 unknown.

660 Since the IgT heavy chain does not possess the additional cysteine residue engaged in the
661 formation of the covalent bond between the two heavy chains in the monomer, the presence
662 of a conserved cysteine residue in all Antarctic species is another peculiarity of the CH2
663 remnant. Cysteine pairs involved in the formation of disulfide bonds are highly conserved
664 relative to unpaired cysteines and to other amino acids [72]. Consequently, whenever
665 disulfide bonds are permanently consolidated in proteins, the cysteine residues involved in
666 the formation of such bonds rarely vary. Since the remnant portion of the CH2 domain just
667 maintained the second of the two canonical cysteines, we speculate that the high
668 conservation of this residue might be linked with a function in: (i) bridging the two heavy
669 chains in the monomer; (ii) keeping the monomer units covalently linked in the multimeric
670 form, creating the so-called “redox forms” observed for other Ig isotypes. In general,
671 variations in disulfide connectivity allow a higher degree of polymerization, influencing the
672 structure, stability and effector functions of Ig molecules, as it typically happens in the case
673 of IgM [73]. Although little information is available concerning the aggregation status of fish
674 IgT, biochemical studies carried out in the rainbow trout have revealed that they are present
675 in the serum as a monomer and that, unlike IgM, the multimeric complexes found in the
676 mucus are non-covalently linked [73]. Considering that teleosts lack the J chain, essential for

677 the formation of Ig polymers, the principles that drive their association in multimers are
678 presently unknown.

679 Teleost mucosal Igs can associate with the polymeric Ig receptor (pIgR), which is also
680 present in *T. bernacchii* (unpublished data) and possibly in other Antarctic fish, and enables
681 their secretion into the gut lumen, similar to mammalian IgA and IgM [74, 75]. Antarctic IgT
682 contain, at the carboxy terminus of the secretory tail, a sequence motif similar to the one
683 found in other teleosts, which may be involved in polymerization.

684 A third factor that might allow IgT polymerization relies on glycosylation, even though
685 the role of glycan moieties in teleost mucosa-associated Igs still needs to be clarified. The
686 IgT of Antarctic species contain a high number of sequons consistent with glycosylation
687 sites. In particular, the presence of two such sites in the C-terminal secretory tail of the
688 Antarctic IgT may suggest a role in the assembly of Ig complexes, as they match the
689 position of conserved glycosylation sites found in mammalian IgM and involved in the
690 polymerization of this Ig isotype [76].

691 In light of these observations, it is tempting to speculate that the remnant CH2 domain
692 may provide an advantage for the activity of this Ig isotype under the thermodynamically
693 unfavorable cold conditions of the Antarctica. Although the prevalent role of notothenioid
694 IgT in mucosal immunity needs to be verified, the transcriptomic analysis we performed
695 indicates that, similar to IgM, IgT is mostly expressed in head kidney, gills and spleen, both
696 in Antarctic and non-Antarctic notothenioids. On the other hand, IgT was only found to be
697 expressed at moderate levels in the intestine, and at nearly non-detectable levels in skin.
698 Interestingly, while IgT are expressed at lower levels than IgM, they appear to be expressed
699 at levels significantly higher in *T. bernacchii* than *E. maclovinus*. The ratios IgT/IgM we
700 found reversed in the same tissues (gills and head kidney) from the Antarctic and
701 non-Antarctic representatives suggest a remarkable role of IgT in the adaptive immune
702 response under subzero conditions, despite almost complete devoid of a CH domain. These
703 observations, hence, provide some preliminary clues about the functional role of IgT in a
704 cold adapted species. However, additional data are still needed to further investigate this
705 issue. Among others, for a better comprehension, it will be worthy to take into account the
706 presence of the different parasite fauna documented in non-Antarctic species so far (Torres
707 and Soto, 2004; Brickle and MacKenzie, 2007) [77, 78].

708 We have reported in previous studies that Antarctic fish show heavy parasitic loads and
709 mount a specific plasma and bile antibody response against a wide spectrum of nematode
710 antigens [79, 80]. However, by that time, the IgT isotype had not yet been discovered. Hence,
711 it will be intriguing to assess the presence of IgT specific for naturally occurring nematode
712 parasites in Antarctic fish. Since a specialization of IgT in controlling pathogens at mucosal
713 surfaces and microbiota homeostasis has been recently uncovered in rainbow trout [81-83], it
714 will be challenging to explore, in this respect, the role played by IgT in cold adapted teleosts,
715 presently unknown.

716

717 **5. Conclusions**

718 The many challenges notothenioid fishes faced during their evolutionary history
719 triggered a wave of genomic changes that led to a number of peculiar features that might
720 have been preserved as adaptive traits. The development of an IgT molecule nearly
721 completely devoid of a CH domain might be just another addition to the list of the highly
722 successful strategies these organisms have used to adapt to a very challenging environment.
723 The evolutionary scenario we propose involves a gradual process, which has shaped the *IgH*
724 gene locus over a long period of time, through the action of contrasting evolutionary forces,
725 contributing to the generation of an Ig molecule with a unique architecture among
726 vertebrates.

727

728 **Acknowledgments**

729 The authors wish to thank Dr Ennio Cocca (IBBR, CNR, Naples, Italy), for providing
730 tissue samples of *E. maclovinus*, *C. rastrospinosus*, and *C. aceratus* specimens; Drs Daniela
731 Giordano, and Cinzia Verde (IBBR, CNR, Naples, Italy) for providing *P. urvillii* and *G.*
732 *gibberifrons* blood samples. The authors are grateful to Prof. Alberto Pallavicini (University
733 of Trieste, Italy) for his critical comments on the manuscript.

734 In memory of Guido di Prisco, for his scientific career entirely devoted to unveil the
735 mystery of evolutionary adaptations for life in Antarctica.

736

737 **Funding:** This research was funded by the Italian National Program for Research in
738 Antarctica [grant number PNRA 16_00099–A1 Project].

739

740

741 **CRedit authorship contribution statement**

742 **Alessia Ametrano:** Conceptualization, Investigation, Formal analysis, Validation,
743 Visualization, Writing - Review & Editing. **Marco Gerdol:** Conceptualization,
744 Investigation, Formal analysis, Funding acquisition, Resources, Writing - Original Draft,
745 Writing - Review & Editing Visualization. **Maria Vitale:** Investigation, Formal analysis,
746 Writing - Review & Editing. **Samuele Greco:** Investigation, Formal analysis, Writing -
747 Review & Editing. **Umberto Oreste:** Visualization, Writing - Review & Editing. **Maria**
748 **Rosaria Coscia:** Conceptualization, Writing - Original Draft, Writing - Review & Editing,
749 Supervision, Project administration, Funding acquisition.

750 All authors have read and agreed to the published version of the manuscript.

751

752

753 **Declaration of competing interest**

754 Authors declare no conflict of interest. The funders had no role in the design of the study;
755 in the collection, analyses, or interpretation of data; in the writing of the manuscript, or in the
756 decision to publish the results.

757

758

759 **References**

- 760 1. Salinas, I.; Zhang, Y.-A.; Sunyer, J. O. Mucosal Immunoglobulins and B Cells of
761 Teleost Fish. *Dev Comp Immunol* **2011**, 35, 1346–1365,
762 doi:10.1016/j.dci.2011.11.009.
- 763
- 764 2. Chen, K.; Cerutti, A. New Insights into the Enigma of Immunoglobulin D.
765 *Immunological Reviews* **2010**, 237, 160–179,
766 doi:10.1111/j.1600-065X.2010.00929.x.
- 767
- 768 3. Perdiguero, P.; Martín-Martín, A.; Benedicenti, O.; Díaz-Rosales, P.; Morel, E.;
769 Muñoz-Atienza, E.; García-Flores, M.; Simón, R.; Soletto, I.; Cerutti, A. et al.
770 Teleost IgD+IgM⁻ B Cells Mount Clonally Expanded and Mildly Mutated
771 Intestinal IgD Responses in the Absence of Lymphoid Follicles. *Cell Reports* **2019**,
772 29, 4223–4235.e5, doi: 10.1016/j.celrep.2019.11.101.
- 773
- 774 4. Danilova, N.; Bussmann, J.; Jekosch, K.; Steiner, L. A. The Immunoglobulin
775 Heavy-Chain Locus in Zebrafish: Identification and Expression of a Previously
776 Unknown Isotype, Immunoglobulin Z. *Nat Immunol* **2005**, 6, 295–302,
777 doi:10.1038/ni1166.
- 778
- 779 5. Hansen, J. D.; Landis, E. D.; Phillips, R. B. Discovery of a Unique Ig Heavy-Chain
780 Isotype (IgT) in Rainbow Trout: Implications for a Distinctive B Cell
781 Developmental Pathway in Teleost Fish. *Proc Natl Acad Sci U S A* **2005**, 102, 6919–
782 6924, doi:10.1073/pnas.0500027102.
- 783
- 784 6. Zhang, Y.-A.; Salinas, I.; Li, J.; Parra, D.; Bjork, S.; Xu, Z.; LaPatra, S. E.;
785 Bartholomew, J.; Sunyer, J. O. IgT, a Primitive Immunoglobulin Class Specialized
786 in Mucosal Immunity. *Nat Immunol* **2010**, 11, 827–835, doi:10.1038/ni.1913.
- 787
- 788 7. Xu, Z.; Parra, D.; Gómez, D.; Salinas, I.; Zhang, Y.-A.; von Gersdorff Jørgensen,
789 L.; Heinecke, R. D.; Buchmann, K.; LaPatra, S.; Sunyer, J. O. Teleost Skin, an
790 Ancient Mucosal Surface That Elicits Gut-like Immune Responses. *Proc Natl Acad*
791 *Sci U S A* **2013**, 110, 13097–13102, doi:10.1073/pnas.1304319110.
- 792
- 793 8. Tacchi, L.; Musharrafieh, R.; Larragoite, E. T.; Crossey, K.; Erhardt, E. B.; Martin,
794 S. A. M.; LaPatra, S. E.; Salinas, I. Nasal Immunity Is an Ancient Arm of the
795 Mucosal Immune System of Vertebrates. *Nat Commun* **2014**, 5, 5205,
796 doi:10.1038/ncomms6205.

797

798

799

800

801

802

803

804

805

806

807

808

809

810

811

812

813

814

815

816

817

818

819

820

821

822

823

824

825

826

827

828

829

830

831

832

833

834

835

836

837

9. Xu, Z.; Takizawa, F.; Parra, D.; Gómez, D.; von Gersdorff Jørgensen, L.; LaPatra, S. E.; Sunyer, J. O. Mucosal Immunoglobulins at Respiratory Surfaces Mark an Ancient Association That Predates the Emergence of Tetrapods. *Nat Commun* **2016**, *7*, 10728, doi:10.1038/ncomms10728.
10. Yu, Y.-Y.; Kong, W.-G.; Xu, H.-Y.; Huang, Z.-Y.; Zhang, X.-T.; Ding, L.-G.; Dong, S.; Yin, G.-M.; Dong, F.; Yu, W.; et al. Convergent Evolution of Mucosal Immune Responses at the Buccal Cavity of Teleost Fish. *iScience* **2019**, *19*, 821–835, doi:10.1016/j.isci.2019.08.034.
11. Yu, Y.-Y.; Kong, W.; Yin, Y.-X.; Dong, F.; Huang, Z.-Y.; Yin, G.-M.; Dong, S.; Salinas, I.; Zhang, Y.-A.; Xu, Z. Mucosal Immunoglobulins Protect the Olfactory Organ of Teleost Fish against Parasitic Infection. *PLoS Pathog* **2018**, *14*, e1007251, doi:10.1371/journal.ppat.1007251.
12. Zhang, N.; Zhang, X.-J.; Chen, D.-D.; Sunyer, J. O.; Zhang, Y.-A. Molecular Characterization and Expression Analysis of Three Subclasses of IgT in Rainbow Trout (*Oncorhynchus Mykiss*). *Dev Comp Immunol* **2017**, *70*, 94–105, doi:10.1016/j.dci.2017.01.001.
13. Buonocore, F.; Stocchi, V.; Nunez-Ortiz, N.; Randelli, E.; Gerdol, M.; Pallavicini, A.; Facchiano, A.; Bernini, C.; Guerra, L.; Scapigliati, G.; Picchietti, S. Immunoglobulin T from Sea Bass (*Dicentrarchus Labrax* L.): Molecular Characterization, Tissue Localization and Expression after Nodavirus Infection. *BMC Mol Biol* **2017**, *18*, 8, doi:10.1186/s12867-017-0085-0.
14. Fillatreau, S.; Six, A.; Magadan, S.; Castro, R.; Sunyer, J. O.; Boudinot, P. The Astonishing Diversity of Ig Classes and B Cell Repertoires in Teleost Fish. *Front Immunol* **2013**, *4*, doi:10.3389/fimmu.2013.00028.
15. Savan, R.; Aman, A.; Sato, K.; Yamaguchi, R.; Sakai, M. Discovery of a New Class of Immunoglobulin Heavy Chain from Fugu. *Eur J Immunol* **2005**, *35*, 3320–3331, doi:10.1002/eji.200535248.
16. Velázquez, J.; Acosta, J.; Lugo, J. M.; Reyes, E.; Herrera, F.; González, O.; Morales, A.; Carpio, Y.; Estrada, M. P. Discovery of Immunoglobulin T in Nile Tilapia (*Oreochromis Niloticus*): A Potential Molecular Marker to Understand Mucosal Immunity in This Species. *Dev Comp Immunol* **2018**, *88*, 124–136, doi:10.1016/j.dci.2018.07.013.

- 838
839 17. Savan, R.; Aman, A.; Nakao, M.; Watanuki, H.; Sakai, M. Discovery of a Novel
840 Immunoglobulin Heavy Chain Gene Chimera from Common Carp (*Cyprinus*
841 *Carpio L.*). *Immunogenetics* **2005**, *57*, 458–463, doi:10.1007/s00251-005-0015-z.
842
- 843 18. Ryo, S.; Wijdeven, R. H. M.; Tyagi, A.; Hermsen, T.; Kono, T.; Karunasagar, I.;
844 Rombout, J. H. W. M.; Sakai, M.; Verburg-van Kemenade, B. M. L.; Savan, R.
845 Common Carp Have Two Subclasses of Bonyfish Specific Antibody IgZ Showing
846 Differential Expression in Response to Infection. *Dev Comp Immunol* **2010**, *34*,
847 1183–1190, doi:10.1016/j.dci.2010.06.012.
848
- 849 19. Giacomelli, S.; Buonocore, F.; Albanese, F.; Scapigliati, G.; Gerdol, M.; Oreste, U.;
850 Coscia, M. R. New Insights into Evolution of IgT Genes Coming from Antarctic
851 Teleosts. *Mar Genomics* **2015**, *24*, 55–68, doi:10.1016/j.margen.2015.06.009.
852
- 853 20. Eastman, J. T. The Nature of the Diversity of Antarctic Fishes. *Polar Biol* **2005**, *28*,
854 93–107, doi:10.1007/s00300-004-0667-4.
855
- 856 21. Antarctic Fish Biology: Evolution in a Unique Environment. Joseph T. Eastman.
857 1993. San Diego: Academic Press, Xiii + 322 p, Illustrated, Hard Cover. ISBN
858 0-12-228140-3. *Polar Record* 1994, *30*, 59–60, doi:10.1017/S0032247400021100.
859
- 860 22. DeVries, A. L.; Wohlschlag, D. E. Freezing Resistance in Some Antarctic Fishes.
861 *Science* **1969**, *163*, 1073–1075, doi:10.1126/science.163.3871.1073
862
- 863
- 864 23. Hofmann, G. E.; Buckley, B. A.; Airaksinen, S.; Keen, J. E.; Somero, G. N.
865 Heat-Shock Protein Expression Is Absent in the Antarctic Fish *Trematomus*
866 *Bernacchii* (Family Nototheniidae). *J Exp Biol* **2000**, *203*, 2331–2339.
867
- 868 24. Barber, D. L.; Westermann, J. E. M.; White, M. G. The Blood Cells of the Antarctic
869 Icefish *Chaenocephalus Aceratus* Lönnberg: Light and Electron Microscopic
870 Observations. *J Fish Biol* **1981**, *19*, 11–28.
871 doi:10.1111/j.1095-8649.1981.tb05807.x.
872
- 873 25. Near, T. J.; Eytan, R. I.; Dornburg, A.; Kuhn, K. L.; Moore, J. A.; Davis, M. P.;
874 Wainwright, P. C.; Friedman, M.; Smith, W. L. Resolution of Ray-Finned Fish
875 Phylogeny and Timing of Diversification. *Proc Natl Acad Sci U S A* **2012**, *109*,
876 13698–13703, doi:10.1073/pnas.1206625109.
877

- 878 26. Ceballos, S. G.; Lessa, E. P.; Victorio, M. F.; Fernández, D. A. Phylogeography of
879 the Sub-Antarctic Notothenioid Fish *Eleginops Maclovinus*: Evidence of Population
880 Expansion. *Mar Biol* **2012**, 159, 499–505, doi:10.1007/s00227-011-1830-4.
881
- 882 27. Near, T. J.; Ghezelayagh, A.; Ojeda, F. P.; Dornburg, A. Recent Diversification in
883 an Ancient Lineage of Notothenioid Fishes (*Bovichtus*: Notothenioidei). *Polar Biol*
884 **2019**, 42, 943–952, doi:10.1007/s00300-019-02489-1.
885
- 886 28. Near, T. J.; Dornburg, A.; Harrington, R. C.; Oliveira, C.; Pietsch, T. W.; Thacker,
887 C. E.; Satoh, T. P.; Katayama, E.; Wainwright, P. C.; Eastman, J. T.; et al.
888 Identification of the Notothenioid Sister Lineage Illuminates the Biogeographic
889 History of an Antarctic Adaptive Radiation. *BMC Evol Biol* **2015**, 15,
890 doi:10.1186/s12862-015-0362-9.
891
- 892
- 893 29. Shin, S. C.; Kim, S. J.; Lee, J. K.; Ahn, D. H.; Kim, M. G.; Lee, H.; Lee, J.; Kim,
894 B.-K.; Park, H. Transcriptomics and Comparative Analysis of Three Antarctic
895 Notothenioid Fishes. *PLoS One* **2012**, 7, e43762,
896 doi:10.1371/journal.pone.0043762.
897
- 898
- 899 30. Near, T. J.; MacGuigan, D. J.; Parker, E.; Struthers, C. D.; Jones, C. D.; Dornburg,
900 A. Phylogenetic Analysis of Antarctic Notothenioids Illuminates the Utility of
901 RADseq for Resolving Cenozoic Adaptive Radiations. *Mol Phylogenet Evol* **2018**,
902 129, 268–279, doi:10.1016/j.ympev.2018.09.001.
903
- 904 31. Bargelloni, L.; Babbucci, M.; Ferrareso, S.; Papetti, C.; Vitulo, N.; Carraro, R.;
905 Pauletto, M.; Santovito, G.; Lucassen, M.; Mark, F. C.; et al. Draft Genome
906 Assembly and Transcriptome Data of the Icefish *Chionodraco Myersi* Reveal the
907 Key Role of Mitochondria for a Life without Hemoglobin at Subzero Temperatures.
908 *Commun Biol* **2019**, 2, 443, doi:10.1038/s42003-019-0685-y.
909
- 910 32. Kim, B.-M.; Amores, A.; Kang, S.; Ahn, D.-H.; Kim, J.-H.; Kim, I.-C.; Lee, J. H.;
911 Lee, S. G.; Lee, H.; Lee, J.; et al. Antarctic Blackfin Icefish Genome Reveals
912 Adaptations to Extreme Environments. *Nat Ecol Evol* **2019**, 3, 469–478,
913 doi:10.1038/s41559-019-0812-7.
914
- 915
- 916 33. Koepfli, K.-P.; Paten, B.; Genome 10K Community of Scientists; O'Brien, S. J. The
917 Genome 10K Project: A Way Forward. *Annu Rev Anim Biosci* **2015**, 3, 57–111,
918 doi:10.1146/annurev-animal-090414-014900.

- 919
920 34. Altschul, S. F.; Gish, W.; Miller, W.; Myers, E. W.; Lipman, D. J. Basic Local
921 Alignment Search Tool. *J Mol Biol* **1990**, 215, 403–410,
922 doi:10.1016/S0022-2836(05)80360-2.
923
- 924 35. Thompson, J. D.; Higgins, D. G.; Gibson, T. J. CLUSTAL W: Improving the
925 Sensitivity of Progressive Multiple Sequence Alignment through Sequence
926 Weighting, Position-Specific Gap Penalties and Weight Matrix Choice. *Nucleic
927 Acids Res* **1994**, 22, 4673–4680,
928 doi:10.1093/nar/22.22.4673.
929
- 930 36. Gupta, R.; Jung, E.; Brunak, S. Prediction of N-glycosylation sites in human
931 proteins. **2004**
932
- 933 37. Kelley, L. A.; Mezulis, S.; Yates, C. M.; Wass, M. N.; Sternberg, M. J. E. The
934 Phyre2 Web Portal for Protein Modeling, Prediction and Analysis. *Nat Protoc* **2015**,
935 10, 845–858, doi:10.1038/nprot.2015.053.
936
- 937 38. Price, A. L.; Jones, N. C.; Pevzner, P. A. De Novo Identification of Repeat Families
938 in Large Genomes. *Bioinformatics* **2005**, 21 Suppl 1, i351-358,
939 doi:10.1093/bioinformatics/bti1018.
940
- 941 39. Smit, A.F.A.; Hubley, R.; Green, P. **1996**. *RepeatMaster Open-3.0*
942
- 943 40. Hubley, R.; Finn, R. D.; Clements, J.; Eddy, S. R.; Jones, T. A.; Bao, W.; Smit, A.
944 F. A.; Wheeler, T. J. The Dfam Database of Repetitive DNA Families. *Nucleic
945 Acids Res* **2016**, 44, D81-89, doi:10.1093/nar/gkv1272.
946
- 947 41. Edgar, R. C. MUSCLE: Multiple Sequence Alignment with High Accuracy and
948 High Throughput. *Nucleic Acids Res* **2004**, 32, 1792–1797, doi:10.1093/nar/gkh340.
949
- 950 42. Talavera, G.; Castresana, J. Improvement of Phylogenies after Removing Divergent
951 and Ambiguously Aligned Blocks from Protein Sequence Alignments. *Syst Biol*
952 **2007**, 56, 564–577, doi:10.1080/10635150701472164.
953
- 954 43. Darriba, D.; Posada, D.; Kozlov, A. M.; Stamatakis, A.; Morel, B.; Flouri, T.
955 ModelTest-NG: A New and Scalable Tool for the Selection of DNA and Protein
956 Evolutionary Models. *Mol Biol Evol* **2020**, 37, 291–294,
957 doi:10.1093/molbev/msz189.
958

- 959 44. Tavaré, S. Some Probabilistic and Statistical Problems in the Analysis of DNA
960 Sequences. Some mathematical questions in biology / DNA sequence analysis
961 edited by Robert M. Miura **1986**.
962
- 963 45. Cavanaugh, J. E. Unifying the Derivations for the Akaike and Corrected Akaike
964 Information Criteria. *Statistics & Probability Letters* **1997**, 33, 201–208,
965 doi:10.1016/S0167-7152(96)00128-9.
966
- 967 46. Huelsenbeck, J. P.; Ronquist, F. MRBAYES: Bayesian Inference of Phylogenetic
968 Trees. *Bioinformatics* **2001**, 17, 754–755, doi:10.1093/bioinformatics/17.8.754.
969
- 970 47. Rambaut, A.; Drummond, A. J.; Xie, D.; Baele, G.; Suchard, M. A. Posterior
971 Summarization in Bayesian Phylogenetics Using Tracer 1.7. *Syst Biol* **2018**, 67,
972 901–904, doi:10.1093/sysbio/syy032.
973
- 974 48. Kumar, S.; Stecher, G.; Li, M.; Knyaz, C.; Tamura, K. MEGA X: Molecular
975 Evolutionary Genetics Analysis across Computing Platforms. *Mol Biol Evol* **2018**,
976 35, 1547–1549, doi:10.1093/molbev/msy096.
977
- 978 49. Weaver, S.; Shank, S. D.; Spielman, S. J.; Li, M.; Muse, S. V.; Kosakovsky Pond,
979 S. L. Datamonkey 2.0: A Modern Web Application for Characterizing Selective and
980 Other Evolutionary Processes. *Mol Biol Evol* **2018**, 35, 773–777,
981 doi:10.1093/molbev/msx335.
982
- 983 50. Kosakovsky Pond, S. L.; Frost, S. D. W. Not So Different After All: A Comparison
984 of Methods for Detecting Amino Acid Sites Under Selection. *Mol Biol Evol* **2005**,
985 22, 1208–1222, doi:10.1093/molbev/msi105.
986
- 987 51. Murrell, B.; Wertheim, J. O.; Moola, S.; Weighill, T.; Scheffler, K.; Pond, S. L. K.
988 Detecting Individual Sites Subject to Episodic Diversifying Selection. *PLOS Genet*
989 **2012**, 8, e1002764, doi:10.1371/journal.pgen.1002764.
990
- 991 52. Bailey, T. L.; Elkan, C. Fitting a Mixture Model by Expectation Maximization to
992 Discover Motifs in Biopolymers. *Proc Int Conf Intell Syst Mol Biol* **1994**, 2, 28–36.
993
- 994 53. Bailey, T. L.; Boden, M.; Buske, F. A.; Frith, M.; Grant, C. E.; Clementi, L.; Ren,
995 J.; Li, W. W.; Noble, W. S. MEME SUITE: Tools for Motif Discovery and
996 Searching. *Nucleic Acids Res* **2009**, 37 (Web Server issue), W202–208,
997 doi:10.1093/nar/gkp335.
998

- 999 54. Hancks, D. C.; Kazazian, H. H. Roles for Retrotransposon Insertions in Human
1000 Disease. *Mobile DNA* **2016**, 7, 65, doi:10.1186/s13100-016-0065-9.
1001
- 1002 55. Davis, M. B.; Dietz, J.; Standiford, D. M.; Emerson, C. P. Transposable Element
1003 Insertions Respecify Alternative Exon Splicing in Three Drosophila Myosin Heavy
1004 Chain Mutants. *Genetics* **1998**, 150, 1105–1114.
1005
- 1006 56. Ejima, Y.; Yang, L. Trans Mobilization of Genomic DNA as a Mechanism for
1007 Retrotransposon-Mediated Exon Shuffling. *Hum Mol Genet* **2003**, 12, 1321–1328,
1008 doi:10.1093/hmg/ddg138.
1009
- 1010 57. Colombo, M.; Damerau, M.; Hanel, R.; Salzburger, W.; Matschiner, M. Diversity
1011 and Disparity through Time in the Adaptive Radiation of Antarctic Notothenioid
1012 Fishes. *J Evol Biol* **2015**, 28, 376–394, doi:10.1111/jeb.12570.
1013
- 1014 58. Gambòn-Deza, F.; Sánchez-Espinel, C.; Magadàn-Mompò, S. Presence of an
1015 unique IgT on the IGH locus in three-spined stickleback fish (*Gasterosteus*
1016 *aculeatus*) and the very recent generation of a repertoire of VH genes. *Dev Comp*
1017 *Immunol* **2010**, 34, 114–122, doi: 10.1016/j.dci.2009.08.011
1018
- 1019 59. Bause, E. Structural Requirements of N-Glycosylation of Proteins. Studies with
1020 Proline Peptides as Conformational Probes. *Biochem J* **1983**, 209, 331–336.
1021
- 1022 60. Coscia, M. R.; Morea, V.; Tramontano, A.; Oreste, U. Analysis of a CDNA
1023 Sequence Encoding the Immunoglobulin Heavy Chain of the Antarctic Teleost
1024 *Trematomus Bernacchii*. *Fish Shellfish Immunol* **2000**, 10, 343–357,
1025 doi:10.1006/fsim.1999.0244.
1026
- 1027 61. Coscia, M. R.; Varriale, S.; Giacomelli, S.; Oreste, U. Antarctic Teleost
1028 Immunoglobulins: More Extreme, More Interesting. *Fish Shellfish Immunol* **2011**,
1029 31, 688–696, doi:10.1016/j.fsi.2010.10.018.
1030
- 1031 62. Coscia, M. R.; Giacomelli, S.; Oreste, U. Allelic Polymorphism of Immunoglobulin
1032 Heavy Chain Genes in the Antarctic Teleost *Trematomus Bernacchii*. *Mar*
1033 *Genomics* **2012**, 8, 43–48, doi:10.1016/j.margen.2012.04.002.
1034
- 1035 63. Flajnik, M. F. A Cold-Blooded View of Adaptive Immunity. *Nat Rev Immunol*
1036 **2018**, 18, 438–453, doi:10.1038/s41577-018-0003-9.
1037
- 1038 64. Bernardi, G.; Bernardi, G. Compositional Constraints and Genome Evolution. *J Mol*
1039 *Evol* **1986**, 24, 1–11, doi:10.1007/BF02099946.

- 1040
1041 65. Jeffares, D. C.; Mourier, T.; Penny, D. The Biology of Intron Gain and Loss. *Trends*
1042 *Genet* **2006**, 22, 16–22, doi:10.1016/j.tig.2005.10.006.
1043
1044
- 1045 66. Glasauer, S. M. K.; Neuhauss, S. C. F. Whole-Genome Duplication in Teleost
1046 Fishes and Its Evolutionary Consequences. *Mol Genet Genomics* **2014**, 289, 1045–
1047 1060, doi:10.1007/s00438-014-0889-2.
1048
- 1049 67. Coscia, M. R.; Varriale, S.; De Santi, C.; Giacomelli, S.; Oreste, U. Evolution of the
1050 Antarctic Teleost Immunoglobulin Heavy Chain Gene. *Mol Phylogenet Evol* **2010**,
1051 55, 226–233, doi:10.1016/j.ympev.2009.09.033.
1052
- 1053 68. Mills, R. E.; Bennett, E. A.; Iskow, R. C.; Devine, S. E. Which Transposable
1054 Elements Are Active in the Human Genome? *Trends Genet* **2007**, 23, 183–191,
1055 doi:10.1016/j.tig.2007.02.006.
1056
- 1057
1058 69. Gelfman, S.; Burstein, D.; Penn, O.; Savchenko, A.; Amit, M.; Schwartz, S.; Pupko,
1059 T.; Ast, G. Changes in Exon–Intron Structure during Vertebrate Evolution Affect
1060 the Splicing Pattern of Exons. *Genome Res* **2012**, 22, 35–50,
1061 doi:10.1101/gr.119834.110.
1062
- 1063
1064 70. Das, S.; Hirano, M.; Tako, R.; McCallister, C.; Nikolaidis, N. Evolutionary
1065 Genomics of Immunoglobulin-Encoding Loci in Vertebrates. *Curr Genomics* **2012**,
1066 13, 95–102, doi:10.2174/138920212799860652.
1067
- 1068 71. Pisano, E.; Coscia, M. R.; Mazzei, F.; Ghigliotti, L.; Coutanceau, J.-P.;
1069 Ozouf-Costaz, C.; Oreste, U. Cytogenetic Mapping of Immunoglobulin Heavy
1070 Chain Genes in Antarctic Fish. *Genetica* **2007**, 130, 9–17,
1071 doi:10.1007/s10709-006-0015-4.
1072
- 1073 72. Wong, J. W. H.; Ho, S. Y. W.; Hogg, P. J. Disulfide Bond Acquisition through
1074 Eukaryotic Protein Evolution. *Mol Biol Evol* **2011**, 28, 327–334,
1075 doi:10.1093/molbev/msq194.
1076
- 1077 73. Kaattari, S.; Evans, D.; Klemer, J. Varied Redox Forms of Teleost IgM: An
1078 Alternative to Isotypic Diversity? *Immunol Rev* **1998**, 166, 133–142,
1079 doi:10.1111/j.1600-065x.1998.tb01258.x.
1080

- 1081 74. Zhang, Y.-A.; Salinas, I.; Sunyer, J. O. Recent Findings on the Structure and
1082 Function of Teleost IgT. *Fish Shellfish Immunol* **2011**, 31, 627–634,
1083 doi:10.1016/j.fsi.2011.03.021.
1084
- 1085 75. Kaetzel, C. S. Coevolution of Mucosal Immunoglobulins and the Polymeric
1086 Immunoglobulin Receptor: Evidence That the Commensal Microbiota Provided the
1087 Driving Force. *Int Sch Res Notices*. Volume **2014**, Article ID 541537, doi:
1088 10.1155/2014/541537.
1089
- 1090 76. Su, Y.-L.; Wang, B.; Hu, M.-D.; Cui, Z.-W.; Wan, J.; Bai, H.; Yang, Q.; Cui, Y.-F.;
1091 Wan, C.-H.; Xiong, L.; Zhang, Y.-A.; Geng, H. Site-Specific N-Glycan
1092 Characterization of Grass Carp Serum IgM. *Front Immunol* **2018**, 9,
1093 doi:10.3389/fimmu.2018.02645.
1094
- 1095 77. Torres, P.; Soto, M. S. *Hysterothylacium Winteri* Sp. n. (Nematoda: Anisakidae), a
1096 Parasite of Chilean Rock Cod, *Eleginops Maclovinus* (Perciformes: Eleginopidae),
1097 from South Chile. *Folia Parasitol (Praha)* **2004**, 51, 55–60,
1098 doi:10.14411/fp.2004.008.
1099
- 1100 78. Brickle, P.; Mackenzie, K. Parasites as Biological Tags for *Eleginops Maclovinus*
1101 (Teleostei: Eleginopidae) around the Falkland Islands. *J Helminthol* **2007**, 81, 147–
1102 153, doi:10.1017/S0022149X07750514.
1103
- 1104 79. Coscia, M. R.; Oreste, U. Presence of antibodies specific for proteins of
1105 *Contracaecum osculatum* (Rudolphi, 1908) in plasma of several Antarctic teleosts.
1106 *Fish Shellfish Immunol* **1998**, 8, 295–302, doi:10.1006/fsim.1998.0140
1107
- 1108 80. Coscia, M. R.; Oreste, U. Plasma and Bile Antibodies of the Teleost *Trematomus*
1109 *bernacchii* Specific for the Nematode *Pseudoterranova Decipiens*. *Dis Aquat Organ*
1110 **2000**, 41, 37–42, doi: 10.3354/dao041037.
1111
- 1112 81. Zhang, X.-T.; Yu, Y.-Y.; Xu, H.-Y.; Huang, Z.-Y.; Liu, X.; Cao, J.-F.; Meng, K.-F.;
1113 Wu, Z.-B.; Han, G.-K.; Zhan, M.-T.; et al. Prevailing Role of Mucosal Igs and B
1114 Cells in Teleost Skin Immune Responses to Bacterial Infection. *J Immunol* **2021**,
1115 206, 1088–1101, doi:10.4049/jimmunol.2001097.
1116
- 1117 82. Kong, W.; Yu, Y.; Dong, S.; Huang, Z.; Ding, L.; Cao, J.; Dong, F.; Zhang, X.; Liu,
1118 X.; Xu, H.; et al. Pharyngeal Immunity in Early Vertebrates Provides Functional
1119 and Evolutionary Insight into Mucosal Homeostasis. *J Immunol* **2019**, 203, 3054–
1120 3067, doi:10.4049/jimmunol.1900863.
1121

- 1122 83. Xu, Z.; Takizawa, F.; Casadei, E.; Shibasaki, Y.; Ding, Y.; Sauters, T. J. C.; Yu, Y.;
 1123 Salinas, I.; Sunyer, J. O. Specialization of Mucosal Immunoglobulins in Pathogen
 1124 Control and Microbiota Homeostasis Occurred Early in Vertebrate Evolution.
 1125 *Science Immunology* **2020**, *5*, doi:10.1126/sciimmunol.aay3254.

1126

Table 1. Length of the IgT CH2 domain in non-Antarctic (in red) and Antarctic (in blue) notothenioid fish. The minimum and maximum length values observed among all species are underlined

Species	Nucleotides	Amino acids
<i>B. diacanthus</i>	<u>285</u>	<u>95</u>
<i>C. gobio</i>	282	94
<i>P. urvillii</i>	282	94
<i>E. maclovinus</i> Ema1	30	10
<i>E. maclovinus</i> Ema2	36	12
<i>T. bernacchii</i> TbeS	33	11
<i>T. bernacchii</i> TbeL	51	17
<i>G. gibberifrons</i> Ggi1	<u>24</u>	<u>8</u>
<i>N. coriiceps</i> Nco1	<u>24</u>	<u>8</u>
<i>H. velifer</i> * Hve1	39	13
<i>G. acuticeps</i> Gac1	39	13

* two *C τ 3* variants

Table 2. Length of the IgT *C τ 1-C τ 2* and *C τ 2-C τ 3* introns in non-Antarctic (in red) and in Antarctic (in blue) notothenioid fish. The minimum and maximum length values observed across all species are underlined

Species	<i>Cτ1-Cτ2</i> intron (nt)	<i>Cτ2-Cτ3</i> intron (nt)
<i>B. diacanthus</i>	242	144
<i>C. gobio</i>	224	147
<i>P. urvillii</i>	222	142
<i>E. maclovinus</i>	<u>206</u>	<u>108</u>
<i>T. bernacchii</i>	318	122
<i>G. gibberifrons</i>	237	115
<i>N. coriiceps</i>	212	126
<i>H. velifer</i>	216	128
<i>G. acuticeps</i>	216	126
<i>T. bernacchii</i>	318	122

	[<i>Ct1</i> exon][<i>Ct2</i> exon	
Ema1	GTGACTCATTCTGGAACCTCAAAAAGTGTGGAATTAAAAA-----TACTCAA---TGT-----CCAAAGA-TAG		80
Ema2	GTGACTCATTCTGGAACCTCAAAAAGTGTGGAATTAAAAA---CAAGA-TACTCAA---TGT-----CCAAAGA-TAG		
Tb30.3	GTGACTCACCTGGAGGACAAAAAAGTTTGTGAGCATTAAAAAGGTTAA-----AGCACCTTCATGTTC		
Tb30.7	GTGACTCACCTGGAGGACAAAAAAGTTTGTGAGCATTAAAAAGGTTATTCAACTTTTTAGGTGCTTTAGCACCTTCATGTCC		
Tb30.8	GTGACTCACCTGGAGGACAAAAAAGTTTGTGAGCATTAAAA-----AGCACCTTCATGTTC		
Nco	GTGGCTCACCTGGAGGACAAAAAAGTTTGAACATTAAAAAN-----CCTTCATGTCC		
Ggi1	GTGACTCACTCTGGAGGATCAACAAGTTTGAACATTAAAAAT-----CCTTCATGTAC		
Hve1	GTGACTCACCTGGAGGACCAACAAGTTTGTGAGCATTAAAAAGGCAATT----CAACA-----AGCACCTTCATGTCC		
Hve2	GTGACTCACCTGGAGGACCAACAAGTTTGTGAGCATTAAAAAGGTAATT----CAACA-----AGCACCTTCATGTCC		
Gac1	GTGACTCACCTGGAGGACCAACAAGTTTGAACATTAAAAAGGCAATT----CAACA-----AGCACCTTCATGTAC		
	.:****.:: ** ***:** :*****		
][<i>Ct3</i> exon		
Ema1	CAT--CGAAAGCTCCCAT-----CACAAATAGAACTGAAAAAATCAAGTCCCAAAGAAGTGTTCATAATAACA		160
Ema2	CAT--CGAAAGCTCCCAT-----CGCAATAGAACTGAAAAAATCAAGTCCCAAAGAAGTGTTCATAATAACA		
Tb30.3	AGAGACTAACGCTACCAT-----CACGATGAAACTGAACAAATCAAGTCCCAAAGAAATGTTCACAACA		
Tb30.7	AGAGACTAACGCTCCCAT-----CACGATGAAACTGAACAAATCAAGTCCCAAAGAAATGTTCACAACA		
Tb30.8	-----CTACCAT-----CACGATGAAACTGAACAAATCAAGTCCCAAAGAAATGTTCACAACA		
Nco	AAAGACTAAAGCTCCCAT-----CACGATGAAACTGAACAAATCAAGTCCCAAAGAAATGTTCACAACA		
Ggi1	AGAGACTAAAGCTCCCAT-----CACGATGAAACTGAACAAATCAAGTCCCAAAGAAATGTACACCAACA		
Hve1	AAAGACTAAAGCTCCCGTTTGTGTAGCTCCCGTCACGATGAAACTGATCATA--AGTCCCAAAGAAATGTTCACAACA		
Hve2	AAAGACTAAAGCTCCCGT-----CACGATGAAACTGATCAAA--AGTCCCAAAGAAATGTTCACAACA		
Gac1	AGAGACTAAAGCTCCCAT-----CACGATGAAACTGATCAGA--AGTCCCAAAGAAATGTTCACAACA		
	:*:* **:*:*:***:* * ***:*****:***.**:*.***		

Fig. 1. Multiple alignment of partial cDNA sequences spanning the 3' end of the *Ct1* exon, and the 5' end of the *Ct3* exon. Nucleotide sequences are representatives of different cDNA clones coding for different variants (wherever found), derived from the non-Antarctic species *E. maclovinus* (clones Ema1 and Ema2), and from the Antarctic species *G. gibberifrons* (clone Ggi1), *H. velifer* (clones Hve1 and Hve2, both coding for the same *Ct2* remnant but differing in *Ct3* length), *G. acuticeps* (clone Gac1). Three cDNA transcripts (Tb30.3, Tb30.7, Tb30.8) encoding the TbeS, TbeL and TbeSts variants in *T. bernacchii* and one cDNA sequence from *N. coriiceps* (Nco), obtained in a previous work [19], have been added for comparison. *Ct* exon boundaries are reported above the alignment. Gaps are indicated by dashes. Below the alignment, identical nucleotides are marked with an asterisk, positions where only one sequence shows a different nucleotide are marked with a dot, positions differing in two nucleotides are marked with a colon. The duplicated 9-nt sequence at the beginning of the *Ct3* exon of the Hve2 transcript is underlined. Since several sequences varying in length at the 5' and/or at 3' were obtained from each species, only the region of each representative sequence that aligned over the same length has been shown here. Full alignments are provided in Figure S1.

gEma1 GTATGTGATTTTATTTACTTGCCATCATGGCTTTCTTTTCTCGTTGACTCCAAATGAATCACGTTCTTAATCTCAGTTTT 80
gEma3 GTATGTGATTTTATTTACTTGCCATCATGGCTTTCTTTTCTCGTTGACTCCAAATGAATCACGTTCTTAATCTCAGTTTT
g30.8 GTACGTGATTTTATTTACTTGCTATCATGATCTTCTTTACTCTTTGAGTCCAAATGAAGCACATT-----CTCAGTTTT
g30.6 GTACGTGATTTTATTTACTTGCTATCATGATCTTCTTTACTCTTTGAGTCCAAATGAAGCACATT-----CTCAGTTTT
gNco1 GTATATGATTTTATTTACTTCTTGCCCGATCATCTTTACTCTTTGAGTCCAAATGCTGATCATT-----CGT-----
gGgi2 GTATGTGATTTTATTTACTTGCTATCATGATCTTCTTTACTCTTTGAGCCCAAATGAAGCACATT-----CTCAGTTTT
gGgi3 GTATGAGATTTTATTTACTTGCTATCAGGATCTTCTT--ACTCTTTGAGCCCAAATGAAGCACAT-----CTCAGTTT--
gHve2 GTACGTGATTTTATTTACTTGCTATCATGATCTTCTTTCTCTTTGAGTCCAAATGAAGCACATT-----CTCAGTTGT
gHve5 GTACGTGATTTTATTTACTTGCTATCATGATCTTCTTTACTCTTTGAGTCCAAATGAAGCACATT-----CTCAGTTGT
gGac1 GTACGTGATTTTATTTACTTGCTATCATGATCTTCTTTACTCTTTGAGTCCAAATGAAGCACATT-----CTCAGATTC
*** .*.*****. *.:..* *::..** *:::***:*****. .:. *:* *..

gEma1 ATCCTGAAAAATGTTTCAGTAGCACTCTTACTTTCTCACGTTTATCGTCCTCCGTCATGTGTCTT----- 160
gEma3 ATCCTGAAAAATGTTTCAGTAGCACTCTTACTTTCTCACGTTTATCGTCCTCCGTCATGTGTCTT-----
g30.8 ATCCTGAAAAATGTTTCATTAACACTCTCACTTTCTCATGTTTATCTTCATCCATCATGTGTCTTACTTAA---ACAAATCT
g30.6 ATCCTGAAAAATGTTTCATTAACACTCTCACTTTCTCAATGTTTATCTCCATCCATCATGTGTCTTACTTAA---ACAAATCT
gNco1 -CCTTAAAAAT-----CACTTTCTCTGTTTATCTACGTCATCATGTGTCTTACTTGA---ACAAATCT
gGgi2 ATCCTGAAAAATGTTTCATTAACACTCCCCTTTCTCATGTTTATCTTCATCCATCATGTGTCTTACTTAA---ACAAATCT
gGgi3 ATCCTGAAAAATGTTTCATTAACACTCTCACTTTCTCATGTT--ATCTTCATCCATCATGTGTCTTACTTACTTTACAAATCT
gHve2 ATCCTGAAAAATGTTTCATTAACAC--TCACTTTCTCATGTTTATCTTCATCCATCATGTGTCTTACTTAA---AAAAATCT
gHve5 ATCCTGAAAAATGTTTCATTAACAC--TCACTTTCTCATGTTTATCTTCATCCATCATGTGTCTTACTTAA---AAAAATCT
gGac1 ATCCTGAAAAATGTTTCATTAACAC--CCACTTTCTCATGTTTATCTTCATCCATCATGTGTCTTACTTAA---AAAAATCA
*. * .***** :***. * .:*** *::: * *:::*****

gEma1 AATATGATGCATTTCAAACCTGATGTTAATG----- 240
gEma3 AATATGATGCATTTTCAGACTGATGTTAATG-----
g30.8 AAAATGATGCATTTTCAGACTGATATCAATAAAAGATATCTATTTGTACTCTTTCTTCCCCATGCTTAGAGGCTAAAGCAC
g30.6 AAAATGATGCATTTTCAGACTGATATCAATAAAAGATATCTATTTGTACTCTTTCTTCCCCATGCTTAGAGGTTAAAGCAC
gNco1 AAAATGATGAATTTCAAACCTGATATCAATT-----
gGgi2 AAAATGATGAATTTTCAGACTGATATCAATT-----
gGgi3 AAAATGATGAATTTTCAGACTGATATCAATT-----
gHve2 AAAATGATGCATTTTCAGACGGATATCATTT-----
gHve5 AAAATGATGCATTTTCAGACGGATATCATTT-----
gGac1 AAAATGATGCATTTTCAGACTGATATCATTT-----
.:***. *.:***:.* *:

gEma1 -----CAAATATCTAATTGTGTTCTTTCTTCCCC 320
gEma3 -----CAAATATCTAATTGTGTTCTTTCTTCCCC
g30.8 CTTTCTCATGTTTATCTTCATCCATCATGTGTCTTACTTAAACAAATCTAAAAAATATCTATTTGTACTCTTTCTTCCCC
g30.6 CTTTCTCATGTTTATCTTCATCCATCATGTGTCTTACTTAAACAAATCTAAAAAATATCTATTTGTACTCTTTCTTCCCC
gNco1 -----AAAAATATCTATTTGTACTCTTTCTTCCCC
gGgi2 -----AAAAAGATCTATCTGACTCTTTCTTCCCC
gGgi3 -----AAAAGTATCTATCTGACTCTTTCTTCCCC
gHve2 -----AAAATATCTATTTGCACCTTTCTTCCCC
gHve5 -----AAAATATCTATTTGCACCTTTCTTCCCC
gGac1 -----AAAATATCTATTTGCACCTTTCTTCCCC
:*. * .*****: * :*****.***

] [Cr2 exon] [

gEma1 ATGCTT-----CAAACAAGATACTC-----AATGTCCAAAGATAGCATCGAAAGGTAGA 400
gEma3 ATGCTT-----AGAACAAGATACTC-----AATGTCCAAAGATAGCATCGAAAGGTAGA
g30.8 ATGCTT-----AGAGGGTTAA-----AGCACCTTCATGTCCAGAGACTAACGGTAAA
g30.6 ATGCTT-----AGAGGGTTATTCAACTTTTAGGTGCTTTAGCACCTTCATGTCCAGAGACTAACGGTAAA
gNco1 ATGCTTGGCCATAAATCAA-CAAG-----ATCCTTCATGTCCAAAGACTAAAGGTAAA
gGgi2 ATGCTTGGACATAAATCAA-CAAG-----ATCCTTCATGTATAGAGACTAAAGGTAAA
gGgi3 ATGCTTGGACATAAATCAA-CAAG-----ATCCTTCATGTACAGAGACTAAAGGTAAA
gHve2 ATGCTT-----AGAGGCAATTCAAC-----AAGCACCTTCATGTCCAAAGACTAAAGGTAAA
gHve5 ATGCTT-----AGAGGCAATTCAAC-----AAGCACCTTCATGTCCAAAGACTAAAGGTAAA
gGac1 ATGCTT-----AGAGGCAATTCAAC-----AAGCACCTTCATGTACAGAGACTAAAGGTAAA
***** ** :*:***: * :*:***:***:

Cr2-Cr3 intron

gEma1 GGGCTAC-AC--TCATTACATAAATACTGATTTACTGTAAGTATAGTATACATTATGCTCAAATATAAGGGATGTACAAT 480
gEma3 GGGCTAC-AC--TCATTACATAAATACTGATTTACTGTAAGTATAGTATACATTATGCTCAAATATAAGGGATGTACAAT
g30.8 GAGCTACTACTGTCAATTACATAAATACTGTTTACTGTAAGTATAGTATACATTATACTCAAATATAAGAGATCTACTGT
g30.6 GAGCTACTACTGTCAATTACATAAATACTGTTTACTGTAAGTATAGTATACATTATACTCAAATATAAGAGATCTACTGT
gNco1 GAGCTACTACTGTCAATTACATAAATACTGTTTACTGTAAGTATAGTATACATTATACTCAAATATAAGAGATCTACTGT
gGgi2 GAGTTACTACTGTCAATTACATAAATACTGTTTACTGTAAGTATAGTATAGTATACATTATACTCAAATATAAGAGATCTACTGT
gGgi3 GAGCTACTACTGTCAATTACATAAATACTGTTTACTGTAAGTATAGTATACATAAATACTCAAATATAAGAGATCTACTGT
gHve2 GAGCTACTACTGTCAATTACATAAATACTGTTTACTGTAAGTATAGTATACATAAATACTCAAATATAAGAGATCTACTGT
gHve5 GAGCTACTACTGTCAATTACATAAATACTGTTTACTGTAAGTATAGTATACATAAATACTCAAATATAAGAGATCTACTGT
gGac1 GAACTACTACTGTCAATTACATAAATACTGTTTACTGTAAGTATAGTATACATTATACTCAAATATAAGAGATCTACTGT
*.:** * .*****.*****:*****

] [Cr3 exon

gEma1 TTTTATTTATTG-TTTGTAT-----TCTTGTCTGTAGCTCCCAT-----CACTATATAACTGAAAA 560
gEma3 TTTTATTTATTG-TTTGTAT-----TCTTGTCTGTAGCTCCCAT-----CACAATAGAACTGAAAA
g30.8 TTGTATTTATTGATTTATTTCTGTTGATCTTGTCTGTAGCTACCAT-----CACGATGAACTGAACA
g30.6 TTGTATTTATTGATTTATTTCTGTTGATCTTGTCTGTAGCTACCAT-----CACGATGAACTGAACA
gNco1 CTGTATGATTGATTTTTTCTGTTGATCTTGTCTGTAGCTCCCAT-----AACGATGAACTGAACA
gGgi2 TTGTATTTATTGATTTATTTCTGTTGATCTTGTCTGTAGCTCCCAT-----CACGATGAACTGAACA
gGgi3 TTGTATTTATTGATTTATTTCTGTTGATCTTGTCTGTAGCTCCCAT-----CACGATGAACTGAACA

gHve2 TTGTATTTATTGATTTGTTTCTGTTGATCTTGTTTCTGTAGCTCCCGT-----CACGATGAAACTGATCA
gHve5 TTCTATTTTATTGATTTTCTTCTCTGATCTTCTTCTCTACCTCCCGCTTCTCTACCTCCCGCTACCATCAAACTCATCA
gGac1 TTCTATTTTATTGATTTTCTTCTCTGATCTTCTTCTCTACCTCCCGCTTCTCTACCTCCCGCTACCATCAAACTCATCA
Journal Pre-proof
.:***.***** ** *.* *****:***:*. ** **: ***** :*

Fig. 2. Multiple alignment of partial genomic sequences encoding the notothenioid IgT CH region. Nucleotide sequences are representatives of different genomic clones coding for different variants (wherever found) from the non-Antarctic species *E. maclovinus* (clones gEma1 and gEma3), and from the Antarctic species *N. coriiceps* (clone gNco1), *G. gibberifrons* (clones gGgi2 and gGgi3/pseudogene), *H. velifer* (clones gHve2 and gHve5, both coding for the same *Cτ2* remnant but differing in *Cτ3* length), and *G. acuticeps* (clone gGac1). The *T. bernacchii* clones g30.8 and g30.6, coding for the respective TbeS and TbeL variants, previously determined [19], have been added for comparison. The exon-intron boundaries are reported above the alignment. Donor and acceptor splicing sites are shaded in black. Gaps are indicated by dashes. Below the alignment, identical nucleotides are marked with an asterisk, positions where only one sequence shows a different nucleotide are marked with a dot, positions differing in two nucleotides are marked with a colon. The two duplicated 46-bp long sequences in the *T. bernacchii* *Cτ1-Cτ2* intron and the one duplicate 9-bp long sequence in the gHve5 *Cτ3* exon are underlined. Since several sequences varying in length at the 5' and/or at 3' ends were obtained for each species, only the region of each sequence that aligned over the same length has been shown. Full alignments are provided in Fig. S2.

nucleotides are marked with a colon. Twenty six nucleotides (underlined, in bold) of a 32-nt long region (shaded in green) at the beginning of the *P. urvillii* *C τ 1-C τ 2* intron were present within the 32-nt long complementary reverse motif (shaded in cyan) in the respective *C τ 2* exon. Since several sequences varying in length at the 5' and/or at 3' ends were obtained for each species, only the region of each sequence that aligned over the same length has been shown. Full alignments are provided in Figure S3.

Journal Pre-proof




Fig. 4. Bayesian phylogeny of notothenioid IgTs, based on the concatenated multiple sequence alignment of the $C\tau 1$ - $C\tau 2$ and $C\tau 2$ - $C\tau 3$ introns. Phylogeny was built with two parallel MCMC analyses, run for 1 million generations. Nodes supported by posterior probability values lower than 50% were collapsed.




Fig. 5. Significantly enriched sequence motifs identified by MEME in the *Ct1-Ct2* intron structure, exemplified by colored boxes. The degree of nucleotide conservation of each of the five motifs identified is reported as a sequence logo. Sequence names design different genomic clones from different species, as follows: *B. diacanthus* (gBdi), *C. gobio* (gCgo), *P. urvillii* (gPur1), *E. maclovinus* (gEma1 and gEma3), *G. gibberifrons* (gGgi2 and the pseudogene gGgi3), *G. acuticeps* (Gac1), *H. velifer* (gHve2 and gHve5), *N. coriiceps* (gNco1), and *T. bernacchii* (g30.6 and g30.8), representing the previously described TbeL and TbeS variants [19].




Fig. 6. Distribution of AT content in the $C\tau 1-C\tau 2$ and $C\tau 2-C\tau 3$ introns. The representative genomic clones encoding the $C\tau$ region are from the non-Antarctic species *B. diacanthus* (clone gBdi), *C. gobio* (gCgo), *P. urvillii* (clone gPur1), *E. maclovinus* (clone gEma1), and from the Antarctic species *T. bernacchii* (clone g30.8), *G. gibberifrons* (clone gGgi2), *N. coriiceps* (clone gNco1), *H. velifer* (clone gHve2), and *G. acuticeps* (clone gGac1).

CH1 Domain

E. coioides YSQTAA-PALFPLVQ**CK**SGTAGTIVTVG**CIA**QDFFPESLTF**QWT**DASGTTQTFKQYPTVMKKDNKYTGVSVLVDSKSAWDSRRSFS**CSV**THPGGSESVTLQKP--
S. aurata SGQTATA-PTLFPLVQ**CK**PGTGDRITVG**CLAL**DFS^PSSLTF**QWT**DARGTSLNSVXYPPVEKKNKYTEVSLVELSKSDWDAMHSFT**CS**AGSMSLM----LQKPSS
T. orientalis SSETPSS-PTLYPLLQ**C**SDTGNKVTVG**CLAR**DFPKSID**FQW**NDARGTRVDSAQYISGQ--NNKYTGVSVVQVSRSDIKS--SY**NC**SV^DDHLGRTMAVTVKLP--
G. aculeatus ----TTS-PTLFPLVK**CE**PD-GDKVTVG**CLAH**HNYPKSLTF**QWT**NACGADLTS^DQYPSILNNNKYTGVSLLKVPKSDWNLRHSFE**CS**VTHAGGSRSVTLGKK--
S. chuatsi SSVTVAS-PTLFPLVQ**CN**SGPADKITVG**CLAR**DFYPKSLTF**QWT**NS^SSGTALTS^ENYPPAEKKNKYTGVSLLVQVSKSDWDSRKS**FKC**SVHHNGSTHDLQVHKP--
S. caurinus STVNAAS-PTLFPLVQ**CT**PGSADKVT**VGCLAK**DYPRSLTF**QWT**DGSGNTQTSYQYPAVEKKNKYTGVSLLQVQKSEWDSK**SFNC**SVTHPGGQKSVTLEKP--
D. labrax SGTSTR--PTVFPLM**QCG**SGSGDKVT**LGCL**ATGFTPSSLT**YSW**SKNGSLLTD**FIQ**YPAVQKKNVYMGVSQIRVSRQDWDAR**QPFQ**CVATHAAGNAQADIVKP--
 Bdi NTV--AS-PTLFPLL**Q**CNPVSAST--VG**CLANN**FYPKSLTF**QWT**DTS^GTTLSSVQYPTVEKKN**NYT**GVSLQVSKSDWDSKKS**F**TC^SVTHDGSQRSVQLSK---
 Ema1TIRVG**CLAN**DFT**PNA**VT**FQW**TASG**STLS**ALQY**PLME**KKNKY**TG**VSVLEVSKSN**WDS**SRK**SFN**CA**VT**HS**GT**SKS**VEI**-K---
 Ema2FPLV**QC**DT**SAN**TI**R**VG**CLAN**DFT**SNA**VT**FR**W**T**DASG**STLS**ALQY**PLME**KKNKY**TG**VSVLEVSKSN**WDS**SRK**SFN**CA**VT**HS**GT**SKS**VEI**-K---
 TbeS SNEIDAS-PTLFPLVQ**CN**PGSAE**IT**VG**CLAN**DFS^PNSVTF**QWT**DASG**TNL**PAVQ**YPP**TVKKNKY**TG**ISV**VN**VPKSDWDLRKS**YK**CA**VT**HPGGQK**SL**SI-K---
 TbeL SNEIDAS-PTLFPLVQ**CN**PGSAE**IT**VG**CLAN**DFS^PNSVTF**QWT**DASG**TNL**PAVQ**YPP**TVKKNKY**TG**ISV**VN**VPKSDWDLRKS**YK**CA**VT**HPGGQK**SL**SI-K---
 TbeSts SNEIDAS-PTLFPLVQ**CN**PGSAE**IT**VG**CLAN**DFS^PNSVTF**QWT**DASG**TNL**PAVQ**YPP**TVKKNKY**TG**ISV**VN**VPKSDWDLRKS**YK**CA**VT**HPGGQK**SL**SI-K---
 Nco SEIVASSP**PTLF**PLV**QCN**PDSA**EIT**AVG**CLAN**AFY**PNS**VT**FQW**TASG**TLS**AVQ**YPS**IEKKNKY**TG**VSVL**NP**KSDWDSRKS**FKCN**VAH**PGG**PK**SL**NI-K---
 GgilS-PTLFPLVQ**CN**PGSAE**IT**VG**CLAN**AFY**PNS**VT**FQW**TASG**TNL**SAVQ**YPP**TVKKNKY**TG**VSV**VK**VPKSDWDSRKS**SFN**CA**VT**HS**GG**ST**SL**NI-K---
 Hve1S-PTLFPLVQ**CN**PGSAE**IT**VG**CLAN**AFY**PNS**VT**FQW**TASG**TLS**AVQ**YPP**TE**EN**NKY**TG**VSV**VN**VKSDW**YS**RKS**SFN**CT**VT**HPGG**P**T**SL**SI-K---
 Hve2S-PTLFPLVQ**CN**PGSAE**IT**VG**CLAN**AFY**PNS**VT**FQW**TASG**TLS**AVQ**YPP**TE**EN**NKY**TG**VSV**VN**VKSDW**YS**RKS**SFN**CT**VT**HPGG**P**T**SL**SI-K---
 Gac1S-PTLFPLVQ**CN**PGSAE**IT**VG**CLAN**AFY**PNS**VT**FQW**TASG**TLS**AVQ**YPP**TE**EN**NKY**TG**VSV**VN**VKSDW**YS**RKS**SFN**CT**VT**HPGG**P**T**SL**NI-K---
 .**.* . : .**.* . * . . : * . * . : . . * : . .

CH2 Domain

E. coioides --PPPKVTLVAVPAGDTQTLV**CT**IEDLPSN**QLS**-V**KWK**KDDNSV**TG**FTDCPPQ**LNG**GVY**TAVS**ILKVT**NSE**WDSKAVY**TCE**V**TN**Q**G**TT**YP**PKV**SK**
S. aurata ----PPKVTL**LS**SVPRGDTQALV**CT**IEDYAPK**ELT**-V**KWK**KNAHDV**PG**STD**WK**PE**SI**GDMFSA**VSIL**NVK**NTD**WNRD**VTCE**V**TH**AG**TQ**Y**IK**K**ASK**
T. orientalis ---SPPRVTL**LS**SVPNGDTQV**LV**CTIE**EF**LP**ETLS**-V**KWK**KNRD**Y**ESD**F**TD**DW**V**PKQ**IGDV**YSA**VSVL**KV**K**ND**W**ES**KAVY**TCE**V**TH**K**GKI**Y**EK**K**ASK**
G. aculeatus -----
S. chuatsi ---IPPKVTL**VS**VPSEDSQALV**CT**IEDGRSG**TLD**SK**FWK**KNGAELND**YIQ**S**PIQ**KT**GEL**HS**AVS**VLK**VK**NTD**W**SKAVY**TCE**V**TY**SG**TQ**Y**KK**K**ASK**
S. caurinus ---TPPKVTL**L**AVPGDDQTLV**CT**IQDLIS**NELS**-V**KWK**KDVND**VTG**ST**TWAA**Q**KTG**SL**YSA**VSVL**KV**R**NI**D**W**SDAVY**TCE**V**M**H**G**T**KH**T**K**K**V**T**K**
D. labrax --RDPPKVTL**VS**VPRGDSQALM**CT**VED**FLPK**DL**T**-IN**W**KKNQ**ND**VAG-T**NW**DP**VL**TGD**LYSA**VSVL**KV**K**NT**D**W**NS**NAV**Y**T**C**V**ATH**R**G**K**Q**Y**K**K**I**ASK**
 Bdi DV**S**FPPK**L**TLVAVPGEDAQALV**CT**IED**FLPK**N**V**L-V**KWK**KNGNSV**TG**CI**D**W**AP**K**IG**N**V**Y**SA**VSVL**KV**K**NN**D**W**SN**AV**Y**T**C**E**V**V**H**Q**G**T**S**Y**T**M**R**ASK**
 Ema1 -----NT**QC**PK-----I**ASK**
 Ema2 -----KQDT**QC**PK-----I**ASK**
 TbeS -----K**V**K-----A**P**S**C**PE-----**TN**
 TbeL -----K**V**I--**Q**LL**GAL**A**P**S**C**PE-----**TN**
 TbeSts -----
 Nco -----**NP**S**C**PK-----**TK**
 Ggil -----**NP**S**C**TE-----**TK**
 Hve1 -----K**A**I**Q**Q-----A**P**S**C**PK-----**TK**
 Hve2 -----K**V**I**Q**Q-----A**P**S**C**PK-----**TK**
 Gac1 -----K**A**I**Q**Q-----A**P**S**C**TE-----**TK**

CH3 Domain

E. colioides -----VPITVTLTQSSPKEIFSNQAKFECVITGTDQGTGP-DFQIIWQVDGQNVTDN--IETKPGSK----KISTMTRAHTDWQSIKVRCSAIRDNMTPIVQELTIQKG
S. aurata -----APITVTLNQPSPKKEIFSNQVKLECVITGQDRSIVDDTKITWQIDGQNIENVTHLATQSSSGQYSKTSTMTRSYTEWLGVTQVRCVSAEGNDMTPVVQDLTVHKG
T. orientalis -----APITVTLNPPSPKFFNNQAELECIIVEGQDNTIVSETEMWQINGKNVAGNMGLLKTAGS--QYSKTNTLTRSLTEWLVQVNTVRCVSAKRKDVTVTKDLTFHKG
G. aculeatus ASCSMIRNITMTLKPTSPKDFINNQAKFECRINGPDQNTINEMKFTWQINGADVTKNSSETTDATGS----KISTLTRDRSEWQIDNVRCVSAKG-PKTHVSKDLHINKG
S. chuatsi -----APITVTLNQPSPKKEIFSNQAELECIITGQDETIVDEIKVWQIDGQVDSDNINETTKSVDG-QRIKTSTMTSRTEWQRVNKRCSAIRDDDT-LIQDLTVHKG
S. caurinus -----APITMTLRHSSPKDVFINNQAKLECVVTVGQDIG---QQITWEIDGQHVTSNNTETTTSGSS----KISTITRSRTDWKGVNTVRCVLAALRDNMTPIVQDPTFNKG
D. labrax -----ATVTVTLNQPSAKEIFSNNEAKLECIIVTGRDKSIVNEIQITWQIDGQTVAEINIESRVSSDGD-QHSKTSTMSRNRTEWQGVNKRCSAAREDMTPVVQDLTVHKG
Bdi -----APITMILNQPSPKEMFNHQAACLECIISGQDQATMNDIQITWQMYGENVTTHNNIETTWSGGD----KVSVITLSVTEWQVNVKRCSAMRDNMTPIVQDLIVHKG
Ema1 -----APITIELKKSSPKVEFNQITFECCIISGEDEATLNAFTISWKNVGNWTKNTENGRSGG-----KSTMKLSVDELKDATNVGCSATGRNGVPVEKELTLQTS
Ema2 -----APIAIELKKSSPKVEFNQITFECCIISGEDEATLNAFTISWKNVGNWTKNTENGRSGG-----KSTMKLSVDELKDATNVGCSATGRNGVPVEKELTLQTS
TbeS -----ATITMKNKSSPKEMFNQARFDCVISGDEETTVDQFQITWQVNGAIRTTSEQ-GRSGG-----IKTSTMRLSLHESLS--NVRCVSAKSD-DVTVFQDLPPPPI
TbeL -----APITMKNKSSPKEMFNQARFDCVISGDEETTVDQFQITWQVNGAIRTTSEQ-GRSGG-----IKTSTMRLSLHESLS--NVRCVSAKSD-DVTVFQDLPPPPI
TbeSts -----TTITMKNKSSPKEMFNQARFDCVISGDEETTVDQFQITWQVNGAIRTTSEQ-GRSGG-----IKTSTMRLSLHESLS--NVRCVSAKSD-DVTVFQDLPPPPI
Nco -----APITMKNKSSPKEMFNQTKFDCVISGQDETTVNFEITWQVDGATRATSEP-TSSGG-----IKTSTMTLTLNESLR--DVRCVSAKSDKVTAVSQNLLTPKT
Ggil -----APITMELNKSSPKEMYTNQARFDCVIRGQDVTTVNLFQITWQVNGAIRTTSEP-RSSGG-----TKTSTMRLSLNESLS--DVRCVSAKSD-NVTVVQDLTPPKT
Hve1 -APVCVAPVTMCLIIS-SKEMFNQQRKIDCVISGQDETTVNKFNITWQVNEAIRPTSEE-TSSGG-----IKTSTMRLSLHESLS--EVCSASNG-DVTVFQNLPPPPI
Hve2 -----APVTMCLIIS-SKEMFNQTKIDCVISGQDETTVNKFNITWQVNEAIRPTSEE-TSSGG-----IKTSTMRLSLHESLS--EVCSASNG-VTVFQNLPPPPI
Gac1 -----APITMELIRS-PKEMFNSQTKIDCVISGQDETTVKKFQITWLVNGAIRPTSEE--KNPGI----IKTSTMWLSLHDSLS--DVCSVSAKSD-DVTVQCNLPPPPI
* . . . * * * * * * * * * *

CH4 Domain

E. colioides DGSDPKV--TVHILPLEDIDKAAQGSSEVTLVCLVSSRAQDYIYAWEEDTGQNTGTYSQDGINFPPQK--IQNRYLVTSIYITTKDKWNTTKF--SCNVW-PAGGNKSMKSRDVSAMSN
S. aurata DGEKPKPK-VHVLQDNDLQEXDS---EITLVCLVTSRKLQDYIYAWESEHTGQKNGEYTDGINFPPQK--TKTYSVTSVYITTKKWN--THKVGCVVW-PAGSAVKVAPKEVSKAKGN
T. orientalis DRSKTTV--TVHILSEEEIRKNSD---VTLVCLVSSSVQDYIYAWSDDAGQNTGNVYDGIPTFPQK--TQHGYSVTSVYITINKEKWNQQRVFNQNVW-LVGSNKSMIIRGVSKATGN
G. aculeatus D-NEPKV--TVHVLPEEDVKEGA---EVTLLCLVSTREPQDFYIAWLEENGSNVSVDYNDGIDFPPSMK--TDERYSVTSLYTTSLKKNW-NFVKFTCKVWSPGG---GTQRDVSKMVF
S. chuatsi DGREPVK--TVHVLTEEDINKGA---EVTLLCLVSSPVLQDYIYAWESED---ETNIYTDGINFPPQK--TQHGYSVTSVYITTKKWN-KFNMFYCNVW-PAGSNDSEMPRGVSKSHGN
S. caurinus DESEPKV--TVHVLPEEDISSNPAK--VTLVCLVSSRLKTDYIYAWEHIGRDVSSYTDGIDSLPQK--NQDGYSVTSIYITTKDKW--KYTMFSCNAW-PATNNNSMRSDVSKAQNN
D. labrax DVSKTKV--TVHVLPEEV-NKEA---EVTLLCLVSPVLQDSYIYAWESENIEKSPGIYKDGINFPQK--TKDGYFVTSVYITTKAEWETPGNVFNCHVR-VAGSNSSMKSRAVSKAQGN
Bdi DQSKSKPKATVHILPEEDIGKASAD-VTLVCLVSSDVQDFYIAWTEQIRELVSNSYSDGINFPPQK--TQHGYSVTSLYTTTKKWNKLNK-MFSCVW-PAGNKEAMVQRQVSKAHSN
Ema1 GGNVPTV-A-VHTLPNNDIKEDTP---ITLVCLVSTVQRDYIYAWNIANKT--INYQDGDIFPAQK--SEKGYVTSIYITTKKWNKLNK--FSCIVWYPGSKENISD---VSSSMDT
Ema2 GGNVPTV-A-VHTLPNNDIKEDAP---ITLVCLVSTVQRDYIYAWNIANKT--INYQDGDIFPAQK--SEKGYVTSIYITTKKWNKLNK--FSCIVWYPGS.....
TbeS DGKEPKV--TVHIVPEEDID-ADPSDEVTLVCLVFSSEK-DYYIYAWAEHIGQNTPIYQDGDIDLPPQK--ANNFYFATSSYTTNKTQWQNH--MFSCNVW-PAGAKKAMAPRTVSNMNN
TbeL DGKEPKV--TVHIVPEEDID-ADPSDEVTLVCLVFSSEK-DYYIYAWAEHIGQNTPIYQDGDIDLPPQK--ANNFYFATSSYTTNKTQWQNH--MFSCNVW-PAGAKKAMAPRTVSNMNN
TbeSts DGKEPKV--TVHIVPEEDID-ADPSDEVTLVCLVFSSEK-DYYIYAWAEHIGQNTPIYQDGDIDLPPQK--ANNFYFATSSYTTNKTQWQNH--MFSCNVW-PAGAKKAMAPRTVSNMNN
Nco DGTDPKV--TVHILPEEDIN-AEPSNEVTLVCLVFSXXX-DYYIYAWTDYRGRKDTPIYLDGIDFPPQK--ANKGYFVTSIYITTKKWNKLNK--IISCNVW-PAGVKKPMITRTVSKMNN
Ggil DGTEPKV--TVHILPEEDIN-ADPSNEVTLVCLVFSVQDYIYAWAEYIEQNTPIYEDGIDFLPKRPTDKGYFVTSIFTNKTQWQSH--MFSCNVW-PAGVKKPMITRTVSNMNN
Hve1 DGTTPKV--TVHILPEEDIN-GNPS-EVTL.....
Hve2 DGTTPKV--TVHILPEEDIN-ENPS-EVTLVCLVFSVQDYIYAWAEHIGQNTPIYQDGINFPVNRPTDKGYSVTSIYITTKKWNKLNK--SFSCNVW-PAGVKKPMITRTVSNMNN
Gac1 DGTQPKV--TVHILPEEDID-ANPS-EVTLCL.....
* *

Secretory tail

```

E. coioides  SVECKK-
S. aurata   SHEC---
T. orientalis SLECDK-
G. aculeatus STECKES
S. chuatsi  SIECRK-
S. caurinus SIYCKK-
D. labrax   SLECDK-
Bdi         STECEHV
TbeS       SIECKK
TbeL       SIECKK
TbeSts     SIECKK
Nco        SIECKK
Ggi1       SIECKK
Hve2       SIECKK
* **

```

Membrane-bound

```

Bdi          STEYFTPESKVSFALSCTDDDIEE-DEYSSLWSTTTSFIFLFISSLVYNMIFTMVKMK
Ema1        ----SFECSKLGFHLSCTEGAIEE-DDYSSLWSTTLSFIFLFISLLYSMIFTMVKMK
Tbe         --SIEYLMSKLSSALSCTDEAIDEEDEYSSLWSTTLSFIFLFISLLYSMIFTMVKMK
G. aculeatus STES-FKLRVAMSCTDDAIDE-DEYSSLWSTTSFIFLFISSLISYSMVFSLVKMK
          . . .***. .*. * .***** *****.** *. *..****

```

Fig. 7. Multiple alignment of deduced amino acid sequences of the IgT CH domains from temperate, Antarctic and non-Antarctic species. Temperate Perciform species: *Epinephelus coioides*, orange-spotted grouper (GU182366), *Sparus aurata*, seabream (KX599200), *Thunnus orientalis*, Pacific bluefin tuna (KF713336), *Gasterosteus aculeatus*, three-spined stickleback [40], *Siniperca chuatsi*, mandarin fish (DQ016660), *Sebastes caurinus*, copper rockfish (GE798008), *Dicentrarchus labrax*, European seabass (KM410929); non-Antarctic species: *B. diacanthus* (Bdi, KP876590), *E. maclovinus* (Ema1 and Ema); Antarctic species: *T. bernacchii* (TbeS, TbeL and TbeSts variants), *N. coriiceps* (Nco, KP876589), *G. gibberifrons* (Ggi1), *H. velifer* (Hve1 and Hve2, both coding for same CH2 but differing in the CH3 domain length) and *G. acuticeps* (Gac1). Gaps are indicated by dashes. Identical amino acids are marked with an asterisk, conservative substitutions with colon, and semiconservative substitutions with a dot. Canonical cysteine and tryptophan residues are marked in bold. Extra cysteines are marked in bold and shaded in dark gray. The putative *N*-glycosylation sites are underlined. Asparagines predicted to be glycosylated are shown in red. Sequons containing a proline are in bold, underlined and shaded in gray. The duplicated 3-aa sequence APV at the beginning of the CH3 domain of the Hve1 variant is underlined. The membrane-bound form obtained only for *E. maclovinus* (Ema1) has been aligned with that of *B. diacanthus* (Bdi) and *T. bernacchii* (Tbe), previously obtained [19], and with that of *G. aculeatus* [58], the closest temperate species to the notothenioid species.




Fig. 8. Sites predicted to evolve under purifying and diversifying selection in the regions surrounding the remnant CH2 domain in Cryonotothenioidea. Sites with statistically significant ω values are indicated by blue (negative selection) and red (positive selection) boxes, as detected by FEL and MEME.




Fig. 9. Gene expression levels of IgM and IgT genes in *E. maclovinus* and *T. bernacchii*, calculated based on available RNA-seq data and reported as counts per million (CPM).




Fig. 10. Predicted 3D molecular model of the IgT CH1, remnant CH2, CH3 and CH4 domains from *E. maclovinus* (sequence Ema2), obtained with Phyre 2 (<http://www.sbg.bio.ic.ac.uk/phyre2>).

Highlights

- All analyzed cryonotothenioid species have an *IgT* gene with a *C τ 2* exonic remnant
- Conserved donor/acceptor splice sites allow the expression of the exonic remnant
- The exon loss event took place prior to the radiation of Cryonotothenioidea
- The exon loss is shared by the non-Antarctic sister lineage of Cryonotothenioidea
- the nearly total loss of the *C τ 2* exon occurred in a stepwise mode

CRedit authorship contribution statement

Alessia Ametrano: Conceptualization, Investigation, Formal analysis, Validation, Visualization, Writing - Review & Editing. **Marco Gerdol:** Conceptualization, Investigation, Formal analysis, Funding acquisition, Resources, Writing - Original Draft, Writing - Review & Editing Visualization. **Maria Vitale:** Investigation, Formal analysis, Writing - Review & Editing. **Samuele Greco:** Investigation, Formal analysis, Writing - Review & Editing. **Umberto Oreste:** Visualization, Writing - Review & Editing. **Maria Rosaria Coscia:** Conceptualization, Writing - Original Draft, Writing - Review & Editing, Supervision, Project administration, Funding acquisition.

All authors have read and agreed to the published version of the manuscript.