

Supplementary Material

Robinia pseudoacacia L. (black locust) leaflets as biomonitors of airborne microplastics

Mehriban Jafarova^{1*}, Lisa Grifoni^{1,2}, Monia Renzi³, Tecla Bentivoglio⁴, Serena Anselmi⁴, Aldo Winkler², Luigi Antonello Di Lella¹, Lilla Spagnuolo², Julian Aherne⁵, Stefano Loppi¹

- ¹ Department of Life Sciences, University of Siena, Siena 53100, Italy; mehriban.jafarova@student.unisi.it (M.J.); l.grifoni2@student.unisi.it (L.G.); luigi.dilella@unisi.it (L.A.L.); stefano.loppi@unisi.it (S.L.)
 - ² Istituto Nazionale di Geofisica e Vulcanologia, 00143 Rome, Italy; aldo.winkler@ingv.it (A.W.); lilla.spagnuolo@ingv.it (L.S.)
 - ³ Department of Life Science, University of Trieste, via L. Giorgieri, 10, Italy; mrenzi@units.it (M.R.)
 - ⁴ Bioscience Research Center, via Aurelia Vecchia, 32, Italy; tecla.bentivoglio@bsrc.it (T.B.); serena.anselmi@bsrc.it (S.A.)
 - ⁵ School of Environment, Trent University, Peterborough, ON K9L 0G2 Canada; jaherne@trentu.ca (J.A.)
- * Correspondence: mehriban.jafarova@student.unisi.it (M.J.)




Figure S1. FTIR spectra example of tire wear particles found during the polymer characterization using a spectral library with a match of 78.39 %.