



Periodic solutions of Hamiltonian systems coupling twist with generalized lower/upper solutions

Alessandro Fonda *, Wahid Ullah

Dipartimento di Matematica e Geoscienze, Università degli Studi di Trieste, P.le Europa 1, 34127 Trieste, Italy

Received 29 August 2022; revised 29 August 2023; accepted 4 October 2023

Abstract

The Hamiltonian systems considered in this paper are obtained by weakly coupling two systems having completely different behaviors. The first one satisfies the twist assumptions usually considered for the application of the Poincaré–Birkhoff Theorem, while the second one presents the existence of some well-ordered lower and upper solutions. In the higher dimensional case, we also treat a coupling situation where the classical Hartman condition is assumed.

© 2023 The Author(s). Published by Elsevier Inc. This is an open access article under the CC BY-NC-ND license (<http://creativecommons.org/licenses/by-nc-nd/4.0/>).

MSC: 34C25

Keywords: Hamiltonian systems; Periodic boundary value problem; Poincaré–Birkhoff theorem; Lower and upper solutions; Hartman condition

1. Main results in low dimension

In the first part of the paper we are interested in the periodic problem associated with a four-dimensional system of the type

* Corresponding author.

E-mail addresses: a.fonda@units.it (A. Fonda), wahid.ullah@phd.units.it (W. Ullah).

$$\begin{cases} \dot{q} = \partial_p H(t, q, p) + \varepsilon \partial_p P(t, q, p, u, v), \\ \dot{p} = -\partial_q H(t, q, p) - \varepsilon \partial_q P(t, q, p, u, v), \\ \dot{u} = f(t, v) + \varepsilon \partial_v P(t, q, p, u, v), \\ \dot{v} = g(t, u) - \varepsilon \partial_u P(t, q, p, u, v). \end{cases} \tag{1.1}$$

The idea is to consider a Poincaré–Birkhoff situation for the system

$$\dot{q} = \partial_p H(t, q, p), \quad \dot{p} = -\partial_q H(t, q, p), \tag{1.2}$$

and the existence of well-ordered lower/upper solutions for the system

$$\dot{u} = f(t, v), \quad \dot{v} = g(t, u). \tag{1.3}$$

The coupling function $P = P(t, q, p, u, v)$ will be assumed to have a bounded gradient with respect to (q, p, u, v) , and ε will be a small parameter. All functions involved are assumed to be continuous, and T -periodic in their first variable t .

In the second part of the paper we will extend our results to higher dimensional systems, both concerning the couple (q, p) and the couple (u, v) . To this aim, for the couple (q, p) we will apply some recent generalizations of the Poincaré–Birkhoff Theorem (see [12,13,18,19]), while for the couple (u, v) the treatment of lower and upper solutions will be based on two different situations. The first one comes from the recent papers [14,17], while the second approach involves a classical condition by Hartman [22].

Now, in order to better understand the spirit of our results, some historical hints may be useful.

Just three months before his death, in 1912, Poincaré published his paper [28] in which he conjectured the existence of at least two fixed points for an area-preserving homeomorphism of a planar circular annulus onto itself, such that the points of the inner circle Γ_1 are moved along Γ_1 in the clockwise sense and the points of the outer circle Γ_2 are moved along Γ_2 in the counter-clockwise sense. The existence of one fixed point was proved by Birkhoff the year later, while the proof of the existence of a second fixed point was provided by Birkhoff himself only in 1925 (see [4] for a modern exposition). The Poincaré–Birkhoff Theorem has then been generalized in several directions (see [15] and the references therein).

In 1983, one of the most brilliant results for the periodic problem associated with a Hamiltonian system was proved by Conley and Zehnder [7], giving a partial answer to a conjecture by Arnold [1,2]. They obtained the multiplicity of periodic solutions for a Hamiltonian system in \mathbb{R}^{2M} assuming the C^2 -smooth Hamiltonian function $H = H(t, q, p)$ to be periodic in t and in the space variables q_1, \dots, q_M , and quadratic in p on a neighborhood of infinity. They also mentioned a possible relation of their result with the Poincaré–Birkhoff Theorem. The results in [7] have been developed by different researchers in several directions (see, e.g., [5,9,21,23–25]).

Recently, a deeper relation between these results and the Poincaré–Birkhoff Theorem has been established by the first author and Ureña [18], replacing the quadratic assumption considered in [7] by a local “twist condition” on the solutions of the system. The first author then extended the results of [18] jointly with Gidoni [12], introducing a very general twist condition in order to find the periodic solutions. The same authors further extended the theory, in a second paper [13], to the case when the Hamiltonian function includes a nonresonant quadratic term. The possibility of resonance has also been studied in [6] by assuming some Ahmad–Lazer–Paul conditions.

On the other hand, the history of lower and upper solutions goes back to the pioneering work of Picard [27] in 1893. The first attempts towards a modern definition of lower and upper solutions were made by Scorza Dragoni [29] in 1939 for the following equation

$$\ddot{u} = g(t, u, \dot{u}). \tag{1.4}$$

A few years later Nagumo [26] provided the classical definition of lower solution α and upper solution β of (1.4) by assuming the inequalities

$$\ddot{\alpha}(t) \geq g(t, \alpha(t), \dot{\alpha}(t)), \quad \ddot{\beta}(t) \leq g(t, \beta(t), \dot{\beta}(t)).$$

He also introduced an extra assumption, which we nowadays call *Nagumo condition*, so to find the existence of a solution. We refer to [8] for more historical information and further developments of the theory.

The notion of lower and upper solutions has recently been extended in [14,17] to planar systems. Moreover, the first author together with Garzon and Sfecci [11] further extended this fertile theory to coupled systems which contain both the periodicity-twist conditions and a pair of well-ordered lower and upper solutions. However, due to some technical problems, they only used *constant* lower and upper solutions, while proposing as an open problem the case of non-constant lower/upper solutions.

In this paper, we provide a partial answer to this open problem and extend the theory to systems which contain the periodicity-twist conditions together with generalized well-ordered lower/upper solutions, coupled by a perturbation term.

The paper is organized as follows.

In Section 2 we state our result in the low dimensional case by coupling “twist” and strict lower/upper solutions. The proof of this result is given in Section 3. In Section 4 we provide some consequences of the main result and an example of application.

In Section 5 we extend our previous theorem to higher dimensions, and provide some variants and an example of application. In Section 6 we prove a result by coupling “twist” with a Hartman-type condition [22] in higher dimensions. This condition extends the concept of *constant* lower and upper solutions to higher dimensions, and has been extensively studied by many authors (see [10] and the references therein).

Finally, in Section 7 we illustrate an application to the theory of perturbations of completely integrable systems.

2. A first multiplicity result

Let us first recall what we know about systems (1.2) and (1.3), separately.

The Poincaré–Birkhoff Theorem. *Here are our assumptions concerning system (1.2).*

- A1. *The function $H(t, q, p)$ is 2π -periodic in q .*
- A2. *There are $a < b$ such that all the solutions (q, p) of system (1.2) starting with $p(0) \in [a, b]$ are defined on $[0, T]$ and*

$$\begin{cases} p(0) = a & \Rightarrow & q(T) - q(0) < 0, \\ p(0) = b & \Rightarrow & q(T) - q(0) > 0. \end{cases}$$

Notice that the original Poincaré–Birkhoff Theorem was stated for functions defined on a planar annulus. However, as explained in [4], it can equivalently be stated on the strip $[a, b] \times \mathbb{R}$, after a change of variables in suitable polar coordinates. In this setting, assumption A2 is usually called a “twist condition”.

The following result was proved in [18].

Theorem 2.1. *Assume that A1 and A2 hold true. Then, system (1.2) has at least two geometrically distinct T -periodic solutions (q, p) such that $p(0) \in]a, b[$.*

Notice that, when a T -periodic solution (q, p) has been found, infinitely many others appear by just adding an integer multiple of 2π to the q -th component. We say that two solutions are geometrically distinct if they cannot be obtained from each other in this way.

We also want to remark here that the period 2π in assumption A1 is inessential; any period would be possible.

Lower and upper solutions. Let us first recall the definitions of lower and upper solutions for the T -periodic problem associated with system (1.3).

Definition 2.2. A T -periodic C^1 -function $\alpha : \mathbb{R} \rightarrow \mathbb{R}$ is said to be a “lower solution” for the T -periodic problem associated with system (1.3) if there exists a T -periodic C^1 -function $v_\alpha : \mathbb{R} \rightarrow \mathbb{R}$ such that

$$\begin{cases} v < v_\alpha(t) & \Rightarrow & f(t, v) < \dot{\alpha}(t), \\ v > v_\alpha(t) & \Rightarrow & f(t, v) > \dot{\alpha}(t), \end{cases} \tag{2.1}$$

and

$$\dot{v}_\alpha(t) \geq g(t, \alpha(t)). \tag{2.2}$$

The lower solution is “strict” if the strict inequality in (2.2) holds.

Definition 2.3. A T -periodic C^1 -function $\beta : \mathbb{R} \rightarrow \mathbb{R}$ is said to be an “upper solution” for the T -periodic problem associated with system (1.3) if there exists a T -periodic C^1 -function $v_\beta : \mathbb{R} \rightarrow \mathbb{R}$ such that

$$\begin{cases} v < v_\beta(t) & \Rightarrow & f(t, v) < \dot{\beta}(t), \\ v > v_\beta(t) & \Rightarrow & f(t, v) > \dot{\beta}(t), \end{cases} \tag{2.3}$$

and

$$\dot{v}_\beta(t) \leq g(t, \beta(t)). \tag{2.4}$$

The upper solution is “strict” if the strict inequality in (2.4) holds.

Notice that, when $f(t, v) = v$, the above definitions reduce to the classical ones for the second order equation (1.4), by choosing $v_\alpha = \dot{\alpha}$ and $v_\beta = \dot{\beta}$.

The following result was proved in [14,17].

Theorem 2.4. Assume that there exist a lower solution α and an upper solution β for the T -periodic problem associated with system (1.3), such that $\alpha \leq \beta$. Then, system (1.3) has a T -periodic solution (u, v) such that $\alpha \leq u \leq \beta$.

Back to the coupled system. Let us state our hypotheses. We will assume A1, A2, and

A3. There exist a strict lower solution α and a strict upper solution β for the T -periodic problem associated with system (1.3), such that $\alpha \leq \beta$.

Moreover, we need the function $f(t, v)$ to be smooth and strictly increasing in its second variable, precisely as follows.

A4. The second order partial derivatives $\partial_{tv}^2 f(t, v)$ and $\partial_{vv}^2 f(t, v)$ exist and are continuous; moreover, there exists $\lambda > 0$ such that

$$\partial_v f(t, v) \geq \lambda, \quad \text{for every } (t, v) \in [0, T] \times \mathbb{R}.$$

Concerning the function $P(t, q, p, u, v)$, besides its periodicity in t and q , we also need some smoothness condition, as specified in the next assumption.

A5. The function $P(t, q, p, u, v)$ is 2π -periodic in q and has a bounded gradient with respect to (q, p, u, v) ; moreover, the partial derivative $\partial_v P$ is independent of q and p , and the map $\partial_v P(t, u, v)$ is continuously differentiable.

Here is the main result of this section.

Theorem 2.5. Assume that A1 – A5 hold true. Then there exists $\bar{\varepsilon} > 0$ such that, if $|\varepsilon| \leq \bar{\varepsilon}$, there are at least two geometrically distinct T -periodic solutions of system (1.1), with $p(0) \in]a, b[$ and $\alpha \leq u \leq \beta$.

The proof of the Theorem 2.5 will be given in Section 3.

Remark 2.6. Theorem 2.5 provides a partial answer to an open problem raised in [11], where only constant lower and upper solutions were considered. However, our result only applies to weakly coupled systems with strict lower and upper solutions. Hence, the problem raised in [11] remains open.

Remark 2.7. As already noticed in [18], instead of using a constant interval $[a, b]$, it is possible to deal with a varying interval $[a(q), b(q)]$, where $a, b : \mathbb{R} \rightarrow \mathbb{R}$ are continuous and 2π -periodic functions. Indeed, if a and b are continuously differentiable, then this case can be reduced to the previous one by the symplectic change of variables

$$\psi(q, p) = \left(\int_0^q \frac{b(s) - a(s)}{2} ds, \frac{2p - b(q) - a(q)}{b(q) - a(q)} \right).$$

On the other hand, if the functions a and b are only continuous, then by the Fejer Theorem they can be replaced by smooth functions. Notice that the new Hamiltonian $\tilde{H}(t, \tilde{q}, \tilde{p}) = H(t, \psi^{-1}(\tilde{q}, \tilde{p}))$ is periodic in \tilde{q} with period $\tau := \frac{1}{2} \int_0^{2\pi} (b(s) - a(s))ds$.

Before going to the proof, we now present a variant of Theorem 2.5 which is more related to Poincaré-Birkhoff Theorem as originally stated by Poincaré [28].

We first recall the definition of “rotation number”. Assume that $t_1 < t_2$ and let $\phi : [t_1, t_2] \rightarrow \mathbb{R}^2$ be a continuous curve such that $\phi(t) \neq (0, 0)$ for every $t \in [t_1, t_2]$. Writing $\phi(t) = \rho(t)(\cos \theta(t), \sin \theta(t))$, where $\rho : \mathbb{R} \rightarrow]0, +\infty[$ and $\theta : \mathbb{R} \rightarrow \mathbb{R}$ are continuous, we define

$$\text{Rot}(\phi; [t_1, t_2]) = -\frac{\theta(t_2) - \theta(t_1)}{2\pi}.$$

In the sequel, $\mathcal{D}(\Gamma)$ denotes the open bounded region delimited by a planar Jordan curve Γ . Here is the statement.

Theorem 2.8. *Let assumptions A3 – A5 hold true. Let k be any integer and assume that there exist $\rho > 0$ and two planar Jordan curves Γ_1, Γ_2 , strictly star-shaped with respect to the origin, with*

$$0 \in \mathcal{D}(\Gamma_1) \subseteq \overline{\mathcal{D}(\Gamma_1)} \subseteq \mathcal{D}(\Gamma_2),$$

such that the solutions of system (1.2) starting with $(q(0), p(0)) \in \overline{\mathcal{D}(\Gamma_2)} \setminus \mathcal{D}(\Gamma_1)$ are defined on $[0, T]$ and satisfy

$$(q(t), p(t)) \neq (0, 0), \quad \text{for every } t \in [0, T];$$

moreover,

$$\begin{cases} (q(0), p(0)) \in \Gamma_1 & \Rightarrow \text{Rot}((q, p); [0, T]) < k, \\ (q(0), p(0)) \in \Gamma_2 & \Rightarrow \text{Rot}((q, p); [0, T]) > k. \end{cases} \tag{2.5}$$

Then system (1.1) has at least two T -periodic solutions (q, p, u, v) such that

$$\alpha \leq u \leq \beta, \\ (q(0), p(0)) \in \mathcal{D}(\Gamma_2) \setminus \overline{\mathcal{D}(\Gamma_1)},$$

and

$$\text{Rot}((q, p); [0, T]) = k.$$

The same is true if (2.5) is replaced by the following:

$$\begin{cases} (q(0), p(0)) \in \Gamma_1 & \Rightarrow \text{Rot}((q, p); [0, T]) > k, \\ (q(0), p(0)) \in \Gamma_2 & \Rightarrow \text{Rot}((q, p); [0, T]) < k. \end{cases}$$

In the above theorem, the Hamiltonian function H is not assumed to be periodic in the variable q . The 2π -periodicity can indeed be recovered when passing to some kind of polar coordinates. The proof is almost the same as in [11, Theorem 10], so we omit it, for brevity.

3. Proof of Theorem 2.5

Let $A = \min \alpha$ and $B = \max \beta$. Then there exists a constant $C > 0$ such that

$$|g(t, u)| \leq C, \quad \text{for every } (t, u) \in [0, T] \times [A, B].$$

Moreover, by A4, there exist positive constants c, d such that $|\dot{\alpha}(t)| < c, |\dot{\beta}(t)| < c$ for every $t \in [0, T]$, and

$$\begin{cases} f(t, v) \geq c, & \text{for } v \geq d, \\ f(t, v) \leq -c, & \text{for } v \leq -d. \end{cases} \tag{3.1}$$

We can find two straight lines $\gamma_{\pm} : \mathbb{R} \rightarrow \mathbb{R}$, whose equations are

$$\gamma_+(u) = \mu u + R, \quad \gamma_-(u) = \mu u - R,$$

where $\mu < -C/c$ and $R > 0$ are chosen in such a way that

$$\gamma_-(u) < -d < d < \gamma_+(u),$$

and

$$\gamma_-(u) \leq \dot{\alpha}(t), v_{\alpha}(t), \dot{\beta}(t), v_{\beta}(t) \leq \gamma_+(u), \tag{3.2}$$

for every $(t, u) \in [0, T] \times [A, B]$.

Let us define the set

$$\mathcal{V} = \{(t, q, p, u, v) \in \mathbb{R}^5 : \alpha(t) \leq u \leq \beta(t), \gamma_-(u) \leq v \leq \gamma_+(u)\}.$$

We can choose a constant $\hat{d} > \max\{c, d, C/|\mu|\}$ such that

$$-\hat{d} < \gamma_-(u) < \gamma_+(u) < \hat{d}, \quad \text{for every } u \in [A, B].$$

Consider the function $\eta : \mathbb{R} \times \mathbb{R} \rightarrow \mathbb{R}$, defined as

$$\eta(t, u) = \begin{cases} \alpha(t), & \text{if } u \leq \alpha(t), \\ u, & \text{if } \alpha(t) \leq u \leq \beta(t), \\ \beta(t) & \text{if } u \geq \beta(t). \end{cases}$$

Now define the functions

$$\tilde{g}(t, u) = g(t, \eta(t, u)) - \eta(t, u) + u, \tag{3.3}$$

and

$$\tilde{f}(t, v) = \begin{cases} v, & \text{if } v \leq -\hat{d} - 1, \\ f(t, v) - (v + \hat{d})(v - f(t, v)), & \text{if } -\hat{d} - 1 \leq v \leq -\hat{d}, \\ f(t, v), & \text{if } -\hat{d} \leq v \leq \hat{d}, \\ f(t, v) + (v - \hat{d})(v - f(t, v)), & \text{if } \hat{d} \leq v \leq \hat{d} + 1, \\ v, & \text{if } v \geq \hat{d} + 1. \end{cases} \tag{3.4}$$

By A3, there exists a $\xi > 0$ such that

$$\dot{v}_\alpha(t) - g(t, \alpha(t)) > \xi, \quad \text{for every } t \in [0, T], \tag{3.5}$$

$$\dot{v}_\beta(t) - g(t, \beta(t)) < -\xi, \quad \text{for every } t \in [0, T]. \tag{3.6}$$

By the global existence assumption in A2, we note that there exists a constant $C_1 > 0$ such that, for any solution (q, p) of (1.2) starting with $p(0) \in [a, b]$, one has that

$$|p(t)| \leq C_1, \quad \text{for every } t \in [0, T].$$

Let $\sigma : \mathbb{R} \rightarrow \mathbb{R}$ be a C^∞ -function such that

$$\sigma(s) = \begin{cases} 1, & \text{if } s \leq C_1, \\ 0, & \text{if } s > C_1 + 1, \end{cases} \tag{3.7}$$

and set $\widehat{H}(t, q, p) = \sigma(|p|)H(t, q, p)$. Then \widehat{H} has a bounded gradient with respect to (q, p) . Now consider the modified system

$$\begin{cases} \dot{q} = \partial_p \widehat{H}(t, q, p) + \varepsilon \partial_p P(t, q, p, u, v), \\ \dot{p} = -\partial_q \widehat{H}(t, q, p) - \varepsilon \partial_q P(t, q, p, u, v), \\ \dot{u} = \tilde{f}(t, v) + \varepsilon \partial_v P(t, q, p, u, v), \\ \dot{v} = \tilde{g}(t, u) - \varepsilon \partial_u P(t, q, p, u, v), \end{cases} \tag{3.8}$$

where the new Hamiltonian function is defined as

$$\tilde{H}(t, q, p, u, v) = \widehat{H}(t, q, p) + \int_0^v \tilde{f}(t, s) ds - \int_0^u \tilde{g}(t, s) ds + \varepsilon P(t, q, p, u, v).$$

We can also write the modified system (3.8) as $\dot{z} = J \nabla \tilde{H}(t, z)$, where $J = \begin{pmatrix} 0 & -1 \\ 1 & 0 \end{pmatrix}$ is the standard symplectic matrix, and $z = (q, p, u, v)$. Notice that

$$\tilde{H}(t, z) = \frac{1}{2}(v^2 - u^2) + K(t, z),$$

where K is a function having a bounded gradient with respect to z . Moreover, by A2, since $\partial_p P$ and $\partial_q P$ are bounded, if $|\varepsilon|$ is small enough, for any solution (q, p, u, v) of (3.8) one still has that

$$\begin{cases} p(0) = a & \Rightarrow & q(T) - q(0) < 0, \\ p(0) = b & \Rightarrow & q(T) - q(0) > 0. \end{cases}$$

Then, by [13, Corollary 2.4], we conclude that the modified system (3.8) has at least two geometrically distinct T -periodic solutions such that $p(0) \in]a, b[$, provided that $|\varepsilon|$ is small enough.

We now need to show that such solutions z are such that $(t, z(t)) \in \mathcal{V}$ for every $t \in [0, T]$. Let us first prove the following five lemmas.

Lemma 3.1. *If $|\varepsilon|$ is small enough, there exist v_α^ε and v_β^ε such that*

$$f(t, v_\alpha^\varepsilon(t)) + \varepsilon \partial_v P(t, \alpha(t), v_\alpha^\varepsilon(t)) = \dot{\alpha}(t), \tag{3.9}$$

$$f(t, v_\beta^\varepsilon(t)) + \varepsilon \partial_v P(t, \beta(t), v_\beta^\varepsilon(t)) = \dot{\beta}(t), \tag{3.10}$$

for every $t \in [0, T]$. Moreover,

$$\lim_{\varepsilon \rightarrow 0} v_\alpha^\varepsilon = v_\alpha, \quad \lim_{\varepsilon \rightarrow 0} \dot{v}_\alpha^\varepsilon = \dot{v}_\alpha, \quad \lim_{\varepsilon \rightarrow 0} v_\beta^\varepsilon = v_\beta, \quad \lim_{\varepsilon \rightarrow 0} \dot{v}_\beta^\varepsilon = \dot{v}_\beta,$$

uniformly in $[0, T]$, i.e., $v_\alpha^\varepsilon \rightarrow v_\alpha$ and $v_\beta^\varepsilon \rightarrow v_\beta$ in $C^1([0, T], \mathbb{R})$, as $\varepsilon \rightarrow 0$.

Proof. We only prove the statement concerning v_α^ε , since the one for v_β^ε can be proved in a similar way. Consider the space $X = C^1([0, T], \mathbb{R})$. By A4 and A5, the functions f and $\partial_v P$ are continuously differentiable, and since

$$f(t, v_\alpha(t)) = \dot{\alpha}(t), \quad \text{for every } t \in \mathbb{R},$$

we have that $\alpha \in C^2([0, T], \mathbb{R})$. We can then define a function $\tilde{F} : X \times \mathbb{R} \rightarrow X$ by

$$\tilde{F}(v, \varepsilon)(t) = f(t, v(t)) + \varepsilon \partial_v P(t, \alpha(t), v(t)) - \dot{\alpha}(t). \tag{3.11}$$

Now, clearly $\tilde{F}(v_\alpha, 0) = 0$, and for all $h \in X$, we have

$$\begin{aligned} \frac{\partial \tilde{F}}{\partial v}(v_\alpha, 0)(h)(t) &= \lim_{\sigma \rightarrow 0} \frac{\tilde{F}(v_\alpha + \sigma h, 0) - \tilde{F}(v_\alpha, 0)}{\sigma}(t) \\ &= \lim_{\sigma \rightarrow 0} \frac{f(t, v_\alpha(t) + \sigma h(t)) - f(t, v_\alpha(t))}{\sigma} \\ &= \partial_v f(t, v_\alpha(t))h(t). \end{aligned}$$

Let us prove that \tilde{F} is differentiable with respect to its first variable at $(v_\alpha, 0)$, with

$$[d_v \tilde{F}(v_\alpha, 0)(h)](t) = \partial_v f(t, v_\alpha(t))h(t).$$

Writing

$$\tilde{F}(v, 0) = \tilde{F}(v_\alpha, 0) + d_v \tilde{F}(v_\alpha, 0)(v - v_\alpha) + r(v), \tag{3.12}$$

we need to prove that

$$v \xrightarrow{C^1} v_\alpha \implies \frac{r(v)}{\|v - v_\alpha\|_{C^1}} \xrightarrow{C^1} 0. \tag{3.13}$$

Substituting (3.11) in (3.12), we obtain

$$f(t, v(t)) = f(t, v_\alpha(t)) + \partial_v f(t, v_\alpha(t))(v(t) - v_\alpha(t)) + r(v)(t).$$

By the Lagrange Mean Value Theorem, for every $t \in [0, T]$ there exists $\zeta(t) \in [v_\alpha(t), v(t)]$ such that

$$f(t, v(t)) - f(t, v_\alpha(t)) = \partial_v f(t, \zeta(t))(v(t) - v_\alpha(t)).$$

Then,

$$\begin{aligned} \frac{|r(v)(t)|}{\|v - v_\alpha\|_{C^1}} &= |\partial_v f(t, \zeta(t)) - \partial_v f(t, v_\alpha(t))| \frac{|v(t) - v_\alpha(t)|}{\|v - v_\alpha\|_{C^1}} \\ &\leq |\partial_v f(t, \zeta(t)) - \partial_v f(t, v_\alpha(t))|, \quad \text{for every } t \in [0, T]. \end{aligned}$$

If $v \rightarrow v_\alpha$ in C^1 , then $v \rightarrow v_\alpha$ uniformly, hence also $\zeta \rightarrow v_\alpha$ uniformly. Since the partial derivative of f with respect to v is continuous, taking a constant $M > \|\partial_v f\|_\infty$, the map $\partial_v f : [0, T] \times [-M, M] \rightarrow \mathbb{R}$ is uniformly continuous. It then follows that, if $v \rightarrow v_\alpha$ in C^1 , then

$$\frac{r(v)(t)}{\|v - v_\alpha\|_{C^1}} \rightarrow 0, \quad \text{uniformly for } t \in [0, T].$$

It remains to be proved that, if $v \rightarrow v_\alpha$ in C^1 , then

$$\frac{d}{dt} \left(\frac{r(v)(t)}{\|v - v_\alpha\|_{C^1}} \right) \rightarrow 0, \quad \text{uniformly for } t \in [0, T].$$

We have

$$\begin{aligned} \frac{d}{dt} r(v)(t) &= \partial_t f(t, v(t)) + \partial_v f(t, v(t))\dot{v}(t) - \partial_t f(t, v_\alpha(t)) - \partial_v f(t, v_\alpha(t))\dot{v}_\alpha(t) \\ &\quad - \left(\partial_{tv}^2 f(t, v_\alpha(t)) + \partial_{vv}^2 f(t, v_\alpha(t))\dot{v}_\alpha(t) \right) (v(t) - v_\alpha(t)) \\ &\quad - \partial_v f(t, v_\alpha(t))(\dot{v}(t) - \dot{v}_\alpha(t)) \\ &= \left(\partial_t f(t, v(t)) - \partial_t f(t, v_\alpha(t)) - \partial_{tv}^2 f(t, v_\alpha(t))(v(t) - v_\alpha(t)) \right) \\ &\quad + (\partial_v f(t, v(t)) - \partial_v f(t, v_\alpha(t)))\dot{v}(t) - \partial_{vv}^2 f(t, v_\alpha(t))\dot{v}_\alpha(t)(v(t) - v_\alpha(t)). \end{aligned}$$

Again by using the Lagrange Mean Value Theorem twice, for every $t \in [0, T]$ there exist $\xi(t)$ and $\eta(t)$ in $[v_\alpha(t), v(t)]$ such that

$$\partial_t f(t, v(t)) - \partial_t f(t, v_\alpha(t)) = \partial_{tv}^2 f(t, \xi(t))(v(t) - v_\alpha(t)), \tag{3.14}$$

$$\partial_v f(t, v(t)) - \partial_v f(t, v_\alpha(t)) = \partial_{vv}^2 f(t, \eta(t))(v(t) - v_\alpha(t)). \tag{3.15}$$

Then,

$$\begin{aligned} \frac{d}{dt}r(v(t)) &= \left(\partial_{tv}^2 f(t, \xi(t)) - \partial_{tv}^2 f(t, v_\alpha(t)) \right) (v(t) - v_\alpha(t)) \\ &\quad + \left(\partial_{vv}^2 f(t, \eta(t))\dot{v}(t) - \partial_{vv}^2 f(t, v_\alpha(t))\dot{v}_\alpha(t) \right) (v(t) - v_\alpha(t)). \end{aligned}$$

If $v \rightarrow v_\alpha$ in C^1 , the first term in the sum converges to 0 when divided by $\|v - v_\alpha\|_{C^1}$, uniformly on $[0, T]$, by the continuity of $\partial_{tv}^2 f$. For the second term, we have

$$\begin{aligned} &\left| \partial_{vv}^2 f(t, \eta(t))\dot{v}(t) - \partial_{vv}^2 f(t, v_\alpha(t))\dot{v}_\alpha(t) \right| \leq \\ &\leq \left| \partial_{vv}^2 f(t, \eta(t)) - \partial_{vv}^2 f(t, v_\alpha(t)) \right| |\dot{v}(t)| + \left| \partial_{vv}^2 f(t, v_\alpha(t)) \right| |\dot{v}(t) - \dot{v}_\alpha(t)|, \end{aligned}$$

which converges uniformly to 0 when $v \rightarrow v_\alpha$ in C^1 , since both $|\dot{v}(t)|$ and $\left| \partial_{vv}^2 f(t, v_\alpha(t)) \right|$ are bounded, $\dot{v} \rightarrow \dot{v}_\alpha$ uniformly on $[0, T]$, and the map $\partial_{vv}^2 f$ is continuous.

We have thus proved (3.13). Therefore,

$$d_v \tilde{F}(v_\alpha, 0) = \partial_v f(\cdot, v_\alpha(\cdot)) \text{Id},$$

where $\text{Id} : X \rightarrow X$ is the identity map. By A4, we have that $\partial_v f(t, v_\alpha(t)) > 0$, for every $t \in [0, T]$, so the map $d_v \tilde{F}(v_\alpha, 0) : X \rightarrow X$ is invertible.

By the Implicit Function Theorem, there exists an $\bar{\varepsilon} > 0$ and a map $\varphi :] - \bar{\varepsilon}, \bar{\varepsilon}[\rightarrow B_X(v_\alpha, \bar{\varepsilon})$, of class C^1 , such that, for every $\varepsilon \in] - \bar{\varepsilon}, \bar{\varepsilon}[$ and $v \in B_X(v_\alpha, \bar{\varepsilon})$,

$$\tilde{F}(v, \varepsilon) = 0 \iff v = \varphi(\varepsilon).$$

Setting $v_\alpha^\varepsilon = \varphi(\varepsilon)$, the proof is completed. \square

Lemma 3.2. *There exists $\tilde{\varepsilon} > 0$ such that, if $|\varepsilon| < \tilde{\varepsilon}$, then for every $t \in [0, T]$ and $u \in [A, B]$ the following inequalities hold:*

$$\begin{cases} \tilde{f}(t, v) + \varepsilon \partial_v P(t, u, v) < \dot{\alpha}(t), & \text{if } v < v_\alpha^\varepsilon(t), \\ \tilde{f}(t, v) + \varepsilon \partial_v P(t, u, v) > \dot{\alpha}(t), & \text{if } v > v_\alpha^\varepsilon(t), \end{cases} \tag{3.16}$$

$$\begin{cases} \tilde{f}(t, v) + \varepsilon \partial_v P(t, u, v) < \dot{\beta}(t), & \text{if } v < v_\beta^\varepsilon(t), \\ \tilde{f}(t, v) + \varepsilon \partial_v P(t, u, v) > \dot{\beta}(t), & \text{if } v > v_\beta^\varepsilon(t), \end{cases} \tag{3.17}$$

$$\begin{cases} \tilde{g}(t, u) - \varepsilon \partial_u P(t, q, p, u, v) < \dot{v}_\alpha^\varepsilon(t), & \text{if } u \leq \alpha(t), \\ \tilde{g}(t, u) - \varepsilon \partial_u P(t, q, p, u, v) > \dot{v}_\beta^\varepsilon(t), & \text{if } u \geq \beta(t). \end{cases} \tag{3.18}$$

Proof. We only prove the first inequality in (3.16), the proof of the second inequality in (3.16) and of the inequalities in (3.17) being similar.

We first want to prove that, for $|\varepsilon|$ small enough, we have

$$v < v_\alpha^\varepsilon(t) \implies f(t, v) + \varepsilon \partial_v P(t, u, v) < \dot{\alpha}(t). \tag{3.19}$$

By A4, there exists $\lambda > 0$ such that $\partial_v f(t, v_\alpha^\varepsilon(t)) \geq \lambda$, and by A5 there exists a constant $\widehat{C} > 0$ such that

$$\left| \partial_{vv}^2 P(t, u, v_\alpha(t)) \right| \leq \widehat{C}, \quad \text{for every } (t, u) \in [0, T] \times [A, B].$$

So, if $2|\varepsilon|\widehat{C} < \lambda$, we have

$$\partial_v (f(t, v_\alpha^\varepsilon(t)) + \varepsilon \partial_v P(t, u, v_\alpha^\varepsilon(t))) \geq \frac{\lambda}{2}, \quad \text{for every } (t, u) \in [0, T] \times [A, B].$$

By continuity, there exists a $\bar{\delta} > 0$ such that

$$|v - v_\alpha^\varepsilon(t)| < \bar{\delta} \implies \partial_v (f(t, v) + \varepsilon \partial_v P(t, u, v)) \geq \frac{\lambda}{4},$$

for every $(t, u) \in [0, T] \times [A, B]$. So, by (3.9), there exists $\tau > 0$ such that

$$\begin{aligned} v \in [v_\alpha^\varepsilon(t) - \tau, v_\alpha^\varepsilon(t)] &\implies f(t, v) + \varepsilon \partial_v P(t, u, v) < \dot{\alpha}(t), \\ v \in [v_\alpha^\varepsilon(t), v_\alpha^\varepsilon(t) + \tau] &\implies f(t, v) + \varepsilon \partial_v P(t, u, v) > \dot{\alpha}(t). \end{aligned}$$

Without loss of generality, we can assume

$$-d < \dot{\alpha}(t), v_\alpha(t), \dot{\beta}(t), v_\beta(t) < d,$$

where the constant d is as in (3.1), and take $|\varepsilon|, \tau$ small enough so that

$$-d < v_\alpha^\varepsilon(t) - \tau < v_\alpha^\varepsilon(t) + \tau < d.$$

By (2.1) and (3.1), there exists $\varrho > 0$ such that

$$\begin{aligned} f(t, v) - \dot{\alpha}(t) &\leq -\varrho, \quad \text{for } v \leq -d, \\ f(t, v) - \dot{\alpha}(t) &\geq \varrho, \quad \text{for } v \geq d. \end{aligned}$$

If $|\varepsilon|$ is small enough, since $\partial_v P$ is bounded, we have that

$$\begin{aligned} f(t, v) + \varepsilon \partial_v P(t, u, v) - \dot{\alpha}(t) &\leq -\frac{\varrho}{2}, \quad \text{for } v \leq -d, \\ f(t, v) + \varepsilon \partial_v P(t, u, v) - \dot{\alpha}(t) &\geq \frac{\varrho}{2}, \quad \text{for } v \geq d. \end{aligned}$$

Now it remains only to check what happens in the intervals $[v_\alpha^\varepsilon(t) + \tau, d]$ and $[-d, v_\alpha^\varepsilon(t) - \tau]$. Let us only consider the first interval, the argument being similar for the other one. If $v \in [v_\alpha^\varepsilon(t) + \tau, d]$, then, if $|\varepsilon|$ is small enough, by Lemma 3.1 and (2.1), using (3.1), we have

$$f(t, v) \geq f(t, v_\alpha^\varepsilon(t) + \tau) \geq f\left(t, v_\alpha(t) + \frac{\tau}{2}\right) > \dot{\alpha}(t).$$

By Weierstrass Theorem, there exists a $m > 0$ such that

$$f(t, v) - \dot{\alpha}(t) \geq m,$$

for every $t \in [0, T]$ and $v \in [v_\alpha^\varepsilon(t) + \tau, d]$. Then, for $|\varepsilon|$ small enough,

$$f(t, v) + \varepsilon \partial_v P(t, u, v) - \dot{\alpha}(t) > 0,$$

for every $t \in [0, T]$ and $v \in [v_\alpha^\varepsilon(t) + \tau, d]$. We have thus proved (3.19).

Now, since $-\hat{d} < v_\alpha^\varepsilon(t) < \hat{d}$ for $|\varepsilon|$ small enough, we have the following three cases.

Case 1. If $-\hat{d} \leq v < v_\alpha^\varepsilon(t)$, then by (3.4) and (3.19) we have

$$\tilde{f}(t, v) + \varepsilon \partial_v P(t, u, v) = f(t, v) + \varepsilon \partial_v P(t, u, v) < \dot{\alpha}(t).$$

Case 2. If $v \leq -\hat{d} - 1$, then by (3.4) we have

$$\begin{aligned} \tilde{f}(t, v) + \varepsilon \partial_v P(t, u, v) &= v + \varepsilon \partial_v P(t, u, v) \\ &\leq -\hat{d} + \varepsilon \partial_v P(t, u, v) < \dot{\alpha}(t), \end{aligned}$$

for $|\varepsilon|$ small enough, since $\partial_v P$ is bounded.

Case 3. If $-\hat{d} - 1 \leq v < -\hat{d}$, then by (3.4) and (2.1) we have

$$\begin{aligned} \tilde{f}(t, v) + \varepsilon \partial_v P(t, u, v) &= f(t, v) - (v + \hat{d})(v - f(t, v)) + \varepsilon \partial_v P(t, u, v) \\ &= (1 + (v + \hat{d}))f(t, v) - (v + \hat{d})v + \varepsilon \partial_v P(t, u, v) \\ &< \dot{\alpha}(t), \end{aligned}$$

for $|\varepsilon|$ small enough, since $-(v + \hat{d}) \in [0, 1]$ and $f(t, v) < \dot{\alpha}(t)$, $v < \dot{\alpha}(t)$.

The proof of the first inequality in (3.16) is thus completed.

We now prove the first inequality in (3.18), the second one being analogous. Suppose $u \leq \alpha(t)$. By (3.3) and (3.5), we have

$$\begin{aligned} \tilde{g}(t, u) - \varepsilon \partial_u P(t, q, p, u, v) &= g(t, \alpha(t)) - \alpha(t) + u - \varepsilon \partial_u P(t, q, p, u, v) \\ &\leq g(t, \alpha(t)) - \varepsilon \partial_u P(t, q, p, u, v) \\ &< \dot{v}_\alpha(t) - \xi - \varepsilon \partial_u P(t, q, p, u, v) \\ &< \dot{v}_\alpha^\varepsilon(t), \end{aligned}$$

for $|\varepsilon|$ small enough, since $\dot{v}_\alpha^\varepsilon \rightarrow \dot{v}_\alpha$ uniformly. The proof of the first inequality in (3.18) is thus completed. \square

Let us define the open sets

$$\begin{aligned} A_{NW} &= \{(t, u, v) \in \mathbb{R}^3 : u < \alpha(t), v > v_\alpha^\varepsilon(t)\}, \\ A_{SW} &= \{(t, u, v) \in \mathbb{R}^3 : u < \alpha(t), v < v_\alpha^\varepsilon(t)\}, \\ A_{NE} &= \{(t, u, v) \in \mathbb{R}^3 : u > \beta(t), v > v_\beta^\varepsilon(t)\}, \\ A_{SE} &= \{(t, u, v) \in \mathbb{R}^3 : u > \beta(t), v < v_\beta^\varepsilon(t)\}. \end{aligned}$$

Lemma 3.3. *For every solution $z = (q, p, u, v)$ of system (3.8), the following assertions hold true:*

$$\begin{aligned} (t_0, u(t_0), v(t_0)) \in A_{NW} &\Rightarrow (t, u(t), v(t)) \in A_{NW} \text{ for every } t < t_0, \\ (t_0, u(t_0), v(t_0)) \in A_{SE} &\Rightarrow (t, u(t), v(t)) \in A_{SE} \text{ for every } t < t_0, \\ (t_0, u(t_0), v(t_0)) \in A_{NE} &\Rightarrow (t, u(t), v(t)) \in A_{NE} \text{ for every } t > t_0, \\ (t_0, u(t_0), v(t_0)) \in A_{SW} &\Rightarrow (t, u(t), v(t)) \in A_{SW} \text{ for every } t > t_0. \end{aligned}$$

Proof. We only prove the first assertion, since the remaining ones can be proved similarly. We suppose on contrary that there exists $t_1 < t_0$ such that

$$\begin{aligned} (t_0, u(t_0), v(t_0)) &\in A_{NW}, \\ (t, u(t), v(t)) &\in A_{NW}, \quad \text{for } t \in]t_1, t_0[, \end{aligned}$$

and

$$(t_1, u(t_1), v(t_1)) \in \partial A_{NW}.$$

Notice that

$$\begin{aligned} \partial A_{NW} &= \{t, u, v\} \in \mathbb{R}^3 : u = \alpha(t), v \geq v_\alpha^\varepsilon(t)\} \\ &\cup \{t, u, v\} \in \mathbb{R}^3 : u \leq \alpha(t), v = v_\alpha^\varepsilon(t)\}. \end{aligned} \tag{3.20}$$

Assume $v(t_1) > v_\alpha^\varepsilon(t_1)$. Without loss of generality, we may assume that there exists $\delta > 0$ such that $[t_1, t_1 + \delta] \subseteq]t_1, t_0[$ and $v(t) > v_\alpha^\varepsilon(t)$ for every $t \in [t_1, t_1 + \delta]$. Now define $w : [t_1, t_1 + \delta] \rightarrow \mathbb{R}$ by $w(t) = u(t) - \alpha(t)$. Then, we have that $w(t_1 + \delta) < 0$ and, by Lemma 3.2,

$$\dot{w}(t) = \dot{u}(t) - \dot{\alpha}(t) = \tilde{f}(t, v) + \varepsilon \partial_v P(t, u(t), v(t)) - \dot{\alpha}(t) > 0,$$

for every $t \in]t_1, t_1 + \delta]$ and $|\varepsilon|$ small enough. Hence, $w(t_1) < 0$, implying that $u(t) < \alpha(t)$ for every $t \in [t_1, t_1 + \delta]$. Then, by (3.20), we necessarily have that $v(t_1) = v_\alpha^\varepsilon(t_1)$. Now if we define the map $G(t) = v(t) - v_\alpha^\varepsilon(t)$, then G is continuous on $[t_1, t_0]$, $G(t_1) = 0$ and $G(t) > 0$ for every $t \in]t_1, t_0]$. But then, using (3.18) and the fact that $u(t) \leq \alpha(t)$ for every $t \in [t_1, t_0]$, we have

$$\dot{G}(t_1) = \dot{v}(t_1) - \dot{v}_\alpha^\varepsilon(t_1) = \tilde{g}(t_1, u(t_1)) - \varepsilon \partial_u P(t, q, p, u, v) - \dot{v}_\alpha^\varepsilon(t_1) < 0,$$

for $|\varepsilon|$ small enough; a contradiction. \square

We now define the sets

$$A_W = \{(t, u, v) \in \mathbb{R}^3 : u < \alpha(t), v = v_\alpha^\varepsilon(t)\},$$

$$A_E = \{(t, u, v) \in \mathbb{R}^3 : u > \beta(t), v = v_\beta^\varepsilon(t)\}.$$

Lemma 3.4. *If $z = (q, p, u, v)$ is a solution of system (3.8) such that $(t_0, u(t_0), v(t_0)) \in A_W$, then there exists a $\delta > 0$ such that*

$$t \in]t_0 - \delta, t_0[\Rightarrow (t, u(t), v(t)) \in A_{NW},$$

$$t \in]t_0, t_0 + \delta[\Rightarrow (t, u(t), v(t)) \in A_{SW}.$$

Similarly, if $(t_0, u(t_0), v(t_0)) \in A_E$, then there exists a $\delta > 0$ such that

$$t \in]t_0 - \delta, t_0[\Rightarrow (t, u(t), v(t)) \in A_{SE},$$

$$t \in]t_0, t_0 + \delta[\Rightarrow (t, u(t), v(t)) \in A_{NE}.$$

Proof. We give only the proof of the first part, the proof of the second part being similar. Let $z = (q, p, u, v)$ be a solution of system (3.8) such that $(t_0, u(t_0), v(t_0)) \in A_W$. Then $v(t_0) = v_\alpha^\varepsilon(t_0)$ and $u(t_0) < \alpha(t_0)$. Let us define a map $G(t) = v(t) - v_\alpha^\varepsilon(t)$. Then, G is continuous with $G(t_0) = 0$, and by (3.18) we have

$$\dot{G}(t_0) = \dot{v}(t_0) - \dot{v}_\alpha^\varepsilon(t_0)$$

$$= \tilde{g}(t_0, u(t_0)) - \varepsilon \partial_u P(t_0, q(t_0), p(t_0), u(t_0), v(t_0)) - \dot{v}_\alpha^\varepsilon(t_0) < 0,$$

for $|\varepsilon|$ small enough. So, there exists $\delta > 0$ such that $G(t) > 0$ for every $t \in]t_0 - \delta, t_0[$, and $u(t) < \alpha(t)$ for every $t \in [t_0 - \delta, t_0 + \delta]$. The conclusion is thus proved. \square

Lemma 3.5. *If $z = (q, p, u, v)$ is a T -periodic solution of system (3.8), then $(t, z(t)) \in \mathcal{V}$, for every $t \in \mathbb{R}$.*

Proof. Let us first prove that, for every $t \in \mathbb{R}$, we have

$$\alpha(t) \leq u(t) \leq \beta(t). \tag{3.21}$$

Suppose that there exists a solution $z = (q, p, u, v)$ of system (3.8) such that $u(t_0) < \alpha(t_0)$ for some $t_0 \in [0, T]$. If $(t_0, u(t_0), v(t_0)) \in A_{NW}$, then from Lemma 3.3 we have that $(t, u(t), v(t)) \in A_{NW}$ for every $t < t_0$. Then, by (3.16), we have

$$\frac{d}{dt}(u - \alpha)(t) = \tilde{f}(t, v(t)) + \varepsilon \partial_v P(t, u(t), v(t)) - \dot{\alpha}(t) > 0,$$

for $|\varepsilon|$ small enough and every $t < t_0$, which is clearly a contradiction, because $u - \alpha$ is a periodic solution. The same reasoning applies if $(t_0, u(t_0), v(t_0)) \in A_{SW}$. Finally, if $(t_0, u(t_0), v(t_0)) \in A_W$, then by Lemma 3.4 we know that the solution will be in A_{SW} or in A_{NW} at some time near t_0 , hence we obtain a contradiction again. Then, $u(t) \geq \alpha(t)$ for every $t \in [0, T]$. In a similar way we can prove that $u(t) \leq \beta(t)$ for every $t \in [0, T]$.

Finally we prove that

$$\gamma_-(u(t)) \leq v(t) \leq \gamma_+(u(t)). \tag{3.22}$$

For such a solution $z = (q, p, u, v)$, by (3.3) and (3.21) we see that $\tilde{g}(t, u(t)) = g(t, u(t))$. Now, define the T -periodic function $H_-(t) = v(t) - \gamma_-(u(t))$. Let $t_m \in [0, T]$ be such that $H_-(t_m) = \min H_-$ and assume by contradiction that $H_-(t_m) < 0$. Then,

$$\begin{aligned} \dot{H}_-(t_m) &= \dot{v}(t_m) - \gamma'_-(u(t_m))\dot{u}(t_m) \\ &= g(t_m, u(t_m)) - \varepsilon \partial_u P(t, q(t_m), p(t_m), u(t_m), v(t_m)) - \mu \dot{u}(t_m) \\ &= g(t_m, u(t_m)) - \mu \tilde{f}(t_m, v(t_m)) \\ &\quad - \varepsilon (\partial_u P(t_m, q(t_m), p(t_m), u(t_m), v(t_m)) + \mu \partial_v P(t_m, u(t_m), v(t_m))). \end{aligned}$$

We now consider the following cases:

Case 1. If $-\hat{d} \leq v(t_m) \leq \gamma_-(u(t_m))$, then $\tilde{f}(t_m, v(t_m)) = f(t_m, v(t_m))$ and so we have

$$\begin{aligned} \dot{H}_-(t_m) &\leq C - \mu \cdot (-c) \\ &\quad - \varepsilon (\partial_u P(t_m, q(t_m), p(t_m), u(t_m), v(t_m)) + \mu \partial_v P(t_m, u(t_m), v(t_m))) \\ &< 0, \end{aligned}$$

for $|\varepsilon|$ small enough, since $g(t_m, u(t_m)) \leq C$ and $\mu < -\frac{C}{c}$.

Case 2. If $v(t_m) < -\hat{d} - 1$, then $\tilde{f}(t_m, v(t_m)) = v(t_m)$ and so we have

$$\begin{aligned} \dot{H}_-(t_m) &\leq C - \mu v(t_m) \\ &\quad - \varepsilon (\partial_u P(t_m, q(t_m), p(t_m), u(t_m), v(t_m)) + \mu \partial_v P(t_m, u(t_m), v(t_m))) \\ &< C - \mu \cdot (-\hat{d} - 1) \\ &\quad - \varepsilon (\partial_u P(t_m, q(t_m), p(t_m), u(t_m), v(t_m)) + \mu \partial_v P(t_m, u(t_m), v(t_m))) \\ &< 0, \end{aligned}$$

for $|\varepsilon|$ small enough, since $g(t_m, u(t_m)) \leq C$ and $\hat{d} > C/|\mu|$.

Case 3. If $-\hat{d} - 1 \leq v(t_m) \leq -\hat{d}$, then $\tilde{f}(t_m, v(t_m))$ is a linear interpolation between $f(t_m, v(t_m))$ and $v(t_m)$, hence

$$\min\{f(t_m, v(t_m)), v(t_m)\} \leq \tilde{f}(t_m, v(t_m)) \leq \max\{f(t_m, v(t_m)), v(t_m)\},$$

so we have

$$\begin{aligned} \dot{H}_-(t_m) &\leq C - \mu \max\{f(t_m, v(t_m)), v(t_m)\} \\ &\quad - \varepsilon (\partial_u P(t_m, q(t_m), p(t_m), u(t_m), v(t_m)) + \mu \partial_v P(t_m, u(t_m), v(t_m))). \end{aligned}$$

If $f(t_m, v(t_m)) \leq v(t_m)$, then

$$\begin{aligned} \dot{H}_-(t_m) &\leq C + \mu \hat{d} \\ &\quad - \varepsilon (\partial_u P(t_m, q(t_m), p(t_m), u(t_m), v(t_m)) + \mu \partial_v P(t_m, u(t_m), v(t_m))) \\ &< 0, \end{aligned}$$

for $|\varepsilon|$ small enough, since $\hat{d} > c$ and $\mu < -C/c$. On the other hand, if $v(t_m) < f(t_m, v(t_m))$, then again

$$\begin{aligned} \dot{H}_-(t_m) &\leq C - \mu \cdot (-c) \\ &\quad - \varepsilon (\partial_u P(t_m, q(t_m), p(t_m), u(t_m), v(t_m)) + \mu \partial_v P(t_m, u(t_m), v(t_m))) \\ &< 0, \end{aligned}$$

for $|\varepsilon|$ small enough.

In all the above three cases we obtain contradictions, hence we have proved that $v(t) \geq \gamma_-(u(t))$, for every $t \in [0, T]$. In a similar way we can prove that $v(t) \leq \gamma_+(u(t))$, for every $t \in [0, T]$. \square

We have thus proved that, if $z = (q, p, u, v)$ is a solution of system (3.8), then $(t, z(t)) \in \mathcal{V}$, for every $t \in \mathbb{R}$, and so z is a solution of system (1.1). This completes the proof of Theorem 2.5. \square

4. Consequences and applications of Theorem 2.5

Let $\phi : \mathbb{R} \rightarrow \mathbb{R}$ be an increasing diffeomorphism with a bounded derivative, such that $\phi(0) = 0$. Consider the system

$$\begin{cases} \dot{q} = \partial_p H(t, q, p) + \varepsilon \partial_p P(t, q, p, u), \\ \dot{p} = -\partial_q H(t, q, p) - \varepsilon \partial_q P(t, q, p, u), \\ \frac{d}{dt}(\phi(\dot{u})) = g(t, u) - \varepsilon \partial_u P(t, q, p, u), \end{cases} \tag{4.1}$$

where $P = P(t, q, p, u)$ is a perturbation term which is 2π -periodic in q and has a bounded gradient with respect to (q, p, u) . As a direct consequence of the Theorem 2.5, we have the following result.

Corollary 4.1. *Let A1 and A2 hold. Moreover, let there exist two T -periodic C^2 -functions $\alpha, \beta : \mathbb{R} \rightarrow \mathbb{R}$ with $\alpha \leq \beta$, such that*

$$\frac{d}{dt}(\phi(\dot{\alpha}))(t) > g(t, \alpha(t)), \quad \frac{d}{dt}(\phi(\dot{\beta}))(t) < g(t, \beta(t)),$$

for every $t \in [0, T]$. Then there exists $\bar{\varepsilon} > 0$ such that, if $|\varepsilon| \leq \bar{\varepsilon}$, system (4.1) has at least two geometrically distinct T -periodic solutions, such that $p(0) \in]a, b[$ and $\alpha \leq u \leq \beta$.

Proof. Define $v_\alpha, v_\beta : \mathbb{R} \rightarrow \mathbb{R}$ as

$$v_\alpha(t) = \phi(\dot{\alpha}(t)), \quad v_\beta(t) = \phi(\dot{\beta}(t)).$$

Setting $f(t, v) = \phi^{-1}(v)$, all the assumptions of Theorem 2.5 are satisfied, and so the conclusion follows. \square

Notice that, taking $\phi(s) = s$ for all $s \in \mathbb{R}$, the last equation in (4.1) becomes

$$\ddot{u} = g(t, u) - \varepsilon \partial_u P(t, q, p, u).$$

Example 4.2. Consider the following system

$$\begin{cases} -\ddot{q} = a \sin q + \varepsilon \partial_q P(t, q, u), \\ -\ddot{u} = -g(t, u) + \varepsilon \partial_u P(t, q, u), \end{cases} \tag{4.2}$$

where $a > 0$. Assume that P is 2π -periodic in q and has a bounded gradient with respect to (q, u) , and the function g satisfies the Landesman–Lazer condition

$$\int_0^T \limsup_{u \rightarrow -\infty} g(t, u) dt < 0 < \int_0^T \liminf_{u \rightarrow +\infty} g(t, u) dt. \tag{4.3}$$

By using (4.3) and [16, Lemma 2], we get a strict lower solution α and a strict upper solution β of the equation $\ddot{u} = g(t, u)$. So, Corollary 4.1 applies, and thus system (4.2) has at least two geometrically distinct solutions. Notice that also Theorem 2.8 could be applied in this case, providing subharmonic solutions of period kT , with any integer $k > 2\pi/(T\sqrt{a})$, in the spirit of [20].

5. The higher dimensional case

We now consider the following system

$$\begin{cases} \dot{q} = \nabla_p H(t, q, p) + \varepsilon \nabla_p P(t, q, p, u, v), \\ \dot{p} = -\nabla_q H(t, q, p) - \varepsilon \nabla_q P(t, q, p, u, v), \\ \dot{u}_j = f_j(t, v_j) + \varepsilon \partial_{v_j} P(t, q, p, u, v), \quad j = 1, \dots, L, \\ \dot{v}_j = g_j(t, u_j) - \varepsilon \partial_{u_j} P(t, q, p, u, v), \quad j = 1, \dots, L. \end{cases} \tag{5.1}$$

For $z = (q, p, u, v) \in \mathbb{R}^N$ we write

$$\begin{aligned} q &= (q_1, \dots, q_M) \in \mathbb{R}^M, & p &= (p_1, \dots, p_M) \in \mathbb{R}^M, \\ u &= (u_1, \dots, u_L) \in \mathbb{R}^L, & v &= (v_1, \dots, v_L) \in \mathbb{R}^L. \end{aligned}$$

We assume all the involved functions to be continuous, and T -periodic in their first variable t .

We first recall the definition of lower and upper solution for the T -periodic problem associated with the system

$$\dot{u}_j = f_j(t, v_j), \quad \dot{v}_j = g_j(t, u_j), \quad j = 1, \dots, L. \tag{5.2}$$

Definition 5.1. A T -periodic C^1 -function $\alpha : \mathbb{R} \rightarrow \mathbb{R}^L$ is said to be a “lower solution” for the T -periodic problem associated with system (5.2) if there exists a T -periodic C^1 -function $v_\alpha : \mathbb{R} \rightarrow \mathbb{R}^L$ such that, for every $j = 1, \dots, L$ we have

$$\begin{cases} s < v_{\alpha,j}(t) & \Rightarrow f_j(t, s) < \dot{\alpha}_j(t), \\ s > v_{\alpha,j}(t) & \Rightarrow f_j(t, s) > \dot{\alpha}_j(t), \end{cases} \tag{5.3}$$

and

$$\dot{v}_{\alpha,j}(t) \geq g_j(t, \alpha_j(t)). \tag{5.4}$$

The lower solution is “strict” if the strict inequalities in (5.4) hold.

Definition 5.2. A T -periodic C^1 -function $\beta : \mathbb{R} \rightarrow \mathbb{R}^L$ is said to be an “upper solution” for the T -periodic problem associated with system (5.2) if there exists a T -periodic C^1 -function $v_\beta : \mathbb{R} \rightarrow \mathbb{R}^L$ such that, for every $j = 1, \dots, L$ we have

$$\begin{cases} s < v_{\beta,j}(t) & \Rightarrow f_j(t, s) < \dot{\beta}_j(t), \\ s > v_{\beta,j}(t) & \Rightarrow f_j(t, s) > \dot{\beta}_j(t), \end{cases} \tag{5.5}$$

and

$$\dot{v}_{\beta,j}(t) \leq g_j(t, \beta_j(t)). \tag{5.6}$$

The upper solution is “strict” if the strict inequalities in (5.6) hold.

We first consider the case when \mathcal{D} is a rectangle in \mathbb{R}^M , i.e.

$$\mathcal{D} = [a_1, b_1] \times \dots \times [a_M, b_M].$$

Let us state our hypotheses in this setting.

A1'. The function $H(t, q, p)$ is 2π -periodic in each variable q_1, \dots, q_M .

A2'. There exists an M -tuple $\sigma = (\sigma_1, \dots, \sigma_M) \in \{-1, 1\}^M$ such that all the solutions (q, p) of system (5.10) starting with $p(0) \in \mathcal{D}$ are defined on $[0, T]$, and, for every $i = 1, \dots, M$, we have

$$\begin{cases} p_i(0) = a_i & \Rightarrow \sigma_i(q_i(T) - q_i(0)) < 0, \\ p_i(0) = b_i & \Rightarrow \sigma_i(q_i(T) - q_i(0)) > 0. \end{cases}$$

In the sequel, inequalities of n -tuples will be meant componentwise.

- A3'. There exist a strict lower solution α and a strict upper solution β for the T -periodic problem associated with system (5.2), such that $\alpha \leq \beta$.
- A4'. The second order partial derivatives $\partial_{ts}^2 f_j(t, s)$ and $\partial_{ss}^2 f_j(t, s)$ exist and are continuous; moreover, there exists $\lambda > 0$ such that

$$\partial_s f_j(t, s) \geq \lambda, \quad \text{for every } (t, s) \in [0, T] \times \mathbb{R},$$

for every $j = 1, \dots, L$.

- A5'. The function $P(t, q, p, u, v)$ is 2π -periodic in q_1, \dots, q_M and has a bounded gradient with respect to (q, p, u, v) ; moreover, the partial derivative $\nabla_v P$ is independent of q and p , and the map $\nabla_v P(t, u, v)$ is continuously differentiable.

Here is our first generalization of Theorem 2.5.

Theorem 5.3. *Assume that A1' – A5' hold true. Then there exists $\bar{\varepsilon} > 0$ such that, if $|\varepsilon| \leq \bar{\varepsilon}$, there are at least $M + 1$ geometrically distinct T -periodic solutions of system (5.1), with $p(0) \in \overset{\circ}{\mathcal{D}}$ and $\alpha \leq u \leq \beta$.*

The proof follows exactly the lines of the proof of Theorem 2.5, working separately on the components (u_j, v_j) of the solutions $z = (q, p, u, v)$ of system (5.1). The main idea is to modify the system so to have a Hamiltonian function of the type

$$\tilde{H}(t, z) = \frac{1}{2}(|v|^2 - |u|^2) + K(t, z),$$

where K is a function having a bounded gradient with respect to z , and then apply [13, Corollary 2.4] again. We avoid the details, for brevity.

Remark 5.4. Based on the Remark 2.7, we could have varying intervals $[a_i(s), b_i(s)]$ instead of the intervals $[a_i, b_i]$ in the rectangle \mathcal{D} , where $a_i, b_i : \mathbb{R} \rightarrow \mathbb{R}$ are 2π -periodic continuous functions.

We can now provide a higher dimensional version of Theorem 2.8. Recall that $\mathcal{D}(\Gamma)$ denotes the open bounded region delimited by a planar Jordan curve Γ .

Theorem 5.5. *Assume that A3' – A5' hold true. Let k_1, k_2, \dots, k_M be integers and assume that, for each $i \in \{1, \dots, M\}$, there exist two planar Jordan curves Γ_1^i, Γ_2^i , strictly star-shaped with respect to the origin, with*

$$0 \in \mathcal{D}(\Gamma_1^i) \subseteq \overline{\mathcal{D}(\Gamma_1^i)} \subseteq \mathcal{D}(\Gamma_2^i),$$

such that the solutions of system (5.10) with $(q_i(0), p_i(0)) \in \overline{\mathcal{D}(\Gamma_2^i)} \setminus \mathcal{D}(\Gamma_1^i)$ for every $i \in \{1, \dots, M\}$ are defined on $[0, T]$ and satisfy

$$(q_i(t), p_i(t)) \neq (0, 0), \quad \forall t \in [0, T],$$

and

$$\begin{cases} (q_i(0), p_i(0)) \in \Gamma_1^i & \Rightarrow \text{Rot}((q_i, p_i); [0, T]) < k_i, \\ (q_i(0), p_i(0)) \in \Gamma_2^i & \Rightarrow \text{Rot}((q_i, p_i); [0, T]) > k_i. \end{cases} \tag{5.7}$$

Then system (5.1) has at least $M + 1$ geometrically distinct T -periodic solutions $z^{(n)} = (q^{(n)}, p^{(n)}, u^{(n)}, v^{(n)})$ for $n = 1, \dots, M + 1$ such that

$$\begin{aligned} \alpha &\leq u^{(n)} \leq \beta, \\ (q_i^{(n)}(0), p_i^{(n)}(0)) &\in \mathcal{D}(\Gamma_2^i) \setminus \overline{\mathcal{D}(\Gamma_1^i)}, \end{aligned}$$

and

$$\text{Rot}((q_i^{(n)}, p_i^{(n)}); [0, T]) = k_i,$$

for $i = 1, \dots, M$. The same is true if for some $i \in \{1, \dots, M\}$ the assumption (5.7) is replaced by the following

$$\begin{cases} (q_i(0), p_i(0)) \in \Gamma_1^i & \Rightarrow \text{Rot}((q_i, p_i); [0, T]) > k_i, \\ (q_i(0), p_i(0)) \in \Gamma_2^i & \Rightarrow \text{Rot}((q_i, p_i); [0, T]) < k_i. \end{cases}$$

Here is an example of application of the above theorems.

Example 5.6. Consider the following system

$$\begin{cases} -\ddot{q}_i = a_i \sin q_i + \varepsilon \partial_{q_i} P(t, q, u), & i = 1, \dots, M, \\ -\ddot{u}_j = -g_j(t, u_j) + \varepsilon \partial_{u_j} P(t, q, u), & j = 1, \dots, L, \end{cases} \tag{5.8}$$

where $a_i > 0$. Assume that P is 2π -periodic in q_1, \dots, q_M and has a bounded gradient with respect to (q, u) , and for each $j \in \{1, \dots, L\}$, the function g_j satisfies the Landesman–Lazer condition

$$\int_0^T \limsup_{s \rightarrow -\infty} g_j(t, s) dt < 0 < \int_0^T \liminf_{s \rightarrow +\infty} g_j(t, s) dt. \tag{5.9}$$

By using (5.9) and [16, Lemma 2], we get a strict lower solution α_j and a strict upper solution β_j of the equation $\ddot{u}_j = g_j(t, u_j)$, with $\alpha_j(t) < \beta_j(t)$. So all the assumptions of Theorem 5.3 hold and thus system (5.8) has at least $M + 1$ geometrically distinct T -periodic solutions. As noticed in Example 4.2, subharmonic solutions with a sufficiently large period may be detected also in this case.

We now consider two variants of Theorem 5.3.

We say that \mathcal{D} is a convex body of \mathbb{R}^M if it is a closed convex bounded subset of \mathbb{R}^M having nonempty interior. By assuming that \mathcal{D} has a smooth boundary, we denote the unit outward

normal at $\zeta \in \partial\mathcal{D}$ by $\nu_{\mathcal{D}}(\zeta)$. Moreover, we say that \mathcal{D} is strongly convex if for any $p \in \partial\mathcal{D}$, the map $\mathcal{F} : \mathcal{D} \rightarrow \mathbb{R}$ defined by $\mathcal{F}(\xi) = \langle \xi - p, \nu_{\mathcal{D}}(p) \rangle$ has a unique maximum point at $\xi = p$.

Let us first state the following “avoiding rays” assumption for the system

$$\dot{q} = \nabla_p H(t, q, p), \quad \dot{p} = -\nabla_q H(t, q, p). \tag{5.10}$$

A2''. There exists a convex body \mathcal{D} of \mathbb{R}^M , having a smooth boundary, such that all the solutions (q, p) of system (5.10) starting with $p(0) \in \mathcal{D}$ are defined on $[0, T]$, and

$$p(0) \in \partial\mathcal{D} \implies q(T) - q(0) \notin \{\lambda \nu_{\mathcal{D}}(p(0)) : \lambda \geq 0\}.$$

Theorem 5.7. *If in the statement of Theorem 5.3 we replace assumption A2' by A2'', the same conclusion holds.*

The only difference in the proof is that instead of [13, Corollary 2.4], we apply [13, Corollary 2.1].

To conclude this section we introduce an “indefinite twist” assumption.

A2'''. There are a strongly convex body \mathcal{D} of \mathbb{R}^M having a smooth boundary and a symmetric regular $M \times M$ matrix \mathbb{A} such that all the solutions (q, p) of system (5.10) starting with $p(0) \in \mathcal{D}$ are defined on $[0, T]$, and

$$p(0) \in \partial\mathcal{D} \implies \langle q(T) - q(0), \mathbb{A} \nu_{\mathcal{D}}(p(0)) \rangle > 0.$$

Theorem 5.8. *If in the statement of Theorem 5.3 we replace assumption A2' by A2''', the same conclusion holds.*

Again the proof is the same, the only difference being that instead of [13, Corollary 2.4], we apply [13, Corollary 2.3].

Remark 5.9. We could also have assumed some very general twist conditions, in the line of [11–13]. These involve the “avoiding cones condition” for rather general domains. However, in this paper we preferred to present our ideas in some more concrete situations. The interested reader will have no difficulties in adapting our results to the more general setting.

6. Twist with Hartman type condition

In this section we consider a system in \mathbb{R}^{2M+2L} of the type

$$\begin{cases} \dot{q} = \nabla_p H(t, q, p) + \varepsilon \nabla_p P(t, q, p, u), \\ \dot{p} = -\nabla_q H(t, q, p) - \varepsilon \nabla_q P(t, q, p, u), \\ \dot{u} = v, \quad \dot{v} = \nabla_u G(t, u) - \varepsilon \nabla_u P(t, q, p, u). \end{cases} \tag{6.1}$$

Here again all functions involved are assumed to be continuous, and T -periodic in t . We will assume the periodicity condition A1' and one of the twist conditions A2', A2'' or A2''', even if in the following statement we concentrate on A2'''. We also assume condition A5' which, in this setting, can be stated in the following simpler form.

A5''. The function $P(t, q, p, u)$ is 2π -periodic in q_1, \dots, q_M and has a bounded gradient with respect to (q, p, u) .

Here is our statement, involving a Hartman-type condition (see [10,22] and the references therein).

Theorem 6.1. *Assume that A1', A2''' and A5'' hold true and that there exists $R > 0$ such that*

$$|u| = R \implies \langle \nabla_u G(t, u), u \rangle > 0. \tag{6.2}$$

Then there exists $\bar{\varepsilon} > 0$ such that for $|\varepsilon| \leq \bar{\varepsilon}$, there are at least $M + 1$ geometrically distinct T -periodic solutions of system (6.1) such that $p(0) \in \mathring{D}$ and $|u(t)| \leq R$ for every $t \in \mathbb{R}$.

Proof. First of all, we modify the function G outside the ball $B_R = \{u : |u| \leq R\}$. By (6.2) and the continuity of the inner product, there exists $\tilde{\rho} > 0$ and $\delta > 0$ such that

$$R \leq |u| \leq R + \tilde{\rho} \implies \langle \nabla_u G(t, u), u \rangle \geq \delta. \tag{6.3}$$

We can assume without loss of generality that

$$G(t, u) \leq 0, \quad \text{when } R \leq |u| \leq R + \tilde{\rho}. \tag{6.4}$$

Indeed, if it is not already the case, it is sufficient to replace $G(t, u)$ by $G(t, u) - M$, where

$$M = \max\{|G(t, u)| : 0 \leq t \leq T, R < |u| \leq R + \tilde{\rho}\}.$$

Its gradient will not be changed.

Moreover, as in the previous proofs, after a truncation we can from now on assume that H has a bounded gradient with respect to (q, p) .

Now choose a C^∞ -function $\eta : \mathbb{R} \rightarrow \mathbb{R}$ satisfying

$$\begin{cases} \eta(s) = 1, & \text{if } s \leq R, \\ \dot{\eta}(s) \leq 0, & \text{if } R \leq s \leq R + \tilde{\rho}, \\ \eta(s) = 0, & \text{if } s > R + \tilde{\rho}, \end{cases}$$

and define the function

$$\tilde{G}(t, u) = \begin{cases} G(t, u), & \text{if } |u| \leq R, \\ \eta(|u|) G(t, u) + (1 - \eta(|u|)) \frac{1}{2} |u|^2, & \text{if } R \leq |u| \leq R + \tilde{\rho}, \\ \frac{1}{2} |u|^2, & \text{if } |u| > R + \tilde{\rho}. \end{cases}$$

Notice that, outside the ball $B_{R+\tilde{\rho}}$, the system becomes almost linear.

We now consider the new system

$$\begin{cases} \dot{q} = \nabla_p H(t, q, p) + \varepsilon \nabla_p P(t, q, p, u), \\ \dot{p} = -\nabla_q H(t, q, p) - \varepsilon \nabla_q P(t, q, p, u), \\ \dot{u} = v, \quad \dot{v} = \nabla_u \tilde{G}(t, u) - \varepsilon \nabla_u P(t, q, p, u), \end{cases} \tag{6.5}$$

where the new Hamiltonian function is

$$\tilde{H}(t, q, p, u, v) = H(t, q, p) + \frac{1}{2}|v|^2 - \tilde{G}(t, u) + \varepsilon P(t, q, p, u).$$

Writing the modified system (6.5) as $\dot{z} = J\nabla\tilde{H}(t, z)$, we see that

$$\tilde{H}(t, z) = \frac{1}{2}(|v|^2 - |u|^2) + K(t, z),$$

with $K(t, z)$ has a bounded gradient with respect to $z = (q, p, u, v)$. Then, by [13, Corollary 2.3], the modified system (6.5) has at least $M + 1$ geometrically distinct T -periodic solutions, such that $p(0) \in \dot{D}$, provided that $|\varepsilon|$ is small enough.

We need to prove that the T -periodic solutions of system (6.5) we have found are such that $|u(t)| \leq R$ for every $t \in [0, T]$, so that they are indeed solutions of system (6.1).

Assume by contradiction that there exists $t_0 \in \mathbb{R}$ such that

$$|u(t_0)| = \max\{|u(t)| : t \in [0, T]\} > R.$$

Consider the function $f(t) = |u(t)|^2$. We have that $\dot{f}(t_0) = 0$ and $\ddot{f}(t_0) \leq 0$. Being $\dot{f}(t) = \langle 2u(t), \dot{u}(t) \rangle$, we compute

$$\begin{aligned} \ddot{f}(t) &= 2\langle \dot{u}(t), \dot{u}(t) \rangle + 2\langle u(t), \ddot{u}(t) \rangle \\ &= 2|\dot{u}(t)|^2 + 2\langle u(t), \nabla_u \tilde{G}(t, u(t)) - \varepsilon \nabla_u P(t, q(t), p(t), u(t)) \rangle \\ &\geq 2\langle u(t), \nabla_u \tilde{G}(t, u(t)) - \varepsilon \nabla_u P(t, q(t), p(t), u(t)) \rangle. \end{aligned} \tag{6.6}$$

We have two cases.

Case 1. If $|u(t_0)| > R + \tilde{\rho}$, then by the Cauchy–Schwartz inequality and the fact that $|\nabla_u P(t, q, p, u)| < C$, the inequality (6.6) implies that

$$\begin{aligned} \ddot{f}(t_0) &\geq 2\langle u(t_0), u(t_0) - \varepsilon \nabla_u P(t_0, q(t_0), p(t_0), u(t_0)) \rangle \\ &\geq 2|u(t_0)|^2 - 2|\varepsilon| |u(t_0)| |\nabla_u P(t_0, q(t_0), p(t_0), u(t_0))| \\ &\geq 2|u(t_0)| (|u(t_0)| - |\varepsilon| |\nabla_u P(t_0, q(t_0), p(t_0), u(t_0))|) \\ &> 2R^2 > 0, \end{aligned}$$

for $|\varepsilon|$ small enough, a contradiction.

Case 2. If $R < |u(t_0)| < R + \tilde{\rho}$, then again (6.6) implies that

$$\begin{aligned} \ddot{f}(t_0) &\geq 2\left\langle u(t_0), \dot{\eta}(|u(t_0)|) \frac{u(t_0)}{|u(t_0)|} G(t_0, u(t_0)) + \eta(|u(t_0)|) \nabla_u G(t_0, u(t_0)) \right\rangle \\ &\quad + 2\left\langle u(t_0), -\dot{\eta}(|u(t_0)|) \frac{u(t_0)}{|u(t_0)|} \frac{1}{2}|u(t_0)|^2 + (1 - \eta(|u(t_0)|)) u(t_0) \right\rangle \\ &\quad - 2|u(t_0)| |\varepsilon| |\nabla_u P(t_0, q(t_0), p(t_0), u(t_0))| \end{aligned}$$

$$\begin{aligned}
 &= 2\dot{\eta}(|u(t_0)|) |u(t_0)|G(t_0, u(t_0)) + 2\eta(|u(t_0)|) \langle u(t_0), \nabla_u G(t_0, u(t_0)) \rangle \\
 &\quad - \dot{\eta}(|u(t_0)|) |u(t_0)|^3 + 2(1 - \eta(|u(t_0)|)) |u(t_0)|^2 \\
 &\quad - 2|u(t_0)| |\varepsilon| |\nabla_u P(t_0, q(t_0), p(t_0), u(t_0))|.
 \end{aligned}$$

By (6.3) and (6.4), since $\dot{\eta}(|u(t_0)|) \leq 0$ and

$$\eta(|u(t_0)|) \langle u(t_0), \nabla_u G(t_0, u(t_0)) \rangle + (1 - \eta(|u(t_0)|)) |u(t_0)|^2 \geq \min\{\delta, R^2\},$$

we have that

$$\ddot{f}(t_0) \geq 2 \min\{\delta, R^2\} - 2(R + \tilde{\rho})|\varepsilon| |\nabla_u P(t_0, q(t_0), p(t_0), u(t_0))| > 0,$$

when $|\varepsilon|$ is small enough, a contradiction. The proof is thus completed. \square

Remark 6.2. In the case $L = 1$, writing $g(t, u) = \nabla_u G(t, u)$, the Hartman condition becomes

$$g(t, -R) < 0 < g(t, R).$$

It is thus seen that $\alpha = -R$ and $\beta = R$ are *constant* strict lower/upper solutions, with $\alpha < \beta$.

7. Perturbations of completely integrable systems

There is a very large literature on the periodic problem for perturbations of completely integrable systems (see, e.g., [3,13] and references therein), starting from Poincaré, who referred to Hamiltonian perturbation theory as the “Problème général de la Dynamique”.

We will add now an extra term to the Hamiltonian function, involving a Hartman-type situation. Consider the system

$$\begin{cases} \dot{\varphi} = \nabla_{\mathcal{K}}(I) + \varepsilon \nabla_I P(t, \varphi, I, u), & \dot{I} = -\varepsilon \nabla_{\phi} P(t, \varphi, I, u), \\ \ddot{u} = \nabla_u G(t, u) - \varepsilon \nabla_u P(t, \varphi, I, u), \end{cases} \tag{7.1}$$

where $(\varphi, I) \in \mathbb{R}^{2M}$ and $u \in \mathbb{R}^L$. As usual we assume that all the involved functions are continuous and T -periodic in t . The perturbation function $P : \mathbb{R} \times \mathbb{R}^{2M+L} \rightarrow \mathbb{R}$ is assumed to have a bounded gradient with respect to (φ, I, u) . Moreover, it is τ_i -periodic in each variable φ_i , i.e.

$$P(t, \dots, \varphi_i + \tau_i, \dots) = P(t, \dots, \varphi_i, \dots),$$

and we assume that there exist $I^0 \in \mathbb{R}^M$ and some integers m_1, \dots, m_M such that

$$T \nabla_{\mathcal{K}}(I^0) = (m_1 \tau_1, \dots, m_M \tau_M).$$

We are thus dealing with a completely resonant torus. Here is our result.

Theorem 7.1. *In the above setting, assume that there exist $I^0 \in \mathbb{R}^M$, a symmetric invertible $M \times M$ matrix \mathbb{A} and $\rho > 0$ such that*

$$0 < |I - I^0| \leq \rho \implies \langle \nabla \mathcal{K}(I) - \nabla \mathcal{K}(I^0), \mathbb{A}(I - I^0) \rangle > 0. \tag{7.2}$$

Moreover, let there exist $R > 0$ such that

$$|u| = R \implies \langle \nabla_u G(t, u), u \rangle > 0.$$

Then, for every $\sigma > 0$, there exists $\tilde{\varepsilon} > 0$ such that, for $|\varepsilon| < \tilde{\varepsilon}$, there are at least $M + 1$ geometrically distinct solutions of system (7.1), with

$$\begin{aligned} \varphi(t + T) &= \varphi(t) + T \nabla \mathcal{K}(I^0), & u(t + T) &= u(t), & I(t + T) &= I(t), \\ |\varphi(t) - \varphi(0) - t \nabla \mathcal{K}(I^0)| &+ |I(t) - I^0| < \sigma, \end{aligned}$$

and

$$|u(t)| \leq R,$$

for every $t \in \mathbb{R}$.

The proof is based on Theorem 6.1, following the same reasoning as in [11, Theorem 24], so we omit it, for brevity.

Remark 7.2. It can easily be seen that assumption (7.2) is satisfied if the function \mathcal{K} is twice continuously differentiable at I^0 , with

$$\det \mathcal{K}''(I^0) \neq 0.$$

It is indeed sufficient to choose $\mathbb{A} = \mathcal{K}''(I^0)$.

Data availability

No data was used for the research described in the article.

References

- [1] V.I. Arnold, Sur une propriété topologique des applications globalement canoniques de la mécanique classique, C. R. Acad. Sci. Paris 261 (1965) 3719–3722.
- [2] V.I. Arnold, The stability problem and ergodic properties for classical dynamical systems, in: Proc. Internat. Congr. Math., Moscow, 1966, Mir, Moscow, 1966, pp. 387–392 (in Russian), English translation in: AMS Transl. Ser. 2, vol. 70, 1968, pp. 5–11.
- [3] D. Bernstein, A. Katok, Birkhoff periodic orbits for small perturbations of completely integrable Hamiltonian systems with convex Hamiltonians, Invent. Math. 88 (1987) 222–241.
- [4] M. Brown, W.D. Neumann, Proof of the Poincaré–Birkhoff fixed point theorem, Mich. Math. J. 24 (1977) 21–31.
- [5] K.C. Chang, On the periodic nonlinearity and the multiplicity of solutions, Nonlinear Anal. 13 (1989) 527–537.
- [6] F. Chen, D. Qian, An extension of the Poincaré–Birkhoff theorem for Hamiltonian systems coupling resonant linear components with twisting components, J. Differ. Equ. 321 (2022) 415–448.

- [7] C.C. Conley, E.J. Zehnder, The Birkhoff–Lewis fixed point theorem and a conjecture of V.I. Arnold, *Invent. Math.* 73 (1983) 33–49.
- [8] C. De Coster, P. Habets, *Two-Point Boundary Value Problems, Lower and Upper Solutions*, Elsevier, Amsterdam, 2006.
- [9] P.L. Felmer, Periodic solutions of spatially periodic Hamiltonian systems, *J. Differ. Equ.* 98 (1992) 143–168.
- [10] G. Feltrin, F. Zanolin, Bound sets for a class of ϕ -Laplacian operators, *J. Differ. Equ.* 297 (2021) 508–535.
- [11] A. Fonda, M. Garzón, A. Sfecci, An extension of the Poincaré–Birkhoff Theorem coupling twist with lower and upper solutions, *J. Math. Anal. Appl.* 528 (2023) 127599.
- [12] A. Fonda, P. Gidoni, An avoiding cones condition for the Poincaré–Birkhoff Theorem, *J. Differ. Equ.* 262 (2017) 1064–1084.
- [13] A. Fonda, P. Gidoni, Coupling linearity and twist: an extension of the Poincaré–Birkhoff Theorem for Hamiltonian systems, *NoDEA Nonlinear Differ. Equ. Appl.* 27 (2020) 55.
- [14] A. Fonda, G. Klun, A. Sfecci, Well-ordered and non-well-ordered lower and upper solutions for periodic planar systems, *Adv. Nonlinear Stud.* 21 (2021) 397–419.
- [15] A. Fonda, M. Sabatini, F. Zanolin, Periodic solutions of perturbed Hamiltonian systems in the plane by the use of the Poincaré–Birkhoff Theorem, *Topol. Methods Nonlinear Anal.* 40 (2012) 29–52.
- [16] A. Fonda, R. Toader, Periodic solutions of radially symmetric perturbations of Newtonian systems, *Proc. Am. Math. Soc.* 140 (4) (2012) 1331–1341.
- [17] A. Fonda, R. Toader, A dynamical approach to lower and upper solutions for planar systems, *Discrete Contin. Dyn. Syst.* 41 (2021) 3683–3708.
- [18] A. Fonda, A.J. Ureña, A higher dimensional Poincaré–Birkhoff theorem for Hamiltonian flows, *Ann. Inst. Henri Poincaré, Anal. Non Linéaire* 34 (2017) 679–698.
- [19] A. Fonda, A.J. Ureña, A Poincaré–Birkhoff theorem for Hamiltonian flows on nonconvex domains, *J. Math. Pures Appl.* 129 (2019) 131–152.
- [20] A. Fonda, F. Zanolin, Periodic oscillations of forced pendulums with a very small length, *Proc. R. Soc. Edinb.* 127A (1997) 67–76.
- [21] G. Fournier, D. Lupo, M. Ramos, M. Willem, *Limit Relative Category and Critical Point Theory*, *Dynamics Reported: Expositions in Dynamical Systems*, vol. 3, Springer, Berlin, 1994, pp. 1–24.
- [22] P. Hartman, On boundary value problems for systems of ordinary, nonlinear, second order differential equations, *Trans. Am. Math. Soc.* 96 (1960) 493–509.
- [23] F.W. Josellis, Lyusternik–Schnirelman theory for flows and periodic orbits for Hamiltonian systems on $\mathbb{T}^n \times \mathbb{R}^n$, *Proc. Lond. Math. Soc.* 68 (1994) 641–672.
- [24] J.Q. Liu, A generalized saddle point theorem, *J. Differ. Equ.* 82 (1989) 372–385.
- [25] J. Mawhin, Forced second order conservative systems with periodic nonlinearity, in: *Analyse Non Linéaire*, Perpignan, 1987, *Ann. Inst. Henri Poincaré, Anal. Non Linéaire* 6 (suppl) (1989) 415–434.
- [26] M. Nagumo, Über die Differentialgleichung $y'' = f(t, y, y')$, *Proc. Phys. Math. Soc. Jpn.* 19 (1937) 861–866.
- [27] E. Picard, Sur l’application des méthodes d’approximations successives à l’étude de certaines équations différentielles ordinaires, *J. Math. Pures Appl.* 9 (1893) 217–271.
- [28] H. Poincaré, Sur un théorème de géométrie, *Rend. Circ. Mat. Palermo* 33 (1912) 375–407.
- [29] G. Scorza Dragoni, Il problema dei valori ai limiti studiato in grande per le equazioni differenziali del secondo ordine, *Math. Ann.* 105 (1931) 133–143.