

Inclusive and differential cross section measurements of $t\bar{t}b\bar{b}$ production in the lepton+jets channel at $\sqrt{s} = 13$ TeV



The CMS collaboration

E-mail: cms-publication-committee-chair@cern.ch

ABSTRACT: Measurements of inclusive and normalized differential cross sections of the associated production of top quark-antiquark and bottom quark-antiquark pairs, $t\bar{t}b\bar{b}$, are presented. The results are based on data from proton-proton collisions collected by the CMS detector at a centre-of-mass energy of 13 TeV, corresponding to an integrated luminosity of 138 fb^{-1} . The cross sections are measured in the lepton+jets decay channel of the top quark pair, using events containing exactly one isolated electron or muon and at least five jets. Measurements are made in four fiducial phase space regions, targeting different aspects of the $t\bar{t}b\bar{b}$ process. Distributions are unfolded to the particle level through maximum likelihood fits, and compared with predictions from several event generators. The inclusive cross section measurements of this process in the fiducial phase space regions are the most precise to date. In most cases, the measured inclusive cross sections exceed the predictions with the chosen generator settings. The only exception is when using a particular choice of dynamic renormalization scale, $\mu_R = \frac{1}{2} \prod_{i=t,\bar{t},b,\bar{b}} m_{T,i}^{1/4}$, where $m_{T,i}^2 = m_i^2 + p_{T,i}^2$ are the transverse masses of top and bottom quarks. The differential cross sections show varying degrees of compatibility with the theoretical predictions, and none of the tested generators with the chosen settings simultaneously describe all the measured distributions.

KEYWORDS: B Physics, Hadron-Hadron Scattering, Top Physics

ARXIV EPRINT: [2309.14442](https://arxiv.org/abs/2309.14442)

Contents

1	Introduction	2
2	The CMS detector	4
3	Simulated samples	4
4	Event reconstruction	7
4.1	Detector-level object reconstruction and identification	7
4.2	Particle-level object definitions	9
5	Event selection and definition of fiducial phase space regions and observables	10
5.1	Event selection	10
5.2	Fiducial phase space regions	11
5.3	Observables	11
6	Signal extraction and unfolding	12
6.1	Ancillary variable	14
6.2	Multivariate algorithm for jet assignment	15
6.3	Unfolding methodology	17
7	Systematic uncertainties	21
7.1	Experimental uncertainties	21
7.2	Modelling uncertainties	23
8	Results	26
8.1	Inclusive cross sections	26
8.2	Normalized differential cross sections	32
9	Summary	41
A	Leading nuisance parameter impacts	44
B	Groups of impacts	48
C	Correlation of POIs	52
D	Variation of renormalization and factorization scales	55
E	Normalized differential cross section compatibility	57
	The CMS collaboration	65

1 Introduction

The associated production of top and bottom quark-antiquark pairs, $t\bar{t}b\bar{b}$, in proton-proton (pp) collisions at the CERN LHC is notoriously challenging to model because of the non-negligible mass of the b quark, and the difference in the typical energy scales of interactions involving top and b quarks [1, 2]. Comparing predictions for $t\bar{t}b\bar{b}$ production cross sections with inclusive and differential measurements is an important test of perturbative quantum chromodynamics (QCD) calculations. Furthermore, $t\bar{t}b\bar{b}$ production is a leading background for searches and other measurements, such as the measurement of the associated production of top quark pairs with Higgs bosons ($t\bar{t}H$), where the Higgs boson decays to a pair of b quarks ($H \rightarrow b\bar{b}$) [3–8], and measurements of the simultaneous production of four top quarks ($t\bar{t}t\bar{t}$) [9–19]. These two processes provide direct access to the top quark Yukawa coupling, a crucial parameter of the standard model (SM) [20, 21]. An improved understanding of $t\bar{t}b\bar{b}$ production will help reduce the uncertainties in these important measurements.

Fixed-order calculations of inclusive and differential cross sections have been obtained at next-to-leading order (NLO) in QCD for $t\bar{t}b\bar{b}$ production [22–26] and for the associated production of $t\bar{t}b\bar{b}$ with one additional jet ($t\bar{t}b\bar{b}j$) [1]. In addition, full NLO QCD corrections to off-shell $t\bar{t}b\bar{b}$ production are available [27, 28]. While these fixed-order calculations provide important insights into the dynamics of the $t\bar{t}b\bar{b}$ process, they cannot easily be compared with measurements because of the large extrapolations to the full parton-level phase space that this would involve. Instead, predictions obtained by matching matrix-element (ME) generators to parton showers (PSs) can be more directly compared to the data and can be used to model the $t\bar{t}b\bar{b}$ background in other measurements. Such predictions have been obtained using different modelling approaches [29–34]. The current state of the art in simulating $t\bar{t}b\bar{b}$ production uses ME calculations at NLO in QCD with massive b quarks and matching to a PS [2, 35, 36]. These calculations with massive b quarks use parton distribution functions (PDFs) of the proton in the four-flavour scheme (4FS), where b quarks are not part of the proton PDF. In particular, in final states with gluon splittings, $g \rightarrow b\bar{b}$, the b quark mass is taken into account. These calculations provide a description of $t\bar{t}b\bar{b}$ production based on the NLO MEs in the entire phase space where the $b\bar{b}$ pair can be resolved as one or two b quark jets.

Measurements of inclusive and differential $t\bar{t}b\bar{b}$ cross sections have previously been performed by the ATLAS and CMS Collaborations in pp collisions at centre-of-mass energies of 7, 8, and 13 TeV in final states with zero, one, or two charged leptons ($\ell = e, \mu$), using samples corresponding to integrated luminosities of up to 41.5 fb^{-1} [37–45]. To date, $t\bar{t}b\bar{b}$ simulations have shown a tendency to underpredict the inclusive $t\bar{t}b\bar{b}$ cross sections. Normalized differential distributions are generally in agreement with the predictions from event generators, within experimental and theoretical uncertainties, although the size of these uncertainties has not yet made it possible to definitively rule out or prefer specific modelling approaches. The dominant uncertainties in previous studies were those related to b tagging calibration, jet energy scale (JES), and the limited precision of the NLO calculations, i.e. the choice of renormalization (μ_R) and factorization scales (μ_F).

This paper reports the measurement of inclusive and normalized differential cross sections of $t\bar{t}b\bar{b}$ production in four fiducial phase space regions in the lepton+jets channel. The measurement uses pp collision data recorded with the CMS detector at the LHC from 2016

to 2018 at $\sqrt{s} = 13$ TeV, corresponding to an integrated luminosity of 138 fb^{-1} . Top quarks almost always decay into a W boson and a b quark. We consider events in which one W boson decays into a pair of quarks, and the other W boson decays into a charged lepton (electron or muon) and a neutrino. Decays of the W boson into a tau lepton decaying into an electron or muon are also implicitly considered. After hadronization of the quarks, and including the b quarks not originating from the top quarks, the considered $t\bar{t}b\bar{b}$ events contain one electron or muon and at least five jets, of which three result from the hadronization of b quarks (b jets).

We measure inclusive cross sections and normalized differential cross sections, where the latter correspond to the ratios of the absolute differential cross sections to the inclusive ones. Measurements are performed in different phase space regions, targeting distinct aspects of $t\bar{t}b\bar{b}$ production. One phase space region targets fully resolved $t\bar{t}b\bar{b}$ events (labelled as $t\bar{t}b\bar{b}$). Another, more inclusive, phase space region requires only three b jets, so as to select $t\bar{t}b\bar{b}$ events in which the pairs of additional b quarks are reconstructed as a single jet, or in which one b jet is outside of the detector acceptance (labelled as $t\bar{t}b$). Finally, we measure cross sections aimed at the study of additional QCD radiation in $t\bar{t}b$ or $t\bar{t}b\bar{b}$ events (labelled as $t\bar{t}bj$ and $t\bar{t}b\bar{b}j$, respectively), as these have been shown to be sensitive to the modelling of $t\bar{t}b\bar{b}$ production [1].

The observed distributions are unfolded to the particle level using binned maximum likelihood fits, whereby the data from muon and electron channels from all data-taking years are fitted simultaneously, and uncertainties are estimated using a nuisance parameter profiling procedure. Differential cross sections are measured for several observables, using two complementary approaches for the event interpretation given below.

In the first approach, no attempt is made at directly identifying the additional b jets in $t\bar{t}b\bar{b}$ events in either data or simulation. The generator-level observables are defined using stable particles exclusively, with no reference to the simulated event history or the origin of b jets from top quark decays or QCD radiation. This ensures that the observable definitions at the generator and detector levels are highly consistent. Furthermore, the distributions unfolded in this way can be compared with predictions from event generators which do not provide any information about the origin of b quarks in the final state. In the phase space region with four b jets ($t\bar{t}b\bar{b}$), observables are defined via the pair of b jets with the smallest angular separation to target b jets produced through the splitting of gluons into $b\bar{b}$.

In the second approach, the b jets not originating from top quark decays are identified at the generator level using the simulated event history, and a multivariate algorithm is developed to identify the resulting reconstructed b jets among all observed jets. This approach is more accurate at identifying additional b jets and thus these observables can be more sensitive to the modelling of additional heavy-quark production in $t\bar{t}$ events, at the cost of restricting future reinterpretations of the results and introducing additional modelling assumptions about the parton history.

The unfolded results are compared to different predictions based on various NLO ME calculations, interfaced with different PS simulations. Tabulated results are provided in the HEPData record for this analysis [46].

This paper is organized as follows. Section 2 describes the CMS detector. The event generation and detector simulation are detailed in section 3. The reconstruction of electrons,

muons, and jets and the identification of b jets, as well as the corresponding definition of particle-level objects, are discussed in section 4. Section 5 describes the event selection and the definition of the measured observables at the generator and detector levels. In section 6, the extraction of the signal, the identification of additional b jets using a multivariate algorithm, and the unfolding of the observables are presented. The treatment of systematic uncertainties is detailed in section 7, and the results, compared to several theoretical predictions, are presented in section 8. Finally, a summary of the results is provided in section 9.

2 The CMS detector

The central feature of the Compact Muon Solenoid (CMS) apparatus is a superconducting solenoid of 6 m internal diameter, providing a magnetic field of 3.8 T. Within the solenoid volume are a silicon pixel and strip tracker, a lead tungstate crystal electromagnetic calorimeter (ECAL), and a brass and scintillator hadron calorimeter (HCAL), each composed of a barrel and two endcap sections. Forward calorimeters extend the pseudorapidity (η) coverage provided by the barrel and endcap detectors. Muons are measured in gas-ionization detectors embedded in the steel flux-return yoke outside the solenoid. A more detailed description of the CMS detector, together with a definition of the coordinate system used and the relevant kinematic variables, can be found in refs. [47, 48].

Events of interest are selected using a two-tiered trigger system. The first level (L1), composed of custom hardware processors, uses information from the calorimeters and muon detectors to select events at a rate of around 100 kHz within a fixed latency of about $4 \mu\text{s}$ [49]. The second level, known as the high-level trigger, consists of a farm of processors running a version of the full event reconstruction software optimized for fast processing, and reduces the event rate to around 1 kHz before data storage [50].

3 Simulated samples

Samples of simulated events, produced with Monte Carlo (MC) event generators, are used in this analysis to estimate the contributions from background processes, model the correspondence between the observables at the generator and detector levels for unfolding, and compare the unfolded results with theoretical predictions.

For all simulated samples, additional pp interactions in the same or neighbouring bunch crossings (pileup) are generated with PYTHIA (v8.240) [51] and overlapped with the simulated hard interactions to match the pileup multiplicity measured in data. The detector response is modelled using a detailed simulation of the CMS detector, based on GEANT4 [52].

The response of the detector and event reconstruction of the $t\bar{t}b\bar{b}$ signal is modelled using a sample of $t\bar{t}b\bar{b}$ events generated using POWHEG-BOX-RES [53] and OPENLOOPS [54], referred to as the nominal $t\bar{t}b\bar{b}$ sample (or POWHEG+OL+P8 $t\bar{t}b\bar{b}$ 4FS), where the $t\bar{t}b\bar{b}$ MEs are calculated at NLO in QCD with massive b quarks [2], and matched with PYTHIA for parton showering and hadronization. The 4FS NNPDF3.1 next-to-NLO (NNLO) PDF set is used for the description of the proton structure. The b quark mass is set to 4.75 GeV and the POWHEG damping parameter that regulates the damping of real emissions in the NLO calculation when matching to the PS is set to a value of $h_{\text{damp}} = 1.379m_t$. Dynamic

μ_F and μ_R scales were chosen as $\mu_F = H_T/4$ and $\mu_R = \frac{1}{2} \prod_{i=t,\bar{t},b,\bar{b}} m_{T,i}^{1/4}$, respectively, where $H_T = \sum_{i=t,\bar{t},b,\bar{b},g} m_{T,i}$ and the transverse mass $m_{T,i} = \sqrt{m_i^2 + p_{T,i}^2}$, following ref. [1]. The choice of renormalization scale is a geometric average of top and bottom quark transverse masses and is a natural choice for taking into account the widely separated energy scales of both particles. The factorization scale is based on the maximum momentum of final state radiation that is still resummed into the PDF and hence is parameterized as the scalar sum of transverse top and bottom quark masses.

The sensitivity of the detector response to the modelling of the $t\bar{t}b\bar{b}$ process is evaluated using an alternative sample of $t\bar{t}b\bar{b}$ events, obtained from the inclusive $t\bar{t}$ simulation with POWHEG (v2) matched to PYTHIA (also referred to as POWHEG+P8 $t\bar{t}$ 5FS). The b quarks are assumed to be massless, and accordingly, five flavour scheme (5FS) proton PDFs (also NNPDF3.1 NNLO) are used in the calculation. In that sample, events with one additional b quark are modelled by the ME at leading order (LO) in QCD, while any further b quarks are generated by the PS. In particular, in $g \rightarrow b\bar{b}$ splittings only the emission of the gluon off initial-state partons or the top quarks is described at the ME-level. Hence, for a description of $t\bar{t}b\bar{b}$, the modelling of the splitting itself is necessarily handled by the PS. Since it has been shown that $g \rightarrow b\bar{b}$ splittings are the dominant mechanism for the production of additional b jets with top quark pairs, both in $t\bar{t}b$ and $t\bar{t}b\bar{b}$ events [2], this results in large uncertainties from the choice of the μ_R scale used for the strong coupling constant in the final-state PS. Conversely, in $t\bar{t}b\bar{b}$ samples generated in the 4FS, the dominant uncertainty comes from the choice of μ_R scale in the MEs, while uncertainties from the PS scale are smaller. The μ_F and μ_R scales in the POWHEG+P8 $t\bar{t}$ 5FS sample were set to $\mu_F = \mu_R = m_{T,t}$.

In addition to the above, we consider several alternative predictions of $t\bar{t}b\bar{b}$ cross sections, to be compared with the measurements. The generator settings used for the POWHEG+OL+P8 $t\bar{t}b\bar{b}$ 4FS and inclusive $t\bar{t}$ simulation, as well as a number of alternative generator setups described below, which are used for comparison of results.

The POWHEG inclusive $t\bar{t}$ generator is also interfaced with HERWIG (v7.13) for parton showering and hadronization [55, 56], using the CH3 underlying event tune [57] (referred to as POWHEG+H7 $t\bar{t}$ 5FS). Two other sets of simulated $t\bar{t}b\bar{b}$ events in the 4FS at NLO in QCD were obtained using MADGRAPH5_aMC@NLO (referred to as MG5_aMC+P8 $t\bar{t}b\bar{b}$ 4FS), and SHERPA (v2.2.4) [35, 58] with OPENLOOPS (referred to as SHERPA+OL $t\bar{t}b\bar{b}$ 4FS). The MG5_aMC+P8 $t\bar{t}b\bar{b}$ 4FS simulation is matched with PYTHIA and uses MADSPIN to decay the top quarks; the μ_F and μ_R scales are set to the sums of the transverse masses m_T of all partons in the final state ($\sum m_T$). In the SHERPA+OL $t\bar{t}b\bar{b}$ 4FS sample, the ME and PS are matched in the MC@NLO scheme, the PDF set used is NNPDF3.0 NNLO in the 4FS, and the scales are set to $\mu_F = H_T/2$ and $\mu_R = \prod_{i=t,\bar{t},b,\bar{b}} m_{T,i}^{1/4}$. Finally, MADGRAPH5_aMC@NLO is used to generate a sample of $t\bar{t}$ +jets events with up to two additional jets described at NLO QCD in the ME calculations, merged in the FxFx scheme [33] and using massless b quarks (referred to as MG5_aMC+P8 $t\bar{t}$ +jets FxFx 5FS). The μ_F and μ_R scales are set to $\mu_F = \mu_R = \sum m_T$, while the jet cutoff in the ME calculations is set to 20 GeV and the merging scale to 40 GeV. The generator settings of all $t\bar{t}b\bar{b}$ simulation approaches are summarized in table 1.

Contributions to the $t\bar{t}b\bar{b}$ phase space include b quarks produced in multiple parton interactions (MPI) as well as those produced in the matrix element or shower of the hard

Generator setup	Process/ME order	Generator/Shower	Tune	PDF set	h_{damp}	Scales
POWHEG+P8 $t\bar{t}$ 5FS	$t\bar{t}$ / NLO	POWHEG v2/ PYTHIA 8.240	CP5	5FS NNPDF3.1 NNLO	$1.379m_t$	$\mu_F = \mu_R = m_{T,t}$
POWHEG+H7 $t\bar{t}$ 5FS	$t\bar{t}$ / NLO	POWHEG v2/ HERWIG 7.13	CH3	5FS NNPDF3.1 NNLO	$1.379m_t$	$\mu_F = \mu_R = m_{T,t}$
POWHEG+OL+P8 $t\bar{t}b\bar{b}$ 4FS	$t\bar{t}b\bar{b}$ / NLO	POWHEG-BOX-RES/ PYTHIA 8.240	CP5	4FS NNPDF3.1 NNLO as 0118	$1.379m_t$	$\mu_R = \frac{1}{2} \prod_{i=t,\bar{t},b,\bar{b}} m_{T,i}^{1/4}$, $\mu_F = H_T/4$
SHERPA+OL $t\bar{t}b\bar{b}$ 4FS	$t\bar{t}b\bar{b}$ / NLO	SHERPA 2.2.4	SHERPA	4FS NNPDF3.0 NNLO as 0118	—	$\mu_R = \prod_{i=t,\bar{t},b,\bar{b}} m_{T,i}^{1/4}$, $\mu_F = H_T/2$
MG5_aMC+P8 $t\bar{t}b\bar{b}$ 4FS	$t\bar{t}b\bar{b}$ / NLO	MADGRAPH5_aMC@NLO v2.4.2/ PYTHIA 8.230	CP5	4FS NNPDF3.1 NNLO as 0118	—	$\mu_F = \mu_R = \sum m_T$
MG5_aMC+P8 $t\bar{t}$ +jets FxFx 5FS	$t\bar{t}$ +jets FxFx/ NLO [≤ 2 jets]	MADGRAPH5_aMC@NLO v2.6.1/ PYTHIA 8.240	CP5	5FS NNPDF3.1 NNLO	—	$\mu_F = \mu_R = \sum m_T$, qCut = 40 GeV, qCutME = 20 GeV

Table 1. Generator settings for different modelling approaches of $t\bar{t}b\bar{b}$ production. The top quark mass value is set to $m_t = 172.5$ GeV for all generator setups, and for the generator setups using massive b quarks, the b quark mass value is set to $m_b = 4.75$ GeV. In the scale settings, H_T corresponds to the scalar m_T sum, $H_T = \sum_{i=t,\bar{t},b,\bar{b},g} m_{T,i}$, and $m_{T,i} = \sqrt{m_i^2 + p_{T,i}^2}$ is the transverse mass. For generators setups using POWHEG the h_{damp} value is specified. Other generator setups do not use this parameter and are marked with (—).

process itself. For the POWHEG+P8 $t\bar{t}$ 5FS sample, this contribution is found to be 8–12% depending on the phase space considered. The dedicated $t\bar{t}b\bar{b}$ simulations have a much smaller contribution (<2% for POWHEG+OL+P8 $t\bar{t}b\bar{b}$ 4FS) because they do not include contributions from $t\bar{t}$ production in the primary parton scattering, where additional b quarks come only from MPI.

The predictions from the exclusive $t\bar{t}b\bar{b}$ samples are normalized using the total $t\bar{t}b\bar{b}$ cross sections obtained at NLO in QCD from the corresponding generators, whereas the inclusive $t\bar{t}$ and $t\bar{t}$ +jets samples are normalized using the total cross section for $t\bar{t}$ production, $\sigma(t\bar{t}) = 833.9$ pb, computed at NNLO in QCD using TOP++ (v2.0) [59], including soft-gluon resummations to next-to-next-to-leading logarithmic accuracy [60], and assuming a top quark mass of $m_t = 172.5$ GeV.

The POWHEG+P8 $t\bar{t}$ 5FS sample is used for the estimation of the inclusive $t\bar{t}$ background. However, using the nominal $t\bar{t}b\bar{b}$ sample for the signal in conjunction with the inclusive $t\bar{t}$ sample for the backgrounds would result in a double counting of the $t\bar{t}b\bar{b}$ contribution. This overlap is avoided by removing from the inclusive sample events containing at least one b jet not originating from a top quark (as defined below in section 4.2), with $p_T > 20$ GeV and $|\eta| < 2.4$. Conversely, in the nominal $t\bar{t}b\bar{b}$ sample only the events with at least one additional b jet are kept. Henceforth the $t\bar{t}B$ process is defined as the production of events passing these criteria. All simulated events in our measured fiducial phase spaces (see section 5.2) are found to pass the $t\bar{t}B$ selection criteria. The remaining events in the inclusive $t\bar{t}$ samples are categorized into a $t\bar{t}C$ contribution, with events containing at least one charm (c) jet (also with $p_T > 20$ GeV and $|\eta| < 2.4$) for which the simulation histories of the matched c hadrons do not include any top quark, and a $t\bar{t}$ +light contribution with all remaining events.

Simulated samples of minor backgrounds include single top quark production in the t and s channels, as well as tW production (collectively Single t); $t\bar{t}W$, $t\bar{t}H$, and $t\bar{t}Z$ production

Process	Group	ME order	Generator	Notes
tW	Single t	NLO	POWHEG v2	
t channel	Single t	NLO	POWHEG v2	MADSPIN for heavy particle decays [67]
s channel	Single t	NLO	MADGRAPH5_aMC@NLO v2.6.1	
t \bar{t} H	t \bar{t} H	NLO	POWHEG v2	
t \bar{t} Z	t \bar{t} V	NLO	MADGRAPH5_aMC@NLO v2.6.1	MADSPIN for heavy particle decays [67]
t \bar{t} W	t \bar{t} V	NLO	MADGRAPH5_aMC@NLO v2.6.1	FxFx merging up to 1 additional jet [33] MADSPIN for heavy particle decays [67]
W+jets	V+jets	LO	MADGRAPH5_aMC@NLO v2.6.5	MLM merging up to 4 additional jets [68]
Z+jets	V+jets	LO	MADGRAPH5_aMC@NLO v2.6.5	MLM merging up to 4 additional jets [68]

Table 2. Generator settings for various minor background samples simulated with POWHEG [34, 61–66] or MADGRAPH5_aMC@NLO [32]. The “Group” column refers to the grouping of processes in the maximum likelihood fits.

(collectively t \bar{t} X); and the production of Z/ γ^* or W in association with jets (collectively V+jets). An overview of the simulation settings used for these backgrounds is given in table 2.

For all of these minor background samples, the proton structure is described by the NNPDF3.1 set of NNLO PDFs [69], and parton showering and hadronization are simulated with PYTHIA, using the CP5 tune [70] for the underlying event description. The value of the Higgs boson mass is assumed to be 125 GeV, while the top quark mass value is set to $m_t = 172.5$ GeV. In the POWHEG samples, the h_{damp} parameter is set to a value of $h_{\text{damp}} = 1.379m_t$ as a part of the CP5 tune.

The cross sections for t \bar{t} , Z+jets, W+jets, and single top quark production are obtained at NNLO in QCD [59, 60, 71]. Samples for t \bar{t} W, t \bar{t} Z, and t \bar{t} H production are normalized to predictions at NLO in QCD [32, 66].

4 Event reconstruction

4.1 Detector-level object reconstruction and identification

A particle-flow (PF) algorithm [72] is applied to reconstruct and identify each individual particle in an event, with an optimized combination of information from the various elements of the CMS detector. The primary vertex (PV) is taken to be the vertex corresponding to the hardest scattering in the event, evaluated using tracking information alone, as described in section 9.4.1 of ref. [73].

The energy of electrons is determined using a multivariate algorithm from a combination of the electron momentum at the PV as determined by the tracker, the energy of the corresponding ECAL cluster, and the energy sum of all bremsstrahlung photons, obtained from the ECAL, spatially compatible with originating from the electron track [74]. Electrons are identified by placing requirements on the cluster shape in the ECAL, the track quality, and the compatibility between the track and the ECAL cluster. Electrons from photon conversions are rejected. The electrons used in this analysis are required to have $|\eta| < 2.5$ and $p_T > 29$ GeV for data collected in 2016. In data collected in 2017–2018, the p_T threshold is raised to 34 GeV, except for electrons with $|\eta| < 2.1$, for which the threshold is 30 GeV. These requirements are

chosen to be as low as possible so as to maximize the signal selection efficiency while staying above the trigger thresholds in the respective data-taking periods. Electrons with a cluster pseudorapidity $|\eta_{\text{SC}}|$ between 1.444 and 1.566 are not considered, in order to avoid the gap between the barrel and endcap ECAL sections. The average identification efficiency for the primary electrons is about 70%, including the isolation requirements described below. Another looser set of requirements with 95% selection efficiency is also considered, with $p_{\text{T}} > 15$ GeV and $|\eta| < 2.5$, and relaxed identification criteria. These define the “veto” electrons, which are used to reject events with more than one lepton.

The p_{T} of muons is obtained from the curvature of the corresponding track, combining information from the inner tracker and the outer muon detector system [75]. The muons are identified based on the quality of the combined track fit and on the number of hits in the different tracking detectors, with an efficiency of about 95%. Muons are required to have $|\eta| < 2.4$ and $p_{\text{T}} > 26$ (29) GeV in 2016 (2017–2018), while “veto” muons are defined with $p_{\text{T}} > 15$ GeV, $|\eta| < 2.4$, and looser identification criteria.

Electron or muon tracks are required to have a longitudinal and transverse distance to the PV smaller than 0.5–5 mm, depending on the lepton flavour and $|\eta|$. In order to suppress backgrounds from hadrons misidentified as leptons, or from leptons produced from hadrons decaying inside of jets, the leptons are required to be isolated from hadronic activity. The lepton isolation is defined as the ratio between the scalar p_{T} sum of all PF candidates in a cone around the lepton excluding the lepton itself, and the lepton p_{T} . The cone size is $\Delta R < 0.4$ (0.3) for electrons (muons). The isolation is corrected by removing contributions from pileup [76]. The maximum primary (veto) electron isolation varies between 0.03–0.08 (0.20–0.27), decreasing with p_{T} and increasing with $|\eta_{\text{SC}}|$. For primary (veto) muons, the isolation is required to be < 0.15 (< 0.25). Residual differences between lepton reconstruction, identification, and isolation efficiencies in data and simulation are corrected. The efficiencies are measured as a function of lepton p_{T} and η in data samples enriched in $Z \rightarrow \ell^+ \ell^-$ events using a “tag-and-probe” method. For electrons (muons), the corrections are between 1–5% ($< 2\%$) with uncertainties of less than 2 (1)% [74, 75].

The energy of charged hadrons is determined from a combination of their momentum measured in the tracker and the matching ECAL and HCAL energy deposits, corrected for the response function of the calorimeters to hadronic showers. The energy of neutral hadrons is obtained from the corresponding corrected ECAL and HCAL energies. Hadronic jets are reconstructed by clustering PF candidates using the anti- k_{T} algorithm [77], as implemented in the FASTJET package [78], with a distance parameter of $R = 0.4$. Jet momentum is determined as the vectorial sum of all particle momenta in the jet, and is found from simulation to be, on average, within 5–10% of the true momentum over the whole p_{T} spectrum and detector acceptance. The jet energy resolution (JER) amounts typically to 10 (15–20)% at 100 (30) GeV [79]. Pileup interactions can contribute additional tracks and calorimetric energy depositions, increasing the apparent jet momentum. To mitigate this effect, tracks identified to be originating from pileup vertices are discarded and an offset correction is applied to correct for remaining contributions [76]. Jet energy corrections are derived from simulation studies so that the average measured energy of jets becomes identical to that of jets at generator level. In-situ measurements of the momentum balance in dijet,

γ +jet, Z+jet, and multijet events are used to determine any residual differences between the JES in data and in simulation, and appropriate corrections are applied to data [79]. Additional selection criteria are applied to each jet to remove jets potentially falsely reconstructed, with dominant contributions from instrumental effects or reconstruction failures. A multivariate algorithm is used to identify and remove jets likely originating from pileup interactions [76]. Jets used in the analysis are required to have $p_T > 30$ GeV and $|\eta| < 2.4$ and to be separated from the selected electron or muon by $\Delta R > 0.4$.

The DEEPJET algorithm is used to identify b jets [80–82]. This algorithm uses a deep neural network (DNN) discriminant to combine information from charged and neutral PF candidates clustered in the jet with features of secondary vertices within the jet. Jets are labelled as b tagged if they pass a “medium” working point of the discriminant, corresponding to an efficiency for correctly identifying b jets in $t\bar{t}$ events of 75–80% and to a misidentification probability of about 15–17% for c jets and about 1.5–2% for other jets. We will also refer to jets failing the medium working point as light-tagged jets. Another, more restrictive, (“tight”) b tagging working point is also used, yielding a misidentification rate of 2.5–3.5 for c jets and 0.15–0.25% for other jets. This improved mistagging rate comes at the cost of reducing the b jet identification efficiency to around 60%.

The b jet (mis)identification probabilities in the simulation are corrected to match the efficiencies measured in data at both the medium and tight working points [80]. The tagging efficiency in data for b jets is obtained from a combination of five different measurements, three of which make use of a sample of multijet events enriched in b jets, by requiring jets to contain a muon, while the remaining two use samples enriched in $t\bar{t}$ events and containing respectively one or two isolated electrons or muons. These three samples are statistically independent of each other and with the events used in this analysis and yield compatible measurements of the b tagging efficiency. The b tagging efficiency correction factors are measured as functions of jet p_T and vary between 0.9–1.0 depending on the tagger working point, the data period, and the jet p_T , and have uncertainties of up to 10%. For c jets, the misidentification efficiency in the simulation is corrected using the same correction factors as for b jets, but with double their uncertainty. This has been shown to cover the true c jet misidentification probability [80]. For other jets, the misidentification probability is measured as a function of jet p_T in an inclusive sample of multijet events. The corresponding simulation-to-data correction factors vary between 0.6–1.9, depending on the tagger working point, the data period, and the jet p_T . Their uncertainties vary from 10–30%.

4.2 Particle-level object definitions

The fiducial phase space regions and observables for the inclusive and differential cross sections reported in this paper are defined based on the properties of stable final-state particles, with proper lifetimes $\tau_0 > 10$ mm. Particles with $|\eta| > 5$ are not considered. The definition of objects at the particle level, which follow closely those at the detector level introduced earlier, is described in the following.

Prompt particle-level electrons are selected with $p_T > 29$ GeV and $|\eta| < 2.5$ and are “dressed” with photons from final-state radiation (FSR) by adding to their four-momentum the momentum of any photon within $\Delta R < 0.1$. Prompt particle-level muons are required to

have $p_T > 26$ GeV and $|\eta| < 2.4$. A similar FSR dressing procedure as for the electrons is applied to the muons. Furthermore, “veto” electrons and muons are defined by relaxing the thresholds to $p_T > 15$ GeV, and are then used to reject events with more than one lepton. The same dressing procedure as described above is also applied for the veto electrons and muons.

Jets are obtained by clustering all particles, excluding any neutrinos, using the anti- k_T algorithm with $R = 0.4$. They are required to have $p_T > 25$ GeV and $|\eta| < 2.4$ and are not considered if they are within $\Delta R < 0.4$ of a prompt electron or muon. At the particle level, the flavour of jets is defined unambiguously by rescaling the momenta of all generated b hadrons to a negligible value and including them in the jet clustering procedure [83]. Jets thus matched to at least one b hadron are considered as b jets, while all remaining jets are labelled as light jets.

For the purpose of defining observables for the second approach introduced in section 1, and the definition of the $t\bar{t}B$ process, we also define as “additional” b jets those particle-level b jets for which the simulation histories of the matched b hadrons do not include any top quark. This definition is therefore not uniquely based on stable particles, since it refers to the simulated parton-level event content, which is specific to each generator and not provided by every generator.

5 Event selection and definition of fiducial phase space regions and observables

5.1 Event selection

Data are collected using a set of triggers requiring the presence of one isolated electron or muon. The lepton p_T thresholds applied by these single-lepton triggers vary between 27–32 (24–27) GeV for electrons (muons), depending on the data-taking period. In addition, triggers selecting events with one isolated electron and a scalar jet p_T sum (H_T) above 150 GeV are used, allowing the electron p_T threshold to be set to 28 GeV for $|\eta| < 2.1$. The trigger selections used in collision data are also applied in the simulation, and residual differences between data and simulation are corrected. For electron triggers, efficiencies are measured in a control region enriched with $t\bar{t}$ events, collected using triggers requiring the presence of missing transverse momentum. These present a negligible correlation with the triggers used in this analysis. Muon trigger efficiencies are measured using the tag-and-probe technique in a control region enriched in $Z \rightarrow \ell^+ \ell^-$ events.

Data from the different data-taking periods (2016, 2017, 2018) are analyzed separately and are only combined at the likelihood level, as will be explained in section 6.3. Additionally, data collected in 2016 are split into two groups, the “2016preVFP” and “2016postVFP” eras, due to substantial changes in the detector conditions between them. At the beginning of the 2016 period, saturation effects in the readout chips of the strip tracker led to lower signal-to-noise ratios and fewer hits on tracks. This issue was mitigated by changing the feedback pre-amplifier bias voltage (VFP) during the 2016 data-taking run [84].

5.2 Fiducial phase space regions

The measured cross sections are derived from the fiducial phase space regions and observables described below. The definitions at the particle and detector levels follow each other as closely as possible in order to minimize extrapolations outside of the detector acceptance. Both at the particle and detector levels, events are first required to have exactly one primary electron or muon, and no additional “veto” electrons or “veto” muons, as defined in section 4. At the particle level, events with electrons and muons are combined, and no requirement is placed on the decay channels of the top quarks, so that the fiducial phase space regions can also contain $t\bar{t}b\bar{b}$ events in which both top quarks decay leptonically, but one lepton is outside the detector acceptance and is not selected as a “veto” lepton. Furthermore, electrons and muons produced indirectly through the decay of a tau lepton are included. At the detector level, events with electrons or muons are classified into two distinct channels in this measurement.

Four different and partially overlapping phase space regions are then considered, each targeting different aspects of $t\bar{t}b\bar{b}$ production, as introduced in section 1. The most inclusive selection considered, labelled as 5j3b (or ≥ 5 jets: $\geq 3b$) and targeting $t\bar{t}b$, requires the presence of at least five jets, of which at least three must be b(-tagged) jets at particle (detector) level. In a second selection, labelled as 6j4b (or ≥ 6 jets: $\geq 4b$) and targeting $t\bar{t}b\bar{b}$, at least six jets are required, of which at least four are b(-tagged) jets. Two additional selections are defined which help target the properties of additional light jets produced in the event, targeting $t\bar{t}bj$ and $t\bar{t}b\bar{b}j$. These are the 6j3b3l phase space (also labelled ≥ 6 jets: $\geq 3b$, ≥ 3 light), requiring at least six jets, including at least three b(-tagged) jets and at least three light(-tagged) jets; and the 7j4b3l phase space (also labelled ≥ 7 jets: $\geq 4b$, ≥ 3 light), requiring at least seven jets, including at least four b(-tagged) jets and at least three light(-tagged) jets.

5.3 Observables

As described in section 1, we consider two classes of observables for unfolding. In the first class, all observables are defined using stable particles exclusively, without reference to any simulated event history. This implies that no strict distinction is made between b jets originating from the decay of top quarks, or from additional radiation. In this way, there is a close correspondence between particle- and detector-level observables (see sections 4.1 and 4.2), and the definition of the observables is independent of any specific event generator. In order to probe different features of $t\bar{t}b\bar{b}$ production, and in particular the properties of b jets produced from QCD radiation, we distinguish different categories of objects using simple kinematic criteria.

In the 5j3b phase space, we focus the measurements on the b jet with the third-largest p_T , which at the generator level is a true additional b jet (not coming from top quark decays) in the nominal signal simulation in approximately 49% of 5j3b events. This identification allows for the study of the properties of the additional b jets even in the case where they cannot be individually resolved. In that phase space, we measure, each independently, the number of jets in the event (N_{jets}), the number of b jets in the event (N_b), the p_T and $|\eta|$ of the 3rd b jet ($p_T(b_3)$, $|\eta(b_3)|$), the scalar p_T sum of all jets (H_T), and the scalar p_T sum of all b jets (H_T^b).

In the 6j4b phase space where there are at least four b jets in the event, in order to measure properties that are expected to be sensitive to the modelling of gluon splittings to $b\bar{b}$,

we define the “extra” b jets as the pair of b jets with the smallest angular separation, defined by

$$\Delta R_{bb} = \sqrt{(\Delta\phi_{bb})^2 + (\Delta\eta_{bb})^2}, \quad (5.1)$$

where $\Delta\phi_{bb}$ and $\Delta\eta_{bb}$ are the angular separations of the pair of b jets in azimuthal angle ϕ (in radians) and η , respectively. We denote that pair by bb^{extra} in the following. With this strategy, in the fiducial generator-level phase space, the two bb^{extra} jets correspond to true additional b jets in $\sim 49\%$ of events in the 6j4b phase space. Other choices, such as picking the pairs of jets with the 3rd and 4th largest p_T , result in smaller probabilities of correctly identifying the additional jets ($\sim 28\%$).

In the 6j4b phase space, we measure, each independently, the number of jets in the event, the p_T and $|\eta|$ of the 3rd and 4th b jets ($p_T(b_3), p_T(b_4), |\eta(b_3)|, |\eta(b_4)|$), the scalar jet p_T sum (H_T), and the scalar b-jet p_T sum (H_T^b). Considering all possible pairs of distinct b jets, we independently measure the average ΔR_{bb} over all pairings ($\Delta R_{bb}^{\text{avg}}$), and the largest invariant mass (m_{bb}^{max}). For the bb^{extra} pair, we measure, each independently, the distance in η between the two b jets of the pair ($\Delta R(bb^{\text{extra}})$), the invariant mass of the pair ($m(bb^{\text{extra}})$), the p_T and $|\eta|$ of the pair ($p_T(bb^{\text{extra}}), |\eta(bb^{\text{extra}})|$), and the p_T and $|\eta|$ of the leading (b_1^{extra}) and sub-leading (b_2^{extra}) b jets in the pair ($p_T(b_1^{\text{extra}}), p_T(b_2^{\text{extra}}), |\eta(b_1^{\text{extra}})|, |\eta(b_2^{\text{extra}})|$).

In the 6j3b3l and 7j4b3l phase space regions, we target the properties of the additional light jets (see sections 4.1 and 4.2). In each of these phase space regions, we measure the scalar p_T sum of the light jets in the event (H_T^{light}). We then remove from consideration the pair of light jets with the invariant mass closest to the W boson mass obtained from a fit to the two detector- or particle-level jets matched to the W boson decay in simulation. We then measure, independently, the p_T of the leading remaining light jet ($p_T(lj_1^{\text{extra}})$), and the $\Delta\phi$ between that light jet and the lowest p_T b jet ($|\Delta\phi(lj_1^{\text{extra}}, b_{\text{soft}})|$). The latter variable probes the amount of recoil against additional QCD radiation in $t\bar{t}b\bar{b}$ events that is absorbed by the softest b jet [1]. In the 6j3b3l and 7j4b3l phase space regions, the leading remaining light jet corresponds to a light jet not from top quark decays in $\sim 94\%$ of cases. The softest b jet is an additional b jet in ~ 50 (65)% of cases in the 6j3b3l (7j4b3l) phase space.

In the second class of observables, we focus on the 6j4b phase space and explicitly target the b jets that do not originate from decaying top quarks. At the particle level, these are the additional b jets as defined in section 4.2, labelled as bb^{add} . In case more than two additional b jets are present, the two additional b jets leading in p_T are selected. At the detector level, the pair of b-tagged jets most consistent with the true additional b jets is identified using a DNN discriminant, described in section 6.2. For the bb^{add} pair defined in this way, we measure the opening angle between the two b jets of the pair ($\Delta R(bb^{\text{add}})$), the invariant mass of the pair ($m(bb^{\text{add}})$), the p_T and $|\eta|$ of the pair ($p_T(bb^{\text{add}}), |\eta(bb^{\text{add}})|$), and the p_T and $|\eta|$ of the leading and subleading b jets in the pair ($p_T(b_1^{\text{add}}), p_T(b_2^{\text{add}}), |\eta(b_1^{\text{add}})|, |\eta(b_2^{\text{add}})|$).

A summary of all observables is given in table 3. In the measurement of each observable we also measure the inclusive cross section in the respective phase space (σ_{fid}).

6 Signal extraction and unfolding

In the selected regions with at least three b-tagged jets, the data are highly enriched in $t\bar{t}$ +jets events, which consist of about 30% $t\bar{t}B$, 20% $t\bar{t}C$, and 50% $t\bar{t}$ +light events, with minor

Observable		5j3b	6j4b	6j3b3l	7j4b3l
σ_{fid}	Inclusive cross section	✓	✓	✓	✓
Global observables					
N_{jets}	Jet multiplicity	✓	✓		
N_{b}	b jet multiplicity	✓			
H_{T}^{j}	Scalar p_{T} sum of all jets	✓	✓		
H_{T}^{b}	Scalar p_{T} sum of all b jets	✓	✓		
$H_{\text{T}}^{\text{light}}$	Scalar p_{T} sum of all light jets			✓	✓
Observables related to b jets					
$p_{\text{T}}(\text{b}_3)$	p_{T} of third hardest b jet	✓	✓		
$ \eta(\text{b}_3) $	$ \eta $ of third hardest b jet	✓	✓		
$p_{\text{T}}(\text{b}_4)$	p_{T} of fourth hardest b jet		✓		
$ \eta(\text{b}_4) $	$ \eta $ of fourth hardest b jet		✓		
Observables considering all pairs of b jets (bb)					
$\Delta R_{\text{bb}}^{\text{avg}}$	Average ΔR of all bb pairs		✓		
$m_{\text{bb}}^{\text{max}}$	Highest invariant mass among all bb pairs		✓		
Observables related to the pair of b jets closest in ΔR (bb^{extra})					
$p_{\text{T}}(\text{b}_1^{\text{extra}})$	p_{T} of leading extra b jet		✓		
$ \eta(\text{b}_1^{\text{extra}}) $	$ \eta $ of leading extra b jet		✓		
$p_{\text{T}}(\text{b}_2^{\text{extra}})$	p_{T} of subleading extra b jet		✓		
$ \eta(\text{b}_2^{\text{extra}}) $	$ \eta $ of subleading extra b jet		✓		
$\Delta R(\text{bb}^{\text{extra}})$	ΔR of bb^{extra} pair		✓		
$ \eta(\text{bb}^{\text{extra}}) $	$ \eta $ of bb^{extra} pair		✓		
$m(\text{bb}^{\text{extra}})$	invariant mass of bb^{extra} pair		✓		
$p_{\text{T}}(\text{bb}^{\text{extra}})$	p_{T} of bb^{extra} pair		✓		
Observables related to the pair of b jets not from $t\bar{t}$ decay ($\text{bb}^{\text{add.}}$)					
$p_{\text{T}}(\text{b}_1^{\text{add.}})$	p_{T} of leading additional b jet		✓*		
$ \eta(\text{b}_1^{\text{add.}}) $	$ \eta $ of leading additional b jet		✓*		
$p_{\text{T}}(\text{b}_2^{\text{add.}})$	p_{T} of subleading additional b jet		✓*		
$ \eta(\text{b}_2^{\text{add.}}) $	$ \eta $ of subleading additional b jet		✓*		
$\Delta R(\text{bb}^{\text{add.}})$	ΔR of $\text{bb}^{\text{add.}}$ pair		✓*		
$ \eta(\text{bb}^{\text{add.}}) $	$ \eta $ of $\text{bb}^{\text{add.}}$ pair		✓*		
$m(\text{bb}^{\text{add.}})$	invariant mass of $\text{bb}^{\text{add.}}$ pair		✓*		
$p_{\text{T}}(\text{bb}^{\text{add.}})$	p_{T} of $\text{bb}^{\text{add.}}$ pair		✓*		
Observables related to extra light jets					
$p_{\text{T}}(\text{l}_{\text{j}1}^{\text{extra}})$	p_{T} of leading extra light jet			✓	✓
$ \Delta\phi(\text{l}_{\text{j}1}^{\text{extra}}, \text{b}_{\text{soft}}) $	$\Delta\phi$ of leading extra light jet and softest b jet			✓	✓

Table 3. Description of all measured observables for each of the four fiducial phase space regions. Observables marked as (✓*) rely on the definition of additional b jets, and do not fully correspond to the 6j4b fiducial phase space defined at the particle level, but also require the presence of b jets without top (anti)quarks in their simulated history.




Figure 1. Jet (left) and b-tagged jet (right) multiplicity with the ≥ 5 jets: $\geq 3b$ selection prior to any fit, shown for both lepton channels and all data periods combined. For the purpose of visualisation, the contributions from simulation have been scaled by a common factor (0.98) to match the yield in data. The $t\bar{t}X$ contribution includes the $t\bar{t}H$, $t\bar{t}W$, and $t\bar{t}Z$ processes. The $t\bar{t}B$ contribution includes the superset of fiducial and out of acceptance $t\bar{t}b\bar{b}$ events, following the definitions used for the measurements. The shaded bands include all a priori uncertainties described in section 7, including the $t\bar{t}B$ cross section uncertainty estimated from the nominal $t\bar{t}b\bar{b}$ simulation. Only effects on the shape of the distributions are considered. The last bins also contain the overflow.

background contributions, in descending order of importance, from single top quark production, $t\bar{t}H$, V +jets, $t\bar{t}Z$, and $t\bar{t}W$ processes. Events from $t\bar{t}H$, $t\bar{t}Z$, or $t\bar{t}W$ processes are referred to as a combined $t\bar{t}X$ contribution if a distinction is not necessary. In the other selected regions with at least four b-tagged jets, the $t\bar{t}B$ contribution is about two thirds of all events. Background contamination from inclusive multijet production, where a jet or a nonisolated lepton is misidentified as a primary lepton, has been verified with simulated multijet events to be negligible at the level of the precision of this measurement ($\leq 1\%$ in the $5j3b$ phase space), and this background is therefore not considered further. The prefit composition of the events as a function of number of jets and number of b-tagged jets at the medium working point is shown in figure 1, with the $5j3b$ selection applied. The events displayed in these distributions are the superset of events used for the measurements presented in section 8.

For each observable, the cross section is measured using a dedicated binned maximum likelihood fit, which simultaneously yields the inclusive cross section in the corresponding fiducial phase space and the unfolded normalized differential cross section of the observable (section 6.3). In order to improve the separation between the signal and background components, thereby better constraining the background contributions using the data, we use an “ancillary” variable that divides the detector-level selections into signal- and background-enriched categories, which are then fitted simultaneously.

6.1 Ancillary variable

We use the number of jets passing the tight b tagging working point as an ancillary variable. The distributions of the number of tight b-tagged jets with the $5j3b$ and $6j4b$ selections applied are shown in figure 2. The regions with three or more tight b-tagged jets are highly




Figure 2. Number of jets b tagged at the tight working point with the ≥ 5 jets: $\geq 3b$ (left) and ≥ 6 jets: $\geq 4b$ selections (right) prior to any fit, shown for all lepton channels and years combined. For the purpose of visualisation, the contributions from simulation have been scaled by a common factor (0.98 on the left, 0.95 on the right) to match the yield in data. The $t\bar{t}B$ contribution includes the superset of fiducial and out of acceptance $t\bar{t}b\bar{b}$ events, following the definitions used for the measurements. The shaded bands include all uncertainties described in section 7, including the $t\bar{t}B$ cross section uncertainty estimated from the nominal $t\bar{t}b\bar{b}$ simulation. Only effects on the shape of the distributions are considered. The last bins also contain the overflow. The vertical dashed lines indicate the ancillary regions.

enriched in $t\bar{t}B$ signal events, as defined in section 3, since they typically feature three or four true b jets after fiducial selection, whereas the regions with 0–2 tight b -tagged jets are dominated by $t\bar{t}C$ or $t\bar{t}$ +light backgrounds where c or light jets have been misidentified as medium b -tagged jets and, therefore, serve as control regions for these backgrounds. In the fits of observables in the 5j3b and 6j3b3l phase space regions we use three ancillary regions containing zero or one, two, or three or more tight b -tagged jets. In the other phase space regions we only use two ancillary regions: one containing events with fewer than three tight b -tagged jets, and another one containing events with at least three tight b -tagged jets. The use of ancillary variables is not only useful for obtaining signal-enriched regions but also allows for the constraint of b tagging related uncertainties in the fit.

6.2 Multivariate algorithm for jet assignment

For the eight observables in the 6j4b phase space with explicit identification of the additional b jets, a multivariate algorithm based on a DNN is used to identify the pair of b -tagged jets most consistent with the true additional b jets as defined by the generator-level information. The four b -tagged jets in an event with the highest p_T , in the following referred to as candidate jets, are grouped into six permutations of candidate jet pairs, depending on their ranking in p_T . Permutations within a pair are not treated separately. A detector-level jet is considered correctly identified as an additional jet if it lies within an angular distance of $\Delta R < 0.4$ to an additional particle-level b jet.

An illustration of the DNN architecture is shown in figure 3. The DNN makes use of two sets of input variables, targeting jet-specific input information and global event information separately.




Figure 3. Structural representation of the neural network used for the assignment of the additional b-jet pair. The neural network uses two sets of input variables: global event information is connected to three dense network layers, and jet-specific information is connected via convolutional network layers (CNN) and a long short-term memory (LSTM) cell. The input sequences are concatenated into one dense layer. The output layer consists of six nodes, each representing one of the six possible candidate jet combinations.

For the input variables targeting jet-specific information, i.e. features of the four candidate jets, an automated feature engineering is performed for each jet using five convolutional neural network (CNN) layers [85] with filter matrices of size 1×1 [81]. The size of the filter matrices is chosen such that they aggregate the different features, separately for each jet, into higher-level features of the consecutive layers. These layers are followed by a long short-term memory (LSTM) cell [86], where the independent jet features are combined into a sequence, allowing the network to learn from the correlations of the jet features. The five features used for each candidate jet are the p_T , η , a flag indicating whether it passes the tight b tagging working point, the angular separation (ΔR) with the charged lepton, and the invariant mass with the charged lepton. The 30 input variables targeting global event information include the properties of the six dijet combinations of the four candidate b jets. These input variables are connected via three dense network layers. The inputs consist of the scalar p_T sum of the four candidate b jets, the p_T , η , and ϕ of the charged lepton, the $\Delta\phi$, $\Delta\eta$, and invariant mass of the dijet combinations, the ΔR of the dijet combinations and the charged lepton, and the jet and b-tagged jet multiplicities. Both input sequences are concatenated into one dense layer, which is connected to an output layer consisting of six nodes, each representing one of the six possible candidate jet combinations. The pair of b-tagged jets with the highest DNN output value per event is chosen as the additional b-jet pair to be used further in the analysis.

The training of the DNN is performed with simulated events passing the fiducial 6j4b phase space definition using the KERAS [87, 88] package with a TENSORFLOW [89] backend and the ADAM optimizer [90]. In the training, the categorical cross-entropy loss function is minimized. To reduce biases from imbalances in the numbers of events for the different dijet categories, events are weighted in the training such that each category has the same number of weighted events. Potential overfitting is mitigated using a dropout percentage of 10% [91] and an early-stopping procedure to stop the training if no decrease in the loss minimization has been achieved for 20 epochs (iterations over the training data set) on a set of spectator

events not used for the training (validation data set). In order not to bias the evaluation of the DNN, events from the POWHEG+P8 $t\bar{t}$ 5FS simulation passing the fiducial 6j4b definition are used in the training, while the evaluation for the measurement is performed with the nominal $t\bar{t}b\bar{b}$ sample. It has been validated that the performance of the DNN is independent of the choice of which of those samples of simulated $t\bar{t}b\bar{b}$ events is used for the training or the evaluation. This implies that the measurements of the DNN-based observables are not biased by the choice of signal model used for training the DNN.

The performance of the DNN is evaluated in the simulation based on the fraction of events in which it selects the correct pair of additional b jets (accuracy). The accuracy of the DNN is determined to be about 49%, which represents a significant increase in identification accuracy compared to choosing the two b jets closest in ΔR (as in section 5), which only yields an accuracy of about 41%.

6.3 Unfolding methodology

The observed distributions are unfolded to the particle level by removing estimated background contributions and correcting for resolution, acceptance and efficiency effects of the detector and event reconstruction so that the measured distributions can be directly compared to theoretical predictions provided by MC event generators.

Unfolding is performed through a maximum likelihood fit by constructing, for each observable, a statistical model that links the distributions at the particle and detector levels. The values of the particle-level cross sections which maximize the agreement between the predicted detector-level distributions and the observed data are determined from the fit. In these models, freely floating parameters of interest determine the total cross section of the signal process in the corresponding phase space, as well as the normalized differential cross section of the signal process in discrete bins of the considered observable. The models are constructed using the simulated signal and background samples described in section 3 and include nuisance parameters that model the effect of systematic uncertainties in the signal and background predictions. The fit is performed by minimizing the negative-log-likelihood (NLL) of the data with respect to the parameters of the model. The calculation of the NLL and its minimization are performed using SMOOFIT [92, 93], itself relying on the JAX package [94, 95] for automatic differentiation of the NLL function, providing fast and numerically stable evaluations of the NLL gradients and of the second derivative (Hessian) matrix. Both the central values and confidence intervals for the absolute and normalized differential cross sections are extracted from the fit.

The electron and muon channels as well as the four data-taking eras (2016preVFP, 2016postVFP, 2017, and 2018) and the ancillary variable regions are combined at the likelihood level. The likelihood used to measure the unfolded distributions can be written as

$$L(\vec{\mu}, \vec{\alpha}) = \left[\prod_{e,i} \text{Poi} \left(D_{e,i} \middle| S_{e,i}(\vec{\mu}, \vec{\alpha}) + \sum_{p \in \text{bkg.}} N_{e,i}^p(\vec{\alpha}) \right) \right] \mathcal{N}(\vec{\alpha}), \quad (6.1)$$

where the parameters $\vec{\mu}$ are freely-floating parameters of interest, $\vec{\alpha}$ are profiled nuisance parameters used to model systematic uncertainties, $D_{e,i}$ are the observed yields in data-taking era e and detector-level bin i (including the lepton channels and ancillary regions), $N_{e,i}^p$

are the predicted yields of background process p in era e and bin i , $S_{e,i}$ are the predicted signal yields in era e and bin i , $\text{Poi}(d|\nu)$ is the Poisson probability mass function for counts d with mean ν , and $\mathcal{N}(\vec{\alpha})$ is the Gaussian constraint term (with mean of zero and width of one) of the nuisance parameters.

We denote by M_{ij}^e the expected number of signal events in the fiducial phase space in the detector-level bin i and generator-level bin j for the data-taking era e . For every generator-level bin there are, therefore, up to

$$(4 \text{ eras}) \times (2 \text{ channels}) \times (2 \text{ or } 3 \text{ anc. regions}) = 16 \text{ or } 24 \text{ detector-level bins}, \quad (6.2)$$

depending on the phase space. In addition, the detector-level binning is chosen to be finer than the generator-level binning, with every bin at the detector level being about half the width of the generator-level bins (except for discrete observables such as the number of jets or b jets, and the observables in the 7j4b3l region where the data yields are the lowest). This improves the separation between signal and background. Including other control regions in the fit or using a finer binning at the detector level has not been found to provide any significant further improvements in the sensitivity of the measurements. The expected event yields in era e can be written as $M_{ij}^e = \mathcal{L}_e \sigma_j^0 K_{ij}^e$, where \mathcal{L}_e is the integrated luminosity in era e , σ_j^0 is the prefit cross section in bin j estimated using the nominal $t\bar{t}b\bar{b}$ sample, and K_{ij}^e are response matrices, i.e. the probability for a simulated event in era e and generator-level bin j to be reconstructed and selected in detector-level bin i . The total expected signal yields S in eq. (6.1) are computed as functions of the parameters of interest as

$$S_{e,i}(\vec{\mu}, \vec{\alpha}) = \mu_{\text{fid}} \sum_{j=1}^n \mu_j M_{ij}^e(\vec{\alpha}), \quad (6.3)$$

where $\mu_{\text{fid}} = \sigma_{\text{fid}}/\sigma_{\text{fid}}^0$ is the signal-strength modifier for the inclusive cross section and μ_j are the parameters varying the fraction of signal events in each generator-level bin j . To preserve unity, the yields in the last generator-level bin n are not scaled independently, but as a function of the other bins as

$$\mu_n(\mu_1 \dots \mu_{n-1}) = \frac{1}{F_n} \left(1 - \sum_{i=1}^{n-1} \mu_i F_i \right). \quad (6.4)$$

where $F_j = \sigma_j^0/\sigma_{\text{fid}}^0 = \sigma_j^0/\sum_{i=1}^n \sigma_i^0$ is the *a priori* fractional cross section in bin j . In this way, the measured inclusive cross section is directly obtained as $\hat{\sigma}_{\text{fid}} = \hat{\mu}_{\text{fid}} \sigma_{\text{fid}}^0$. Furthermore, the measured normalized differential cross section in bin j is extracted as $1/\hat{\sigma}_{\text{fid}} d\hat{\sigma}_j/dX = \hat{\mu}_j F_j/w_j$, where w_j is the width of generator-level bin j . Thus, the yields in the last bin n are constrained through eq. (6.4) to conserve the total signal normalization for a given value of μ_{fid} . The covariance matrix of all fit parameters is obtained from the inverse Hessian of the NLL at the minimum and is used to compute confidence intervals on the measured cross sections. We have verified that the resulting intervals are equivalent to those obtained by finding the level crossings of the profiled NLL, and we have validated the frequentist coverage properties of the confidence intervals using pseudo-experiments.

Figure 4 shows the response matrix for an example observable, both with and without ancillary variables. For the purpose of visualization, the response matrices shown are averaged

across eras and lepton channels and are normalized so that the values sum to 100% across each column (generator-level bin). Most response matrices are highly diagonal, which can be explained by the close correspondence between the particle- and detector-level definitions of the observables. Compared to these purely particle-level based observables, there is considerably more migration for the observables based on the additional b jets, which are defined via the simulated history at the particle level but only identified with limited accuracy by the DNN at the detector level.

Despite a careful definition of the fiducial phase space at the generator and detector level, a fraction of the events selected at the detector level consists of simulated $t\bar{t}B$ events that do not pass the corresponding generator-level fiducial phase space definitions. There is an inherent ambiguity on how to treat this contribution, labelled as “out of acceptance” (OOA). While technically part of the $t\bar{t}b\bar{b}$ process being measured, these events are outside of the fiducial phase space and should not be included in the measured cross sections. Therefore, we treat this OOA contribution as a background and assign to it theoretical systematic uncertainties estimated using the nominal $t\bar{t}b\bar{b}$ simulation. This OOA contribution constitutes about 40% of selected $t\bar{t}B$ events in the 6j4b and 7j4b3l regions and reduces to about 20 and 12% in the more inclusive 6j3b3l and 5j3b regions, respectively.

Two distinct mechanisms explain the presence of OOA events in the detector-level selection. First, mismeasurements of the energy or direction of leptons or jets can cause events that are outside the generator-level phase space to pass the detector-level selection. This class of events features a similar topology to the fiducial signal events. Second, the misidentification of light or c jets as b jets can lead to events with only two or three b jets within the fiducial acceptance to be selected in the regions requiring the presence of three or four b-tagged jets, respectively. This latter mechanism is dominant in the two regions with four b jets, 6j4b and 7j4b3l, and explains why the OOA contribution is larger in these regions, while the fractions of OOA events are reduced in the signal-enriched ancillary regions. However, $t\bar{t}B$ events with only three b jets selected within the generator-level fiducial volume feature a distinct topology from those with four b-tagged jets, since they might either contain a soft (low- p_T) or forward (high- $|\eta|$) b jet, or a b jet resulting from the hadronization of a collinear $b\bar{b}$ quark pair. Due to this difference, the two OOA configurations are treated as separate background sources in the 6j4b and 7j4b3l regions and are labelled as $t\bar{t}b$ OOA and $t\bar{t}b\bar{b}$ OOA, respectively, based on the presence of one or two “additional” b jets (in the sense of section 4.2).

For some of the unfolded distributions, there is no well-defined upper bound on the measured observable. In order to include all observed events in the measurement, while keeping the upper edge of the unfolded distributions at a reasonable value, we normalize the differential cross sections in these bins by the width of the bin as it appears in the histogram, while at the same time including any overflow events, both in the data and in the simulation, into these last bins.

In order to assess the presence of biases due to the choice of the nominal signal model (POWHEG+OL+P8 $t\bar{t}b\bar{b}$ 4FS), bias tests were performed with the POWHEG+P8 $t\bar{t}$ 5FS simulation. These tests consisted of unfolding pseudo-data generated using signal predictions from the POWHEG+P8 $t\bar{t}$ 5FS simulation, while using response matrices constructed from





Figure 4. Response matrix for $\Delta R(bb^{\text{extra}})$ in the ≥ 6 jets: $\geq 4b$ phase space. The x (y) axes show the generator- (detector-)level observables. The upper figure includes the ancillary variable, unrolled on the same axis as the detector-level observable, so that the binning of the detector-level observable, stacked vertically, is repeated twice. For the lower figure, the ancillary variables are projected out to more easily visualize the correspondence between true and reconstructed values. The coloured bins show the finer binning used at reconstructed level (bins split in two), while the numbers show the values one would obtain when using the same binning at the generator and detector level.

the POWHEG+OL+P8 $t\bar{t}b\bar{b}$ 4FS samples, thereby verifying whether the differences in signal modelling and response matrices resulted in any biases in the measurements. These studies showed that the unfolded results were compatible with the expectations of the POWHEG+P8 $t\bar{t}$ 5FS signal prediction within systematic uncertainties, supporting the validity of using the response matrices of the POWHEG+OL+P8 $t\bar{t}b\bar{b}$ 4FS model for unbiased unfolding.

7 Systematic uncertainties

Systematic uncertainties are evaluated by appropriate variations of the signal and background simulations. The uncertainty sources may affect background yields and distributions, as well as the selection efficiency and the kinematic distributions of the signals. These uncertainties are taken into account via nuisance parameters in the likelihood fits.

In certain cases where the statistical uncertainty of a systematic variation is comparable to or larger than the size of the systematic variation or there is significant bin-to-bin variation, the systematic templates need to be smoothed. This is done via a lowess-based smoothing algorithm, which constrains the shape differences between the upward and downward fluctuations if they have a shape effect (smoothing) or, if not, converts them to rate-only effects. If, on the other hand, the systematic uncertainty in question is determined not to have a shape or rate effect, then it is removed altogether.

A summary of all systematic uncertainties is given in table 4 and in the following sections, grouped by experimental uncertainties in section 7.1 and modelling uncertainties in section 7.2.

7.1 Experimental uncertainties

Integrated luminosity: the integrated luminosities of the data-taking periods are individually measured with uncertainties of 1.2, 2.3, and 2.5% for 2016, 2017, and 2018 data-taking periods, respectively [96–98]. The uncertainty in the integrated luminosity of the combined data set is 1.6% when taking into account the correlations between the periods.

Pileup reweighting: the prediction of the number of pileup interactions in simulation is performed by assuming a total inelastic pp cross section of 69.2 mb. Changes in the assumed pileup multiplicity are estimated by varying the total inelastic cross section by $\pm 4.6\%$ [76]. This uncertainty is treated as fully correlated between the data-taking periods.

Lepton reconstruction and identification: separate uncertainties are assigned to the corrections applied to the reconstruction and identification efficiencies for electrons in the simulation. Similarly, the muon tracking, identification, and isolation efficiencies are estimated and the associated uncertainties are propagated to signal and background distributions used in the fits. These uncertainties are taken to be fully correlated across the data-taking periods.

Trigger efficiencies: the trigger efficiency scale factors are varied within their uncertainties, separately for electron and muon triggers and for each data-taking period.

L1 prefring: during the 2016–2017 data-taking periods, a gradual shift in the timing of the inputs of the ECAL L1 trigger in the forward endcap region ($|\eta| > 2.4$) led to a specific

	Source	Corr. (period)	Corr. (process)
Experimental	Integrated luminosity	~	✓
	Pileup reweighting	✓	✓
	Electron reconstruction and identification	✓	✓
	Muon reconstruction and identification	✓	✓
	Trigger efficiencies	×	✓
	L1 prefiring	✓	✓
	JES	~	✓
	JER	×	✓
	b tagging	~	✓
	Theoretical	μ_R scale	✓
μ_F scale		✓	~
Top quark p_T modelling		✓	✓
PDF		✓	✓
PS scales: ISR		✓	×
PS scales: FSR		✓	×
ME-PS matching (h_{damp})		✓	~
Underlying-event tune		✓	✓
Colour reconnection		✓	✓
b quark fragmentation		✓	✓
Inclusive $t\bar{t}C$ cross section		✓	—

Table 4. Summary of the systematic uncertainty sources in the inclusive and differential $t\bar{t}b\bar{b}$ cross section measurements. The first column lists the source of the uncertainty. The second (third) column indicates the treatment of correlations of the uncertainties between different data-taking periods (processes), where ✓ means fully correlated, ~ means partially correlated (i.e. contains sub-sources that are either fully correlated or uncorrelated), × means uncorrelated, and — means not applicable.

inefficiency, known as “prefiring”. A similar effect is present for the muon system due to its limited time resolution, most pronounced in 2016, but also impacting data collected in 2017–2018. Corrections of this effect are applied to simulated events, and 20% of the corrections are assigned as the associated uncertainties.

Jet energy scale and resolution: uncertainties in the determination of the JES are taken into account by shifting the jet momenta in the simulation up and down, separately for several sources of uncertainty such as the overall energy scale, differences in flavour response, and residual differences between energy scale measurements. Some of these sources are treated separately per data-taking period, while some are correlated for all periods. Uncertainties in the JER are evaluated by increasing or decreasing the variation of jet energies between the reconstructed and particle levels, or by smearing the measured jet energy in case no matching particle-level jet could be found [79]. This uncertainty is uncorrelated between data-taking periods.

b tagging: differences in the b tagging efficiency between data and simulation are corrected by applying correction factors to simulated events, derived as a function of jet p_T and $|\eta|$. Systematic uncertainties are considered separately for light jets and b/c jets. For b and c jets, nine different sources of uncertainties from the measurements of the correction factors are considered [80]. Included amongst these uncertainties are effects from variations in b fragmentation and gluon to $b\bar{b}$ splitting. Scale factors and their uncertainties on the b tagging efficiency and mistag rates are split based on the flavour of the initiating quark, but do not discriminate based on the origin of the b quark as coming, for example, from a gluon splitting or top quark decay. Statistical uncertainties in the scale factor measurements are treated independently for the medium and tight b tagging working points, and for the four data-taking periods. All other sources of uncertainties are correlated between the data-taking periods and the b tagging working points.

7.2 Modelling uncertainties

This analysis is affected by uncertainties in the modelling of the background processes and the migration matrices linking the measured particle-level observables with the detector-level observables. Variations applied to the signal process are defined such that the predicted yields at generator level remain constant for all values of the corresponding nuisance parameters, independently for each bin of the generator-level distributions. This procedure ensures that the modelling uncertainties only have an effect on the signal selection efficiency, due to shape variations within each generator-level bin. All modelling uncertainties are correlated between the data-taking eras. Some uncertainties are not correlated between processes. In the phase space regions where the OOA contribution is separated into $t\bar{t}b$ OOA and $t\bar{t}b\bar{b}$ OOA contributions, the two components can have different correlations with the signal. However, by repeating the fits with different correlation assumptions for the signal and the OOA processes, it has been verified that the results were not sensitive to this choice. While most modelling uncertainties are assumed to be correlated between ancillary regions, various alternative correlation assumptions were tested for the scale uncertainties, with no significant effect on the result.

Renormalization and factorization scales: uncertainties covering the choice of μ_R and μ_F scales in the ME generators are considered by shifting the scales independently up and down by a factor of two. These uncertainties are treated separately for each process but correlated for $t\bar{t}+\text{light}$ and $t\bar{t}C$, as these contributions are estimated using the same POWHEG+P8 $t\bar{t}$ 5FS simulation. Where separate $t\bar{t}b\bar{b}$ OOA and $t\bar{t}b$ OOA contributions are considered, the uncertainty in the former contribution is correlated with the signal while the latter taken as uncorrelated. For the $t\bar{t}B$ OOA processes, shape and normalization effects of the μ_R scale are decorrelated, and both are uncorrelated with the signal.

Top quark p_T modelling: because of discrepancies between the observed and simulated p_T spectrum of top quarks in $t\bar{t}$ events, simulated events are reweighted to match the top quark p_T distribution predicted at NNLO [99]. This procedure improves the agreement of the simulated predictions with the data. We consider as uncertainty in this reweighting the full effect of the reweighting itself. The uncertainty is applied to all $t\bar{t}$ subprocesses (including the signal and all OOA processes), but not to single top quark or $t\bar{t}X$ production.

PDF: the uncertainties in the PDFs are evaluated by using replicas of the NNPDF set [69]. For the 4FS set used for the $t\bar{t}b\bar{b}$ signal simulation, uncertainties are estimated using the root-mean-square of all residuals in the predictions obtained using the PDF replicas, as these are defined by sampling from the covariance matrix of the PDF fit. For the 5FS set used in all the other simulated samples, the replicas correspond to the leading eigenvectors of the PDF fit covariance matrix, and the uncertainties are obtained as the quadratic sum of the residuals in the predictions across the replicas. This uncertainty is treated as correlated for all processes. An additional uncertainty from the value of the strong coupling constant, α_S , used in the PDF, is also included.

PS scales: the uncertainty from the choice of scale at which the strong coupling constant is evaluated in the PS is estimated by varying the scale up and down by a factor of two, independently for initial-state radiation (ISR) and final state radiation (FSR). These uncertainties are considered as uncorrelated for all processes. Accordingly, the $t\bar{t}B$ OOA processes is considered uncorrelated with the signal. Where separate $t\bar{t}b\bar{b}$ OOA and $t\bar{t}b$ OOA contributions are considered, the $t\bar{t}b\bar{b}$ OOA contribution is correlated with the signal, and $t\bar{t}b$ OOA is treated separately.

ME-PS matching (h_{damp}): in the POWHEG generator, the scale that separates the phase space of the first QCD emission into soft and hard parts is controlled by the h_{damp} parameter. The nominal value of the h_{damp} parameter in the CP5 tune [70] is $h_{\text{damp}} = 1.379m_t$ and the uncertainties are estimated with varied values of $h_{\text{damp}} = 2.305m_t$ and $0.874m_t$ for the $t\bar{t}$ and $t\bar{t}b\bar{b}$ samples. This uncertainty is treated as correlated for $t\bar{t}$ +light and $t\bar{t}C$ processes but decorrelated from the $t\bar{t}B$ contributions. This uncertainty is only applied to $t\bar{t}$ subprocesses, which does not include single top or $t\bar{t}X$ production.

Underlying event tune: the simulation of the underlying event is based on the CP5 tune of the PYTHIA event generator, and uncertainties are estimated via varied tune settings [70], applied to the $t\bar{t}$ samples. This uncertainty is only applied to $t\bar{t}B$, $t\bar{t}$ +light, and $t\bar{t}C$ processes and is treated as correlated.

Colour reconnection: the default colour reconnection model in the PYTHIA PS simulation is replaced by three alternative models [100, 101]. These uncertainties are treated as correlated for $t\bar{t}B$, $t\bar{t}$ +light, and $t\bar{t}C$ processes. This uncertainty is only applied to $t\bar{t}$ subprocesses, which does not include single top or $t\bar{t}X$ production.

b quark fragmentation: we include a theoretical uncertainty for the fragmentation of the b quarks into hadrons. The fragmentation of the initiating parton into observable hadrons is subject to modelling uncertainties. We estimate this effect by varying simultaneously the generator- and detector-level momenta of b jets up and down by 1% in the simulated samples. This is consistent with the effect from reasonable variations of the fragmentation functions on the b jet p_T distribution in ref. [102]. Residual effects on the efficiency to identify b jets, due to a possible mismodelling of the fragmentation of b quarks, are already accounted for by the calibration of the b tagging efficiency in the simulation.

Inclusive $t\bar{t}C$ cross section: for the $t\bar{t}C$ process an additional 20% normalization uncertainty is applied, corresponding to the precision of the inclusive $t\bar{t}C$ cross section measurement by CMS [45]. This measurement found that the $t\bar{t}C$ cross section agreed with the POWHEG+P8 $t\bar{t}$ 5FS prediction.

No additional cross section uncertainties are considered for the backgrounds since the uncertainties listed above already result in variations in the predicted event yields that cover the uncertainties in the theoretical cross sections used to normalize the background contributions.

The finite size of the simulated MC samples is taken into account as a systematic uncertainty, following a method similar to the one proposed by Barlow and Beeston [103, 104]. For every bin of the detector-level distributions, a single Gaussian-constrained nuisance parameter varies the predicted yields, summed over all processes including the signal, within their statistical uncertainty.

The impact of a group of k nuisance parameters $\alpha_{n_1} \dots \alpha_{n_k}$ on the parameter of interest p is computed as

$$I_p = \sqrt{\sum_{i,j=1}^k C_{pn_i} \tilde{C}_{n_i n_j}^{-1} C_{n_j p}}, \tag{7.1}$$

where C is the covariance matrix between all parameters, and \tilde{C} is the covariance matrix restricted to the parameters $n_1 \dots n_k$. In this way, the effects of a set of nuisance parameters are combined while taking into account their correlation in the fit. The total systematic uncertainties are calculated by considering all nuisance parameters in the sum in eq. (7.1), and the statistical uncertainties as the difference in quadrature between the total uncertainties and the total systematic uncertainties.

Table 5 shows the contributions of various sources of uncertainties to the total uncertainties in the inclusive cross sections, obtained from the combined impacts $I_{\mu_{\text{fid}}}$ on the parameters scaling the inclusive cross sections in the fits of the representative observables discussed in section 8.1. For these fits, the contributions of the 20 single nuisance parameters with the largest contributions to the uncertainty in the inclusive cross section can be found in appendix A. Figure 5 shows the effect of the sources of uncertainty on the normalized differential cross section measurements for the H_T of b jets in the 5j3b phase space. For the four representative observables, corresponding figures are shown in appendix B. The uncertainties in the inclusive cross sections are dominated by systematic sources, while the precision in the differential measurements is mainly limited by the statistical uncertainty in the data since the rate-based effects of many systematic uncertainties cancel out in the ratio between the absolute differential cross sections and the inclusive cross sections. The leading systematic uncertainties originate from the calibration of the b tagging and of the JES, the choice of μ_R scale in the signal $t\bar{t}b\bar{b}$ and background $t\bar{t}$ processes, and, for the differential measurements only, the finite number of simulated events. Previous measurements of the inclusive $t\bar{t}b\bar{b}$ cross section by the ATLAS [39] and CMS [42–44] Collaborations had the same leading sources of systematic uncertainty.

Uncertainty source	Relative uncertainty (%)			
	5j3b	6j3b3l	6j4b	7j4b3l
Integrated luminosity	1.6	1.6	2.0	1.8
Pileup reweighting	0.2	0.8	0.4	0.5
Lepton and trigger	1.1	0.9	1.9	1.8
JES, JER	2.1	1.6	3.5	5.7
b tagging	4.5	3.9	7.0	9.1
μ_R and μ_F scales	2.8	6.8	8.2	12
Top quark p_T modelling	0.3	1.0	0.6	1.3
PDF	0.2	0.7	1.0	1.9
PS scales	2.8	2.7	2.4	1.5
ME-PS matching (h_{damp})	0.4	0.9	1.3	2.8
Underlying event	0.4	<0.1	0.4	0.4
Colour reconnection	1.1	1.5	1.9	4.5
b quark fragmentation	0.3	0.4	0.4	0.4
Inclusive $t\bar{t}C$ cross section	0.5	0.3	1.9	2.6
MC statistical	0.8	1.6	2.4	2.8
Total systematic uncertainty	6.0	8.7	13	17
Statistical uncertainty	0.6	1.2	2.2	3.3
Total uncertainty	6.0	8.8	13	17

Table 5. Contributions of the considered sources of uncertainty to the total uncertainty in the inclusive cross sections. For each group of uncertainty sources, the impacts of the corresponding nuisance parameters on the total cross section are combined, taking into account their correlation in the fit. The numbers show relative uncertainties (in %). The statistical uncertainty is obtained as the difference, in quadrature, between the total uncertainty and the sum of all systematic uncertainties.

The nuisance parameter associated with the normalization of the $t\bar{t}C$ process is not found to be constrained significantly beyond its prefit expectation and also shows no significant deviation from its expected value, which is consistent with the results of ref. [45]. The correlation of that nuisance parameter with the inclusive $t\bar{t}b\bar{b}$ cross section in the fits to different observables is below 5 (20)% in the phase space regions with at least three (four) b jets.

8 Results

The results are obtained with the statistical procedures described in section 6.3. Inclusive and differential cross section results are presented in sections 8.1 and 8.2, respectively.

8.1 Inclusive cross sections

The inclusive $t\bar{t}b\bar{b}$ cross sections, measured in each of the fiducial phase space regions, are shown in figure 6 and listed in table 6, along with the predictions obtained from the $t\bar{t}b\bar{b}$




Figure 5. Effect of the considered sources of uncertainties on the measurement of the normalized differential cross section of the H_T of b jets in the ≥ 5 jets: $\geq 3b$ phase space, obtained by combining the impacts of associated nuisance parameters according to eq. (6.1). The ordering of the various sources is similar for other observables and in the other phase space regions. The last bin of the distribution is not shown, since it has no associated parameter of interest but is constrained by the other bins as described in section 6.3. The category “other theory” includes b quark fragmentation, top quark p_T modelling, PDF, h_{damp} , colour reconnection, and underlying event uncertainties. The category “other experimental” includes pileup and the integrated luminosity uncertainties.

generator setups described in section 3. In each phase space, the representative observable for which we report the measured inclusive cross section is that for which the measured value is closest to the mean of all measured inclusive cross sections in order to provide a value representative of the ensemble of measurements. These observables are $|\eta(b_3)|$ in the 5j3b phase space, H_T^{light} in the 6j3b3l phase space, $|\eta(b_2^{\text{extra}})|$ in the 6j4b phase space, and $|\Delta\phi(l_{j_1}^{\text{extra}}, b_{\text{soft}})|$ in the 7j4b3l phase space. Their observed distributions at the detector level, combined for both lepton channels and all data-taking periods, are shown in figures 7 and 8 after the fits to data. The correlations between the parameters $\vec{\mu}$ in the fit of $|\eta(b_3)|$ in the 5j3b phase space are shown in figure 9. For the other representative observables, corresponding figures are shown in appendix C. The inclusive cross section results for all observables are provided in the HEPData record for this analysis [46].

The measured cross sections in all phase space regions are larger than the theoretical predictions, as was also observed in previous measurements of the inclusive $t\bar{t}b\bar{b}$ cross section (with somewhat different fiducial definitions) by the CMS [42–44] and ATLAS [39] Collaborations, with the notable exception of the POWHEG+OL+P8 $t\bar{t}b\bar{b}$ 4FS generator setup. The choice of reduced central μ_R and μ_F scales in the POWHEG+OL+P8 $t\bar{t}b\bar{b}$ 4FS sample, justified by the previous measurements and by studies of fixed-order NLO QCD corrections to the $t\bar{t}b\bar{b}j$ process [1], results in significantly larger cross sections in all phase




Figure 6. Measured inclusive cross sections for each considered phase space, compared to predictions from different $t\bar{t}b\bar{b}$ simulation approaches shown as coloured symbols. The predictions include uncertainties (horizontal bars) due to the limited number of simulated events. The blue colour is reserved for models using massive b quarks and NLO QCD $t\bar{t}b\bar{b}$ MEs, while red is used for the inclusive $t\bar{t}$ generators at NLO in QCD with massless b quarks. The right panel shows the ratios between the predicted and measured cross sections, with the black bars showing the relative uncertainties in the measurements.

Fiducial phase space	5j3b	6j3b3l	6j4b	7j4b3l
Measured cross section	2367	1037	291	147
	± 142 (syst)	± 90 (syst)	± 36 (syst)	± 24 (syst)
	± 14 (stat)	± 12 (stat)	± 6 (stat)	± 5 (stat)
POWHEG+OL+P8 $t\bar{t}b\bar{b}$ 4FS	2361	1183	361	197
μ_R variation	+1161/-737	+826/-433	+183/-113	+121/-67
μ_F variation	+126 /-100	+97 /-78	+23 /-18	+16 /-13
POWHEG+P8 $t\bar{t}$ 5FS	1791	899	240	129
POWHEG+H7 $t\bar{t}$ 5FS	1665	762	197	95
SHERPA+OL $t\bar{t}b\bar{b}$ 4FS	1391	677	216	116
MG5_aMC+P8 $t\bar{t}b\bar{b}$ 4FS	1024	524	187	101
MG5_aMC+P8 $t\bar{t}$ +jets FxFx 5FS	1560	712	203	101

Table 6. Measured and predicted inclusive cross sections in the four considered phase space regions (in fb).




Figure 7. The $|\eta|$ of the third-hardest b jet in p_T ($|\eta(b_3)|$) in the ≥ 5 jets: $\geq 3b$ phase space (upper) and the $|\eta|$ of the subleading additional b jet ($|\eta(b_2^{\text{extra}})|$) in the ≥ 6 jets: $\geq 4b$ phase space (lower) after the fit to data, shown for both lepton channels and all data periods combined. The distributions are shown separately for each ancillary region, as defined in section 6.1. In the ≥ 5 jets: $\geq 3b$ (≥ 6 jets: $\geq 4b$) phase space the ancillary regions are defined as ≤ 2 , 2 , and ≥ 3 (≤ 3 and ≥ 3) tight b-tagged jets. The shaded bands include all uncertainties described in section 7 after profiling the nuisance parameters in the fit, estimated by sampling the predicted yields from the fit covariance matrix. The blue line shows the sum of the predicted yields for all processes before the fit to data, using the nominal $t\bar{t}b\bar{b}$ samples and its corresponding cross section for the signal. In the ratio panel the expected yields before the fit to data are shown relative to the predicted yields after the fit to data. The last bins contain the overflow.




Figure 8. The H_T of all light jets in the ≥ 6 jets: $\geq 3b$, ≥ 3 light phase space (upper) and the azimuthal angle between the hardest remaining light jet and the softest b jet ($|\Delta\phi(lj_1^{\text{extra}}, b_{\text{soft}})|$) in the ≥ 7 jets: $\geq 4b$, ≥ 3 light phase space (lower) after the fit to data, shown for both lepton channels and all data periods combined. The distributions are shown separately for each ancillary region, as defined in section 6.1. In the ≥ 6 jets: $\geq 3b$, ≥ 3 light (≥ 7 jets: $\geq 4b$, ≥ 3 light) phase space the ancillary regions are defined as ≤ 2 , 2, and ≥ 3 (≤ 3 and ≥ 3) tight b-tagged jets. The shaded bands include all uncertainties described in section 7 after profiling the nuisance parameters in the fit, estimated by sampling the predicted yields from the fit covariance matrix. The blue line shows the sum of the predicted yields for all processes before the fit to data, using the nominal $t\bar{t}b\bar{b}$ samples and its corresponding cross section for the signal. In the ratio panel the expected yields before the fit to data are shown relative to the predicted yields after the fit to data. The last bins contain the overflow.




Figure 9. Correlations between the parameters of interest $\vec{\mu}$ in the fit for $|\eta(b_3)|$ in the ≥ 5 jets: $\geq 3b$ phase space.

space regions. This prediction agrees well with the measurement in the 5j3b phase space, but overestimates the cross section by 10–35% in the other phase space regions. Nevertheless, the POWHEG+OL+P8 $t\bar{t}b\bar{b}$ 4FS predictions agree with the measurements when considering their μ_R scale uncertainties of about 50%, estimated by varying the μ_R scale by a factor of two. When considering μ_R scale uncertainties in both the ME and in the PS, the POWHEG+P8 $t\bar{t}$ 5FS predictions agree with the measurements in the phase space regions targeting additional light radiation in $t\bar{t}b$ and $t\bar{t}b\bar{b}$ events. The cross sections based on $t\bar{t}b\bar{b}$ matrix element calculations are expected to increase if $t\bar{t}$ events with additional b quarks produced in MPI are included. For the POWHEG+OL+P8 $t\bar{t}b\bar{b}$ 4FS prediction, this increase is estimated using the POWHEG+P8 $t\bar{t}$ 5FS simulation to be 6–10% depending on the phase space.

The SHERPA+OL $t\bar{t}b\bar{b}$ 4FS predictions are between 20 and 50% lower than the measured cross sections. This can mostly be attributed to the μ_R and μ_F scale choices in this SHERPA+OL $t\bar{t}b\bar{b}$ 4FS setup, which are a factor of two higher than the scale choices of the POWHEG+OL+P8 $t\bar{t}b\bar{b}$ 4FS simulation. With common scale settings between the SHERPA+OL $t\bar{t}b\bar{b}$ 4FS and POWHEG+OL+P8 $t\bar{t}b\bar{b}$ 4FS simulation approaches, compatible inclusive and differential cross section predictions are achieved in the studies presented in ref. [105]. Increased cross

section predictions could similarly be achieved in the other simulations by lowering the μ_R and μ_F scale settings.

8.2 Normalized differential cross sections

The normalized differential cross section is measured in four different fiducial phase space regions for 29 observables that use exclusively stable-particle level information without reference to any simulated event history, and eight observables targeting explicitly the b jets that do not originate from decaying top quarks. For each observable, customized bin sizes are chosen, depending on the resolution of the observables and the statistical uncertainty in the measured event yields.

The resulting normalized differential cross sections are shown in figure 10 for the observables of the 5j3b phase space, in figures 11–13 for the observables of the 6j4b phase space, for the observables targeting the b jets that do not originate from decaying top quarks in figures 14 and 15, and finally in figure 16 for the observables of the 6j3b3l and 7j4b3l phase space regions. The measurements are compared to six cross section predictions of the $t\bar{t}b\bar{b}$ process, obtained at the particle level, produced with the different combinations of event generators and PSs introduced in section 3. The various predictions are shown as symbols distinguished by colour and shape. It should be noted that predictions from SHERPA+OL $t\bar{t}b\bar{b}$ 4FS cannot be compared to the observables related to the additional b jets, since that generator does not provide the necessary information (parton-level top quarks) to assign b jets to the decaying top quarks. The calculations using $t\bar{t}b\bar{b}$ in the matrix element do not include contributions coming from the production of $t\bar{t}$ in the primary parton scattering with additional b quarks coming only from MPI.

The compatibility of the predictions with the unfolded data in each of the phase space regions is quantified using χ^2 tests, which are converted to z scores [106], which quantify the p -value in terms of the equivalent number of standard deviations by which each theoretical prediction differs from the mean of a normal distribution centered at the normalized cross-section measurements. These are shown in figure 17 and as tables in appendix E. The χ^2 test statistics are computed using the predicted and measured normalized cross sections, as well as the estimated covariance between the bins of the unfolded distributions, but do not include the inclusive cross sections. No uncertainties in the predictions are considered in these tests. Varying levels of agreement between the predicted and measured distributions are observed, whereby no event generator setup describes all observables well since they each result in $z \geq 2$ for several observables.

In the 5j3b phase space (Figure 10), with a few exceptions, none of the distributions are described well by any of the considered generator setups, quantified by the z scores without any consideration of the uncertainties on the predictions. Most generator setups predict a higher number of inclusive jets than what is observed, with the exception of POWHEG+H7 $t\bar{t}$ 5FS and MG5_aMC+P8 $t\bar{t}$ +jets FxFx 5FS showing better agreement. The b jet multiplicity distribution, which can be interpreted as a measure of the ratio between the cross sections of $t\bar{t}b\bar{b}$ and $t\bar{t}b$, is only well described by the inclusive POWHEG $t\bar{t}$ simulations matched to PYTHIA or HERWIG. The NLO $t\bar{t}b\bar{b}$ simulations all predict a higher ratio of events with at least four b jets over exactly three b jets. The H_T^j distribution shows a significant




Figure 10. Predicted and observed normalized differential cross sections in the ≥ 5 jets; $\geq 3b$ fiducial phase space, for the inclusive jet multiplicity (upper left), the b jet multiplicity (upper right), the inclusive jet H_T (middle left, H_T^j), the H_T of b jets (middle right, H_T^b), the $|\eta|$ of the third b jet (lower left), and the p_T of the third b jet (lower right). The data are represented by points, with inner (outer) vertical bars indicating the systematic (total) uncertainties, also represented as blue (grey) bands. Cross section prediction obtained at the particle level from different simulation approaches are shown, including their statistical uncertainties, as coloured symbols with different shapes. For N_{jets} , N_b , H_T^j , H_T^b , and $p_T(b_3)$, the last bins contain the overflow.




Figure 11. Predicted and observed normalized differential cross sections in the ≥ 6 jets: $\geq 4b$ fiducial phase space, for the inclusive jet H_T (upper left), the H_T of b jets (upper right), the H_T of b jets (upper right), the $|\eta|$ of the third b jet (middle left), the p_T of the third b jet (middle right), the $|\eta|$ of the fourth b jet (lower left), and the p_T of the fourth b jet (lower right). The data are represented by points, with inner (outer) vertical bars indicating the systematic (total) uncertainties, also represented as blue (grey) bands. Cross section predictions obtained at the particle level from different simulation approaches are shown, including their statistical uncertainties, as coloured symbols with different shapes. For H_T and p_T , the last bins contain the overflow.




Figure 12. Predicted and observed normalized differential cross sections in the ≥ 6 jets: $\geq 4b$ fiducial phase space, for the average ΔR of all possible bb pairs (upper left), the largest invariant mass of any bb pair (upper right), the invariant mass (middle left), ΔR (middle right), p_T (lower left), and $|\eta|$ (lower right) of the extra b-jet pair. The data are represented by points, with inner (outer) vertical bars indicating the systematic (total) uncertainties, also represented as blue (grey) bands. Cross section predictions obtained at the particle level from different simulation approaches are shown, including their statistical uncertainties, as coloured symbols. For m_{bb} and p_T , the last bins contain the overflow.




Figure 13. Predicted and observed normalized differential cross sections in the ≥ 6 jets: $\geq 4b$ fiducial phase space, for the $|\eta|$ (upper left) and p_T (upper right) of the first extra b jet, the $|\eta|$ (middle left) and p_T (middle right) of the second extra b jet, and the inclusive jet multiplicity (lower left). The data are represented by points, with inner (outer) vertical bars indicating the systematic (total) uncertainties, also represented as blue (grey) bands. Cross section predictions obtained at the particle level from different simulation approaches are shown, including their statistical uncertainties, as coloured symbols. For N_{jets} and p_T , the last bins contain the overflow.




Figure 14. Predicted and observed normalized differential cross sections in the ≥ 6 jets: $\geq 4b$ fiducial phase space, for the invariant mass (upper left), ΔR (upper right), p_T (lower left), and $|\eta|$ (lower right) of the additional b-jet pair not originating from decaying top quarks. The data are represented by points, with inner (outer) vertical bars indicating the systematic (total) uncertainties, also represented as blue (grey) bands. Cross section predictions obtained at the particle level from different simulation approaches are shown, including their statistical uncertainties, as coloured symbols. For m_{bb} and p_T , the last bins contain the overflow.

trend towards higher values than what is observed in data for MG5_aMC+P8 $t\bar{t}b\bar{b}$ 4FS, and towards lower values for POWHEG+H7 $t\bar{t}$ 5FS, while the other generator setups better describe the distribution. The H_T^b distribution shows a trend towards lower values in data compared to most predictions, with the exception of POWHEG+H7 $t\bar{t}$ 5FS and SHERPA+OL $t\bar{t}b\bar{b}$ 4FS. The predictions of the different simulations for $|\eta(b_3)|$ are all very similar to each other and have z scores around 2. All generator setups predict somewhat larger values of the p_T of the third b jet.

Observables which are not fully described by the NLO $t\bar{t}b\bar{b}$ ME, such as observables related to the radiation of light jets, are expected to have strong dependencies on the μ_R and μ_F scale choices when described with a NLO $t\bar{t}b\bar{b}$ ME simulation setup [2]. Hence,




Figure 15. Predicted and observed normalized differential cross sections in the ≥ 6 jets: $\geq 4b$ fiducial phase space, for the $|\eta|$ (left) and p_T (right) of the first (upper row) and second (lower row) additional b of the b -jet pair not originating from decaying top quarks. The data are represented by points, with inner (outer) vertical bars indicating the systematic (total) uncertainties, also represented as blue (grey) bands. Cross section predictions obtained at the particle level from different simulation approaches are shown, including their statistical uncertainties, as coloured symbols. For p_T , the last bins contain the overflow.

in appendix D, the measured normalized differential cross sections of the N_{jets} and H_T^j observables are compared with alternative μ_R and μ_F scale choices of the POWHEG+OL+P8 $t\bar{t}b\bar{b}$ 4FS simulation approach. These comparisons show that increased μ_R and μ_F scales relative to the nominal scale choices (see table 1) tend to better describe these observables.

The agreement between data and predictions is generally better in the 6j4b phase space (figures 11–13), at least in part due to the larger uncertainties in the measurements. With a few exceptions, the predictions between the various generator setups are also closer to each other than in the 5j3b phase space. H_T^b and H_T^j are generally modelled well, while for the POWHEG+H7 $t\bar{t}$ 5FS simulation a trend toward lower values is observed, similar to the 5j3b phase space. However, most generator setups underpredict the threshold region at low values




Figure 16. Predicted and observed normalized differential cross sections in the ≥ 6 jets: $\geq 3b$, ≥ 3 light (left) and ≥ 7 jets: $\geq 4b$, ≥ 3 light (right) fiducial phase space regions, for the H_T of light jets (upper row), the p_T of the extra light jet (middle row), and the $\Delta\phi$ between the extra light jet and the softest b jet (lower row). The data are represented by points, with inner (outer) vertical bars indicating the systematic (total) uncertainties, also represented as blue (grey) bands. Cross section predictions obtained at the particle level from different simulation approaches are shown, including their statistical uncertainties, as coloured symbols with different shapes. For H_T and p_T , the last bins contain the overflow.




Figure 17. Observed z score for each of the theoretical predictions, given the unfolded normalized differential cross sections and their covariances. A lower value indicates a better agreement between prediction and measurement. The dashed line at $z = 2$ indicates a p -value of 5%. Predictions for which the z score exceeds the visible range of the figure are marked with arrows (\rightarrow).

of H_T . The predictions for $p_T(b_3)$, $|\eta(b_3)|$, $p_T(b_4)$, and $|\eta(b_4)|$ are generally compatible with the data, with only a slight trend visible in data towards more central (lower $|\eta|$) and softer (lower p_T) b jets. Similarly, predictions for the p_T and $|\eta|$ for the extra b jets all describe the data well within the measurement uncertainties. The total number of jets is reasonably described by all predictions and does not show any clear trend.

The measured ΔR_{bb}^{avg} shows a small trend towards lower values for all simulation approaches, but is compatible with the data within the uncertainties of the measurement, as is the maximum invariant mass of any $b\bar{b}$ pair.

The observables related to the $b\bar{b}^{extra}$ pair are modelled well by most of the generator setups, except for POWHEG+H7 $t\bar{t}$ 5FS, which predicts a softer $m(b\bar{b}^{extra})$, and a smaller $\Delta R(b\bar{b}^{extra})$. The variables related to the $b\bar{b}^{add.}$ pair (figures 14–15), i.e. the b jets not part of the top quark decay chain, show similar behaviour, whereby $\Delta R(b\bar{b}^{add.})$, $m(b\bar{b}^{add.})$ and $p_T(b\bar{b}^{add.})$ are not described by POWHEG+H7 $t\bar{t}$ 5FS. Predictions for the p_T and $|\eta|$ of the individual additional b jets all tend to describe the data well.

In the 6j3b3l and 7j4b3l phase space regions (figure 16), the H_T^{light} and $|\Delta\phi(lj_1^{extra}, b_{soft})|$ observables are modelled well by most generator setups, with only POWHEG+OL+P8 $t\bar{t}b\bar{b}$ 4FS predicting a higher H_T^{light} . With the exception of POWHEG+H7 $t\bar{t}$ 5FS, $p_T(lj_1^{extra})$ is observed to be softer than what is predicted by most generator setups. In appendix D, the measured normalized differential cross sections of the $p_T(lj_1^{extra})$ and H_T^{light} observables in the 6j3b3l phase space are compared with alternative μ_R and μ_F scale choices of the POWHEG+OL+P8 $t\bar{t}b\bar{b}$ 4FS simulation approach. These comparisons show that increased μ_R and μ_F scales relative to the nominal scale choices (see table 1) tend to better describe these observables.

9 Summary

Measurements of inclusive and normalized differential cross sections of the associated production of top quark-antiquark and bottom quark-antiquark pairs, $t\bar{t}b\bar{b}$, for events containing an electron or a muon, have been presented. These measurements use proton-proton collision data recorded by the CMS detector at $\sqrt{s} = 13$ TeV and correspond to an integrated luminosity of 138 fb^{-1} .

The inclusive cross sections are measured in four fiducial phase space regions requiring different jet, b jet, and light jet multiplicities. With total uncertainties of 6–17%, depending on the phase space, these are the most precise measurements of the $t\bar{t}b\bar{b}$ cross section to date. The uncertainties are dominated by systematic sources, with the leading uncertainties originating from the calibration of the b tagging and of the jet energy scale, and from the choice of renormalization scale in the signal $t\bar{t}b\bar{b}$ and background $t\bar{t}$ processes. In most cases, the measured inclusive cross sections exceed the predictions with the chosen generator settings. The only exception is when using a particular choice of dynamic renormalization scale, $\mu_R = \frac{1}{2} \prod_{i=t,\bar{t},b,\bar{b}} m_{T,i}^{1/4}$, where $m_{T,i}^2 = m_i^2 + p_{T,i}^2$ are the transverse masses of top and bottom quarks.

Differential cross section measurements are performed as a function of several observables in the aforementioned phase space regions. These observables mainly target b jets as well as additional light jets produced in association with the top quark pairs. In the phase space containing events with at least six jets, of which at least four are b tagged, the additional b-jet

radiation is probed with two different approaches. The first approach uses observables defined purely at the particle level, without any reference to the top quark decay chains, by selecting the two b jets with the smallest angular separation. The second approach uses explicitly the b jets at the generator level that do not originate from top quark decays and identifies those jets at the detector level with a neural network discriminant. The differential measurements have relative uncertainties in the range of 2–50%, depending on the phase space and the observable.

The results are compared to the predictions of several event generator setups, and it is found that none of them simultaneously describe all measured distributions in the various phase space regions. In the more inclusive phase space with five jets and three b jets, the agreement between data and predictions is generally poor, while in the phase space with six jets and four b jets, corresponding to the case in which the two additional b jets in $t\bar{t}b\bar{b}$ production are resolved, most predictions are compatible with the data within the larger experimental uncertainties. These measurements will help to further tune and refine the theoretical predictions and better assess the validity of the theoretical uncertainties estimated from the various $t\bar{t}b\bar{b}$ event generators.

Acknowledgments

We congratulate our colleagues in the CERN accelerator departments for the excellent performance of the LHC and thank the technical and administrative staffs at CERN and at other CMS institutes for their contributions to the success of the CMS effort. In addition, we gratefully acknowledge the computing centres and personnel of the Worldwide LHC Computing Grid and other centres for delivering so effectively the computing infrastructure essential to our analyses. Finally, we acknowledge the enduring support for the construction and operation of the LHC, the CMS detector, and the supporting computing infrastructure provided by the following funding agencies: SC (Armenia), BMBWF and FWF (Austria); FNRS and FWO (Belgium); CNPq, CAPES, FAPERJ, FAPERGS, and FAPESP (Brazil); MES and BNSF (Bulgaria); CERN; CAS, MoST, and NSFC (China); MINCIENCIAS (Colombia); MSES and CSF (Croatia); RIF (Cyprus); SENESCYT (Ecuador); MoER, ERC PUT and ERDF (Estonia); Academy of Finland, MEC, and HIP (Finland); CEA and CNRS/IN2P3 (France); SRNSF (Georgia); BMBF, DFG, and HGF (Germany); GSRI (Greece); NKFIH (Hungary); DAE and DST (India); IPM (Iran); SFI (Ireland); INFN (Italy); MSIP and NRF (Republic of Korea); MES (Latvia); LAS (Lithuania); MOE and UM (Malaysia); BUAP, CINVESTAV, CONACYT, LNS, SEP, and UASLP-FAI (Mexico); MOS (Montenegro); MBIE (New Zealand); PAEC (Pakistan); MES and NSC (Poland); FCT (Portugal); MESTD (Serbia); MCIN/AEI and PCTI (Spain); MOSTR (Sri Lanka); Swiss Funding Agencies (Switzerland); MST (Taipei); MHEI and NSTDA (Thailand); TUBITAK and TENMAK (Turkey); NASU (Ukraine); STFC (United Kingdom); DOE and NSF (U.S.A.).

Individuals have received support from the Marie-Curie programme and the European Research Council and Horizon 2020 Grant, contract Nos. 675440, 724704, 752730, 758316, 765710, 824093, and COST Action CA16108 (European Union); the Leventis Foundation; the Alfred P. Sloan Foundation; the Alexander von Humboldt Foundation; the Science Committee, project no. 22rl-037 (Armenia); the Belgian Federal Science Policy Office; the Fonds pour la Formation à la Recherche dans l'Industrie et dans l'Agriculture (FRIA-Belgium); the

Agentschap voor Innovatie door Wetenschap en Technologie (IWT-Belgium); the F.R.S.-FNRS and FWO (Belgium) under the “Excellence of Science — EOS” — be.h project n. 30820817; the Beijing Municipal Science & Technology Commission, No. Z191100007219010 and Fundamental Research Funds for the Central Universities (China); the Ministry of Education, Youth and Sports (MEYS) of the Czech Republic; the Shota Rustaveli National Science Foundation, grant FR-22-985 (Georgia); the Deutsche Forschungsgemeinschaft (DFG), under Germany’s Excellence Strategy — EXC 2121 “Quantum Universe” — 390833306, and under project number 400140256 — GRK2497; the Hellenic Foundation for Research and Innovation (HFRI), Project Number 2288 (Greece); the Hungarian Academy of Sciences, the New National Excellence Program — ÚNKP, the NKFIH research grants K 124845, K 124850, K 128713, K 128786, K 129058, K 131991, K 133046, K 138136, K 143460, K 143477, 2020-2.2.1-ED-2021-00181, and TKP2021-NKTA-64 (Hungary); the Council of Science and Industrial Research, India; the Latvian Council of Science; the Ministry of Education and Science, project no. 2022/WK/14, and the National Science Center, contracts Opus 2021/41/B/ST2/01369 and 2021/43/B/ST2/01552 (Poland); the Fundação para a Ciência e a Tecnologia, grant CEECIND/01334/2018 (Portugal); the National Priorities Research Program by Qatar National Research Fund; MCIN/AEI/10.13039/501100011033, ERDF “a way of making Europe”, and the Programa Estatal de Fomento de la Investigación Científica y Técnica de Excelencia María de Maeztu, grant MDM-2017-0765 and Programa Severo Ochoa del Principado de Asturias (Spain); the Chulalongkorn Academic into Its 2nd Century Project Advancement Project, and the National Science, Research and Innovation Fund via the Program Management Unit for Human Resources & Institutional Development, Research and Innovation, grant B05F650021 (Thailand); the Kavli Foundation; the Nvidia Corporation; the SuperMicro Corporation; the Welch Foundation, contract C-1845; and the Weston Havens Foundation (U.S.A.).

A Leading nuisance parameter impacts




Figure 18. Post-fit nuisance parameter values and relative impacts on the fiducial cross section, for the fit of $|\eta|$ of the b jet with third-highest p_T in the ≥ 5 jets: $\geq 3b$ phase space. The nuisance parameters are defined such that the prefit value is zero with unity uncertainty.




Figure 19. Post-fit nuisance parameter values and relative impacts on the fiducial cross section, for the fit of $|\eta|$ of the subleading extra b jet in the ≥ 6 jets: $\geq 4b$ phase space. The nuisance parameters are defined such that the prefit value is zero with unity uncertainty.




Figure 20. Post-fit nuisance parameter values and relative impacts on the fiducial cross section, for the fit of H_T of light jets in the ≥ 6 jets: $\geq 3b$, ≥ 3 light phase space. The nuisance parameters are defined such that the prefit value is zero with unity uncertainty.




Figure 21. Post-fit nuisance parameter values and relative impacts on the fiducial cross section, for the fit of $\Delta\phi$ between leading light jet and softest b jet in the ≥ 7 jets: $\geq 4b$, ≥ 3 light phase space. The nuisance parameters are defined such that the prefit value is zero with unity uncertainty.

B Groups of impacts




Figure 22. Effect of the considered sources of uncertainties on the measurement of the normalized differential cross section of the $|\eta|$ of the b jet with third-highest p_T in the ≥ 5 jets: $\geq 3b$ phase space, obtained by combining the impacts of associated nuisance parameters. The last bin of the distribution is not shown, since it has no associated parameter of interest but is constrained by the other bins. The category “other theory” includes b quark fragmentation, top quark p_T modelling, PDF, h_{damp} , colour reconnection, and underlying event uncertainties. The category “other experimental” includes pileup and the integrated luminosity uncertainties.




Figure 23. Effect of the considered sources of uncertainties on the measurement of the normalized differential cross section of the $|\eta|$ of the subleading extra b jet in the ≥ 6 jets: $\geq 4b$ phase space, obtained by combining the impacts of associated nuisance parameters. The last bin of the distribution is not shown, since it has no associated parameter of interest but is constrained by the other bins. The category “other theory” includes b quark fragmentation, top quark p_T modelling, PDF, h_{damp} , colour reconnection, and underlying event uncertainties. The category “other experimental” includes pileup and the integrated luminosity uncertainties.




Figure 24. Effect of the considered sources of uncertainties on the measurement of the normalized differential cross section of the H_T of light jets in the ≥ 6 jets: $\geq 3b$, ≥ 3 light phase space, obtained by combining the impacts of associated nuisance parameters. The last bin of the distribution is not shown, since it has no associated parameter of interest but is constrained by the other bins. The category “other theory” includes b quark fragmentation, top quark p_T modelling, PDF, h_{damp} , colour reconnection, and underlying event uncertainties. The category “other experimental” includes pileup and the integrated luminosity uncertainties.




Figure 25. Effect of the considered sources of uncertainties on the measurement of the normalized differential cross section of the $\Delta\phi$ between leading light jet and softest b jet in the ≥ 7 jets: $\geq 4b$, ≥ 3 light phase space, obtained by combining the impacts of associated nuisance parameters. The last bin of the distribution is not shown, since it has no associated parameter of interest but is constrained by the other bins. The category “other theory” includes b quark fragmentation, top quark p_T modelling, PDF, h_{damp} , colour reconnection, and underlying event uncertainties. The category “other experimental” includes pileup and the integrated luminosity uncertainties.

C Correlation of POIs




Figure 26. Correlations between the parameters of interest $\vec{\mu}$ in the fit of the $|\eta|$ of the subleading extra b jet in the ≥ 6 jets: $\geq 4b$ phase space.




Figure 27. Correlations between the parameters of interest $\vec{\mu}$ in the fit of the H_T of light jets in the ≥ 6 jets: $\geq 3b$, ≥ 3 light phase space.




Figure 28. Correlations between the parameters of interest $\vec{\mu}$ in the fit of the $\Delta\phi$ between leading light jet and softest b jet in the ≥ 7 jets: $\geq 4b$, ≥ 3 light phase space.

D Variation of renormalization and factorization scales




Figure 29. Ratio of normalized differential cross section predictions of the POWHEG+OL+P8 $t\bar{t}b\bar{b}$ 4FS modelling approach with different μ_R and μ_F scale settings relative to the measured normalized differential cross sections for the number of jets (upper) and H_T of jets (lower) in the ≥ 5 jets: $\geq 3b$ phase space. The systematic (total) uncertainties of the measurement are represented as grey (blue) bands. Variations of the μ_R (μ_F) scale relative to the nominal scale setting are shown in orange (purple). The last bin in the distributions contains the overflow.




Figure 30. Ratio of normalized differential cross section predictions of the POWHEG+OL+P8 $t\bar{t}b\bar{b}$ 4FS modelling approach with different μ_R and μ_F scale settings relative to the measured normalized differential cross sections for the extra light jet (upper) and H_T of light jets (lower) in the ≥ 6 jets: $\geq 3b$, ≥ 3 light phase space. The systematic (total) uncertainties of the measurement are represented as grey (blue) bands. The systematic (total) uncertainties of the measurement are represented as grey (blue) bands. Variations of the μ_R (μ_F) scale relative to the nominal scale setting are shown in orange (purple). The last bin in the distributions contains the overflow.

E Normalized differential cross section compatibility

5j3b phase space	Observed z score					
	H_T^b	H_T^j	N_b	N_{jets}	$p_T(b_3)$	$ \eta(b_3) $
MG5_aMC+P8 $t\bar{t}$ +jets FxFx 5FS	4.88	1.15	2.43	5.50	4.54	2.12
MG5_aMC+P8 $t\bar{t}b\bar{b}$ 4FS	>6	5.07	>6	>6	>6	2.01
POWHEG+H7 $t\bar{t}$ 5FS	2.00	4.10	0.46	4.45	1.53	2.27
POWHEG+OL+P8 $t\bar{t}b\bar{b}$ 4FS	3.14	4.56	4.43	>6	2.19	2.61
POWHEG+P8 $t\bar{t}$ 5FS	3.01	3.05	1.73	5.17	1.88	2.52
SHERPA+OL $t\bar{t}b\bar{b}$ 4FS	3.14	0.16	5.18	5.37	3.08	2.31

Table 7. Observed z score for each of the theoretical predictions in the 5j3b phase space, given the unfolded data and covariance matrix. For the determination of the z score, only the measurement uncertainties are considered.

6j4b phase space	Observed z score								
	H_T^b	H_T^j	N_{jets}	$\Delta R_{bb}^{\text{avg}}$	m_{bb}^{max}	$p_T(b_3)$	$p_T(b_4)$	$ \eta(b_3) $	$ \eta(b_4) $
MG5_aMC+P8 $t\bar{t}$ +jets FxFx 5FS	0.41	-0.83	1.91	-1.30	1.63	1.19	0.27	0.10	0.98
MG5_aMC+P8 $t\bar{t}b\bar{b}$ 4FS	0.79	0.81	2.50	-0.02	1.30	1.54	1.11	0.69	0.83
POWHEG+H7 $t\bar{t}$ 5FS	0.92	2.44	1.80	-1.46	0.04	1.24	0.68	0.50	-0.09
POWHEG+OL+P8 $t\bar{t}b\bar{b}$ 4FS	0.60	0.42	1.00	-1.29	0.04	1.36	0.69	0.19	0.15
POWHEG+P8 $t\bar{t}$ 5FS	0.68	0.10	1.08	-0.33	0.34	1.34	0.56	0.38	0.18
SHERPA+OL $t\bar{t}b\bar{b}$ 4FS	0.52	0.17	0.86	0.29	0.65	1.48	0.43	0.52	0.17

Table 8. Observed z score for each of the theoretical predictions in the 6j4b phase space, given the unfolded data and covariance matrix. For the determination of the z score, only the measurement uncertainties are considered.

6j4b phase space	Observed z score							
	$\Delta R(bb^{\text{extra}})$	$m(bb^{\text{extra}})$	$p_T(bb^{\text{extra}})$	$p_T(b_1^{\text{extra}})$	$p_T(b_2^{\text{extra}})$	$ \eta(b_1^{\text{extra}}) $	$ \eta(b_2^{\text{extra}}) $	$ \eta(bb^{\text{extra}}) $
MG5_aMC+P8 $t\bar{t}$ +jets FxFx 5FS	1.31	0.56	0.42	0.27	0.63	0.76	0.17	0.51
MG5_aMC+P8 $t\bar{t}b\bar{b}$ 4FS	0.23	-0.02	0.50	0.52	-0.56	0.38	-0.95	0.62
POWHEG+H7 $t\bar{t}$ 5FS	3.17	1.60	1.00	1.41	0.26	0.47	-0.39	0.88
POWHEG+OL+P8 $t\bar{t}b\bar{b}$ 4FS	-0.79	-0.69	0.60	1.05	-0.19	-0.31	-1.15	0.38
POWHEG+P8 $t\bar{t}$ 5FS	1.33	0.07	0.85	1.24	-0.19	0.03	-1.25	1.07
SHERPA+OL $t\bar{t}b\bar{b}$ 4FS	0.03	-0.99	0.95	1.12	-0.52	0.13	-1.02	0.40

Table 9. Observed z score for each of the theoretical predictions in the 6j4b phase space of the observables related to the bb^{extra} pair, given the unfolded data and covariance matrix. For the determination of the z score, only the measurement uncertainties are considered.

6j4b phase space	$\Delta R(\text{bb}^{\text{add.}})$	$m(\text{bb}^{\text{add.}})$	$p_T(\text{bb}^{\text{add.}})$	Observed z score				
				$p_T(\text{b}_1^{\text{add.}})$	$p_T(\text{b}_2^{\text{add.}})$	$ \eta(\text{bb}^{\text{add.}}) $	$ \eta(\text{b}_1^{\text{add.}}) $	$ \eta(\text{b}_2^{\text{add.}}) $
MG5_aMC+P8 $t\bar{t}$ +jets FxFx 5FS	1.09	0.81	0.83	1.14	0.86	0.30	1.51	0.27
MG5_aMC+P8 $t\bar{t}b\bar{b}$ 4FS	-1.86	-0.14	0.72	1.69	1.27	0.51	1.49	-0.09
POWHEG+H7 $t\bar{t}$ 5FS	3.62	2.45	2.58	2.61	1.05	0.19	1.25	-0.33
POWHEG+OL+P8 $t\bar{t}b\bar{b}$ 4FS	-2.24	-1.43	1.48	2.17	0.90	0.29	1.09	-0.36
POWHEG+P8 $t\bar{t}$ 5FS	-1.21	-0.68	2.24	2.52	0.83	0.95	1.44	-0.57

Table 10. Observed z score for each of the theoretical predictions in the 6j4b phase space of the observables related to the $\text{bb}^{\text{add.}}$ pair, given the unfolded data and covariance matrix. For the determination of the z score, only the measurement uncertainties are considered.

6j3b3l phase space	H_T^{light}	Observed z score	
		$p_T(\text{lj}_1^{\text{extra}})$	$ \Delta\phi(\text{lj}_1^{\text{extra}}, \text{b}_{\text{soft}}) $
MG5_aMC+P8 $t\bar{t}$ +jets FxFx 5FS	1.71	-0.14	0.81
MG5_aMC+P8 $t\bar{t}b\bar{b}$ 4FS	3.55	4.14	1.62
POWHEG+H7 $t\bar{t}$ 5FS	2.02	1.84	1.03
POWHEG+OL+P8 $t\bar{t}b\bar{b}$ 4FS	5.39	1.62	-0.17
POWHEG+P8 $t\bar{t}$ 5FS	1.52	0.24	0.24
SHERPA+OL $t\bar{t}b\bar{b}$ 4FS	0.33	0.78	0.30

Table 11. Observed z score for each of the theoretical predictions in the 6j3b3l phase space, given the unfolded data and covariance matrix. For the determination of the z score, only the measurement uncertainties are considered.

7j4b3l phase space	H_T^{light}	Observed z score	
		$p_T(\text{lj}_1^{\text{extra}})$	$ \Delta\phi(\text{lj}_1^{\text{extra}}, \text{b}_{\text{soft}}) $
MG5_aMC+P8 $t\bar{t}$ +jets FxFx 5FS	1.14	1.66	0.10
MG5_aMC+P8 $t\bar{t}b\bar{b}$ 4FS	-0.92	2.12	0.16
POWHEG+H7 $t\bar{t}$ 5FS	0.49	-0.06	0.03
POWHEG+OL+P8 $t\bar{t}b\bar{b}$ 4FS	2.00	2.30	-0.03
POWHEG+P8 $t\bar{t}$ 5FS	0.89	1.83	0.13
SHERPA+OL $t\bar{t}b\bar{b}$ 4FS	0.31	0.24	0.00

Table 12. Observed z score for each of the theoretical predictions in the 7j4b3l phase space, given the unfolded data and covariance matrix. For the determination of the z score, only the measurement uncertainties are considered.

Open Access. This article is distributed under the terms of the Creative Commons Attribution License ([CC-BY4.0](https://creativecommons.org/licenses/by/4.0/)), which permits any use, distribution and reproduction in any medium, provided the original author(s) and source are credited.

References

- [1] F. Buccioni, S. Kallweit, S. Pozzorini and M.F. Zoller, *NLO QCD predictions for $t\bar{t}b\bar{b}$ production in association with a light jet at the LHC*, *JHEP* **12** (2019) 015 [[arXiv:1907.13624](#)] [[INSPIRE](#)].
- [2] T. Ježo, J.M. Lindert, N. Moretti and S. Pozzorini, *New NLOPS predictions for $t\bar{t} + b$ -jet production at the LHC*, *Eur. Phys. J. C* **78** (2018) 502 [[arXiv:1802.00426](#)] [[INSPIRE](#)].
- [3] ATLAS collaboration, *Search for the Standard Model Higgs boson decaying into $b\bar{b}$ produced in association with top quarks decaying hadronically in pp collisions at $\sqrt{s} = 8$ TeV with the ATLAS detector*, *JHEP* **05** (2016) 160 [[arXiv:1604.03812](#)] [[INSPIRE](#)].
- [4] ATLAS collaboration, *Search for the standard model Higgs boson produced in association with top quarks and decaying into a $b\bar{b}$ pair in pp collisions at $\sqrt{s} = 13$ TeV with the ATLAS detector*, *Phys. Rev. D* **97** (2018) 072016 [[arXiv:1712.08895](#)] [[INSPIRE](#)].
- [5] ATLAS collaboration, *Observation of Higgs boson production in association with a top quark pair at the LHC with the ATLAS detector*, *Phys. Lett. B* **784** (2018) 173 [[arXiv:1806.00425](#)] [[INSPIRE](#)].
- [6] CMS collaboration, *Search for $t\bar{t}H$ production in the all-jet final state in proton-proton collisions at $\sqrt{s} = 13$ TeV*, *JHEP* **06** (2018) 101 [[arXiv:1803.06986](#)] [[INSPIRE](#)].
- [7] CMS collaboration, *Observation of $t\bar{t}H$ production*, *Phys. Rev. Lett.* **120** (2018) 231801 [[arXiv:1804.02610](#)] [[INSPIRE](#)].
- [8] CMS collaboration, *Search for $t\bar{t}H$ production in the $H \rightarrow b\bar{b}$ decay channel with leptonic $t\bar{t}$ decays in proton-proton collisions at $\sqrt{s} = 13$ TeV*, *JHEP* **03** (2019) 026 [[arXiv:1804.03682](#)] [[INSPIRE](#)].
- [9] ATLAS collaboration, *Search for four-top-quark production in the single-lepton and opposite-sign dilepton final states in pp collisions at $\sqrt{s} = 13$ TeV with the ATLAS detector*, *Phys. Rev. D* **99** (2019) 052009 [[arXiv:1811.02305](#)] [[INSPIRE](#)].
- [10] ATLAS collaboration, *Evidence for $t\bar{t}t\bar{t}$ production in the multilepton final state in proton-proton collisions at $\sqrt{s} = 13$ TeV with the ATLAS detector*, *Eur. Phys. J. C* **80** (2020) 1085 [[arXiv:2007.14858](#)] [[INSPIRE](#)].
- [11] ATLAS collaboration, *Measurement of the $t\bar{t}t\bar{t}$ production cross section in pp collisions at $\sqrt{s} = 13$ TeV with the ATLAS detector*, *JHEP* **11** (2021) 118 [[arXiv:2106.11683](#)] [[INSPIRE](#)].
- [12] CMS collaboration, *Search for standard model production of four top quarks in the lepton + jets channel in pp collisions at $\sqrt{s} = 8$ TeV*, *JHEP* **11** (2014) 154 [[arXiv:1409.7339](#)] [[INSPIRE](#)].
- [13] CMS collaboration, *Search for standard model production of four top quarks with same-sign and multilepton final states in proton-proton collisions at $\sqrt{s} = 13$ TeV*, *Eur. Phys. J. C* **78** (2018) 140 [[arXiv:1710.10614](#)] [[INSPIRE](#)].
- [14] CMS collaboration, *Search for standard model production of four top quarks in proton-proton collisions at $\sqrt{s} = 13$ TeV*, *Phys. Lett. B* **772** (2017) 336 [[arXiv:1702.06164](#)] [[INSPIRE](#)].
- [15] CMS collaboration, *Search for the production of four top quarks in the single-lepton and opposite-sign dilepton final states in proton-proton collisions at $\sqrt{s} = 13$ TeV*, *JHEP* **11** (2019) 082 [[arXiv:1906.02805](#)] [[INSPIRE](#)].
- [16] CMS collaboration, *Search for production of four top quarks in final states with same-sign or multiple leptons in proton-proton collisions at $\sqrt{s} = 13$ TeV*, *Eur. Phys. J. C* **80** (2020) 75 [[arXiv:1908.06463](#)] [[INSPIRE](#)].

- [17] CMS collaboration, *Evidence for four-top quark production in proton-proton collisions at $\sqrt{s} = 13$ TeV*, *Phys. Lett. B* **844** (2023) 138076 [[arXiv:2303.03864](#)] [[INSPIRE](#)].
- [18] ATLAS collaboration, *Observation of four-top-quark production in the multilepton final state with the ATLAS detector*, *Eur. Phys. J. C* **83** (2023) 496 [Erratum *ibid.* **84** (2024) 156] [[arXiv:2303.15061](#)] [[INSPIRE](#)].
- [19] CMS collaboration, *Observation of four top quark production in proton-proton collisions at $\sqrt{s} = 13$ TeV*, *Phys. Lett. B* **847** (2023) 138290 [[arXiv:2305.13439](#)] [[INSPIRE](#)].
- [20] Q.-H. Cao, S.-L. Chen and Y. Liu, *Probing Higgs width and top quark Yukawa coupling from $t\bar{t}H$ and $t\bar{t}H$ productions*, *Phys. Rev. D* **95** (2017) 053004 [[arXiv:1602.01934](#)] [[INSPIRE](#)].
- [21] Q.-H. Cao et al., *Limiting top quark-Higgs boson interaction and Higgs-boson width from multitop productions*, *Phys. Rev. D* **99** (2019) 113003 [[arXiv:1901.04567](#)] [[INSPIRE](#)].
- [22] M. Worek, *Next-to-leading order QCD corrections to $t\bar{t}b\bar{b}$ production at the LHC*, *Acta Phys. Polon. B* **40** (2009) 2937 [[arXiv:0910.4080](#)] [[INSPIRE](#)].
- [23] G. Bevilacqua et al., *Assault on the NLO wishlist: $pp \rightarrow t\bar{t}b\bar{b}$* , *JHEP* **09** (2009) 109 [[arXiv:0907.4723](#)] [[INSPIRE](#)].
- [24] A. Bredenstein, A. Denner, S. Dittmaier and S. Pozzorini, *NLO QCD corrections to $pp \rightarrow t\bar{t}b\bar{b} + X$ at the LHC*, *Phys. Rev. Lett.* **103** (2009) 012002 [[arXiv:0905.0110](#)] [[INSPIRE](#)].
- [25] A. Bredenstein, A. Denner, S. Dittmaier and S. Pozzorini, *NLO QCD corrections to top anti-top bottom anti-bottom production at the LHC: 2. Full hadronic results*, *JHEP* **03** (2010) 021 [[arXiv:1001.4006](#)] [[INSPIRE](#)].
- [26] M. Worek, *On the next-to-leading order QCD K-factor for top $t\bar{t}b\bar{b}$ production at the Tevatron*, *JHEP* **02** (2012) 043 [[arXiv:1112.4325](#)] [[INSPIRE](#)].
- [27] A. Denner, J.-N. Lang and M. Pellen, *Full NLO QCD corrections to off-shell $t\bar{t}b\bar{b}$ production*, *Phys. Rev. D* **104** (2021) 056018 [[arXiv:2008.00918](#)] [[INSPIRE](#)].
- [28] G. Bevilacqua et al., *$t\bar{t}b\bar{b}$ at the LHC: on the size of corrections and b-jet definitions*, *JHEP* **08** (2021) 008 [[arXiv:2105.08404](#)] [[INSPIRE](#)].
- [29] S. Höche et al., *Next-to-leading order QCD predictions for top-quark pair production with up to two jets merged with a parton shower*, *Phys. Lett. B* **748** (2015) 74 [[arXiv:1402.6293](#)] [[INSPIRE](#)].
- [30] M.V. Garzelli, A. Kardos and Z. Trócsányi, *Hadroproduction of $t\bar{t}b\bar{b}$ final states at LHC: predictions at NLO accuracy matched with parton shower*, *JHEP* **03** (2015) 083 [[arXiv:1408.0266](#)] [[INSPIRE](#)].
- [31] S. Höche et al., *Next-to-leading order QCD predictions for top-quark pair production with up to three jets*, *Eur. Phys. J. C* **77** (2017) 145 [[arXiv:1607.06934](#)] [[INSPIRE](#)].
- [32] J. Alwall et al., *The automated computation of tree-level and next-to-leading order differential cross sections, and their matching to parton shower simulations*, *JHEP* **07** (2014) 079 [[arXiv:1405.0301](#)] [[INSPIRE](#)].
- [33] R. Frederix and S. Frixione, *Merging meets matching in MC@NLO*, *JHEP* **12** (2012) 061 [[arXiv:1209.6215](#)] [[INSPIRE](#)].
- [34] S. Frixione, P. Nason and G. Ridolfi, *A positive-weight next-to-leading-order Monte Carlo for heavy flavour hadroproduction*, *JHEP* **09** (2007) 126 [[arXiv:0707.3088](#)] [[INSPIRE](#)].
- [35] F. Cascioli et al., *NLO matching for $t\bar{t}b\bar{b}$ production with massive b-quarks*, *Phys. Lett. B* **734** (2014) 210 [[arXiv:1309.5912](#)] [[INSPIRE](#)].

- [36] G. Bevilacqua, M.V. Garzelli and A. Kardos, $t\bar{t}b\bar{b}$ hadroproduction with massive bottom quarks with *PowHel*, [arXiv:1709.06915](#) [INSPIRE].
- [37] ATLAS collaboration, *Study of heavy-flavor quarks produced in association with top-quark pairs at $\sqrt{s} = 7$ TeV using the ATLAS detector*, *Phys. Rev. D* **89** (2014) 072012 [[arXiv:1304.6386](#)] [INSPIRE].
- [38] ATLAS collaboration, *Measurements of fiducial cross-sections for $t\bar{t}$ production with one or two additional b-jets in pp collisions at $\sqrt{s} = 8$ TeV using the ATLAS detector*, *Eur. Phys. J. C* **76** (2016) 11 [[arXiv:1508.06868](#)] [INSPIRE].
- [39] ATLAS collaboration, *Measurements of inclusive and differential fiducial cross-sections of $t\bar{t}$ production with additional heavy-flavour jets in proton-proton collisions at $\sqrt{s} = 13$ TeV with the ATLAS detector*, *JHEP* **04** (2019) 046 [[arXiv:1811.12113](#)] [INSPIRE].
- [40] CMS collaboration, *Measurement of $t\bar{t}$ production with additional jet activity, including b quark jets, in the dilepton decay channel using pp collisions at $\sqrt{s} = 8$ TeV*, *Eur. Phys. J. C* **76** (2016) 379 [[arXiv:1510.03072](#)] [INSPIRE].
- [41] CMS collaboration, *Measurement of the cross section ratio $\sigma_{t\bar{t}b\bar{b}}/\sigma_{t\bar{t}jj}$ in pp collisions at $\sqrt{s} = 8$ TeV*, *Phys. Lett. B* **746** (2015) 132 [[arXiv:1411.5621](#)] [INSPIRE].
- [42] CMS collaboration, *Measurements of $t\bar{t}$ cross sections in association with b jets and inclusive jets and their ratio using dilepton final states in pp collisions at $\sqrt{s} = 13$ TeV*, *Phys. Lett. B* **776** (2018) 355 [[arXiv:1705.10141](#)] [INSPIRE].
- [43] CMS collaboration, *Measurement of the $t\bar{t}b\bar{b}$ production cross section in the all-jet final state in pp collisions at $\sqrt{s} = 13$ TeV*, *Phys. Lett. B* **803** (2020) 135285 [[arXiv:1909.05306](#)] [INSPIRE].
- [44] CMS collaboration, *Measurement of the cross section for $t\bar{t}$ production with additional jets and b jets in pp collisions at $\sqrt{s} = 13$ TeV*, *JHEP* **07** (2020) 125 [[arXiv:2003.06467](#)] [INSPIRE].
- [45] CMS collaboration, *First measurement of the cross section for top quark pair production with additional charm jets using dileptonic final states in pp collisions at $\sqrt{s} = 13$ TeV*, *Phys. Lett. B* **820** (2021) 136565 [[arXiv:2012.09225](#)] [INSPIRE].
- [46] *HEPData record for this analysis*, CMS-TOP-22-009, (2023).
- [47] CMS collaboration, *The CMS experiment at the CERN LHC*, 2008 *JINST* **3** S08004 [INSPIRE].
- [48] CMS TRACKER GROUP collaboration, *The CMS phase-1 pixel detector upgrade*, 2021 *JINST* **16** P02027 [[arXiv:2012.14304](#)] [INSPIRE].
- [49] CMS collaboration, *Performance of the CMS level-1 trigger in proton-proton collisions at $\sqrt{s} = 13$ TeV*, 2020 *JINST* **15** P10017 [[arXiv:2006.10165](#)] [INSPIRE].
- [50] CMS collaboration, *The CMS trigger system*, 2017 *JINST* **12** P01020 [[arXiv:1609.02366](#)] [INSPIRE].
- [51] T. Sjöstrand et al., *An introduction to PYTHIA 8.2*, *Comput. Phys. Commun.* **191** (2015) 159 [[arXiv:1410.3012](#)] [INSPIRE].
- [52] GEANT4 collaboration, *GEANT4 — a simulation toolkit*, *Nucl. Instrum. Meth. A* **506** (2003) 250 [INSPIRE].
- [53] T. Ježo and P. Nason, *On the treatment of resonances in next-to-leading order calculations matched to a parton shower*, *JHEP* **12** (2015) 065 [[arXiv:1509.09071](#)] [INSPIRE].
- [54] F. Buccioni et al., *OpenLoops 2*, *Eur. Phys. J. C* **79** (2019) 866 [[arXiv:1907.13071](#)] [INSPIRE].


- [55] M. Bähr et al., *Herwig++ physics and manual*, *Eur. Phys. J. C* **58** (2008) 639 [[arXiv:0803.0883](#)] [[INSPIRE](#)].
- [56] J. Bellm et al., *Herwig 7.0/Herwig++ 3.0 release note*, *Eur. Phys. J. C* **76** (2016) 196 [[arXiv:1512.01178](#)] [[INSPIRE](#)].
- [57] CMS collaboration, *Development and validation of Herwig 7 tunes from CMS underlying-event measurements*, *Eur. Phys. J. C* **81** (2021) 312 [[arXiv:2011.03422](#)] [[INSPIRE](#)].
- [58] T. Gleisberg et al., *Event generation with SHERPA 1.1*, *JHEP* **02** (2009) 007 [[arXiv:0811.4622](#)] [[INSPIRE](#)].
- [59] M. Czakon and A. Mitov, *Top++: a program for the calculation of the top-pair cross-section at hadron colliders*, *Comput. Phys. Commun.* **185** (2014) 2930 [[arXiv:1112.5675](#)] [[INSPIRE](#)].
- [60] N. Kidonakis, *Top quark production*, in the proceedings of the *Helmholtz international summer school on physics of heavy quarks and hadrons*, (2014) [[DOI:10.3204/DESY-PROC-2013-03/Kidonakis](#)] [[arXiv:1311.0283](#)] [[INSPIRE](#)].
- [61] P. Nason, *A new method for combining NLO QCD with shower Monte Carlo algorithms*, *JHEP* **11** (2004) 040 [[hep-ph/0409146](#)] [[INSPIRE](#)].
- [62] S. Frixione, P. Nason and C. Oleari, *Matching NLO QCD computations with parton shower simulations: the POWHEG method*, *JHEP* **11** (2007) 070 [[arXiv:0709.2092](#)] [[INSPIRE](#)].
- [63] S. Alioli, P. Nason, C. Oleari and E. Re, *A general framework for implementing NLO calculations in shower Monte Carlo programs: the POWHEG BOX*, *JHEP* **06** (2010) 043 [[arXiv:1002.2581](#)] [[INSPIRE](#)].
- [64] S. Alioli, P. Nason, C. Oleari and E. Re, *NLO single-top production matched with shower in POWHEG: s- and t-channel contributions*, *JHEP* **09** (2009) 111 [*Erratum ibid.* **02** (2010) 011] [[arXiv:0907.4076](#)] [[INSPIRE](#)].
- [65] E. Re, *Single-top Wt-channel production matched with parton showers using the POWHEG method*, *Eur. Phys. J. C* **71** (2011) 1547 [[arXiv:1009.2450](#)] [[INSPIRE](#)].
- [66] H.B. Hartanto, B. Jäger, L. Reina and D. Wackerth, *Higgs boson production in association with top quarks in the POWHEG BOX*, *Phys. Rev. D* **91** (2015) 094003 [[arXiv:1501.04498](#)] [[INSPIRE](#)].
- [67] P. Artoisenet, R. Frederix, O. Mattelaer and R. Rietkerk, *Automatic spin-entangled decays of heavy resonances in Monte Carlo simulations*, *JHEP* **03** (2013) 015 [[arXiv:1212.3460](#)] [[INSPIRE](#)].
- [68] J. Alwall et al., *Comparative study of various algorithms for the merging of parton showers and matrix elements in hadronic collisions*, *Eur. Phys. J. C* **53** (2008) 473 [[arXiv:0706.2569](#)] [[INSPIRE](#)].
- [69] NNPDF collaboration, *Parton distributions for the LHC run II*, *JHEP* **04** (2015) 040 [[arXiv:1410.8849](#)] [[INSPIRE](#)].
- [70] CMS collaboration, *Extraction and validation of a new set of CMS PYTHIA8 tunes from underlying-event measurements*, *Eur. Phys. J. C* **80** (2020) 4 [[arXiv:1903.12179](#)] [[INSPIRE](#)].
- [71] Y. Li and F. Petriello, *Combining QCD and electroweak corrections to dilepton production in FEWZ*, *Phys. Rev. D* **86** (2012) 094034 [[arXiv:1208.5967](#)] [[INSPIRE](#)].
- [72] CMS collaboration, *Particle-flow reconstruction and global event description with the CMS detector*, *2017 JINST* **12** P10003 [[arXiv:1706.04965](#)] [[INSPIRE](#)].

- [73] D. Contardo et al., *Technical proposal for the phase-II upgrade of the CMS detector*, CERN-LHCC-2015-010, CERN, Geneva, Switzerland (2015) [DOI:10.17181/CERN.VU8I.D59J].
- [74] CMS collaboration, *Electron and photon reconstruction and identification with the CMS experiment at the CERN LHC*, 2021 *JINST* **16** P05014 [arXiv:2012.06888] [INSPIRE].
- [75] CMS collaboration, *Performance of the CMS muon detector and muon reconstruction with proton-proton collisions at $\sqrt{s} = 13$ TeV*, 2018 *JINST* **13** P06015 [arXiv:1804.04528] [INSPIRE].
- [76] CMS collaboration, *Pileup mitigation at CMS in 13 TeV data*, 2020 *JINST* **15** P09018 [arXiv:2003.00503] [INSPIRE].
- [77] M. Cacciari, G.P. Salam and G. Soyez, *The anti- k_t jet clustering algorithm*, *JHEP* **04** (2008) 063 [arXiv:0802.1189] [INSPIRE].
- [78] M. Cacciari, G.P. Salam and G. Soyez, *FastJet user manual*, *Eur. Phys. J. C* **72** (2012) 1896 [arXiv:1111.6097] [INSPIRE].
- [79] CMS collaboration, *Jet energy scale and resolution in the CMS experiment in pp collisions at 8 TeV*, 2017 *JINST* **12** P02014 [arXiv:1607.03663] [INSPIRE].
- [80] CMS collaboration, *Identification of heavy-flavour jets with the CMS detector in pp collisions at 13 TeV*, 2018 *JINST* **13** P05011 [arXiv:1712.07158] [INSPIRE].
- [81] E. Bols et al., *Jet flavour classification using DeepJet*, 2020 *JINST* **15** P12012 [arXiv:2008.10519] [INSPIRE].
- [82] CMS collaboration, *Performance summary of AK4 jet b tagging with data from proton-proton collisions at 13 TeV with the CMS detector*, CMS-DP-2023-005, CERN, Geneva, Switzerland (2023).
- [83] M. Cacciari and G.P. Salam, *Pileup subtraction using jet areas*, *Phys. Lett. B* **659** (2008) 119 [arXiv:0707.1378] [INSPIRE].
- [84] CMS collaboration, *Simulation of the silicon strip tracker pre-amplifier in early 2016 data*, CMS-DP-2020-045, CERN, Geneva, Switzerland (2020).
- [85] Y. LeCun et al., *Handwritten digit recognition with a back-propagation network*, in *Proc. 2nd Int. Conf. on Advances in Neural Information Processing Systems (NIPS'89)*, <https://dblp.org/db/conf/nips/nips1989.html>, Denver, CO, U.S.A., 27–30 November 1989, p. 396.
- [86] S. Hochreiter and J. Schmidhuber, *Long short-term memory*, *Neural Comput.* **9** (1997) 1735 [INSPIRE].
- [87] F. Chollet et al., *KERAS: deep learning for humans*, <https://github.com/keras-team/keras>, (2023).
- [88] *Keras webpage*, <http://keras.io>, accessed 19 September 2023.
- [89] M. Abadi et al., *TensorFlow: large-scale machine learning on heterogeneous distributed systems*, arXiv:1603.04467 [INSPIRE].
- [90] D.P. Kingma and J. Ba, *Adam: a method for stochastic optimization*, in *Proc. 3rd Int. Conf. on Learning Representations (ICLR 2015)*, San Diego, CA, U.S.A., 7–9 May 2015 [arXiv:1412.6980] [INSPIRE].
- [91] N. Srivastava et al., *Dropout: a simple way to prevent neural networks from overfitting*, *J. Machine Learning Res.* **15** (2014) 1929 [INSPIRE].



















- [92] S. Wertz, *SMOOFIT: a package for smooth binned likelihood fits*, <https://gitlab.cern.ch/swertz/smoofit>, (2022).
- [93] *Smoofit webpage*, <https://smoofit.readthedocs.io>, accessed 19 September 2023.
- [94] J. Bradbury et al., *JAX: AUTOGRAD and XLA*, <https://github.com/google/jax>, (2023).
- [95] *JAX webpage*, <http://jax.readthedocs.io>, accessed 19 September 2023.
- [96] CMS collaboration, *Precision luminosity measurement in proton-proton collisions at $\sqrt{s} = 13$ TeV in 2015 and 2016 at CMS*, *Eur. Phys. J. C* **81** (2021) 800 [[arXiv:2104.01927](https://arxiv.org/abs/2104.01927)] [[INSPIRE](#)].
- [97] CMS collaboration, *CMS luminosity measurement for the 2017 data-taking period at $\sqrt{s} = 13$ TeV*, *CMS-PAS-LUM-17-004*, CERN, Geneva, Switzerland (2018).
- [98] CMS collaboration, *CMS luminosity measurement for the 2018 data-taking period at $\sqrt{s} = 13$ TeV*, *CMS-PAS-LUM-18-002*, CERN, Geneva, Switzerland (2019).
- [99] M. Czakon et al., *Top-pair production at the LHC through NNLO QCD and NLO EW*, *JHEP* **10** (2017) 186 [[arXiv:1705.04105](https://arxiv.org/abs/1705.04105)] [[INSPIRE](#)].
- [100] S. Argyropoulos and T. Sjöstrand, *Effects of color reconnection on $t\bar{t}$ final states at the LHC*, *JHEP* **11** (2014) 043 [[arXiv:1407.6653](https://arxiv.org/abs/1407.6653)] [[INSPIRE](#)].
- [101] J.R. Christiansen and P.Z. Skands, *String formation beyond leading colour*, *JHEP* **08** (2015) 003 [[arXiv:1505.01681](https://arxiv.org/abs/1505.01681)] [[INSPIRE](#)].
- [102] ATLAS collaboration, *Measurement of b-quark fragmentation properties in jets using the decay $B^\pm \rightarrow J/\psi K^\pm$ in pp collisions at $\sqrt{s} = 13$ TeV with the ATLAS detector*, *JHEP* **12** (2021) 131 [[arXiv:2108.11650](https://arxiv.org/abs/2108.11650)] [[INSPIRE](#)].
- [103] R.J. Barlow and C. Beeston, *Fitting using finite Monte Carlo samples*, *Comput. Phys. Commun.* **77** (1993) 219 [[INSPIRE](#)].
- [104] J.S. Conway, *Incorporating nuisance parameters in likelihoods for multisource spectra*, in the proceedings of the *PHYSTAT 2011*, (2011) [[DOI:10.5170/CERN-2011-006.115](https://doi.org/10.5170/CERN-2011-006.115)] [[arXiv:1103.0354](https://arxiv.org/abs/1103.0354)] [[INSPIRE](#)].
- [105] L. Ferencz et al., *Study of $t\bar{t}b\bar{b}$ and $t\bar{t}W$ background modelling for $t\bar{t}H$ analyses*, [arXiv:2301.11670](https://arxiv.org/abs/2301.11670) [[INSPIRE](#)].
- [106] R.D. Cousins, J.T. Linnemann and J. Tucker, *Evaluation of three methods for calculating statistical significance when incorporating a systematic uncertainty into a test of the background-only hypothesis for a Poisson process*, *Nucl. Instrum. Meth. A* **595** (2008) 480 [[physics/0702156](https://arxiv.org/abs/physics/0702156)] [[INSPIRE](#)].

The CMS collaboration

Yerevan Physics Institute, Yerevan, Armenia

A. Hayrapetyan, A. Tumasyan ¹




Institut für Hochenergiephysik, Vienna, Austria

W. Adam , J.W. Andrejkovic, T. Bergauer , S. Chatterjee , K. Damanakis , M. Dragicevic ,
A. Escalante Del Valle , P.S. Hussain , M. Jeitler ², N. Krammer , D. Liko , I. Mikulec ,
J. Schieck ², R. Schöfbeck , D. Schwarz , M. Sonawane , S. Templ , W. Waltenberger ,
C.-E. Wulz ²















Universiteit Antwerpen, Antwerpen, Belgium

M.R. Darwish ³, T. Janssen , P. Van Mechelen 

Vrije Universiteit Brussel, Brussel, Belgium

E.S. Bols , J. D'Hondt , S. Dansana , A. De Moor , M. Delcourt , H. El Faham ,
S. Lowette , I. Makarenko , A. Morton , D. Müller , A.R. Sahasransu , S. Tavernier ,
M. Tytgat ⁴, S. Van Putte , D. Vannerom 

Université Libre de Bruxelles, Bruxelles, Belgium

B. Clerbaux , G. De Lentdecker , L. Favart , D. Hohov , J. Jaramillo , A. Khalilzadeh,
K. Lee , M. Mahdavihorrani , A. Malara , S. Paredes , L. Pétré , N. Postiau, L. Thomas ,
M. Vanden Bemden , C. Vander Velde , P. Vanlaer 






Ghent University, Ghent, Belgium

M. De Coen , D. Dobur , Y. Hong , J. Knolle , L. Lambrecht , G. Mestdach, C. Rendón,
A. Samalan, K. Skovpen , N. Van Den Bossche , L. Wezenbeek 




















Université Catholique de Louvain, Louvain-la-Neuve, Belgium

A. Benecke , G. Bruno , C. Caputo , C. Delaere , I.S. Donertas , A. Giammanco , K. Jaffel ,
Sa. Jain , V. Lemaitre, J. Lidrych , P. Mastrapasqua , K. Mondal , T.T. Tran , S. Wertz 



Centro Brasileiro de Pesquisas Físicas, Rio de Janeiro, Brazil

G.A. Alves , E. Coelho , C. Hensel , T. Menezes De Oliveira, A. Moraes , P. Rebello Teles ,
M. Soeiro







Universidade do Estado do Rio de Janeiro, Rio de Janeiro, Brazil

W.L. Aldá Júnior , M. Alves Gallo Pereira , M. Barroso Ferreira Filho ,
H. Brandao Malbouisson , W. Carvalho , J. Chinellato⁵, E.M. Da Costa , G.G. Da Silveira ⁶,
D. De Jesus Damiao , S. Fonseca De Souza , J. Martins ⁷, C. Mora Herrera ,
K. Mota Amarilo , L. Mundim , H. Nogima , A. Santoro , S.M. Silva Do Amaral ,
A. Sznajder , M. Thiel , A. Vilela Pereira 

Universidade Estadual Paulista, Universidade Federal do ABC, São Paulo, Brazil

C.A. Bernardes ⁶, L. Calligaris , T.R. Fernandez Perez Tomei , E.M. Gregores ,
P.G. Mercadante , S.F. Novaes , B. Orzari , Sandra S. Padula 

Institute for Nuclear Research and Nuclear Energy, Bulgarian Academy of Sciences, Sofia, Bulgaria

A. Aleksandrov , G. Antchev , R. Hadjiiska , P. Iaydjiev , M. Misheva , M. Shopova , G. Sultanov 




University of Sofia, Sofia, Bulgaria

A. Dimitrov , T. Ivanov , L. Litov , B. Pavlov , P. Petkov , A. Petrov , E. Shumka 

Instituto De Alta Investigación, Universidad de Tarapacá, Casilla 7 D, Arica, Chile

S. Keshri , S. Thakur 













Beihang University, Beijing, China

T. Cheng , Q. Guo, T. Javaid , M. Mittal , L. Yuan 










Department of Physics, Tsinghua University, Beijing, China

G. Bauer⁸, Z. Hu , K. Yi ^{8,9}

Institute of High Energy Physics, Beijing, China

G.M. Chen ¹⁰, H.S. Chen ¹⁰, M. Chen ¹⁰, F. Iemmi , C.H. Jiang, A. Kapoor , H. Liao , Z.-A. Liu ¹¹, F. Monti , R. Sharma , J.N. Song¹¹, J. Tao , J. Wang , H. Zhang 


State Key Laboratory of Nuclear Physics and Technology, Peking University, Beijing, China

A. Agapitos , Y. Ban , A. Levin , C. Li , Q. Li , X. Lyu, Y. Mao, S.J. Qian , X. Sun , D. Wang , H. Yang, C. Zhou 




Sun Yat-Sen University, Guangzhou, China

Z. You 

University of Science and Technology of China, Hefei, China

N. Lu 

Institute of Modern Physics and Key Laboratory of Nuclear Physics and Ion-beam Application (MOE) — Fudan University, Shanghai, China

X. Gao ¹², D. Leggat, H. Okawa , Y. Zhang 

Zhejiang University, Hangzhou, Zhejiang, China

Z. Lin , C. Lu , M. Xiao 



Universidad de Los Andes, Bogota, Colombia

C. Avila , D.A. Barbosa Trujillo, A. Cabrera , C. Florez , J. Fraga , J.A. Reyes Vega

Universidad de Antioquia, Medellin, Colombia

J. Mejia Guisao , F. Ramirez , M. Rodriguez , J.D. Ruiz Alvarez 

University of Split, Faculty of Electrical Engineering, Mechanical Engineering and Naval Architecture, Split, Croatia

D. Giljanovic , N. Godinovic , D. Lelas , A. Sculac 









University of Split, Faculty of Science, Split, Croatia

M. Kovac , T. Sculac 

Institute Rudjer Boskovic, Zagreb, Croatia

P. Bargassa , V. Brigljevic , B.K. Chitroda , D. Ferencek , S. Mishra , A. Starodumov ¹³, T. Susa 

University of Cyprus, Nicosia, Cyprus

A. Attikis , K. Christoforou , S. Konstantinou , J. Mousa , C. Nicolaou, F. Ptochos , P.A. Razis , H. Rykaczewski, H. Saka , A. Stepennov 

Charles University, Prague, Czech Republic

M. Finger , M. Finger Jr. , A. Kveton 


Escuela Politecnica Nacional, Quito, Ecuador

E. Ayala 

Universidad San Francisco de Quito, Quito, Ecuador

E. Carrera Jarrin 










Academy of Scientific Research and Technology of the Arab Republic of Egypt, Egyptian Network of High Energy Physics, Cairo, Egypt

H. Abdalla ¹⁴, Y. Assran^{15,16}

Center for High Energy Physics (CHEP-FU), Fayoum University, El-Fayoum, Egypt

M.A. Mahmoud , Y. Mohammed 

















National Institute of Chemical Physics and Biophysics, Tallinn, Estonia

R.K. Dewanjee ¹⁷, K. Ehataht , M. Kadastik, T. Lange , S. Nandan , C. Nielsen , J. Pata , M. Raidal , L. Tani , C. Veelken 

Department of Physics, University of Helsinki, Helsinki, Finland

H. Kirschenmann , K. Osterberg , M. Voutilainen 















Helsinki Institute of Physics, Helsinki, Finland

S. Bharthuar , E. Brücken , F. Garcia , J. Havukainen , K.T.S. Kallonen , M.S. Kim , R. Kinnunen, T. Lampén , K. Lassila-Perini , S. Lehti , T. Lindén , M. Lotti, L. Martikainen , M. Myllymäki , M.m. Rantanen , H. Siikonen , E. Tuominen , J. Tuominiemi 






























Lappeenranta-Lahti University of Technology, Lappeenranta, Finland

P. Luukka , H. Petrow , T. Tuuva[†]














IRFU, CEA, Université Paris-Saclay, Gif-sur-Yvette, France

M. Besancon , F. Couderc , M. Dejardin , D. Denegri, J.L. Faure, F. Ferri , S. Ganjour ,
 P. Gras , G. Hamel de Monchenault , V. Lohezic , J. Malcles , J. Rander, A. Rosowsky ,
 M.Ö. Sahin , A. Savoy-Navarro ¹⁸, P. Simkina , M. Titov 


















Laboratoire Leprince-Ringuet, CNRS/IN2P3, Ecole Polytechnique, Institut Polytechnique de Paris, Palaiseau, France

C. Baldenegro Barrera , F. Beaudette , A. Buchot Perraguin , P. Busson , A. Cappati ,
 C. Charlot , F. Damas , O. Davignon , G. Falmagne , B.A. Fontana Santos Alves ,
 S. Ghosh , A. Gilbert , R. Granier de Cassagnac , A. Hakimi , B. Harikrishnan ,
 L. Kalipoliti , G. Liu , J. Motta , M. Nguyen , C. Ochando , L. Portales , R. Salerno ,
 U. Sarkar , J.B. Sauvan , Y. Sirois , A. Tarabini , E. Vernazza , A. Zabi , A. Zghiche 

Université de Strasbourg, CNRS, IPHC UMR 7178, Strasbourg, France

J.-L. Agram ¹⁹, J. Andrea , D. Apparu , D. Bloch , J.-M. Brom , E.C. Chabert ,
 C. Collard , S. Falke , U. Goerlach , C. Grimault, R. Haeberle , A.-C. Le Bihan ,
 M.A. Sessini , P. Van Hove 

Institut de Physique des 2 Infinis de Lyon (IP2I), Villeurbanne, France

S. Beauceron , B. Blancon , G. Boudoul , N. Chanon , J. Choi , D. Contardo , P. Depasse ,
 C. Dozen ²⁰, H. El Mamouni, J. Fay , S. Gascon , M. Gouzevitch , C. Greenberg, G. Grenier ,
 B. Ille , I.B. Laktineh, M. Lethuillier , L. Mirabito, S. Perries, M. Vander Donckt , P. Verdier ,
 J. Xiao 




























Georgian Technical University, Tbilisi, Georgia

I. Lomidze , T. Toriashvili ²¹, Z. Tsamalaidze ¹³










RWTH Aachen University, I. Physikalisches Institut, Aachen, Germany

V. Botta , L. Feld , K. Klein , M. Lipinski , D. Meuser , A. Pauls , N. Röwert ,
 M. Teroerde 











RWTH Aachen University, III. Physikalisches Institut A, Aachen, Germany

S. Diekmann , A. Dodonova , N. Eich , D. Eliseev , F. Engelke , M. Erdmann ,
 P. Fackeldey , B. Fischer , T. Hebbeker , K. Hoepfner , F. Ivone , A. Jung , M.y. Lee ,
 L. Mastrolorenzo, M. Merschmeyer , A. Meyer , S. Mukherjee , D. Noll , A. Novak ,
 F. Nowotny, A. Pozdnyakov , Y. Rath, W. Redjeb , F. Rehm, H. Reithler , V. Sarkisovi ,
 A. Schmidt , S.C. Schuler, A. Sharma , A. Stein , F. Torres Da Silva De Araujo ²², L. Vigilante,
 S. Wiedenbeck , S. Zaleski

RWTH Aachen University, III. Physikalisches Institut B, Aachen, Germany







































C. Dziwok , G. Flügge , W. Haj Ahmad ²³, T. Kress , A. Nowack , O. Pooth , A. Stahl ,
 T. Ziemons , A. Zotz 

Deutsches Elektronen-Synchrotron, Hamburg, Germany































H. Aarup Petersen , M. Aldaya Martin , J. Alimena , S. Amoroso, Y. An , S. Baxter ,
 M. Bayatmakou , H. Becerril Gonzalez , O. Behnke , A. Belvedere , S. Bhattacharya 

F. Blekman ²⁴, K. Borras ²⁵, D. Brunner , A. Campbell , A. Cardini , C. Cheng, F. Colombina , S. Consuegra Rodríguez , G. Correia Silva , M. De Silva , G. Eckerlin, D. Eckstein , L.I. Estevez Banos , O. Filatov , E. Gallo ²⁴, A. Geiser , A. Giraldi , G. Greau, V. Guglielmi , M. Guthoff , A. Hinzmann , A. Jafari ²⁶, L. Jeppe , N.Z. Jomhari , B. Kaech , M. Kasemann , H. Kaveh , C. Kleinwort , R. Kogler , M. Komm , D. Krücker , W. Lange, D. Leyva Pernia , K. Lipka ²⁷, W. Lohmann ²⁸, R. Mankel , I.-A. Melzer-Pellmann , M. Mendizabal Morentin , J. Metwally, A.B. Meyer , G. Milella , A. Mussgiller , A. Nürnberg , Y. Otarid, D. Pérez Adán , E. Ranken , A. Raspereza , B. Ribeiro Lopes , J. Rübenach, A. Saggio , M. Scham ^{29,25}, V. Scheurer, S. Schnake ²⁵, P. Schütze , C. Schwanenberger ²⁴, M. Shchedrolosiev , R.E. Sosa Ricardo , L.P. Sreelatha Pramod , D. Stafford, F. Vazzoler , A. Ventura Barroso , R. Walsh , Q. Wang , Y. Wen , K. Wichmann, L. Wiens ²⁵, C. Wissing , S. Wuchterl , Y. Yang , A. Zimmermann Castro Santos 

University of Hamburg, Hamburg, Germany

A. Albrecht , S. Albrecht , M. Antonello , S. Bein , L. Benato , M. Bonanomi , P. Connor , M. Eich, K. El Morabit , Y. Fischer , A. Fröhlich, C. Garbers , E. Garutti , A. Grohsjean , M. Hajheidari, J. Haller , H.R. Jabusch , G. Kasieczka , P. Keicher, R. Klanner , W. Korcaric , T. Kramer , V. Kutzner , F. Labe , J. Lange , A. Lobanov , C. Matthies , A. Mehta , L. Moureaux , M. Mrowietz, A. Nigamova , Y. Nissan, A. Paasch , K.J. Pena Rodriguez , T. Quadfasel , B. Raciti , M. Rieger , D. Savoie , J. Schindler , P. Schleper , M. Schröder , J. Schwandt , M. Sommerhalder , H. Stadie , G. Steinbrück , A. Tews, M. Wolf 









Karlsruher Institut fuer Technologie, Karlsruhe, Germany

S. Brommer , M. Burkart, E. Butz , T. Chwalek , A. Dierlamm , A. Droll, N. Faltermann , M. Giffels , A. Gottmann , F. Hartmann ³⁰, M. Horzela , U. Husemann , M. Klute , R. Koppenhöfer , M. Link, A. Lintuluoto , S. Maier , S. Mitra , M. Mormile , Th. Müller , M. Neukum, M. Oh , E. Pfeffer, G. Quast , K. Rabbertz , I. Shvetsov , H.J. Simonis , N. Trevisani , R. Ulrich , J. van der Linden , R.F. Von Cube , M. Wassmer , S. Wieland , F. Wittig, R. Wolf , S. Wunsch, X. Zuo 

Institute of Nuclear and Particle Physics (INPP), NCSR Demokritos, Aghia Paraskevi, Greece

G. Anagnostou, P. Assiouras , G. Daskalakis , A. Kyriakis, A. Papadopoulos³⁰, A. Stakia 








National and Kapodistrian University of Athens, Athens, Greece

D. Karasavvas, P. Kontaxakis , G. Melachroinos, A. Panagiotou, I. Papavergou , I. Paraskevas , N. Saoulidou , K. Theofilatos , E. Tziaferi , K. Vellidis , I. Zisopoulos 










National Technical University of Athens, Athens, Greece

G. Bakas , T. Chatzistavrou, G. Karapostoli , K. Kousouris , I. Papakrivopoulos , E. Siamarkou, G. Tsiapolitis, A. Zacharopoulou

University of Ioánnina, Ioánnina, Greece

K. Adamidis, I. Bestintzanos, I. Evangelou , C. Foudas, P. Gianneios , C. Kamtsikis, P. Katsoulis, P. Kokkas , P.G. Kosmoglou Kioseoglou , N. Manthos , I. Papadopoulos , J. Strologas 

MTA-ELTE Lendület CMS Particle and Nuclear Physics Group, Eötvös Loránd University, Budapest, Hungary

M. Csanád , K. Farkas , M.M.A. Gadallah ³¹, Á. Kadlecik , P. Major , K. Mandal , G. Pásztor , A.J. Rádl ³², G.I. Veres 

Wigner Research Centre for Physics, Budapest, Hungary

M. Bartók ³³, C. Hajdu , D. Horvath ^{34,35}, F. Sikler , V. Veszpremi 

Institute of Nuclear Research ATOMKI, Debrecen, Hungary

G. Bencze, S. Czellar, J. Karancsi ³³, J. Molnar, Z. Szillasi

















Institute of Physics, University of Debrecen, Debrecen, Hungary

P. Raics, B. Ujvari ³⁶, G. Zilizi 









Karoly Robert Campus, MATE Institute of Technology, Gyongyos, Hungary

T. Csorgo ³², F. Nemes ³², T. Novak 

Panjab University, Chandigarh, India

J. Babbar , S. Bansal , S.B. Beri, V. Bhatnagar , G. Chaudhary , S. Chauhan , N. Dhingra ³⁷, R. Gupta, A. Kaur , A. Kaur , H. Kaur , M. Kaur , S. Kumar , P. Kumari , M. Meena , K. Sandeep , T. Sheokand, J.B. Singh ³⁸, A. Singla 
















University of Delhi, Delhi, India

A. Ahmed , A. Bhardwaj , A. Chhetri , B.C. Choudhary , A. Kumar , M. Naimuddin , K. Ranjan , S. Saumya 

Saha Institute of Nuclear Physics, HBNI, Kolkata, India

S. Baradia , S. Barman ³⁹, S. Bhattacharya , D. Bhowmik, S. Dutta , S. Dutta, B. Gomber ⁴⁰, P. Palit , G. Saha , B. Sahu ⁴⁰, S. Sarkar










Indian Institute of Technology Madras, Madras, India

P.K. Behera , S.C. Behera , S. Chatterjee , P. Jana , P. Kalbhor , J.R. Komaragiri ⁴¹, D. Kumar ⁴¹, M. Mohammad Mobassir Ameen , L. Panwar ⁴¹, R. Pradhan , P.R. Pujahari , N.R. Saha , A. Sharma , A.K. Sikdar , S. Verma 









Tata Institute of Fundamental Research-A, Mumbai, India




































































































T. Aziz, I. Das , S. Dugad, M. Kumar , G.B. Mohanty , P. Suryadevara

Tata Institute of Fundamental Research-B, Mumbai, India












A. Bala , S. Banerjee , R.M. Chatterjee, M. Guchait , S. Karmakar , S. Kumar , G. Majumder , K. Mazumdar , S. Mukherjee , A. Thachayath 

National Institute of Science Education and Research, An OCC of Homi Bhabha National Institute, Bhubaneswar, Odisha, India




























S. Bahinipati ⁴², A.K. Das, C. Kar , D. Maity ⁴³, P. Mal , T. Mishra , V.K. Muraleedharan Nair Bindhu ⁴³, K. Naskar ⁴³, A. Nayak ⁴³, P. Sadangi, P. Saha , S.K. Swain , S. Varghese ⁴³, D. Vats ⁴³

Indian Institute of Science Education and Research (IISER), Pune, IndiaA. Alpana , S. Dube , B. Kansal , A. Laha , A. Rastogi , S. Sharma **Isfahan University of Technology, Isfahan, Iran**H. Bakhshiansohi ⁴⁴, E. Khazaie , M. Zeinali ⁴⁵**Institute for Research in Fundamental Sciences (IPM), Tehran, Iran**S. Chenarani ⁴⁶, S.M. Etesami , M. Khakzad , M. Mohammadi Najafabadi **University College Dublin, Dublin, Ireland**M. Grunewald **INFN Sezione di Bari^a, Università di Bari^b, Politecnico di Bari^c, Bari, Italy**M. Abbrescia ^{a,b}, R. Aly ^{a,c,47}, A. Colaleo ^a, D. Creanza ^{a,c}, B. D' Anzi ^{a,b},
N. De Filippis ^{a,c}, M. De Palma ^{a,b}, A. Di Florio ^{a,c}, W. Elmetenawee ^{a,b,47}, L. Fiore ^a,
G. Iaselli ^{a,c}, G. Maggi ^{a,c}, M. Maggi ^a, I. Margjeka ^{a,b}, V. Mastrapasqua ^{a,b}, S. My ^{a,b},
S. Nuzzo ^{a,b}, A. Pellecchia ^{a,b}, A. Pompili ^{a,b}, G. Pugliese ^{a,c}, R. Radogna ^a,
G. Ramirez-Sanchez ^{a,c}, D. Ramos ^a, A. Ranieri ^a, L. Silvestris ^a, F.M. Simone ^{a,b},
Ü. Sözbilir ^a, A. Stamerra ^a, R. Venditti ^a, P. Verwilligen ^a, A. Zaza ^{a,b}**INFN Sezione di Bologna^a, Università di Bologna^b, Bologna, Italy**G. Abbiendi ^a, C. Battilana ^{a,b}, L. Borgonovi ^a, R. Campanini ^{a,b}, P. Capiluppi ^{a,b},
A. Castro ^{a,b}, F.R. Cavallo ^a, M. Cuffiani ^{a,b}, G.M. Dallavalle ^a, T. Diotallevi ^{a,b}, F. Fabbri ^a,
A. Fanfani ^{a,b}, D. Fasanella ^{a,b}, P. Giacomelli ^a, L. Giommi ^{a,b}, C. Grandi ^a, L. Guiducci ^{a,b},
S. Lo Meo ^{a,48}, L. Lunerti ^{a,b}, S. Marcellini ^a, G. Masetti ^a, F.L. Navarra ^{a,b}, A. Perrotta ^a,
F. Primavera ^{a,b}, A.M. Rossi ^{a,b}, T. Rovelli ^{a,b}, G.P. Siroli ^{a,b}**INFN Sezione di Catania^a, Università di Catania^b, Catania, Italy**S. Costa ^{a,b,49}, A. Di Mattia ^a, R. Potenza ^{a,b}, A. Tricomi ^{a,b,49}, C. Tuve ^{a,b}**INFN Sezione di Firenze^a, Università di Firenze^b, Firenze, Italy**G. Barbagli ^a, G. Bardelli ^{a,b}, B. Camaiani ^{a,b}, A. Cassese ^a, R. Ceccarelli ^a, V. Ciulli ^{a,b},
C. Civinini ^a, R. D'Alessandro ^{a,b}, E. Focardi ^{a,b}, G. Latino ^{a,b}, P. Lenzi ^{a,b}, M. Lizzo ^{a,b},
M. Meschini ^a, S. Paoletti ^a, A. Papanastassiou ^{a,b}, G. Sguazzoni ^a, L. Viliani ^a**INFN Laboratori Nazionali di Frascati, Frascati, Italy**L. Benussi , S. Bianco , S. Meola ⁵⁰, D. Piccolo **INFN Sezione di Genova^a, Università di Genova^b, Genova, Italy**P. Chatagnon ^a, F. Ferro ^a, E. Robutti ^a, S. Tosi ^{a,b}**INFN Sezione di Milano-Bicocca^a, Università di Milano-Bicocca^b, Milano, Italy**A. Benaglia ^a, G. Boldrini ^a, F. Brivio ^a, F. Ceteorelli ^a, F. De Guio ^{a,b}, M.E. Dinardo ^{a,b},
P. Dini ^a, S. Gennai ^a, A. Ghezzi ^{a,b}, P. Govoni ^{a,b}, L. Guzzi ^a, M.T. Lucchini ^{a,b},
M. Malberti ^a, S. Malvezzi ^a, A. Massironi ^a, D. Menasce ^a, L. Moroni ^a, M. Paganoni ^{a,b},
D. Pedrini ^a, B.S. Pinolini ^a, S. Ragazzi ^{a,b}, N. Redaelli ^a, T. Tabarelli de Fatis ^{a,b}, D. Zuolo ^a












**INFN Sezione di Napoli^a, Università di Napoli ‘Federico II’^b, Napoli, Italy;
Università della Basilicata^c, Potenza, Italy; Università G. Marconi^d, Roma, Italy**

S. Buontempo ^a, A. Cagnotta ^{a,b}, F. Carnevali ^{a,b}, N. Cavallo ^{a,c}, A. De Iorio ^{a,b},
F. Fabozzi ^{a,c}, A.O.M. Iorio ^{a,b}, L. Lista ^{a,b,51}, P. Paolucci ^{a,30}, B. Rossi ^a, C. Sciacca ^{a,b}
















INFN Sezione di Padova^a, Università di Padova^b, Padova, Italy; Università di Trento^c, Trento, Italy

R. Ardino ^a, P. Azzi ^a, N. Bacchetta ^{a,52}, D. Bisello ^{a,b}, P. Bortignon ^a, A. Bragagnolo ^{a,b},
R. Carlin ^{a,b}, P. Checchia ^a, T. Dorigo ^a, F. Gasparini ^{a,b}, U. Gasparini ^{a,b}, G. Grosso ^a,
L. Layer ^{a,53}, E. Lusiani ^a, M. Margoni ^{a,b}, A.T. Meneguzzo ^{a,b}, M. Migliorini ^{a,b}, J. Pazzini ^{a,b},
P. Ronchese ^{a,b}, R. Rossin ^{a,b}, F. Simonetto ^{a,b}, G. Strong ^a, M. Tosi ^{a,b}, A. Triossi ^{a,b},
S. Ventura ^a, H. Yarar ^{a,b}, M. Zanetti ^{a,b}, P. Zotto ^{a,b}, A. Zucchetta ^{a,b}, G. Zumerle ^{a,b}
































INFN Sezione di Pavia^a, Università di Pavia^b, Pavia, Italy

S. Abu Zeid ^{a,54}, C. Aimè ^{a,b}, A. Braghieri ^a, S. Calzaferri ^{a,b}, D. Fiorina ^{a,b},
P. Montagna ^{a,b}, V. Re ^a, C. Riccardi ^{a,b}, P. Salvini ^a, I. Vai ^{a,b}, P. Vitulo ^{a,b}





















INFN Sezione di Perugia^a, Università di Perugia^b, Perugia, Italy

S. Ajmal ^{a,b}, P. Asenov ^{a,55}, G.M. Bilei ^a, D. Ciangottini ^{a,b}, L. Fanò ^{a,b}, M. Magherini ^{a,b},
G. Mantovani ^{a,b}, V. Mariani ^{a,b}, M. Menichelli ^a, F. Moscatelli ^{a,55}, A. Piccinelli ^{a,b},
M. Presilla ^{a,b}, A. Rossi ^{a,b}, A. Santocchia ^{a,b}, D. Spiga ^a, T. Tedeschi ^{a,b}






























**INFN Sezione di Pisa^a, Università di Pisa^b, Scuola Normale Superiore di Pisa^c,
Pisa, Italy; Università di Siena^d, Siena, Italy**












P. Azzurri ^a, G. Bagliesi ^a, R. Bhattacharya ^a, L. Bianchini ^{a,b}, T. Boccali ^a, E. Bossini ^a,
D. Bruschini ^{a,c}, R. Castaldi ^a, M.A. Ciocci ^{a,b}, M. Cipriani ^{a,b}, V. D’Amante ^{a,d},
R. Dell’Orso ^a, S. Donato ^a, A. Giassi ^a, F. Ligabue ^{a,c}, D. Matos Figueiredo ^a,
A. Messineo ^{a,b}, M. Musich ^{a,b}, F. Palla ^a, S. Parolia ^a, A. Rizzi ^{a,b}, G. Rolandi ^{a,c},
S. Roy Chowdhury ^a, T. Sarkar ^a, A. Scribano ^a, P. Spagnolo ^a, R. Tenchini ^{a,b},
G. Tonelli ^{a,b}, N. Turini ^{a,d}, A. Venturi ^a, P.G. Verdini ^a

INFN Sezione di Roma^a, Sapienza Università di Roma^b, Roma, Italy







P. Barria ^a, M. Campana ^{a,b}, F. Cavallari ^a, L. Cunqueiro Mendez ^{a,b}, D. Del Re ^{a,b},
E. Di Marco ^a, M. Diemoz ^a, F. Errico ^{a,b}, E. Longo ^{a,b}, P. Meridiani ^a, J. Mijuskovic ^{a,b},
G. Organtini ^{a,b}, F. Pandolfi ^a, R. Paramatti ^{a,b}, C. Quaranta ^{a,b}, S. Rahatlou ^{a,b},
C. Rovelli ^a, F. Santanastasio ^{a,b}, L. Soffi ^a, R. Tramontano ^{a,b}

**INFN Sezione di Torino^a, Università di Torino^b, Torino, Italy; Università del
Piemonte Orientale^c, Novara, Italy**













N. Amapane ^{a,b}, R. Arcidiacono ^{a,c}, S. Argiro ^{a,b}, M. Arneodo ^{a,c}, N. Bartosik ^a,
R. Bellan ^{a,b}, A. Bellora ^{a,b}, C. Biino ^a, N. Cartiglia ^a, M. Costa ^{a,b}, R. Covarelli ^{a,b},
N. Demaria ^a, L. Finco ^a, M. Grippo ^{a,b}, B. Kiani ^{a,b}, F. Legger ^a, F. Luongo ^{a,b},
C. Mariotti ^a, S. Maselli ^a, A. Mecca ^{a,b}, E. Migliore ^{a,b}, M. Monteno ^a, R. Mulargia ^a,
M.M. Obertino ^{a,b}, G. Ortona ^a, L. Pacher ^{a,b}, N. Pastrone ^a, M. Pelliccioni ^a, M. Ruspa ^{a,c},

F. Siviero ^{a,b}, V. Sola ^{a,b}, A. Solano ^{a,b}, D. Soldi ^{a,b}, A. Staiano ^a, C. Tarricone ^{a,b},
M. Tornago ^{a,b}, D. Trocino ^a, G. Umoret ^{a,b}, A. Vagnerini ^{a,b}, E. Vlasov ^{a,b}


INFN Sezione di Trieste^a, Università di Trieste^b, Trieste, Italy

S. Belforte ^a, V. Candelise ^{a,b}, M. Casarsa ^a, F. Cossutti ^a, K. De Leo ^{a,b}, G. Della Ricca ^{a,b}

Kyungpook National University, Daegu, Korea

S. Dogra ^b, J. Hong ^b, C. Huh ^b, B. Kim ^b, D.H. Kim ^b, J. Kim, H. Lee, S.W. Lee ^b, C.S. Moon ^b,
Y.D. Oh ^b, S.I. Pak ^b, M.S. Ryu ^b, S. Sekmen ^b, Y.C. Yang ^b





Chonnam National University, Institute for Universe and Elementary Particles, Kwangju, Korea

G. Bak ^b, P. Gwak ^b, H. Kim ^b, D.H. Moon ^b

Hanyang University, Seoul, Korea

E. Asilar ^b, D. Kim ^b, T.J. Kim ^b, J.A. Merlin, J. Park ^b, J. Song


Korea University, Seoul, Korea

S. Choi ^b, S. Han, B. Hong ^b, K. Lee, K.S. Lee ^b, J. Park, S.K. Park, J. Yoo ^b














Kyung Hee University, Department of Physics, Seoul, Korea

J. Goh ^b









Sejong University, Seoul, Korea

H. S. Kim ^b, Y. Kim, S. Lee


Seoul National University, Seoul, Korea

J. Almond, J.H. Bhyun, J. Choi ^b, S. Jeon ^b, W. Jun ^b, J. Kim ^b, J.S. Kim, S. Ko ^b, H. Kwon ^b,
H. Lee ^b, J. Lee ^b, J. Lee ^b, S. Lee, B.H. Oh ^b, S.B. Oh ^b, H. Seo ^b, U.K. Yang, I. Yoon ^b

University of Seoul, Seoul, Korea

W. Jang ^b, D.Y. Kang, Y. Kang ^b, S. Kim ^b, B. Ko, J.S.H. Lee ^b, Y. Lee ^b, I.C. Park ^b, Y. Roh,
I.J. Watson ^b, S. Yang ^b


Yonsei University, Department of Physics, Seoul, Korea

S. Ha ^b, H.D. Yoo ^b

Sungkyunkwan University, Suwon, Korea
















M. Choi ^b, M.R. Kim ^b, H. Lee, Y. Lee ^b, I. Yu ^b

College of Engineering and Technology, American University of the Middle East (AUM), Dasman, Kuwait














T. Beyrouthy, Y. Maghrbi ^b

Riga Technical University, Riga, Latvia

K. Dreimanis ^b, A. Gaile ^b, G. Pikurs, A. Potrebko ^b, M. Seidel ^b, V. Veckalns ^b⁵⁶

University of Latvia (LU), Riga, LatviaN.R. Strautnieks **Vilnius University, Vilnius, Lithuania**M. Ambrozias , A. Juodagalvis , A. Rinkevicius , G. Tamulaitis **National Centre for Particle Physics, Universiti Malaya, Kuala Lumpur, Malaysia**N. Bin Norjoharuddeen , I. Yusuff ⁵⁷, Z. Zolkapli**Universidad de Sonora (UNISON), Hermosillo, Mexico**J.F. Benitez , A. Castaneda Hernandez , H.A. Encinas Acosta, L.G. Gallegos Maríñez, M. León Coello , J.A. Murillo Quijada , A. Sehrawat , L. Valencia Palomo **Centro de Investigacion y de Estudios Avanzados del IPN, Mexico City, Mexico**G. Ayala , H. Castilla-Valdez , E. De La Cruz-Burelo , I. Heredia-De La Cruz ⁵⁸, R. Lopez-Fernandez , C.A. Mondragon Herrera, D.A. Perez Navarro , A. Sánchez Hernández **Universidad Iberoamericana, Mexico City, Mexico**C. Oropeza Barrera , M. Ramírez García **Benemerita Universidad Autonoma de Puebla, Puebla, Mexico**I. Bautista , I. Pedraza , H.A. Salazar Ibarguen , C. Uribe Estrada **University of Montenegro, Podgorica, Montenegro**I. Bubanja, N. Raicevic **University of Canterbury, Christchurch, New Zealand**P.H. Butler **National Centre for Physics, Quaid-I-Azam University, Islamabad, Pakistan**A. Ahmad , M.I. Asghar, A. Awais , M.I.M. Awan, H.R. Hoorani , W.A. Khan **AGH University of Krakow, Faculty of Computer Science, Electronics and Telecommunications, Krakow, Poland**V. Avati, L. Grzanka , M. Malawski **National Centre for Nuclear Research, Swierk, Poland**H. Bialkowska , M. Bluj , B. Boimska , M. Górski , M. Kazana , M. Szeleper , P. Zalewski **Institute of Experimental Physics, Faculty of Physics, University of Warsaw, Warsaw, Poland**K. Bunkowski , K. Doroba , A. Kalinowski , M. Konecki , J. Krolikowski , A. Muhammad **Warsaw University of Technology, Warsaw, Poland**K. Pozniak , W. Zabolotny 

Laboratório de Instrumentação e Física Experimental de Partículas, Lisboa, Portugal

M. Araujo , D. Bastos , C. Beirão Da Cruz E Silva , A. Boletti , M. Bozzo , P. Faccioli , M. Gallinaro , J. Hollar , N. Leonardo , T. Niknejad , M. Pisano , J. Seixas , J. Varela 




























Faculty of Physics, University of Belgrade, Belgrade, Serbia

P. Adzic , P. Milenovic 

VINCA Institute of Nuclear Sciences, University of Belgrade, Belgrade, Serbia

M. Dordevic , J. Milosevic , V. Rekovic














Centro de Investigaciones Energéticas Medioambientales y Tecnológicas (CIEMAT), Madrid, Spain

M. Aguilar-Benitez, J. Alcaraz Maestre , M. Barrio Luna, Cristina F. Bedoya , M. Cepeda , M. Cerrada , N. Colino , B. De La Cruz , A. Delgado Peris , D. Fernández Del Val , J.P. Fernández Ramos , J. Flix , M.C. Fouz , O. Gonzalez Lopez , S. Goy Lopez , J.M. Hernandez , M.I. Josa , J. León Holgado , D. Moran , C. M. Morcillo Perez , Á. Navarro Tobar , C. Perez Dengra , A. Pérez-Calero Yzquierdo , J. Puerta Pelayo , I. Redondo , D.D. Redondo Ferrero , L. Romero, S. Sánchez Navas , L. Urda Gómez , J. Vazquez Escobar , C. Willmott




















Universidad Autónoma de Madrid, Madrid, Spain

J.F. de Trocóniz 

Universidad de Oviedo, Instituto Universitario de Ciencias y Tecnologías Espaciales de Asturias (ICTEA), Oviedo, Spain

B. Alvarez Gonzalez , J. Cuevas , J. Fernandez Menendez , S. Folgueras , I. Gonzalez Caballero , J.R. González Fernández , E. Palencia Cortezon , C. Ramón Álvarez , V. Rodríguez Bouza , A. Soto Rodríguez , A. Trapote , C. Vico Villalba , P. Vischia 

Instituto de Física de Cantabria (IFCA), CSIC-Universidad de Cantabria, Santander, Spain

S. Bhowmik , S. Blanco Fernández , J.A. Brochero Cifuentes , I.J. Cabrillo , A. Calderon , J. Duarte Campderros , M. Fernandez , C. Fernandez Madrazo , G. Gomez , C. Lasiosa García , C. Martinez Rivero , P. Martinez Ruiz del Arbol , F. Matorras , P. Matorras Cuevas , E. Navarrete Ramos , J. Piedra Gomez , C. Prieels, L. Scodellaro , I. Vila , J.M. Vizan Garcia 


University of Colombo, Colombo, Sri Lanka

M.K. Jayananda , B. Kailasapathy ⁵⁹, D.U.J. Sonnadara , D.D.C. Wickramarathna 

University of Ruhuna, Department of Physics, Matara, Sri Lanka













W.G.D. Dharmaratna ⁶⁰, K. Liyanage , N. Perera , N. Wickramage 

CERN, European Organization for Nuclear Research, Geneva, Switzerland

D. Abbaneo , C. Amendola , E. Auffray , G. Auzinger , J. Baechler, D. Barney , A. Bermúdez Martínez , M. Bianco , B. Bilin , A.A. Bin Anuar , A. Bocci , E. Brondolin 

C. Caillol , T. Camporesi , G. Cerminara , N. Chernyavskaya , D. d'Enterria ,
 A. Dabrowski , A. David , A. De Roeck , M.M. Defranchis , M. Deile , M. Dobson ,
 F. Fallavollita ⁶¹, L. Forthomme , G. Franzoni , W. Funk , S. Giani, D. Gigi, K. Gill ,
 F. Glege , L. Gouskos , M. Haranko , J. Hegeman , V. Innocente , T. James , P. Janot ,
 J. Kieseler , S. Laurila , P. Lecoq , E. Leutgeb , C. Lourenço , B. Maier , L. Malgeri ,
 M. Mannelli , A.C. Marini , F. Meijers , S. Mersi , E. Meschi , V. Milosevic , F. Moortgat ,
 M. Mulders , S. Orfanelli, F. Pantaleo , M. Peruzzi , A. Petrilli , G. Petrucciani ,
 A. Pfeiffer , M. Pierini , D. Piparo , H. Qu , D. Rabady , G. Reales Gutiérrez, M. Rovere ,
 H. Sakulin , S. Scarfi , M. Selvaggi , A. Sharma , K. Shchelina , P. Silva , P. Sphicas ⁶²,
 A.G. Stahl Leitner , A. Steen , S. Summers , D. Treille , P. Tropea , A. Tsiros, D. Walter ,
 J. Wanczyk ⁶³, K.A. Wozniak ⁶⁴, P. Zehetner , P. Zejdl , W.D. Zeuner
























Paul Scherrer Institut, Villigen, Switzerland

T. Bevilacqua ⁶⁵, L. Caminada ⁶⁵, A. Ebrahimi , W. Erdmann , R. Horisberger , Q. Ingram ,
 H.C. Kaestli , D. Kotlinski , C. Lange , M. Missiroli ⁶⁵, L. Noehte ⁶⁵, T. Rohe 




ETH Zurich — Institute for Particle Physics and Astrophysics (IPA), Zurich, Switzerland

T.K. Aarrestad , K. Androsov ⁶³, M. Backhaus , A. Calandri , C. Cazzaniga , K. Datta ,
 A. De Cosa , G. Dissertori , M. Dittmar, M. Donegà , F. Eble , M. Galli , K. Gedia ,
 F. Glessgen , C. Grab , D. Hits , W. Lustermann , A.-M. Lyon , R.A. Manzoni ,
 M. Marchegiani , L. Marchese , C. Martin Perez , A. Mascellani ⁶³, F. Nessi-Tedaldi ,
 F. Pauss , V. Perovic , S. Pigazzini , M.G. Ratti , M. Reichmann , C. Reissel ,
 T. Reitenspiess , B. Ristic , F. Riti , D. Ruini, D.A. Sanz Becerra , R. Seidita ,
 J. Steggemann ⁶³, D. Valsecchi , R. Wallny 










Universität Zürich, Zurich, Switzerland

C. Amsler ⁶⁶, P. Bärtschi , C. Botta , D. Brzhechko, M.F. Canelli , K. Cormier , A. De Wit ,
 R. Del Burgo, J.K. Heikkilä , M. Huwiler , W. Jin , A. Jofrehei , B. Kilminster ,
 S. Leontsinis , S.P. Liechti , A. Macchiolo , P. Meiring , V.M. Mikuni , U. Molinatti ,
 I. Neutelings , A. Reimers , P. Robmann, S. Sanchez Cruz , K. Schweiger , M. Senger ,
 Y. Takahashi 

National Central University, Chung-Li, Taiwan

C. Adloff ⁶⁷, C.M. Kuo, W. Lin, P.K. Rout , P.C. Tiwari ⁴¹, S.S. Yu 



















National Taiwan University (NTU), Taipei, Taiwan

L. Ceard, Y. Chao , K.F. Chen , P.s. Chen, Z.g. Chen, W.-S. Hou , T.h. Hsu, Y.w. Kao,
 R. Khurana, G. Kole , Y.y. Li , R.-S. Lu , E. Paganis , A. Psallidas, X.f. Su,
 J. Thomas-Wilsker , H.y. Wu, E. Yazgan 



High Energy Physics Research Unit, Department of Physics, Faculty of Science, Chulalongkorn University, Bangkok, Thailand

C. Asawatrangkuldee , N. Srimanobhas , V. Wachirapusanand 

Çukurova University, Physics Department, Science and Art Faculty, Adana, Turkey

D. Agyel , F. Boran , Z.S. Demiroglu , F. Dolek , I. Dumanoglu ⁶⁸, E. Eskut , Y. Guler ⁶⁹, E. Gurpinar Guler ⁶⁹, C. Isik , O. Kara, A. Kayis Topaksu , U. Kiminsu , G. Onengut , K. Ozdemir ⁷⁰, A. Polatoz , B. Tali ⁷¹, U.G. Tok , S. Turkcapar , E. Uslan , I.S. Zorbakir 

Middle East Technical University, Physics Department, Ankara, Turkey

K. Ocalan ⁷², M. Yalvac ⁷³














Bogazici University, Istanbul, Turkey

B. Akgun , I.O. Atakisi , E. Gülmez , M. Kaya ⁷⁴, O. Kaya ⁷⁵, S. Tekten ⁷⁶

Istanbul Technical University, Istanbul, Turkey

A. Cakir , K. Cankocak ⁶⁸, Y. Komurcu , S. Sen ⁷⁷

Istanbul University, Istanbul, Turkey

O. Aydilek , S. Cerci ⁷¹, V. Epshteyn , B. Hacisahinoglu , I. Hos ⁷⁸, B. Isildak ⁷⁹, B. Kaynak , S. Ozkorucuklu , O. Potok , H. Sert , C. Simsek , D. Sunar Cerci ⁷¹, C. Zorbilmez 
















Institute for Scintillation Materials of National Academy of Science of Ukraine, Kharkiv, Ukraine

A. Boyaryntsev , B. Grynyov 



















National Science Centre, Kharkiv Institute of Physics and Technology, Kharkiv, Ukraine

L. Levchuk 

























University of Bristol, Bristol, United Kingdom








D. Anthony , J.J. Brooke , A. Bundock , F. Bury , E. Clement , D. Cussans , H. Flacher , M. Glowacki, J. Goldstein , H.F. Heath , L. Kreczko , B. Krikler , S. Paramesvaran , S. Seif El Nasr-Storey, V.J. Smith , N. Stylianou ⁸⁰, K. Walkingshaw Pass, R. White 

Rutherford Appleton Laboratory, Didcot, United Kingdom

A.H. Ball, K.W. Bell , A. Belyaev ⁸¹, C. Brew , R.M. Brown , D.J.A. Cockerill , C. Cooke , K.V. Ellis, K. Harder , S. Harper , M.-L. Holmberg ⁸², Sh. Jain , J. Linacre , K. Manolopoulos, D.M. Newbold , E. Olaiya, D. Petyt , T. Reis , G. Salvi , T. Schuh, C.H. Shepherd-Themistocleous , I.R. Tomalin , T. Williams 

Imperial College, London, United Kingdom














R. Bainbridge , P. Bloch , C.E. Brown , O. Buchmuller, V. Cacchio, C.A. Carrillo Montoya , G.S. Chahal ⁸³, D. Colling , J.S. Dancu, P. Dauncey , G. Davies , J. Davies, M. Della Negra , S. Fayer, G. Fedi , G. Hall , M.H. Hassanshahi , A. Howard, G. Iles , M. Knight , J. Langford , L. Lyons , A.-M. Magnan , S. Malik, A. Martelli , M. Mieskolainen , J. Nash ⁸⁴, M. Pesaresi, B.C. Radburn-Smith , A. Richards, A. Rose , C. Seez , R. Shukla 

A. Tapper , K. Uchida , G.P. Uttley , L.H. Vage, T. Virdee ³⁰, M. Vojinovic , N. Wardle ,
D. Winterbottom 






Brunel University, Uxbridge, United Kingdom

K. Coldham, J.E. Cole , A. Khan, P. Kyberd , I.D. Reid 

Baylor University, Waco, Texas, U.S.A.

S. Abdullin , A. Brinkerhoff , B. Caraway , J. Dittmann , K. Hatakeyama , J. Hiltbrand ,
A.R. Kanuganti , B. McMaster , M. Saunders , S. Sawant , C. Sutantawibul , M. Toms ⁸⁵,
J. Wilson 
















Catholic University of America, Washington, DC, U.S.A.

R. Bartek , A. Dominguez , C. Huerta Escamilla, A.E. Simsek , R. Uniyal ,
A.M. Vargas Hernandez 

















The University of Alabama, Tuscaloosa, Alabama, U.S.A.

R. Chudasama , S.I. Cooper , S.V. Gleyzer , C.U. Perez , P. Rumerio ⁸⁶, E. Usai ,
C. West , R. Yi 





















Boston University, Boston, Massachusetts, U.S.A.

A. Akpinar , A. Albert , D. Arcaro , C. Cosby , Z. Demiragli , C. Erice , E. Fontanesi ,
D. Gastler , J. Rohlf , K. Salyer , D. Sperka , D. Spitzbart , I. Suarez , A. Tsatsos ,
S. Yuan 













Brown University, Providence, Rhode Island, U.S.A.

G. Benelli , X. Coubez²⁵, D. Cutts , M. Hadley , U. Heintz , J.M. Hogan ⁸⁷, T. Kwon ,
G. Landsberg , K.T. Lau , D. Li , J. Luo , S. Mondal , M. Narain [†], N. Pervan ,
S. Sagir ⁸⁸, F. Simpson , W.Y. Wong, X. Yan , W. Zhang

University of California, Davis, Davis, California, U.S.A.

S. Abbott , J. Bonilla , C. Brainerd , R. Breedon , M. Calderon De La Barca Sanchez ,
M. Chertok , M. Citron , J. Conway , P.T. Cox , R. Erbacher , G. Haza , F. Jensen ,
O. Kukral , G. Mocellin , M. Mulhearn , D. Pellett , B. Regnery , W. Wei , Y. Yao ,
F. Zhang 

University of California, Los Angeles, California, U.S.A.

M. Bachtis , R. Cousins , A. Datta , J. Hauser , M. Ignatenko , M.A. Iqbal , T. Lam ,
E. Manca , W.A. Nash , D. Saltzberg , B. Stone , V. Valuev 

University of California, Riverside, Riverside, California, U.S.A.



















R. Clare , M. Gordon, G. Hanson , W. Si , S. Wimpenny [†]

University of California, San Diego, La Jolla, California, U.S.A.









J.G. Branson , S. Cittolin , S. Cooperstein , D. Diaz , J. Duarte , R. Gerosa , L. Giannini ,
J. Guiang , R. Kansal , V. Krutelyov , R. Lee , J. Letts , M. Masciovecchio , F. Mokhtar 

M. Pieri , M. Quinnan , B.V. Sathia Narayanan , V. Sharma , M. Tadel , E. Vourliotis ,
F. Würthwein , Y. Xiang , A. Yagil 

University of California, Santa Barbara — Department of Physics, Santa Barbara, California, U.S.A.

L. Brennan , C. Campagnari , G. Collura , A. Dorsett , J. Incandela , M. Kilpatrick ,
J. Kim , A.J. Li , P. Masterson , H. Mei , M. Oshiro , J. Richman , U. Sarica ,
R. Schmitz , F. Setti , J. Sheplock , D. Stuart , S. Wang 














California Institute of Technology, Pasadena, California, U.S.A.

A. Bornheim , O. Cerri, A. Latorre, J.M. Lawhorn , J. Mao , H.B. Newman , T. Q. Nguyen ,
M. Spiropulu , J.R. Vlimant , C. Wang , S. Xie , R.Y. Zhu 






Carnegie Mellon University, Pittsburgh, Pennsylvania, U.S.A.

J. Alison , S. An , M.B. Andrews , P. Bryant , V. Dutta , T. Ferguson , A. Harilal ,
C. Liu , T. Mudholkar , S. Murthy , M. Paulini , A. Roberts , A. Sanchez , W. Terrill 









































































University of Colorado Boulder, Boulder, Colorado, U.S.A.

J.P. Cumalat , W.T. Ford , A. Hassani , G. Karathanasis , E. MacDonald, N. Manganelli ,
F. Marini , A. Perloff , C. Savard , N. Schonbeck , K. Stenson , K.A. Ulmer ,
S.R. Wagner , N. Zipper 




















Cornell University, Ithaca, New York, U.S.A.

J. Alexander , S. Bright-Thonney , X. Chen , D.J. Cranshaw , J. Fan , X. Fan ,
D. Gadkari , S. Hogan , J. Monroy , J.R. Patterson , J. Reichert , M. Reid , A. Ryd ,
J. Thom , P. Wittich , R. Zou 















Fermi National Accelerator Laboratory, Batavia, Illinois, U.S.A.

M. Albrow , M. Alyari , O. Amram , G. Apollinari , A. Apresyan , L.A.T. Bauerdick ,
D. Berry , J. Berryhill , P.C. Bhat , K. Burkett , J.N. Butler , A. Canepa , G.B. Cerati ,
H.W.K. Cheung , F. Chlebana , G. Cummings , J. Dickinson , I. Dutta , V.D. Elvira ,
Y. Feng , J. Freeman , A. Gandrakota , Z. Gecse , L. Gray , D. Green, S. Grünendahl ,
D. Guerrero , O. Gutsche , R.M. Harris , R. Heller , T.C. Herwig , J. Hirschauer ,
L. Horyn , B. Jayatilaka , S. Jindariani , M. Johnson , U. Joshi , T. Klijnsma , B. Klima ,
K.H.M. Kwok , S. Lammel , D. Lincoln , R. Lipton , T. Liu , C. Madrid , K. Maeshima ,
C. Mantilla , D. Mason , P. McBride , P. Merkel , S. Mrenna , S. Nahn , J. Ngadiuba ,
D. Noonan , V. Papadimitriou , N. Pastika , K. Pedro , C. Pena ⁸⁹, F. Ravera ,
A. Reinsvold Hall ⁹⁰, L. Ristori , E. Sexton-Kennedy , N. Smith , A. Soha , L. Spiegel ,
S. Stoynev , J. Strait , L. Taylor , S. Tkaczyk , N.V. Tran , L. Uplegger ,
E.W. Vaandering , I. Zoi 







University of Florida, Gainesville, Florida, U.S.A.

C. Aruta , P. Avery , D. Bourilkov , L. Cadamuro , P. Chang , V. Cherepanov , R.D. Field,
E. Koenig , M. Kolosova , J. Konigsberg , A. Korytov , K.H. Lo, K. Matchev ,
N. Menendez , G. Mitselmakher , A. Muthirakalayil Madhu , N. Rawal , D. Rosenzweig ,
S. Rosenzweig , K. Shi , J. Wang 























Florida State University, Tallahassee, Florida, U.S.A.

T. Adams , A. Al Kadhim , A. Askew , N. Bower , R. Habibullah , V. Hagopian ,
R. Hashmi , R.S. Kim , S. Kim , T. Kolberg , G. Martinez, H. Prosper , P.R. Prova,
O. Viazlo , M. Wulansatiti , R. Yohay , J. Zhang












Florida Institute of Technology, Melbourne, Florida, U.S.A.

B. Alsufyani, M.M. Baarmand , S. Butalla , T. Elkafrawy ⁵⁴, M. Hohlmann ,
R. Kumar Verma , M. Rahmani, F. Yumiceva 












University of Illinois Chicago, Chicago, U.S.A., Chicago, U.S.A.

M.R. Adams , C. Bennett, R. Cavanaugh , S. Dittmer , R. Escobar Franco , O. Evdokimov ,
C.E. Gerber , D.J. Hofman , J.h. Lee , D. S. Lemos , A.H. Merrit , C. Mills , S. Nanda ,
G. Oh , B. Ozek , D. Pilipovic , T. Roy , S. Rudrabhatla , M.B. Tonjes , N. Varelas ,
X. Wang , Z. Ye , J. Yoo 
























The University of Iowa, Iowa City, Iowa, U.S.A.

M. Alhousseini , D. Blend, K. Dilsiz ⁹¹, L. Emediato , G. Karaman , O.K. Köseyan ,
J.-P. Merlo, A. Mestvirishvili ⁹², J. Nachtman , O. Neogi, H. Ogul ⁹³, Y. Onel , A. Penzo ,
C. Snyder, E. Tiras ⁹⁴

Johns Hopkins University, Baltimore, Maryland, U.S.A.

B. Blumenfeld , L. Corcodilos , J. Davis , A.V. Gritsan , L. Kang , S. Kyriacou ,
P. Maksimovic , M. Roguljic , J. Roskes , S. Sekhar , M. Swartz , T.Á. Vámi 

The University of Kansas, Lawrence, Kansas, U.S.A.

A. Abreu , L.F. Alcerro Alcerro , J. Anguiano , P. Baringer , A. Bean , Z. Flowers ,
D. Grove, J. King , G. Krintiras , M. Lazarovits , C. Le Mahieu , C. Lindsey, J. Marquez ,
N. Minafra , M. Murray , M. Nickel , M. Pitt , S. Popescu ⁹⁵, C. Rogan , C. Royon ,
R. Salvatico , S. Sanders , C. Smith , Q. Wang , G. Wilson 



















Kansas State University, Manhattan, Kansas, U.S.A.

B. Allmond , A. Ivanov , K. Kaadze , A. Kalogeropoulos , D. Kim, Y. Maravin , K. Nam,
J. Natoli , D. Roy , G. Sorrentino 












Lawrence Livermore National Laboratory, Livermore, California, U.S.A.














F. Rebassoo , D. Wright 

University of Maryland, College Park, Maryland, U.S.A.













E. Adams , A. Baden , O. Baron, A. Belloni , A. Bethani , Y.M. Chen , S.C. Eno ,
N.J. Hadley , S. Jabeen , R.G. Kellogg , T. Koeth , Y. Lai , S. Lascio , A.C. Mignerey ,
S. Nabili , C. Palmer , C. Papageorgakis , M.M. Paranjpe, L. Wang , K. Wong 

Massachusetts Institute of Technology, Cambridge, Massachusetts, U.S.A.

J. Bendavid , W. Busza , I.A. Cali , Y. Chen , M. D'Alfonso , J. Eysermans , C. Freer ,
G. Gomez-Ceballos , M. Goncharov, P. Harris, D. Hoang, D. Kovalskyi , J. Krupa , L. Lavezzo 

Y.-J. Lee , K. Long , C. Mironov , C. Paus , D. Rankin , C. Roland , G. Roland ,
S. Rothman , Z. Shi , G.S.F. Stephans , J. Wang, Z. Wang , B. Wyslouch , T. J. Yang 









University of Minnesota, Minneapolis, Minnesota, U.S.A.

B. Crossman , B.M. Joshi , C. Kapsiak , M. Krohn , D. Mahon , J. Mans , B. Marzocchi ,
S. Pandey , M. Revering , R. Rusack , R. Saradhy , N. Schroeder , N. Strobbe ,
M.A. Wadud 












University of Mississippi, Oxford, Mississippi, U.S.A.

L.M. Cremaldi 

















University of Nebraska-Lincoln, Lincoln, Nebraska, U.S.A.

K. Bloom , M. Bryson, D.R. Claes , C. Fangmeier , F. Golf , J. Hossain , C. Joo ,
I. Kravchenko , I. Reed , J.E. Siado , G.R. Snow[†], W. Tabb , A. Wightman , F. Yan ,
D. Yu , A.G. Zecchinelli 









State University of New York at Buffalo, Buffalo, New York, U.S.A.

G. Agarwal , H. Bandyopadhyay , L. Hay , I. Iashvili , A. Kharchilava , C. McLean ,
M. Morris , D. Nguyen , J. Pekkanen , S. Rappoccio , H. Rejeb Sfar, A. Williams 

























Northeastern University, Boston, Massachusetts, U.S.A.

G. Alverson , E. Barberis , Y. Haddad , Y. Han , A. Krishna , J. Li , M. Lu ,
G. Madigan , D.M. Morse , V. Nguyen , T. Orimoto , A. Parker , L. Skinnari ,
A. Tishelman-Charny , B. Wang , D. Wood 










Northwestern University, Evanston, Illinois, U.S.A.

S. Bhattacharya , J. Bueghly, Z. Chen , K.A. Hahn , Y. Liu , Y. Miao , D.G. Monk ,
M.H. Schmitt , A. Taliercio , M. Velasco



















University of Notre Dame, Notre Dame, Indiana, U.S.A.

R. Band , R. Bucci, S. Castells , M. Cremonesi, A. Das , R. Goldouzian , M. Hildreth ,
K.W. Ho , K. Hurtado Anampa , C. Jessop , K. Lannon , J. Lawrence , N. Loukas ,
L. Lutton , J. Mariano, N. Marinelli, I. Mcalister, T. McCauley , C. Mcgrady , K. Mohrman ,
C. Moore , Y. Musienko ¹³, H. Nelson , M. Osherson , R. Ruchti , A. Townsend ,
M. Wayne , H. Yockey, M. Zarucki , L. Zygala 

The Ohio State University, Columbus, Ohio, U.S.A.

A. Basnet , B. Bylsma, M. Carrigan , L.S. Durkin , C. Hill , M. Joyce , A. Lesauvage ,
M. Nunez Ornelas , K. Wei, B.L. Winer , B. R. Yates 






















Princeton University, Princeton, New Jersey, U.S.A.

F.M. Addesa , H. Bouchamaoui , P. Das , G. Dezoort , P. Elmer , A. Frankenthal ,
B. Greenberg , N. Haubrich , S. Higginbotham , G. Kopp , S. Kwan , D. Lange ,
A. Loeliger , D. Marlow , I. Ojalvo , J. Olsen , D. Stickland , C. Tully 

University of Puerto Rico, Mayaguez, Puerto Rico, U.S.A.

S. Malik 









Purdue University, West Lafayette, Indiana, U.S.A.

A.S. Bakshi , V.E. Barnes , S. Chandra , R. Chawla , S. Das , A. Gu , L. Gutay, M. Jones , A.W. Jung , D. Kondratyev , A.M. Koshy, M. Liu , G. Negro , N. Neumeister , G. Paspalaki , S. Piperov , A. Purohit , J.F. Schulte , M. Stojanovic , J. Thieman , A. K. Viridi , F. Wang , W. Xie 

Purdue University Northwest, Hammond, Indiana, U.S.A.

J. Dolen , N. Parashar , A. Pathak 


Rice University, Houston, Texas, U.S.A.

D. Acosta , A. Baty , T. Carnahan , S. Dildick , K.M. Ecklund , P.J. Fernández Manteca , S. Freed, P. Gardner, F.J.M. Geurts , A. Kumar , W. Li , O. Miguel Colin , B.P. Padley , R. Redjimi, J. Rotter , E. Yigitbasi , Y. Zhang 

















University of Rochester, Rochester, New York, U.S.A.

A. Bodek , P. de Barbaro , R. Demina , J.L. Dulemba , C. Fallon, A. Garcia-Bellido , O. Hindrichs , A. Khukhunaishvili , P. Parygin ⁸⁵, E. Popova ⁸⁵, R. Taus , G.P. Van Onsem 

The Rockefeller University, New York, New York, U.S.A.

K. Goulianos 















Rutgers, The State University of New Jersey, Piscataway, New Jersey, U.S.A.

B. Chiarito, J.P. Chou , Y. Gershtein , E. Halkiadakis , A. Hart , M. Heindl , D. Jaroslowski , O. Karacheban ²⁸, I. Laflotte , A. Lath , R. Montalvo, K. Nash, H. Routray , S. Salur , S. Schnetzer, S. Somalwar , R. Stone , S.A. Thayil , S. Thomas, J. Vora , H. Wang 












University of Tennessee, Knoxville, Tennessee, U.S.A.

H. Acharya, D. Ally , A.G. Delannoy , S. Fiorendi , T. Holmes , N. Karunarathna , L. Lee , E. Nibigira , S. Spanier 

Texas A&M University, College Station, Texas, U.S.A.

D. Aebi , M. Ahmad , O. Bouhali ⁹⁶, M. Dalchenko , R. Eusebi , J. Gilmore , T. Huang , T. Kamon ⁹⁷, H. Kim , S. Luo , S. Malhotra, R. Mueller , D. Overton , D. Rathjens , A. Safonov 









Texas Tech University, Lubbock, Texas, U.S.A.

N. Akchurin , J. Damgov , V. Hegde , A. Hussain , Y. Kazhykarim, K. Lamichhane , S.W. Lee , A. Mankel , T. Mengke, S. Muthumuni , T. Peltola , I. Volobouev , A. Whitbeck 






















Vanderbilt University, Nashville, Tennessee, U.S.A.

E. Appelt , S. Greene, A. Gurrola , W. Johns , R. Kunnawalkam Elayavalli , A. Melo , F. Romeo , P. Sheldon , S. Tuo , J. Velkovska , J. Viinikainen 

University of Virginia, Charlottesville, Virginia, U.S.A.

B. Cardwell , B. Cox , J. Hakala , R. Hirosky , A. Ledovskoy , A. Li , C. Neu , C.E. Perez Lara 

Wayne State University, Detroit, Michigan, U.S.A.P.E. Karchin **University of Wisconsin — Madison, Madison, Wisconsin, U.S.A.**

A. Aravind, S. Banerjee , K. Black , T. Bose , S. Dasu , I. De Bruyn , P. Everaerts ,
 C. Galloni, H. He , M. Herndon , A. Herve , C.K. Koraka , A. Lanaro, R. Loveless ,
 J. Madhusudanan Sreekala , A. Mallampalli , A. Mohammadi , S. Mondal, G. Parida ,
 D. Pinna, A. Savin, V. Shang , V. Sharma , W.H. Smith , D. Teague, H.F. Tsoi , W. Vetens ,
 A. Warden 

Authors affiliated with an institute or an international laboratory covered by a cooperation agreement with CERN

S. Afanasiev , V. Andreev , Yu. Andreev , T. Aushev , M. Azarkin , A. Babaev ,
 A. Belyaev , V. Blinov⁹⁸, E. Boos , V. Borshch , D. Budkouski , V. Bunichev ,
 M. Chadeeva , V. Chekhovsky, A. Dermenev , T. Dimova , D. Druzhkin , M. Dubinin ,⁸⁹
 L. Dudko , A. Ershov , G. Gavrilo , V. Gavrilo , S. Gninenko , V. Golovtsov ,
 N. Golubev , I. Golutvin , I. Gorbunov , A. Gribushin , Y. Ivanov , V. Kachanov ,
 L. Kardapoltsev ,⁹⁸ V. Karjavine , A. Karneyeu , V. Kim ,⁹⁸ M. Kirakosyan, D. Kirpichnikov ,
 M. Kirsanov , V. Klyukhin , D. Konstantinov , V. Korenkov , A. Kozyrev ,⁹⁸ N. Krasnikov ,
 A. Lanev , P. Levchenko ,¹⁰⁰ N. Lychkovskaya , V. Makarenko , A. Malakhov ,
 V. Matveev ,⁹⁸ V. Murzin , A. Nikitenko ,^{101,102} S. Obraztsov , V. Oreshkin , A. Oskin,
 V. Palichik , V. Perelygin , M. Perfilov, S. Polikarpov ,⁹⁸ V. Popov, O. Radchenko ,⁹⁸
 V. Rusinov, M. Savina , V. Savrin , V. Shalaev , S. Shmatov , S. Shulha , Y. Skovpen ,⁹⁸
 S. Slabospitskii , V. Smirnov , D. Sosnov , V. Sulimov , E. Tcherniaev , A. Terkulov ,
 O. Teryaev , I. Tlisova , A. Toropin , L. Uvarov , A. Uzunian , P. Volkov , A. Vorobyev[†],
 G. Vorotnikov , N. Voytishin , B.S. Yuldashev¹⁰³, A. Zarubin , I. Zhizhin , A. Zhokin 

[†] *Deceased*¹ *Also at Yerevan State University, Yerevan, Armenia*² *Also at TU Wien, Vienna, Austria*³ *Also at Institute of Basic and Applied Sciences, Faculty of Engineering, Arab Academy for Science, Technology and Maritime Transport, Alexandria, Egypt*⁴ *Also at Ghent University, Ghent, Belgium*⁵ *Also at Universidade Estadual de Campinas, Campinas, Brazil*⁶ *Also at Federal University of Rio Grande do Sul, Porto Alegre, Brazil*⁷ *Also at UFMS, Nova Andradina, Brazil*⁸ *Also at Nanjing Normal University, Nanjing, China*⁹ *Now at The University of Iowa, Iowa City, Iowa, U.S.A.*¹⁰ *Also at University of Chinese Academy of Sciences, Beijing, China*¹¹ *Also at University of Chinese Academy of Sciences, Beijing, China*¹² *Also at Université Libre de Bruxelles, Bruxelles, Belgium*¹³ *Also at an institute or an international laboratory covered by a cooperation agreement with CERN*¹⁴ *Also at Cairo University, Cairo, Egypt*¹⁵ *Also at Suez University, Suez, Egypt*¹⁶ *Now at British University in Egypt, Cairo, Egypt*¹⁷ *Also at Birla Institute of Technology, Mesra, Mesra, India*¹⁸ *Also at Purdue University, West Lafayette, Indiana, U.S.A.*¹⁹ *Also at Université de Haute Alsace, Mulhouse, France*

- ²⁰ Also at *Department of Physics, Tsinghua University, Beijing, China*
- ²¹ Also at *Tbilisi State University, Tbilisi, Georgia*
- ²² Also at *The University of the State of Amazonas, Manaus, Brazil*
- ²³ Also at *Erzincan Binali Yildirim University, Erzincan, Turkey*
- ²⁴ Also at *University of Hamburg, Hamburg, Germany*
- ²⁵ Also at *RWTH Aachen University, III. Physikalisches Institut A, Aachen, Germany*
- ²⁶ Also at *Isfahan University of Technology, Isfahan, Iran*
- ²⁷ Also at *Bergische University Wuppertal (BUW), Wuppertal, Germany*
- ²⁸ Also at *Brandenburg University of Technology, Cottbus, Germany*
- ²⁹ Also at *Forschungszentrum Jülich, Juelich, Germany*
- ³⁰ Also at *CERN, European Organization for Nuclear Research, Geneva, Switzerland*
- ³¹ Also at *Physics Department, Faculty of Science, Assiut University, Assiut, Egypt*
- ³² Also at *Wigner Research Centre for Physics, Budapest, Hungary*
- ³³ Also at *Institute of Physics, University of Debrecen, Debrecen, Hungary*
- ³⁴ Also at *Institute of Nuclear Research ATOMKI, Debrecen, Hungary*
- ³⁵ Now at *Universitatea Babeş-Bolyai — Facultatea de Fizica, Cluj-Napoca, Romania*
- ³⁶ Also at *Faculty of Informatics, University of Debrecen, Debrecen, Hungary*
- ³⁷ Also at *Punjab Agricultural University, Ludhiana, India*
- ³⁸ Also at *UPES — University of Petroleum and Energy Studies, Dehradun, India*
- ³⁹ Also at *University of Visva-Bharati, Santiniketan, India*
- ⁴⁰ Also at *University of Hyderabad, Hyderabad, India*
- ⁴¹ Also at *Indian Institute of Science (IISc), Bangalore, India*
- ⁴² Also at *IIT Bhubaneswar, Bhubaneswar, India*
- ⁴³ Also at *Institute of Physics, Bhubaneswar, India*
- ⁴⁴ Also at *Deutsches Elektronen-Synchrotron, Hamburg, Germany*
- ⁴⁵ Also at *Sharif University of Technology, Tehran, Iran*
- ⁴⁶ Also at *Department of Physics, University of Science and Technology of Mazandaran, Behshahr, Iran*
- ⁴⁷ Also at *Helwan University, Cairo, Egypt*
- ⁴⁸ Also at *Italian National Agency for New Technologies, Energy and Sustainable Economic Development, Bologna, Italy*
- ⁴⁹ Also at *Centro Siciliano di Fisica Nucleare e di Struttura Della Materia, Catania, Italy*
- ⁵⁰ Also at *Università degli Studi Guglielmo Marconi, Roma, Italy*
- ⁵¹ Also at *Scuola Superiore Meridionale, Università di Napoli ‘Federico II’, Napoli, Italy*
- ⁵² Also at *Fermi National Accelerator Laboratory, Batavia, Illinois, U.S.A.*
- ⁵³ Also at *Università di Napoli ‘Federico II’, Napoli, Italy*
- ⁵⁴ Also at *Ain Shams University, Cairo, Egypt*
- ⁵⁵ Also at *Consiglio Nazionale delle Ricerche — Istituto Officina dei Materiali, Perugia, Italy*
- ⁵⁶ Also at *Riga Technical University, Riga, Latvia*
- ⁵⁷ Also at *Department of Applied Physics, Faculty of Science and Technology, Universiti Kebangsaan Malaysia, Bangi, Malaysia*
- ⁵⁸ Also at *Consejo Nacional de Ciencia y Tecnología, Mexico City, Mexico*
- ⁵⁹ Also at *Trincomalee Campus, Eastern University, Sri Lanka, Nilaveli, Sri Lanka*
- ⁶⁰ Also at *Saegis Campus, Nugegoda, Sri Lanka*
- ⁶¹ Also at *INFN Sezione di Pavia, Università di Pavia, Pavia, Italy*
- ⁶² Also at *National and Kapodistrian University of Athens, Athens, Greece*
- ⁶³ Also at *Ecole Polytechnique Fédérale Lausanne, Lausanne, Switzerland*
- ⁶⁴ Also at *University of Vienna Faculty of Computer Science, Vienna, Austria*
- ⁶⁵ Also at *Universität Zürich, Zurich, Switzerland*
- ⁶⁶ Also at *Stefan Meyer Institute for Subatomic Physics, Vienna, Austria*
- ⁶⁷ Also at *Laboratoire d’Annecy-le-Vieux de Physique des Particules, IN2P3-CNRS, Annecy-le-Vieux, France*
- ⁶⁸ Also at *Near East University, Research Center of Experimental Health Science, Mersin, Turkey*
- ⁶⁹ Also at *Konya Technical University, Konya, Turkey*
- ⁷⁰ Also at *Izmir Bakircay University, Izmir, Turkey*

- ⁷¹ Also at Adiyaman University, Adiyaman, Turkey
- ⁷² Also at Necmettin Erbakan University, Konya, Turkey
- ⁷³ Also at Bozok Universitetesi Rektörlüğü, Yozgat, Turkey
- ⁷⁴ Also at Marmara University, Istanbul, Turkey
- ⁷⁵ Also at Milli Savunma University, Istanbul, Turkey
- ⁷⁶ Also at Kafkas University, Kars, Turkey
- ⁷⁷ Also at Hacettepe University, Ankara, Turkey
- ⁷⁸ Also at Istanbul University — Cerrahpasa, Faculty of Engineering, Istanbul, Turkey
- ⁷⁹ Also at Yildiz Technical University, Istanbul, Turkey
- ⁸⁰ Also at Vrije Universiteit Brussel, Brussel, Belgium
- ⁸¹ Also at School of Physics and Astronomy, University of Southampton, Southampton, United Kingdom
- ⁸² Also at University of Bristol, Bristol, United Kingdom
- ⁸³ Also at IPPP Durham University, Durham, United Kingdom
- ⁸⁴ Also at Monash University, Faculty of Science, Clayton, Australia
- ⁸⁵ Now at an institute or an international laboratory covered by a cooperation agreement with CERN
- ⁸⁶ Also at Università di Torino, Torino, Italy
- ⁸⁷ Also at Bethel University, St. Paul, Minnesota, U.S.A.
- ⁸⁸ Also at Karamanoğlu Mehmetbey University, Karaman, Turkey
- ⁸⁹ Also at California Institute of Technology, Pasadena, California, U.S.A.
- ⁹⁰ Also at United States Naval Academy, Annapolis, Maryland, U.S.A.
- ⁹¹ Also at Bingol University, Bingol, Turkey
- ⁹² Also at Georgian Technical University, Tbilisi, Georgia
- ⁹³ Also at Sinop University, Sinop, Turkey
- ⁹⁴ Also at Erciyes University, Kayseri, Turkey
- ⁹⁵ Also at Horia Hulubei National Institute of Physics and Nuclear Engineering (IFIN-HH), Bucharest, Romania
- ⁹⁶ Also at Texas A&M University at Qatar, Doha, Qatar
- ⁹⁷ Also at Kyungpook National University, Daegu, Korea
- ⁹⁸ Also at another institute or international laboratory covered by a cooperation agreement with CERN
- ⁹⁹ Also at Universiteit Antwerpen, Antwerpen, Belgium
- ¹⁰⁰ Also at Northeastern University, Boston, Massachusetts, U.S.A.
- ¹⁰¹ Also at Imperial College, London, United Kingdom
- ¹⁰² Now at Yerevan Physics Institute, Yerevan, Armenia
- ¹⁰³ Also at Institute of Nuclear Physics of the Uzbekistan Academy of Sciences, Tashkent, Uzbekistan