

Research Paper

Nutritional Support in Cancer patients: update of the Italian Intersociety Working Group practical recommendations

Riccardo Caccialanza¹✉, Paolo Cotogni², Emanuele Cereda¹, Paolo Bossi³, Giuseppe Aprile⁴, Paolo Delrio⁵, Patrizia Gnagnarella⁶, Annalisa Mascheroni⁷, Taira Monge⁸, Ettore Corradi⁹, Michele Grieco¹⁰, Sergio Riso¹¹, Francesco De Lorenzo¹², Francesca Traclò¹², Elisabetta Iannelli¹², Giordano Domenico Beretta¹³, Michela Zanetti¹⁴, Saverio Cinieri¹⁵, Vittorina Zagonel¹⁶, and Paolo Pedrazzoli¹⁷, on behalf of the Intersociety (AIOM-SINPE-FAVO-SICO-ASAND) Italian Working Group for Nutritional Support in Cancer Patients[#]

1. Clinical Nutrition and Dietetics Unit, Fondazione IRCCS Policlinico San Matteo, Pavia, Italy
2. Pain Management and Palliative Care, Department of Anesthesia, Intensive Care and Emergency, Molinette Hospital, University of Turin, Turin, Italy;
3. Medical Oncology Unit, ASST Spedali Civili di Brescia, and Department of Medical and Surgical Specialties, Radiological Sciences, and Public Health, University of Brescia, Brescia, Italy
4. Department of Oncology, San Bortolo General Hospital, Vicenza, Italy
5. Colorectal Surgical Oncology-Abdominal Oncology Department, Istituto Nazionale per lo Studio e la Cura dei Tumori, Fondazione Giovanni Pascale IRCCS, Naples, Italy
6. Division of Epidemiology and Biostatistics, IEO European Institute of Oncology IRCCS, Milan, Italy
7. Clinical Nutrition and Dietetics Unit, ASST Melegnano-Martesana, 20077 Melegnano (MI), Italy
8. Clinical Nutrition Unit, S. Giovanni Battista Hospital, Torino, Italy
9. Clinical Nutritional Unit, ASST Grande Ospedale Metropolitano Niguarda, Milano, Italy
10. Department of Surgery, Sant' Eugenio Hospital, Rome, Italy
11. Clinical Nutrition and Dietetics Unit, Maggiore della Carità Hospital, Novara, Italy
12. Italian Federation of Volunteer-based Cancer Organizations, Rome, Italy
13. Department of Oncology, Humanitas Gavazzeni, Bergamo, Italy
14. Department of Medical, Surgical and Health Sciences - University of Trieste, and Azienda Sanitaria Universitaria Giuliano Isontina (ASUGI), Trieste, Italy
15. Medical Oncology Division and Breast Unit, Senatore Antonio Perrino Hospital, ASL Brindisi, Brindisi, Italy
16. Oncology Unit 1, Department of Oncology, Veneto Institute of Oncology-IRCCS, 35128 Padova, Italy
17. Medical Oncology Unit, Fondazione IRCCS Policlinico San Matteo and Department of Internal Medicine, University of Pavia, Pavia, Italy

[#]The Italian Intersociety (AIOM-SINPE-FAVO-SICO-ASAND) Working Group for Nutritional Support in Cancer Patients is listed in the Acknowledgments

✉ Corresponding author: Dr. Riccardo Caccialanza, Clinical Nutrition and Dietetics Unit, Fondazione IRCCS Policlinico San Matteo, Viale Golgi 19, 27100 Pavia, Italy. Tel.: + 39 0382 501615. E-mail: r.caccialanza@smatteo.pv.it

© The author(s). This is an open access article distributed under the terms of the Creative Commons Attribution License (<https://creativecommons.org/licenses/by/4.0/>). See <http://ivyspring.com/terms> for full terms and conditions.

Received: 2022.03.22; Accepted: 2022.05.15; Published: 2022.05.21

Abstract

Malnutrition is a frequent problem in cancer patients, which leads to prolonged and repeated hospitalizations, increased treatment-related toxicity, reduced response to cancer treatment, impaired quality of life, a worse overall prognosis and the avoidable waste of health care resources.

Despite being perceived as a limiting factor in oncologic treatments by both oncologists and patients, there is still a considerable gap between need and actual delivery of nutrition care, and attitudes still vary considerably among health care professionals.

In the last 5 years, the Italian Intersociety Working Group for Nutritional Support in Cancer Patients (WG), has repeatedly revisited this issue and has concluded that some improvement in nutritional care in Italy has occurred, at least with regard to awareness and institutional activities. In the same period, new international guidelines for the management of malnutrition and cachexia have been released.

Despite these valuable initiatives, effective structural strategies and concrete actions aimed at facing the challenging issues of nutritional care in oncology are still needed, requiring the active participation of scientific societies and health authorities.

As a continuation of the WG's work, we have reviewed available data present in the literature from January 2016 to September 2021, together with the most recent guidelines issued by scientific societies and health authorities, thus providing an update of the 2016 WG practical recommendations, with suggestions for new areas/issues for possible improvement and implementation.

Key words: nutritional support, cancer patients, malnutrition, practical recommendations, nutritional care

Introduction

Although malnutrition is recognized by both oncologists and patients as a limiting factor in oncologic treatments, it remains poorly managed [1]. The consequences are serious, leading to reduced anticancer treatment tolerance, poorer prognosis, impaired quality of life (QoL) and the avoidable waste of health care resources associated with prolonged and repeated hospitalizations [2]. Nevertheless, adherence to international guidelines and recommendations is still low, which limits access to high quality nutrition therapy both during and following cancer treatment [3].

Despite the abundance of scientific literature highlighting the problem, and the availability of international guidelines for managing nutritional care in cancer patients, many patients do not receive adequate nutritional support [2-4]. Beyond the obvious clinical consequences, overlooking nutrition care incurs billions in healthcare costs [5-8].

The Italian Association of Medical Oncology, the Italian Society of Artificial Nutrition and Metabolism and the Italian Federation of Volunteer-based Cancer Organizations implemented in 2016 a collaborative Working Group (WG) and initiated a structured project named "Integrating Nutritional Therapy in Oncology", with the aim to increase the awareness of nutritional issues among oncologists and, consequently, to improve the nutritional care of cancer patients in Italy [9]. In 2019, the Italian Society of Surgical Oncology and the Technical Scientific Association of Food, Nutrition and Dietetics joined the WG, which was named "Italian Intersociety Working Group for Nutritional Support in Cancer Patients".

Among its activities, in 2016 the WG issued the first inter-society consensus document in order to provide suitable, concise and practical recommendations for appropriate nutrition in cancer patients [10]. This publication was not meant to be a surrogate for international guidelines, but its aim was to provide oncologists, other professionals involved in cancer care and the patients themselves, with a concise, easily accessible and updated summary of the main recommendations needed to appropriately manage nutritional care in oncology.

In the last 5 years, several further initiatives have been undertaken by the WG [11], which has

concluded that some improvement in nutritional care in Italy has occurred, at least as far as awareness [2] and institutional practices are concerned [12]. In the same period, new international guidelines for the management of malnutrition and related syndromes – such as cachexia – have been released [13-15].

While this represents progress, nutritional care in oncology is still inadequate and needs the involvement and cooperation of scientific societies, the Ministry of Health and the Ministry of Education. Consequently, the WG decided to update the 2016 recommendations, which are presented here. The aim of this document is to: 1) stimulate the national and international Oncology Scientific and Clinical Community; 2) to increase the awareness on nutritional care; 3) to improve the clinical nutrition management of patients with cancer through the provision of simple but mandatory nutrition protocols for daily oncological practice.

Methodology

The WG included physicians (nutrition specialists, oncologists and surgeons), dietitians and patient representatives. We reviewed available data on the nutritional management of patients with cancer, which appeared in the literature from January 2016 to September 2021, including the evidence-based recommendations released in the guidelines issued by scientific societies and health authorities. Authors were also asked to identify further references from their personal collection of literature or other sources and to choose the most relevant ones to be included in the manuscript. After critical evaluation of literature, the original 2016 WG recommendations have been implemented along with accompanying commentaries. Compared to the 2016 paper, we chose to modify the structure, focusing still on nutritional risk and malnutrition recognition, nutritional counseling and oral supplementation, but then, also, on the different phases of the disease, together with current critical issues and future perspectives.

The drafting process was based on a consensus discussion followed by Delphi rounds and votes until agreement was reached. A final version of the paper was circulated and approved by the scientific board of the endorsing scientific societies, which exclusively funded the present project.

Early Recognition of Nutritional Risk and Malnutrition

Screening is key to identifying the risk of malnutrition [16]. If nutrition risk is not assessed at the first oncologic visit, nutritional deficiency will be missed in half the patients, and appropriate measures to counteract it will not be implemented [17,18].

A number of techniques have been used to assess nutrition status in cancer patients although no 'gold standard' has emerged as superior for sensitivity or specificity. The most frequently employed tools are: the Nutritional Risk Screening 2002 (NRS 2002), the Malnutrition Universal Screening Tool (MUST), the Malnutrition Screening Tool (MST), the patient-generated subjective global assessment (PG-SGA), and the Mini Nutritional Assessment (MNA) [17].

They all showed a moderate to substantial agreement with one another and should be employed as tools to guide corrective measures. There is no comprehensive evaluation of their comparative predictive and/or prognostic value on patient outcomes [19].

More recently, the Global Leadership Initiative on Malnutrition (GLIM) criteria, based on a consensus of experts, provides a diagnostic and operational tool to identify and treat malnutrition in several settings [20]. They consider phenotypic and etiological criteria and could be helpful in sharing standardized data worldwide.

Independently of the selected criteria/parameters, nutritional status should be considered a dynamic concept, particularly in oncology; therefore, nutritional screening tests should be administered early and periodically repeated, preferably by nurses, during the whole of the patient's journey - at each outpatient visit and within 48 hours of hospital admission.

As stated by all the available guidelines and recommendations, patients at risk of malnutrition should be referred to a clinical nutrition service/unit/professional for nutritional assessment and treatment. However, due to the foreseeable clinical course, it is reasonable to suggest that patients with certain cancer type (head&neck [H&N], gastrointestinal [GI], lung), advanced disease stage or undergoing more aggressive treatments (high-dose chemotherapy [CT], radical radiotherapy [RT], major abdominal surgery or multimodal [either combined or sequential]), all of which are expected to affect nutritional status, should be immediately referred to clinical nutrition specialists for early comprehensive nutritional assessment, counseling/support and a strict monitoring program, independently of risk

evaluation.

The assessment of nutritional status should preferably include tools to identify both malnutrition and to measure body composition, with particular reference to sarcopenia and muscle mass determination [20-25].

The nutritional evaluation should include the combination of different parameters [20]: anthropometric measurements (body weight, height, body mass index [BMI]), unintentional weight loss enquiry, biochemical data related to metabolic and inflammatory status, the assessment of nutritional intake, QoL, and physical function tests (gait speed, grip strength) to assess muscle performance [21].

Scientific literature suggests that the exclusive use of anthropometric measures is not sufficient to identify body composition alterations, particularly with respect to muscle mass loss [24]. Body composition assessment in cancer patients can be performed by Dual-Energy X-ray Absorptiometry (DEXA) or Bioelectrical Impedance Vectorial Analysis (BIVA), the latter also providing information on hydration and cell mass integrity [26]. In particular, low phase angle is a predictor of compromised nutritional status, impaired muscle function, increased risk of morbidity, and reduced survival [26,27].

Computed Tomography and Magnetic Resonance Imaging are the gold standard techniques to assess body composition and their imaging of lumbar vertebra L3 correlates well with whole-body skeletal muscle mass [22,28].

Nutritional Counseling and Oral Supplementation

Nutritional support should be provided to malnourished patients and those at nutritional risk, in particular when oral energy intake is already insufficient or expected to be inadequate (<60% of estimated caloric requirements) for more than 7 days [13,29,30]. The aim of nutritional counseling is to maintain or improve food intake through a diet enriched in calories, proteins and fluids that are better tolerated, and to favour the management of the nutrition impact symptoms (i.e. anorexia, nausea, vomiting, diarrhea, and dysphagia). It should be the first type of support proposed and should be carried out by a dietitian with documented skills in cancer patient care [10,12] for appropriate dietary intervention and its monitoring [31,32]. As reported in **Table 1**, this process includes a few steps [33] and aims at providing patients with a thorough understanding of nutritional topics that can lead to long-lasting changes in their eating habits, taking into account individual preferences, ethnicity, culture,

estimated nutritional requirements and cancer treatment side effects.

Table 1: Nutritional counseling process in cancer patients

Nutrition Assessment and Reassessment:	<ul style="list-style-type: none"> • body weight assessment / changes / body composition; • biochemical data, medical tests and procedures; • energy, macro and micronutrient requirements; • actual food consumption (preferences and habits), and food and nutrition-related history; • estimated nutritional requirements; • cancer treatment side effects; • preferences, ethnicity, culture.
Nutrition Diagnosis:	<ul style="list-style-type: none"> • problems, difficulties and symptoms related to treatments that limit the consumption or absorption of nutrients; • obstacles to change (inconvenience, social problems, food preferences, lack of knowledge or time, costs).
Nutrition Intervention:	<ul style="list-style-type: none"> • definition of objectives; • meal set-up plan that emphasizes increasing meal frequency by distribution of foods to several small meals; • enriching dishes with energy- and protein-dense ingredients oral nutritional supplements; • food preparation and/or modifying of texture or nutrient content; • specific indication for mucositis and other symptoms, digestion (e.g. pancreatic enzymes) or absorption (e.g. slowing of rapid gastrointestinal transit), antiemetic, and other relevant conditions; • alliances with caregivers.
Nutrition Monitoring/Evaluation:	<ul style="list-style-type: none"> • monitoring and re-evaluation to determine if the patients has achieved, or is making progress toward, the planned goals.

Practical suggestions for managing common symptoms related to cancer treatment, leading to impaired food intake or malabsorption, should be foreseen to optimize patients' diets, in order to cope with nutritional deficiencies and possible swallowing difficulties.

Nutritional interventions should compensate for inadequate energy intake with the objective of improving clinical outcomes. So far, numerous reviews have been published [34-40] in malnourished hospitalized and community-dwelling adults with cancer.

Multiple nutrition interventions have been proposed, including dietary counseling or advice, oral nutritional supplements (ONS) and enteral nutrition (EN). The evidence for nutritional counseling to improve clinical outcomes is heterogeneous. According to the most recent review, nutrition interventions were found able to improve body weight and BMI, nutritional status, protein and energy intake, QoL and response to cancer treatments

[40]. Inconclusive results were found regarding body composition, functional status, complications, unplanned hospital readmissions and survival. Interestingly, Richards and colleagues found that early nutrition intervention, that is initiated within the first week of cancer treatment, can improve patient prognosis and outcomes [40].

When dietary measures fail to meet patients' protein-calorie requirements as detected by nutritional monitoring, the prescription of energy-dense ONS should be considered, due to their proven efficacy in increasing protein-calorie intake and to fill nutritional gaps [13,41].

In patients with cancer, systemic inflammation inhibits nutrient utilization and promotes catabolism, thus leading to muscle breakdown. Calorie and protein fortification of regular foods, even with standard ONS, does not reduce systemic inflammation. Updated nutritional strategies now suggest considering nutrition with anti-catabolic and inflammation-suppressing ingredients. Studies have indicated that ONS with addition of essential amino acids or high-dose leucine may improve muscle protein synthesis even in the presence of inflammation, although results have not been fully consistent [42,43].

Fish oil, a source of long chain omega-3 fatty acids, is currently suggested to improve appetite, oral intake, lean body mass, and body weight in patients with advanced cancer and at risk of malnutrition [13,44].

The European Society of Clinical Nutrition and Metabolism (ESPEN) guidelines on nutrition in cancer patients recommend supplementation with fish oil, a source of long chain omega-3 fatty acids, to stabilize or improve appetite, food intake, lean body mass, and body weight for patients with advanced cancer undergoing CT, but the level of evidence is still low [13].

Studies included in the previously mentioned review, evaluated a sole nutrition intervention of ONS enriched in omega-3 fatty acids (ONS- ω 3) vs. placebo, an isocaloric diet, or an isocaloric ONS: they found significantly reduced weight loss and loss of fat free mass, and significantly increased skeletal muscle mass and lean body mass, QoL, and treatment tolerance in the groups receiving ONS- ω 3.

In a recent pragmatic randomized controlled-trial conducted in 159 H&N cancer patients undergoing RT and CT + RT and receiving nutritional counseling, the use systematic use of ONS- ω 3 resulted in better weight maintenance, increased protein-calorie intake, improved QoL and was associated with better anti-cancer treatment tolerance [45], with no additional costs for the healthcare

system [46].

One limitation of most of the available studies on nutritional counseling and oral supplementation is their reliance on small sample size, different outcome measures, variable nutrition interventions, different timings and follow-up.

Of note, however, is a recent well-conducted multicenter study on 506 hospitalized cancer patients at nutritional risk, which showed that individualized nutritional support reduced the risk of mortality and improved functional and quality of life outcomes [47].

The optimal duration and timing of nutrition interventions should also be explored further, but the general impression that the early approach may be the most effective is gaining support [48].

Nutritional Support in the Different Phases of the Disease

Active Systemic Treatment

The often underdiagnosed unintentional weight loss and progressive skeletal muscle depletion are frequently present and worsen during CT and targeted therapies [22,23,49].

Regardless of cancer type, frailty level or age, patients may be similarly affected by malnutrition, [50-56], that is often linked with a range of different side effects, either due to the disease itself or the antineoplastic treatment, including oral mucositis, dysgeusia or smell alterations, fatigue, dyspepsia, nausea and vomiting. These events, by compromising adherence to therapies, may negatively affect outcome and limit the access to novel therapies.

Overall, the management of weight-losing cancer patients during active oncologic treatment relies on nutritional counseling, non-pharmacological (ONS and physical activity) interventions and pharmacological support, including artificial nutrition (AN) [10, 13-15, 23,57,58].

A proactive assessment of nutritional status is essential for selecting the most cost-effective intervention to apply: for example, oral nutritional support might reduce weight loss and hospitalization, and dronabinol may help to counteract nausea and vomiting or increase food intake. Ongoing randomized trials will make clear if treatment-naïve cancer patients at risk for malnutrition might also profit from an early, short-term, supplemental parenteral nutrition [59]. Whether starving cancer cells may help increasing the activity of chemotherapeutic agents is currently unclear and prospective, well designed, randomized trials on caloric restriction or caloric restriction mimetics are missing [60]. When tested under strict protocols, the preliminary clinical results are promising [61].

However, due to the lack of solid clinical evidence, fasting and fasting-mimicking diets during active treatment are still not recommended, even in cancer types associated with a lower risk of malnutrition (e.g. breast and prostate cancer).

Novel preclinical data have suggested that a relationship between cell metabolism and susceptibility to immunotherapy may exist and uncontrolled studies have reported a correlation between BMI and efficacy of immunotherapeutic agents in several cancers [62-64].

Nevertheless, since conflicting data regarding any benefit from immunomodulators according to baseline BMI have been reported in different cancer types, this issue remains unresolved and the study of a more comprehensive or dynamic nutritional pattern has been suggested [65,66].

A secondary analysis of the data from the Swiss prospective, randomized-controlled, multicenter trial EFFORT compared the outcomes of a protocol-guided individualized nutritional support regime (intervention group) to standard hospital food (control group) in 506 patients with a main admission diagnosis of cancer and characterized by a broad spectrum of cancer sites, treatment types and disease severities. Individualized nutritional support reduced the risk of mortality and improved functional and QoL outcomes in cancer patients with increased nutritional risk, further supporting the inclusion of nutritional care in cancer management guidelines (47).

The efficacy of an early nutritional intervention was also highlighted in the study of Lu et al. (67), in which patients with metastatic esophagogastric cancer, who received an early interdisciplinary supportive care, provided by a team of gastrointestinal oncologists, nurse specialists, dietitians, and psychologists, integrated into standard oncologic care, was associated with an improved overall survival, compared with patients in the standard oncologic care-alone group (14.8 vs 11.9 months).

Overall, even if the above mentioned and other issues require further research-based evidence, we can affirm with confidence that prompt and strictly monitored nutritional support during oncologic active treatment is essential, in order to improve clinical outcomes and provide patients with the most innovative and effective therapeutic options.

Although activating a formal nutritional team in every hospital setting may be difficult, many successful examples may contribute in shaping a shared, cooperative awareness between patient communities and healthcare professionals [68].

The adoption of validated care pathways can

ensure compliance with updated scientific recommendations, guarantee access to treatments backed by clinical evidence, contain clinical risks, rationalize expenditures, and increase equity.

Surgical Setting

Nutritional care is a keystone for clinical outcomes in oncologic surgery.

For now, early oral feeding should always be the preferred nutrition mode after oncologic surgery, and this includes gastro-intestinal procedures [69].

The surgical operation causes a trauma that produces systemic inflammation, stress response and metabolic negative effects [70]. The outcome of oncological surgery is not only related to the technical effectiveness of the surgical procedure itself, but also to the perioperative management. Lack of nutritional pre-operative screening and unnecessary post-operative fasting, raise the risk of underfeeding and surgical complications, especially after major surgery [71].

The perioperative management of surgical patients has been studied in depth in recent years and all these measures have been included in the Enhanced Recovery After Surgery (ERAS) programs [72]: a scheduled perioperative pathway to minimize the surgical stress and improve the postoperative functional restoration [73]. The ERAS protocol can be applied in many surgical fields such as colorectal surgery [74], esophageal surgery [75], pancreatic surgery [76], gynecological surgery [77] and many others [72].

Every patient undergoing major surgery should follow a personalized perioperative ERAS program, comprising scheduled steps, including nutritional care. Most of them are common to many different surgical specialties and procedures, such as:

- Preoperative nutritional screening with validated tools should be always performed before oncologic surgery, because severe under-nutrition has long been known to be detrimental to surgical outcome [78-80].

- Nutritional support should be started if the patient is malnourished or at nutritional risk before surgery. An appropriate nutritional support package should be applied during the hospital stay and following discharge [81]. Nutritional interventions in the perioperative period includes counseling, ONS, EN and parenteral nutrition (PN) for all patients malnourished or at risk of malnutrition [69].

- In patients undergoing gastrointestinal surgery, the use of oral/enteral immunonutrition should be encouraged for a reduction in post-operative infectious complications [82].

- Preoperative fasting from midnight should be always avoided. Patients with no specific risk of aspiration can drink clear fluids until 2 hours prior to surgery and eat solid food until 6 hours before surgery [83].

- Oral feeding should be started as soon as possible after surgery, adapting oral intake according to type of surgery and individual tolerance. Special caution should be dedicated to elderly patients [84].

- Risk of postoperative ileus should be minimized and an opioid-sparing pathway should be applied, especially the use short-acting anesthetics during surgery and applying multimodal analgesia in combination with epidural analgesia post-operatively [85].

Palliative Care

The role of nutritional support for oncology patients in palliative care is still a controversial issue.

However, in advanced cancer patients, preserving nutritional status may be an important goal also during the palliative care phase, because even when the disease can no longer be cured, patients may survive for several months or years. In this context, malnutrition may jeopardize performance status, QoL, tolerance to palliative treatments and survival.

AN can be integrated within a palliative care program when the risk of dying from malnutrition is greater than due to cancer progression [10,13]. ESPEN guidelines suggest that EN should be considered first when the gastrointestinal tract is functional but oral food intake remains inadequate despite nutritional counseling and ONS, whereas, if EN is not sufficient or possible, PN is recommended [13].

Nevertheless, there are many factors that may adversely affect the provision of EN in these patients, e.g. high output ileostomy or intestinal fistulas, large bowel resections, and the presence of nutrition impact symptoms (abdominal pain, nausea, vomiting, diarrhea, or constipation due to peritoneal carcinomatosis). Also, ESPEN guidelines recommend home artificial nutrition (HAN), either EN or PN, in eligible cancer patients with persistent insufficient oral food intake or malabsorption [13].

However, in incurable patients nutritional support should be proposed considering the expected benefit on QoL and the potential benefit on survival [13]. According to the principles of bioethics and ESPEN guidelines [13], the prescription of HAN should be discussed with the patient respecting his/her autonomy and, as also required by law, his/her choice or advance directive to refuse it. Concerning clinical appropriateness, HAN is not recommended in patients with short estimated life

expectancy, severe organ dysfunction or uncontrolled symptoms, Karnofsky performance status <50 or Eastern Cooperative Oncology Group (ECOG) score ≥ 3 , and in the presence of patient refusal [86].

Finally, an important question is whether there is evidence of a potential survival benefit due to home parenteral nutrition (HPN) in palliative cancer patients. Recently, a prospective study compared the survival of malnourished cancer patients in palliative care, eligible for HPN according to ESPEN recommendations, who received HPN with a homogenous group of patients, equally eligible for HPN, who did not receive HPN but artificial hydration for logistic reasons or because of patient refusal [87]. Survival of the two groups showed a statistically significant difference in favor of the HPN group, which had a median overall survival three times higher than that receiving artificial hydration (4.3 versus 1.5 months, respectively).

Critical Issues and Perspectives

Malnutrition in oncology is still under-diagnosed and untreated in an unacceptable proportion of patients [2-4] despite the clinical [45, 88,89] and economic [5,7] benefits resulting from early and appropriate nutritional support. The quality of nutritional cancer care in Italy, and to our knowledge throughout Europe [3], remains poor and far from the standards indicated by the international [13-15] and national [12] guidelines, and the Cancer Patients' Bill of Right for Prompt and Appropriate Nutritional Support [90].

One major issue is that the number of clinical nutrition units/services is inadequate and most oncology units are even devoid of dedicated dietitians. In Italy, according to the available data, there are around 50 clinical nutrition structured hospital units around the country, which is an insufficient number when compared to the number of oncology units (~330) [91] and these are unevenly distributed among the Italian regions.

In addition, clinical nutrition is an overlooked topic at every university teaching level, so that even the basic knowledge is lacking in young physicians and health care professionals.

Nowadays, the general education and training offered in clinical nutrition, which should be tailored to satisfy specific professional requirements, is qualitatively and quantitatively inappropriate in the majority of Western countries [92]. Teaching of clinical nutrition is generally insufficient even within medical education, which is paradoxical, considering the relevance of nutritional issues in relation to both prevention and disease treatment [93,94].

The timing of nutritional intervention is fundamental in order to improve the therapeutic chances, with the "early approach" being the most effective [95,48]. However, it is reasonable to think that not enough attention is still paid to this issue, as the most recent guidelines continue to be focused mainly on cachexia and advanced cancer patients [14,15].

Similarly, nutritional parameters are still not systematically evaluated and considered as potential confounders in outcome assessment and study design in clinical oncology trials.

Last but not least, disinformation remains a critical issue with regards to nutrition for cancer patients. Despite the lack of evidence-based data, hundreds of books and web sites still promote anti-cancer diets, fasting, fasting-mimicking diets and nutritional supplements, which are widely used without medical supervision [96]. The autonomous use of dietary supplements should be discouraged, as they could interfere with oncologic treatments [97]. They should only be prescribed by clinical nutrition specialists according to documented deficiencies and clinical conditions, and their efficacy and patients' compliance should be regularly monitored and reassessed.

Inadequate nutritional management for cancer patients should be considered ethically unacceptable and the lack of standard "high level" evidence cannot be a justification for overlooking nutritional care [96]. However, providing additional evidence of its benefits from properly designed and sized clinical trials should be a key clinical objective in overcoming the barriers hampering the provision of proper nutritional support to cancer patients [2,10,98]. Additional efforts for improving awareness on nutrition in cancer and counter the world of fake news is a priority and will require time and considerable effort, particularly at the institutional level, and for healthcare personnel. Cancer patient associations can play a fundamental role in sensitizing public opinion and Institutions in this regard. In order to avoid an excessive additional workload for oncologists and cancer care managers, it is reasonable to argue that the institution of multi-disciplinary teams or the inclusion of a clinical nutrition specialist in existing local tumor boards (at least for the cancer types associated with the highest nutritional risk [head&neck, gastrointestinal, lung]) is strongly recommended, as it can be beneficial to comprehensive patient management, by encouraging early referral and continuous nutritional support implementation and monitoring.

Table 2. Summary of the Intersociety Italian Working Group for Nutritional Support in Cancer Patients updated recommendations

- Nutritional screening should be performed, preferably by nurses, using validated tools (NRS 2002, MUST, MST, MNA, PG-SGA) upon diagnosis, systematically repeated at each outpatient visit and within 48 hours since hospital admission.
- Patients at nutritional risk should be referred promptly for comprehensive nutritional assessment, possibly including the evaluation of body composition, and support to clinical nutrition services or medical personnel with documented skills in clinical nutrition.
- Patients with cancer types expected to affect nutritional status (head & neck, gastrointestinal, lung), advanced stage or treatment (high-dose chemotherapy, radical radiotherapy, major abdominal surgery or multimodal [either combined or sequential]) should be referred directly to clinical nutrition specialists for early comprehensive nutritional assessment, counseling/support and a strict monitoring program.
- Nutritional support should be initiated swiftly and targeted for each patient according to nutritional and clinical conditions, planned treatment and expected outcome. It should comprise nutritional counseling with the possible use of oral nutritional supplements and/or artificial nutrition (enteral nutrition, total or supplemental parenteral nutrition) according to the assessment and ensure the strict monitoring of spontaneous food intake, tolerance and effectiveness.
- Nutritional support and dietary modifications should aim to assist the maintenance or recovery of nutritional status by increasing or preserving protein and calorie intake. "Alternative hypocaloric anti-cancer diets" (e.g. macrobiotic or vegan diets), fasting and fasting-mimicking diets are not recommended.
- The autonomous use of dietary supplements should be discouraged. They should be prescribed by clinical nutrition specialists according to documented deficiencies and clinical conditions. Their efficacy and patients' compliance should be regularly monitored and reassessed.
- Every cancer patient undergoing major surgery should follow a personalized perioperative "Enhanced Recovery After Surgery" program that should comprise scheduled steps, including nutritional assessment and support.
- Nutritional support should be integrated into palliative care programs when the risk of dying from malnutrition is greater than from cancer progression, according to individual-based evaluations, quality of life implications, life expectancy and patients' will. Nutritional counseling aimed at alleviating nutrition-related symptoms should be provided to cancer patients receiving palliative care.
- Home artificial nutrition should be prescribed - even in the early phase if needed - and regularly monitored using defined protocols shared by all the healthcare professionals involved in patient care at institutional or, ideally regional/national level.
- Nutritional parameters should be always evaluated and considered as potential confounders in outcome assessment and study design in clinical oncology research.
- Adequately sized and designed clinical and cost-effectiveness trials, possibly involving cancer patient associations' representatives in the design process, are still needed in order to improve the evidence in favour of nutritional support in different care settings. The lack of standard "high level" evidence should not be a justification for overlooking nutritional care.
- The introduction of multi-disciplinary nutritional teams or the inclusion of clinical nutrition specialists in the existing local tumor boards (at least for the cancer types associated with the highest nutritional risk [head & neck, gastrointestinal, lung]) is strongly recommended.

Another critical issue is the quality and the clinical relevance of the available nutritional studies in the oncologic field. Unfortunately, many trials report only nutritional outcomes rather than clinical outcomes (i.e. treatment response, toxicity, survival,

QoL, etc.). Consequently, the effectiveness of nutritional support may result of limited interest for oncologists and the other health care professionals. Hence, the implementation of properly designed nutritional trials focusing on primary relevant clinical endpoints is key for the future development of clinical nutrition in oncology.

Among the topics of future studies, calorie and protein needs assessment for cancer patients should be of particular interest in light of the potential parallelism with the intensive care setting debate [99].

Preliminary experience adopting indirect calorimetry in the oncologic setting yield promising results [100] and further studies are ongoing. Waiting for the results, calorie and protein goals of patients with cancer should be still set according to the ESPEN guidelines [13], i.e. 25-30 kcal/kg and 1.3-1.5 g/kg, respectively.

From an economic point of view, quality improvement programs (QIP) involving nutrition support, such as the free provision of ONS at home, showed the potential for saving substantial resources [5,7,101]. QIP in the nutritional care of cancer patients, should be developed and promoted, and reimbursement policies should be reassessed or modulated, accordingly.

The combination of nutritional support with personalized physical activity programs represents another promising and relevant issue, which deserves further exploration and implementation, in order to preserve muscle mass more effectively [102,103].

Finally, in the era of new technologies, telemedicine implementation may be extremely helpful for patient monitoring and support, leaving them safely at home, but its real applicability still requires proper assessment in the oncologic setting.

Conclusions

Compared to the 2016 document, the updated Italian Intersociety WG for Nutritional Support in Cancer Patients makes additional practical recommendations, summarized in **Table 2**, and has identified some new areas/issues for possible improvement and implementation.

In order to implement these recommendations across the whole country, the WG will continue its work by establishing stable collaborations with national and international scientific societies/organizations, health authorities and academic institutions.

Acknowledgments

The Italian Intersociety Working Group for Nutritional Support in Cancer Patients is composed by:

AIOM (Italian Association of Medical Oncology): Giuseppe Aprile, Department of Oncology, San Bortolo General Hospital, Vicenza, Italy; Paolo Bossi, Medical Oncology Unit, ASST Spedali Civili di Brescia and Department of Medical and Surgical Specialties, Radiological Sciences, and Public Health, University of Brescia, Brescia, Italy; Gabriella Farina, Department of Oncology, ASST Fatebenefratelli Sacco, Milan, Italy; Paolo Pedrazzoli, Medical Oncology Unit, Fondazione IRCCS Policlinico San Matteo and Department of Internal Medicine, University of Pavia, Pavia Italy; Silvia Stragliotto, Medical Oncology 1, Veneto Institute of Oncology-IRCCS, Padova, Italy.

SINPE (Italian Society of Artificial Nutrition and Metabolism): Riccardo Caccialanza, Clinical Nutrition and Dietetics Unit, Fondazione IRCCS Policlinico San Matteo, Pavia, Italy; Ettore Corradi, Clinical Nutrition and Dietetics Unit, ASST Grande Ospedale Metropolitano Niguarda, Milan, Italy; Paolo Cotogni, Pain Management and Palliative Care, Department of Anesthesia, Intensive Care and Emergency, Molinette Hospital, University of Turin, Turin, Italy; Annalisa Mascheroni, Clinical Nutrition Service, ASST Melegnano-Martesana, Melegnano (MI), Italy; Sergio Riso, Clinical Nutrition and Dietetics Unit, Maggiore della Carità Hospital, University of Eastern Piedmont, Novara, Italy.

FAVO (Italian Federation of Volunteer-based Cancer Organizations), Rome, Italy: Francesco De Lorenzo, Elisabetta Iannelli, Francesca Tracò.

SICO (Italian Society of Oncological Surgery): Paolo Delrio, Colorectal Surgical Oncology, Istituto Nazionale per lo Studio e la Cura dei Tumori, Fondazione Giovanni Pascale IRCCS-Italia, Naples, Italy; Michele Grieco, General Surgery Unit, Sant'Eugenio Hospital, Rome, Italy.

ASAND (Technical Scientific Association of Food, Nutrition and Dietetics): Patrizia Gnagnarella, Division of Epidemiology and Biostatistics, IEO European Institute of Oncology IRCCS, Milan, Italy; Taira Monge, Clinical Nutrition Unit, S. Giovanni Battista Hospital, Turin, Italy.

We are grateful to Dr. Paul Baines for assistance in editing the manuscript.

Author Contributions

Conceptualization: R.C., P.P., P.C., F. DL. Writing original draft: R.C., P.P., P.C., P.B., G.A., P.D., P.G., A.M., T.M., E.C., M.G., S.R., F. DL., F.T. Supervision: R.C., P.P., F.DL., E.I., P.C., P.B., G.A., P.D., G.D.B., V.Z., S.C., M.Z., P.G. Critical revision: all authors and the Italian Intersociety Working Group for Nutritional Support in Cancer Patients. All authors and the Italian Intersociety Working Group for Nutritional Support in Cancer Patients have read

and agreed to the published version of the manuscript.

Funding

The article was partially supported by Ricerca Corrente funding 2019, Fondazione IRCCS Policlinico San Matteo, Pavia, Italy and by the scientific societies constituting the WG.

Competing Interests

The authors have declared that no competing interest exists.

References

- Caccialanza R, Lobascio F, Cereda E, Aprile G, Farina G, Tracò F, Borioli V, Caraccia M, Turri A, De Lorenzo F, Pedrazzoli P. Cancer-related malnutrition management: A survey among Italian Oncology Units and Patients' Associations. *Curr Probl Cancer*. 2020;44:100554.
- Baracos VE. Cancer-associated malnutrition. *Eur J Clin Nutr*. 2018;72:1255-1259.
- Caccialanza R, Goldwasser F, Marschal O, Ottery F, Schiefke J, Tilleul P, Zalcman G, Pedrazzoli P. Unmet needs in clinical nutrition in oncology: a multinational analysis of real-world evidence. *Ther Adv Med Oncol*. 2020;12:1758835919899852.
- Hébuterne X, Lemarié E, et al. Prevalence of malnutrition and current use of nutrition support in patients with cancer. *JPEN*. 2014;38:196-204.
- Tyler R, Barocas A, Guenter P, Araujo Torres K, Bechtold ML, Chan LN, Collier B, Collins NA, Evans DC, Godamunne K, Hamilton C, Hernandez BJD, Mirtallo JM, Nadeau WJ, Partridge J, Perugini M, Valladares A; ASPEN Value Project Scientific Advisory Council. Value of Nutrition Support Therapy: Impact on Clinical and Economic Outcomes in the United States. *JPEN J Parenter Enteral Nutr*. 2020;44:395-406.
- Goates S, Du K, Braunschweig CA, Arensberg MB. Economic Burden of Disease-Associated Malnutrition at the State Level. *PLoS One*. 2016 Sep 21;11(9):e0161833.
- Riley K, Sulo S, Dabbous F, Partridge J, Kozmic S, Landow W, VanDerBosch G, Falson MK, Sriram K. Reducing Hospitalizations and Costs: A Home Health Nutrition-Focused Quality Improvement Program. *JPEN J Parenter Enteral Nutr*. 2020;44:58-68.
- Mullin GE, Fan L, Sulo S, Partridge J. The Association between Oral Nutritional Supplements and 30-Day Hospital Readmissions of Malnourished Patients at a US Academic Medical Center. *J Acad Nutr Diet*. 2019;19:1168-1175.
- Caccialanza R, De Lorenzo F, Pedrazzoli P. The integrating nutritional therapy in oncology (INTO) project: rationale, structure and preliminary results. *ESMO Open*. 2017;2:e000221.
- Caccialanza R, Pedrazzoli P, Cereda E, et al. Nutritional support in cancer patients: a position paper from the Italian Society of Medical Oncology (AIOM) and the Italian Society of Artificial Nutrition and Metabolism (SINPE). *J Cancer*. 2016;7:131-5.
- Caccialanza R, De Lorenzo F, Lobascio F, Gnagnarella P, Iannelli E, Tracò F, Delrio P, Tancredi R, Pedrazzoli P; Italian Intersociety Working Group for Nutritional Support in Cancer Patients. Nutritional care in cancer patients: Initiatives and perspectives of the Italian Intersociety Working Group for Nutritional Support in Cancer Patients. *Nutrition*. 2021;91-92:111358.
- Accordo Stato-Regioni su "Linee di indirizzo sui percorsi nutrizionali nei pazienti oncologici" (rep. atti n. 224/csr; 14/12/2017). https://www.salute.gov.it/imgs/C_17_pubblicazioni_2682_allegato.pdf
- Arends J, Bachmann P, Baracos V, Barthelemy N, Bertz H, Bozzetti F, Fearon K, Hütterer E, Isenring E, Kaasa S, Krznaric Z, Laird B, Larsson M, Laviano A, Mühlebach S, Muscaritoli M, Oldervoll L, Ravasco P, Solheim T, Strasser F, de van der Schueren M, Preiser JC. ESPEN guidelines on nutrition in cancer patients. *Clin Nutr*. 2017;36:11-48.
- Roeland EJ, Bohlke K, Baracos VE, Bruera E, Del Fabbro E, Dixon S, Fallon M, Herrstedt J, Lau H, Platek M, Rugo HS, Schnipper HH, Smith TJ, Tan W, Loprinzi CL. Management of Cancer Cachexia: ASCO Guideline. *J Clin Oncol*. 2020;38:2438-2453.
- Arends J, Strasser F, Gonella S, Solheim TS, Madeddu C, Ravasco P, Buonaccorso L, de van der Schueren MAE, Baldwin C, Chasen M, Ripamonti CI; ESMO Guidelines Committee. Cancer cachexia in adult patients: ESMO Clinical Practice Guidelines. *ESMO Open*. 2021;6:100092.

16. Arends J, Baracos V, Bertz H, Bozzetti F, Calder PC, Deutz NEP, Erickson N, Laviano A, Lisanti MP, Lobo DN, McMillan DC, Muscaritoli M, Ockenga J, Pirllich M, Strasser F, de van der Schueren M, Van Gossum A, Vaupel P, Weimann A. ESPEN expert group recommendations for action against cancer-related malnutrition. *Clin Nutr.* 2017;36:1187-1196.
17. Reber E, Schönenberger KA, Vasiloglou MF, Stanga Z. Nutritional Risk Screening in Cancer Patients: The First Step Toward Better Clinical Outcome. *Front Nutr.* 2021;8:603936.
18. Muscaritoli M, Lucia S, Farcomeni A, Lorusso V, Saracino V, Barone C, Plastino F, Gori S, Magarotto R, Carteni G, Chiurazzi B, Pavese I, Marchetti L, Zagonel V, Bergo E, Tonini G, Imperatori M, Iacono C, Maiorana L, Pinto C, Rubino D, Cavanna L, Di Cicilia R, Gamucci T, Quadri S, Palazzo S, Minardi S, Merlano M, Colucci G, Marchetti P; PreMiO Study Group. Prevalence of malnutrition in patients at first medical oncology visit: the PreMiO study. *Oncotarget.* 2017;8:79884-79896.
19. Van Bokhorst-de van der Schueren MA, Guaitoli PR, Jansma EP, de Vet HC. Nutrition screening tools: does one size fit all? A systematic review of screening tools for the hospital setting. *Clin Nutr.* 2014;33:39-58.
20. Cederholm T, Jensen GL, Correia MITD, Gonzalez MC, Fukushima R, Higashiguchi T, Baptista G, Barazzoni R, Blaauw R, Coats A, Crivelli A, Evans DC, Gramlich L, Fuchs-Tarlovsky V, Keller H, Llido L, Malone A, Mogensen KM, Morley JE, Muscaritoli M, Nyulasi I, Pirllich M, Pisprasert V, de van der Schueren MAE, Siltharm S, Singer P, Tappenden K, Velasco N, Waitzberg D, Yamwong P, Yu J, Van Gossum A, Compher C; GLIM Core Leadership Committee; GLIM Working Group. GLIM criteria for the diagnosis of malnutrition - A consensus report from the global clinical nutrition community. *Clin Nutr.* 2019;38:1-9.
21. Cruz-Jentoft AJ, Bahat G, Bauer J, Boirie Y, Bruyère O, Cederholm T, Cooper C, Landi F, Rolland Y, Sayer AA, Schneider SM, Sieber CC, Topinkova E, Vandewoude M, Visser M, Zamboni M; Writing Group for the European Working Group on Sarcopenia in Older People 2 (EWGSOP2), and the Extended Group for EWGSOP2. Sarcopenia: revised European consensus on definition and diagnosis. *Age Ageing.* 2019;48:16-31.
22. Pamoukdjian F, Bouillet T, Lévy V, Soussan M, Zelek L, Paillaud E. Prevalence and predictive value of pre-therapeutic sarcopenia in cancer patients: A systematic review. *Clin. Nutr.* 2018; 37:1101-1113.
23. Prado CM, Purcell SA, Laviano A. Nutrition interventions to treat low muscle mass in cancer. *J Cachexia Sarcopenia Muscle.* 2020;11:366-380.
24. Brown JC, Cespedes Feliciano EM, Caan BJ. The evolution of body composition in oncology-epidemiology, clinical trials, and the future of patient care: Facts and numbers. *J. Cachexia Sarcopenia Muscle* 2018;9:1200-1208.
25. Kamarajah SK, Bundred J, Tan BHL. Body composition assessment and sarcopenia in patients with gastric cancer: A systematic review and meta-analysis. *Gastric Cancer* 2019;22:10-22.
26. Grundmann O, Yoon SL, Williams JJ. The value of bioelectrical impedance analysis and phase angle in the evaluation of malnutrition and quality of life in cancer patients—a comprehensive review. *Eur J Clin Nutr.* 2015;69:1290-1297.
27. Lukaski HC, Kyle UG, Kondrup J. Assessment of adult malnutrition and prognosis with bioelectrical impedance analysis: phase angle and impedance ratio. *Curr Opin Clin Nutr Metab Care.* 2017;20:330-339.
28. Ní Bhuachalla ÉB, Daly LE, Power DG, Cushen SJ, MacEneaney P, Ryan AM. Computed tomography diagnosed cachexia and sarcopenia in 725 oncology patients: Is nutritional screening capturing hidden malnutrition? *J Cachexia Sarcopenia Muscle.* 2018;9:295-305.
29. August DA, Huhmann MB; American Society for Parenteral, Enteral Nutrition (ASPEN) Board of Directors. ASPEN clinical guidelines: nutrition support therapy during adult anticancer treatment and in hematopoietic cell transplantation. *JPEN J Parenter Enteral Nutr.* 2009;33:472-500.
30. French Speaking Society of Clinical Nutrition, Metabolism (SFNEP). Clinical nutrition guidelines of the French Speaking Society of Clinical Nutrition and Metabolism (SFNEP): Summary of recommendations for adults undergoing non-surgical anticancer treatment. *Dig Liver Dis.* 2014;46:667-74.
31. Hakel-Smith N, Lewis NM. A standardized nutrition care process and language are essential components of a conceptual model to guide and document nutrition care and patient outcomes. *J Am Diet Assoc.* 2004;104:1878-84.
32. Lacey K, Pritchett E. Nutrition Care Process and Model: ADA adopts road map to quality care and outcomes management. *J Am Diet Assoc.* 2003;103:1061-72.
33. <https://www.eatrightpro.org/practice/quality-management/nutrition-care-process>.
34. Sauer AC, Li J, Partridge J, Sulo S. Assessing the impact of nutrition interventions on health and nutrition outcomes of community-dwelling adults: a systematic review. *Nutrition and Dietary Supplements.* 2018;10:45-57.
35. Baldwin C, Spiro A, Ahern R, Emery PW. Oral nutritional interventions in malnourished patients with cancer: a systematic review and meta-analysis. *J Natl Cancer Inst.* 2012;104:371-85.
36. Blackwood HA, Hall CC, Balstad TR, Solheim TS, Fallon M, Haraldsdottir E, Laird BJ. A systematic review examining nutrition support interventions in patients with incurable cancer. *Support Care Cancer.* 2020;28:1877-1889.
37. de van der Schueren MAE, Laviano A, Blanchard H, Jourdan M, Arends J, Baracos VE. Systematic review and meta-analysis of the evidence for oral nutritional intervention on nutritional and clinical outcomes during chemo(radio)therapy: current evidence and guidance for design of future trials. *Ann Oncol.* 2018;29:1141-1153.
38. Lee JLC, Leong LP, Lim SL. Nutrition intervention approaches to reduce malnutrition in oncology patients: a systematic review. *Support Care Cancer.* 2016;24:469-480.
39. Rinninella E, Fagotti A, Cintoni M, Raoul P, Scaletta G, Quagliozzi L, Miggiano GAD, Scambia G, Gasbarrini A, Mele MC. Nutritional Interventions to Improve Clinical Outcomes in Ovarian Cancer: A Systematic Review of Randomized Controlled Trials. *Nutrients.* 2019;11:1404.
40. Richards J, Arensberg MB, Thomas S, Kerr KW, Hegazi R, Bastasch M. Impact of Early Incorporation of Nutrition Interventions as a Component of Cancer Therapy in Adults: A Review. *Nutrients.* 2020;12:3403.
41. Hubbard GP, Elia M, Holdoway A, Stratton RJ. A systematic review of compliance to oral nutritional supplements. *Clin Nutr.* 2012;31:293-312.
42. Engelen MPKJ, Safar AM, Barter T, Koeman F, Deutz NEP. High anabolic potential of essential amino acid mixtures in advanced nonsmall cell lung cancer. *Ann Oncol.* 2015;26:1960-1966.
43. Deutz NE, Safar A, Schützler S, Memelink R, Ferrando A, Spencer H, van Helvoort A, Wolfe RR. Muscle protein synthesis in cancer patients can be stimulated with a specially formulated medical food. *Clin Nutr.* 2011;30:759-68.
44. Klassen P, Cervantes M, Mazurak VC. N-3 fatty acids during chemotherapy: toward a higher level of evidence for clinical application. *Curr Opin Clin Nutr Metab Care.* 2020;23:82-88.
45. Cereda E, Cappello S, Colombo S, Klersy C, Imarisio I, Turri A, Caraccia M, Borioli V, Monaco T, Benazzo M, Pedrazzoli P, Corbella F, Caccialanza R. Nutritional counseling with or without systematic use of oral nutritional supplements in head and neck cancer patients undergoing radiotherapy. *Radiother Oncol.* 2018;126:81-88.
46. Martin B, Cereda E, Caccialanza R, Pedrazzoli P, Tarricone R, Ciani O. Cost-effectiveness analysis of oral nutritional supplements with nutritional counselling in head and neck cancer patients undergoing radiotherapy. *Cost Eff Resour Alloc.* 2021;19:35.
47. Bargetzi L, Brack C, Herrmann J, Bargetzi A, Hersberger L, Bargetzi M, Kaegi-Braun N, Tribolet P, Gomes F, Hoess C, Pavlicek V, Bilz S, Sigrist S, Brändle M, Henzen C, Thomann R, Rutishauser J, Aujesky D, Rodondi N, Donzé J, Laviano A, Stanga Z, Mueller B, Schuetz P. Nutritional support during the hospital stay reduces mortality in patients with different types of cancers: secondary analysis of a prospective randomized trial. *Ann Oncol.* 2021;32:1025-1033.
48. Ho YW, Yeh KY, Hsueh SW, Hung CY, Lu CH, Tsang NM, Wang HM, Hung YS, Chou WC. Impact of early nutrition counseling in head and neck cancer patients with normal nutritional status. *Support Care Cancer.* 2021;29:2777-2785.
49. Antoun S, Lanoy E, Iacovelli R, Albiges-Sauvin L, Loriot Y, Merad-Taoufik M, Fizazi K, di Palma M, Baracos VE, Escudier B. Skeletal muscle density predicts prognosis in patients with metastatic renal cell carcinoma treated with targeted therapies. *Cancer.* 2013 S119:3377-84.
50. Yavuzsen T, Davis MP, Walsh D, LeGrand S, Lagman R. Systematic review of the treatment of cancer-associated anorexia and weight loss. *J Clin Oncol.* 2005;23:8500-11.
51. Blauwhoff-Buskermolen S, Versteeg KS, de van der Schueren MA, den Braver NR, Berkhof J, Langius JA, Verheul HM. Loss of Muscle Mass During Chemotherapy Is Predictive for Poor Survival of Patients With Metastatic Colorectal Cancer. *J Clin Oncol.* 2016;34:1339-44.
52. Xue H, Sawyer MB, Wischmeyer PE, Baracos VE. Nutrition modulation of gastrointestinal toxicity related to cancer chemotherapy: from preclinical findings to clinical strategy. *JPEN J Parenter Enteral Nutr.* 2011;35:74-90.
53. Álvaro Sanz E, Abilés J, Garrido Siles M, Pérez Ruíz E, Alcaide García J, Rueda Domínguez A. Impact of weight loss on cancer patients' quality of life at the beginning of the chemotherapy. *Support Care Cancer.* 2021;29:627-634.
54. Lin T, Yang J, Hong X, Yang Z, Ge T, Wang M. Nutritional status in patients with advanced lung cancer undergoing chemotherapy: a prospective observational study. *Nutr Cancer.* 2020;72:1225-1230.

55. Neoh MK, Abu Zaid Z, Mat Daud ZA, Md Yusop NB, Ibrahim Z, Abdul Rahman Z, Jamhuri N. Changes in Nutrition Impact Symptoms, Nutritional and Functional Status during Head and Neck Cancer Treatment. *Nutrients*. 2020;12:1225.
56. Lodewijckx E, Kenis C, Flamaing J, Debruyne P, De Groof I, Focan C, Cornélis F, Verschaeve V, Bachmann C, Bron D, Luce S, Debugne G, Van den Bulck H, Goeminne JC, Schrijvers D, Geboers K, Petit B, Langenaeken C, Van Rijswijk R, Specenier P, Jerusalem G, Praet JP, Vandendorre K, Lobeke JP, Milisen K, Wildiers H, Decoster L. Unplanned hospitalizations in older patients with cancer: Occurrence and predictive factors. *J Geriatr Oncol*. 2021;12:368-374.
57. Tuca A, Jimenez-Fonseca P, Gascón P. Clinical evaluation and optimal management of cancer cachexia. *Crit Rev Oncol Hematol*. 2013;88:625-36.
58. Argilés JM, López-Soriano FJ, Busquets S. Mechanisms and treatment of cancer cachexia. *Nutr Metab Cardiovasc Dis*. 2013;23:519-24.
59. Caccialanza R, Cereda E, Klersy C, Brugnatelli S, Borioli V, Ferrari A, Caraccia M, Lobascio F, Pagani A, Delfanti S, Aprile G, Reni M, Rimassa L, Melisi D, Cascinu S, Battistini L, Candiloro F, Pedrazzoli P. Early intravenous administration of nutritional support (IVANS) in metastatic gastric cancer patients at nutritional risk, undergoing first-line chemotherapy: study protocol of a pragmatic, randomized, multicenter, clinical trial. *Ther Adv Med Oncol*. 2020;12:1758835919890281.
60. Pistollato F, Forbes-Hernandez TY, Iglesias RC, Ruiz R, Elexpuru Zabaleta M, Dominguez I, Cianciosi D, Quiles JL, Giampieri F, Battino M. Effects of caloric restriction on immunosurveillance, microbiota and cancer cell phenotype: Possible implications for cancer treatment. *Semin Cancer Biol*. 2021;73:45-57.
61. Vidoni C, Ferraresi A, Esposito A, Maheshwari C, Dhanasekaran DN, Mollace V, Isidoro C. Calorie Restriction for Cancer Prevention and Therapy: Mechanisms, Expectations, and Efficacy. *J Cancer Prev*. 2021;26:224-236.
62. Chu MP, Li Y, Ghosh S, Sass S, Smylie M, Walker J, Sawyer MB. Body composition is prognostic and predictive of ipilimumab activity in metastatic melanoma. *J Cachexia Sarcopenia Muscle*. 2020;11:748-755.
63. Roch B, Coffy A, Jean-Baptiste S, Palaysi E, Daires JP, Pujol JL, Bommar S. Cachexia - sarcopenia as a determinant of disease control rate and survival in non-small lung cancer patients receiving immune-checkpoint inhibitors. *Lung Cancer*. 2020;143:19-26.
64. Floris G, Richard F, Hamy AS, Jongen L, Wildiers H, Ardui J, Punie K, Smeets A, Berteloot P, Vergote I, De Croze D, Meseure D, Salomon A, Laé M, Reyat F, Biganzoli E, Neven P, Desmedt C. Body Mass Index and Tumor-Infiltrating Lymphocytes in Triple-Negative Breast Cancer. *J Natl Cancer Inst*. 2021;113:146-153.
65. Johannet P, Sawyers A, Qian Y, Kozloff S, Gulati N, Donnelly D, Zhong J, Osman I. Baseline prognostic nutritional index and changes in pretreatment body mass index associate with immunotherapy response in patients with advanced cancer. *J Immunother Cancer*. 2020;8:e001674.
66. Shoji F, Takeoka H, Kozuma Y, Toyokawa G, Yamazaki K, Ichiki M, Takeo S. Pretreatment prognostic nutritional index as a novel biomarker in non-small cell lung cancer patients treated with immune checkpoint inhibitors. *Lung Cancer*. 2019;136:45-51.
67. Lu Z, Fang Y, Liu C, Zhang X, Xin X, He Y, Cao Y, Jiao X, Sun T, Pang Y, Wang Y, Zhou J, Qi C, Gong J, Wang X, Li J, Tang L, Shen L. Early Interdisciplinary Supportive Care in Patients With Previously Untreated Metastatic Esophagogastric Cancer: A Phase III Randomized Controlled Trial. *J Clin Oncol*. 2021;39:748-756.
68. Rossi R, Serra P, Suzzi M, Guerra D, Bilotta S, Ricci M, Pallotti MC, Ibrahim T, Frassinetti GL, Zavou V, Nanni C, Altini M, Maltoni M. The challenge for nutritional care in a cancer center: The need for integration between clinical nutritionist, oncologist, and palliative care physician. *Curr Probl Cancer*. 2020;44:100618.
69. Weimann A, Braga M, Carli F, Higashiguchi T, Hübner M, Klek S, Laviano A, Ljungqvist O, Lobo DN, Martindale R, Waitzberg DL, Bischoff SC, Singer P. ESPEN guideline: Clinical nutrition in surgery. *Clin Nutr*. 2017;36:623-650.
70. Soeters PB, Schols AM. Advances in understanding and assessing malnutrition. *Curr Opin Clin Nutr Metab Care* 2009;12:487e94.
71. Yeh DD, Fuentes E, QURashi SA, Cropano C, Kaafarani H, Lee J, et al. Adequate nutrition may get you home: effect of caloric/protein deficits on the discharge destination of critically ill surgical patients. *JPEN J Parenter Enteral Nutr*. 2016;40:37e44.
72. <https://erasociety.org/guidelines/>
73. Grieco M, Lorenzon L, Pernazza G, Carlini M, Brescia A, Santoro R, Crucitti A, Palmieri RM, Santoro E, Stipa F, Sacchi M, Persiani R. Impact of implementation of the ERAS program in colorectal surgery: a multi-center study based on the "Lazio Network" collective database. *Int J Colorectal Dis*. 2020;35:445-453.
74. Gustafsson UO, Scott MJ, Hubner M, Nygren J, Demartines N, Francis N, Rockall TA, Young-Fadok TM, Hill AG, Soop M, de Boer HD, Urman RD, Chang GJ, Fichera A, Kessler H, Grass F, Whang EE, Fawcett WJ, Carli F, Lobo DN, Rollins KE, Balfour A, Baldini G, Riedel B, Ljungqvist O. Guidelines for Perioperative Care in Elective Colorectal Surgery: Enhanced Recovery After Surgery (ERAS®) Society Recommendations: 2018. *World J Surg*. 2019;43:659-695.
75. Low DE, Allum W, De Manzoni G, Ferri L, Immanuel A, Kuppusamy M, Law S, Lindblad M, Maynard N, Neal J, Pramesh CS, Scott M, Mark Smithers B, Addor V, Ljungqvist O. Guidelines for Perioperative Care in Esophagectomy: Enhanced Recovery After Surgery (ERAS®) Society Recommendations. *World J Surg*. 2019;43:299-330.
76. Melloul E, Lassen K, Roulin D, Grass F, Perinel J, Adham M, Wellge EB, Kunzler F, Besselink MG, Asbun H, Scott MJ, Dejoch CHC, Vrochides D, Aloia T, Izbickei JR, Demartines N. Guidelines for Perioperative Care for Pancreatoduodenectomy: Enhanced Recovery After Surgery (ERAS) Recommendations 2019. *World J Surg*. 2020;44:2056-2084.
77. Nelson G, Bakum-Gamez J, Kalogera E, Glaser G, Altman A, Meyer LA, Taylor JS, Iniesta M, Lasala J, Mena G, Scott M, Gillis C, Elias K, Wijk L, Huang J, Nygren J, Ljungqvist O, Ramirez PT, Dowdy SC. Guidelines for perioperative care in gynecologic/oncology: Enhanced Recovery After Surgery (ERAS) Society recommendations-2019 update. *Int J Gynecol Cancer*. 2019;29:651-668.
78. Schwegler I, von Holzen A, Gutzwiller JP, Schlumpf R, Muhlebach S, Stanga S. Nutritional risk is a clinical predictor of postoperative mortality and morbidity in surgery for colorectal cancer. *Br J Surg*. 2010;97:92e7.
79. Durkin MT, Mercer KG, McNulty MF, Phipps L, Upperton J, Giles M, et al. Vascular surgical society of great britain and Ireland: contribution of malnutrition to postoperative morbidity in vascular surgical patients. *Br J Surg*. 1999;86:702.
80. Pikul J, Sharpe MD, Lowndes R, Ghent CN. Degree of preoperative malnutrition is predictive of postoperative morbidity and mortality in liver trans-plant recipients. *Transplantation*. 1994;57:469e72.
81. Beattie AH, Prach AT, Baxter JP, Pennington CR. A randomised controlled trial evaluating the use of enteral nutritional supplements postoperatively in malnourished surgical patients. *Gut* 2000;46:813e8.
82. Marimuthu K, Varadhan KK, Ljungqvist O, Lobo DN. A meta-analysis of the effect of combinations of immune modulating nutrients on outcome in patients undergoing major open gastrointestinal surgery. *Ann Surg*. 2012;255:1060e8.
83. Lambert E, Carey S. Practice guideline recommendations on perioperative fasting. A systematic review. *JPEN J Parenter Enteral Nutr*. 2016;40:1158e65.
84. Osland E, Yunus RM, Khan S, Memon MA. Early versus traditional post-operative feeding in patients undergoing resectional gastrointestinal surgery: a meta-analysis. *JPEN J Parenter Enteral Nutr*. 2011;35:473e87.
85. Ljungqvist O, Scott M, Fearon KC. Enhanced Recovery After Surgery: A Review. *JAMA Surg*. 2017;152:292-298.
86. Cotogni P. Enteral versus parenteral nutrition in cancer patients: Evidences and controversies. *Ann. Palliat. Med*. 2016;1:42-49.
87. Cotogni P, Ossola M, Passera R, Monge T, Fadda M, De Francesco A, Bozzetti F. Home parenteral nutrition versus artificial hydration in malnourished patients with cancer in palliative care: a prospective, cohort survival study. *BMJ Support Palliat Care*. 2020;bmjspcare-2020-002343.
88. Gavazzi C, Colatruoglio S, Valoriani F, Mazzaferro V, Sabbatini A, Biffi R, Mariani L, Miceli R. Impact of home enteral nutrition in malnourished patients with upper gastrointestinal cancer: A multicentre randomised clinical trial. *Eur J Cancer*. 2016;64:107-12.
89. Qiu M, Zhou YX, Jin Y, Wang ZX, Wei XL, Han HY, Ye WF, Zhou ZW, Zhang DS, Wang FH, Li YH, Yang DJ, Xu RH. Nutrition support can bring survival benefit to high nutrition risk gastric cancer patients who received chemotherapy. *Support Care Cancer*. 2015; 23:1933-9.
90. Caccialanza R, De Lorenzo F, Gianotti L, Zagonel V, Gavazzi C, Farina G, Cotogni P, Cinieri S, Cereda E, Marchetti P, Nardi M, Iannelli E, Santangelo C, Tracò F, Pinto C, Pedrazzoli P. Nutritional support for cancer patients: still a neglected right? *Support Care Cancer*. 2017;25:3001-4.
91. Libro Bianco AIOM 2020 - X edizione; <https://www.aiom.it/libro-bianco-2020-x-edizione/>
92. Donini LM, Leonardi F, Rondanelli M, Banderali G, Battino M, Bertoli E, Bordini A, Brighenti F, Caccialanza R, Cairella G, Caretto A, Cena H, Gambarara M, Gentile MG, Giovannini M, Lucchin L, Migliaccio P, Nicastro F, Pisanis F, Piretta L, Radrizzani D, Roggi C, Rotilio G, Scalfi L, Vettor R, Vignati F, Battistini NC, Muscaritoli M. The Domains of Human Nutrition: The Importance of Nutrition Education in Academia and Medical Schools. *Front Nutr*. 2017;4:2.
93. Hark LA. Lessons learned from nutrition curricular enhancements. *Am J Clin Nutr*. 2006;83:968S-970S.

94. Ball L, Crowley J, Laur C, Rajput-Ray M, Gillam S, Ray S. Nutrition in medical education: reflections from an initiative at the University of Cambridge. *J Multidiscip Healthc.* 2014;7:209-15.
95. Cotogni P, Pedrazzoli P, De Waele E, Aprile G, Farina G, Stragliotto S, De Lorenzo F, Caccialanza R. Nutritional Therapy in Cancer Patients Receiving Chemoradiotherapy: Should We Need Stronger Recommendations to Act for Improving Outcomes? *J Cancer.* 2019;10:4318-4325.
96. Mohty R, Savani M, Brissot E, Mohty M. Nutritional Supplements and Complementary/Alternative Medications in Patients With Hematologic Diseases and Hematopoietic Stem Cell Transplantation. *Transplant Cell Ther.* 2021;27:467-473.
97. Wieland LS, Moffet I, Shade S, Emadi A, Knott C, Gorman EF, D'Adamo C. Risks and benefits of antioxidant dietary supplement use during cancer treatment: protocol for a scoping review. *BMJ Open.* 2021;11:e047200.
98. Caccialanza R, Cereda E, Pinto C, Cotogni P, Farina G, Gavazzi C, Gandini C, Nardi M, Zagonel V, Pedrazzoli P. Awareness and consideration of malnutrition among oncologists: Insights from an exploratory survey. *Nutrition.* 2016;32:1028-32.
99. Vidoni C, Ferraresi A, Esposito A, Maheshwari C, Dhanasekaran DN, Mollace V, Isidoro C. Calorie Restriction for Cancer Prevention and Therapy: Mechanisms, Expectations, and Efficacy. *J Cancer Prev.* 2021;26:224-236
100. De Waele E, Mattens S, Honoré PM, Spapen H, De Grève J, Pen JJ. Nutrition therapy in cachectic cancer patients. The Tight Caloric Control (TiCaCo) pilot trial. *Appetite.* 2015;91:298-301.
101. Arensberg MB, Richards J, Benjamin J, Kerr K, Hegazi R. Opportunities for Quality Improvement Programs (QIPs) in the Nutrition Support of Patients with Cancer. *Healthcare (Basel).* 2020;8:227.
102. Bland KA, Kouw IWK, van Loon LJC, Zopf EM, Fairman CM. Exercise-Based Interventions to Counteract Skeletal Muscle Mass Loss in People with Cancer: Can We Overcome the Odds? *Sports Med.* 2022 Feb 4. doi: 10.1007/s40279-021-01638-z. Epub ahead of print.
103. Prado CM, Laviano A, Gillis C, Sung AD, Gardner M, Yalcin S, Dixon S, Newman SM, Bastasch MD, Sauer AC, Hegazi R, Chasen MR. Examining guidelines and new evidence in oncology nutrition: a position paper on gaps and opportunities in multimodal approaches to improve patient care. *Support Care Cancer.* 2021 Nov 23. doi: 10.1007/s00520-021-06661-4. Epub ahead of print.