

The BEHOMO project: Λ Lemaître-Tolman-Bondi N -body simulations[★]

V. Marra^{1,2,3} , T. Castro^{1,2,4} , D. Camarena⁵ , S. Borgani^{1,2,4,6} , and A. Ragagnin^{7,1,2} 

¹ INAF – Osservatorio Astronomico di Trieste, 34131 Trieste, Italy
e-mail: valerio.marra@me.com

² IFPU – Institute for Fundamental Physics of the Universe, 34151 Trieste, Italy

³ Núcleo Cosmo-ufes & Departamento de Física, Universidade Federal do Espírito Santo, 29075-910, Vitória, ES, Brazil

⁴ INFN – Sezione di Trieste, 34100 Trieste, Italy

⁵ PPGCosmo, Universidade Federal do Espírito Santo, 29075-910 Vitória, ES, Brazil

⁶ Dipartimento di Fisica, Sezione di Astronomia, Università di Trieste, 34143 Trieste, Italy

⁷ Dipartimento di Fisica e Astronomia “Augusto Righi”, Alma Mater Studiorum Università di Bologna, 40129 Bologna, Italy

Received 14 March 2022 / Accepted 25 May 2022

ABSTRACT

Context. Our universe may feature large-scale inhomogeneities and anisotropies that cannot be explained by the standard model of cosmology, that is, the homogeneous and isotropic Friedmann-Lemaître-Robertson-Walker metric, on which the Λ cold dark matter model is built, may not accurately describe observations. Currently, there is not a satisfactory understanding of the evolution of the large-scale structure on an inhomogeneous background.

Aims. We have launched the cosmology beyond homogeneity and isotropy (BEHOMO) project to study the inhomogeneous Λ Lemaître-Tolman-Bondi model with the methods of numerical cosmology. Understanding the evolution of the large-scale structure is a necessary step in constraining inhomogeneous models with present and future observables and placing the standard model on more solid ground.

Methods. We perform Newtonian N -body simulations, whose accuracy in describing the background evolution is checked against the general relativistic solution. The large-scale structure of the corresponding Λ cold dark matter simulation is also validated.

Results. We obtain the first set of simulations of the Λ Lemaître-Tolman-Bondi model ever produced. The data products consist of 11 snapshots between redshift 0 and 3.7 for each of the 68 simulations that have been performed, together with halo catalogs and lens planes relative to 21 snapshots, between redshift 0 and 4.2, for a total of approximately 180 TB of data.

Conclusions. We plan to study the growth of perturbations at the linear and nonlinear level, gravitational lensing, and cluster abundances and properties.

Key words. large-scale structure of Universe – gravitation – cosmology: theory – cosmological parameters

1. Introduction

Several anomalous signals in cosmological observables have emerged since the establishment of the Λ cold dark matter (CDM) model as the standard model of cosmology more than two decades ago. Particularly relevant here are the *Hubble* crisis, the cosmic-microwave-background (CMB) anomalies, and the cosmic dipoles and bulk flows (see [Perivolaropoulos & Skara 2022](#), and references therein). Such signals indicate anomalies that challenge the Λ CDM model and its foundations. One may then ask if the universe features large-scale inhomogeneities and anisotropies that cannot be explained by the standard paradigm, or, equivalently, if the homogeneous and isotropic Friedmann-Lemaître-Robertson-Walker (FLRW) metric, on which the Λ CDM model is built, does not accurately describe observations. This constitutes the motivation for studying the universe without assuming homogeneity and isotropy, trying instead to reconstruct the metric directly from observations ([Stebbins 2012](#)).

According to the standard reasoning, the validity of the FLRW metric is a consequence of the observed isotropy of the

universe and the Copernican principle, which states that humans are not special observers. Here, however, we are not advocating that the universe is inhomogeneous and humans are special, rather that the scale at which there is homogeneity and isotropy could be larger than the commonly thought ≈ 100 Mpc ([Scrimgeour et al. 2012](#); [Laurent et al. 2016](#); [Ntelis et al. 2017](#)), that is, the cosmological principle may be valid at grander scales (see Sect. 8 of [Abdalla et al. 2022](#)). We note that this scenario is not necessarily at odds with the observed approximate isotropy of the CMB (see the discussion of the Ehlers-Geren-Sachs theorem in [Rasanen 2009](#)).

Inhomogeneous cosmology is undeniably a challenging subject as it would require a considerable theoretical and numerical effort to study its phenomenology. The absence of an FLRW background makes it particularly difficult to study early universe physics and predict, for instance, the CMB power spectrum. Therefore, in order to present a viable program, here we consider a subclass of inhomogeneous cosmologies. The basic requirement is that, at early times, one recovers a near-FLRW metric such that the standard inflationary paradigm is maintained and the physics that leads to the CMB remains basically unchanged. In other words, we consider a standard cosmology endowed with a nonstandard large-scale structure that is dominated by

[★] Data can be obtained upon request. Further information is available at <https://valerio-marra.github.io/BEHOMO-project>

growing modes. This requirement effectively imposes restrictions on the free functions that characterize inhomogeneous metrics, considerably simplifying both analysis and statistical inference. We name these inhomogeneous models “early-FLRW cosmologies”.

Clearly, early-FLRW cosmologies are constrained by CMB observations. Indeed, as shown by Valkenburg (2012a), perturbations at the last scattering surface of present-day contrast ≈ 0.1 and size ≈ 1 Gpc would produce temperature fluctuations of $\Delta T \approx 50 \mu\text{K}$ on a scale of $\approx 5^\circ$. Similarly, too strong structures along the line of sight at $z \lesssim 1$ would be detected via the integrated Sachs-Wolfe (ISW) effect: a present-day contrast ≈ 0.1 and size ≈ 300 Mpc would produce temperature fluctuations of order $\Delta T \approx 20\text{--}30 \mu\text{K}$ (see, for instance, Zibin 2021; Nadathur et al. 2014). To put this figure in perspective, the famous cold spot of the CMB features $\Delta T \approx 70 \mu\text{K}$ across a 5° region (Vielva 2010). Therefore, early-FLRW cosmologies are, at most, mildly nonlinear large-scale perturbations of the FLRW metric.

The inhomogeneities of early-FLRW cosmologies may be regarded as a particular type of primordial non-Gaussianity. Their distinguishing features are nonstandard amplitudes and phases. Indeed, they are characterized by bulk flows and coherent perturbations in the energy content of the universe at arbitrarily large scales. In other words, large-scale homogeneity and isotropy are violated by the phases of these extra modes, so observations depend on the position of the observer and the notion of an average FLRW observer ceases to be meaningful (Kolb et al. 2010). Specifically, large-scale inhomogeneities alter observations both because they affect photon geodesics and because the observer’s local space-time is perturbed. Of course, this is also true within the Λ CDM model, but there the size of this effect is constrained by the standard perturbation spectrum. For example, cosmic variance on local measurements of H_0 is expected to be at most 1% within Λ CDM (Camarena & Marra 2018).

The background evolution of early-FLRW cosmologies, that is, the evolution neglecting standard primordial perturbations, can be studied via exact solutions of general relativity. If considering more general scenarios, one can use linear-perturbation theory or simulations via codes that use general relativistic (GR) perturbation theory such as *gevolution* (Adamek et al. 2016) and *CONCEPT* (Dakin et al. 2021). A general consequence of spatial gradients is the occurrence of background shear, that is, the fact that the universe expands in an anisotropic way.

The scenario becomes more involved once primordial perturbations are added to the inhomogeneous background. First, there is the issue of the backreaction of small-scale perturbations on the average dynamics of the (possibly inhomogeneous) universe. Here, we assume that backreaction gives a negligible effect, as tested via GR simulations (Giblin et al. 2016; Bentivegna & Bruni 2016; Adamek et al. 2019; Macpherson et al. 2019). An overview of the backreaction proposal is given in Sect. 2.12. Second, because of spatial gradients, the standard primordial perturbations are coupled at first order so that standard perturbation theory does not hold in an inhomogeneous background. Within the spherical Lemaître-Tolman-Bondi (LTB) space-time, this issue has been tackled by Zibin (2008), Clarkson et al. (2009), Dunsby et al. (2010), February et al. (2014), and Meyer et al. (2015) via the numerical integration of the system of coupled equations and by Nishikawa et al. (2012) via second-order perturbation theory. It was concluded that the effect of spatial gradients could have an impact on the growth of perturbations. However, a full per-

turbation theory and modeling is missing, hampering comparison with perturbation observables – the focus of current and next-generation surveys such as the Dark Energy Survey (DES, Abbott et al. 2022)¹, the Dark Energy Spectroscopic Instrument (DESI, Aghamousa et al. 2016)², the Javalambre Physics of the Accelerating universe Astrophysical Survey (J-PAS, Bonoli et al. 2021)³, the Legacy Survey of Space and Time (LSST, Abate et al. 2012)⁴, *Euclid* (Amendola et al. 2018)⁵, and the Square Kilometre Array (SKA, Braun et al. 2015)⁶.

Here we introduce the cosmology beyond homogeneity and isotropy (BEHOMO) project. We propose a program that aims at addressing the modeling of linear and nonlinear perturbations and understanding the rich phenomenology of early-FLRW cosmologies. In order to do so, the basic idea is to apply the methods of numerical cosmology, as pioneered by Alonso et al. (2010, 2012). The ultimate goal is to confront arbitrarily early-FLRW inhomogeneous models with data from next-generation surveys. The idea is to adopt Newtonian N -body simulations, whose accuracy in describing the background evolution shall be checked with GR codes. The basic methodology is to feed state-of-the-art N -body codes such as *GADGET* (Springel et al. 2021) with special early-FLRW initial conditions so that early-FLRW cosmologies can reach the same resolution of standard Λ CDM simulations in approximately the same CPU time (except for the most nonlinear cases). This program will bring the field of inhomogeneous cosmologies into the era of precision cosmology, on par with the Λ CDM model.

In this paper we present the first suite of simulations for the simplest possible early-FLRW cosmologies: spherically symmetric ALTB models. We use the LTB metric to model a spherical inhomogeneity on top of the standard Λ CDM model. Though still a toy model, on a first approximation one can regard the spatial gradients of the ALTB model as an archetype for more realistic structures with background shear. We consider a set of high-resolution simulations with varying inhomogeneity size and depth, the two main physical parameters that describe such a structure. This will allow us to understand and model the effect of spatial gradients on the evolution of perturbations, which is necessary to confront inhomogeneous cosmologies with perturbation observables such as redshift-space distortions, weak lensing, and cluster abundances. As said earlier, the grand goal is to study and then constrain the phenomenology of these beyond- Λ CDM inhomogeneities with observations (Valkenburg et al. 2014; Redlich et al. 2014; Camarena et al. 2021).

In this presentation paper we review the ALTB model in Sect. 2, discuss the numerical details of the inhomogeneous N -body simulations and their data products in Sect. 3, present the results of the simulations in Sect. 4, and discuss the road map of the BEHOMO project⁷ in Sect. 5. Regarding notation: we use “LTB metric” as opposed to “FLRW metric” but “ALTB model” as opposed to “ Λ CDM model”; quantities without explicit radial dependence are relative to the FLRW background if pertinent; bold denotes vectors; and $c = 1$ is assumed unless stated otherwise.

¹ <https://www.darkenergysurvey.org>

² <https://www.desi.lbl.gov>

³ <http://www.j-pas.org>

⁴ <https://www.lsst.org>

⁵ <https://www.euclid-ec.org>

⁶ <https://www.skatelescope.org>

⁷ [valerio-marra.github.io/BEHOMO-project](https://github.com/valerio-marra/BEHOMO-project)

2. The ALTB model

We consider early-FLRW ALTB models, that is, the Λ CDM model endowed with a spherical inhomogeneity, which is described via the exact LTB solution of Einstein's equations. As we are considering early-FLRW cosmologies, this model is fully specified by the radial profile function, whose basic parameters are the effective radius and depth of the inhomogeneity.

In this section, after reviewing the formalism and dynamics of the LTB metric, we connect with the more standard Newtonian-perturbed FLRW metric and discuss the historical relevance of LTB models, putting coherently together results from many different papers.

2.1. Metric

In the comoving and synchronous gauge, the spherically symmetric LTB metric can be written as

$$ds^2 = -dt^2 + \frac{a_{\parallel}^2(t, r)}{1 - k(r)r^2} dr^2 + a_{\perp}^2(t, r)r^2 d\Omega^2, \quad (1)$$

where the longitudinal (a_{\parallel}) and perpendicular (a_{\perp}) scale factors are related by $a_{\parallel} = (a_{\perp}r)'$, and a prime denotes partial derivation with respect to the coordinate radius r . We also adopt the alternative notation $Y(t, r) \equiv a_{\perp}r$ so that $Y' \equiv a_{\parallel}$. In the limit $k \rightarrow \text{const}$ and $a_{\perp} = a_{\parallel} = a$, we recover the FLRW metric, but here $k(r)$ is a free function named the LTB curvature function.

The two scale factors define two different Hubble rates:

$$H_{\perp}(t, r) \equiv \frac{\dot{a}_{\perp}}{a_{\perp}} = \frac{\dot{Y}}{Y}, \quad (2)$$

$$H_{\parallel}(t, r) \equiv \frac{\dot{a}_{\parallel}}{a_{\parallel}} = \frac{\dot{Y}'}{Y'}, \quad (3)$$

where a dot denotes partial derivation with respect to the coordinate time t . This has important implications when confronting these models with observations. For example, cosmic chronometers probe dz/dt and so H_{\parallel} (see Eq. (26)); radial baryon acoustic oscillations (BAOs) also probe H_{\parallel} , but angular BAOs and supernovae probe the angular and luminosity distance, respectively, and so a_{\perp} and H_{\perp} (see Eqs. (26)–(29)). Combining these observables can then place interesting constraints on the background shear (García-Bellido & Haugboelle 2009):

$$\Sigma(t, r) = \frac{2}{3} [H_{\parallel}(t, r) - H_{\perp}(t, r)]. \quad (4)$$

As said earlier, the spatial gradient of the ALTB model is an archetype for more realistic structures.

2.2. Dynamics

By solving Einstein's equations for an irrotational dust source in the presence of a cosmological constant Λ , one obtains the equivalent of the Friedmann equation, which can be written as (Enqvist 2008; Marra & Paakkonen 2012, Appendix B)

$$H_{\perp}^2(t, r) = \frac{8\pi G}{3} \rho_m^e(t, r) + \frac{8\pi G}{3} \rho_{\Lambda} - \frac{k(r)}{a_{\perp}^2(t, r)}, \quad (5)$$

where $\rho_{\Lambda} = \Lambda/8\pi G$, and the last term is the Euclidean average of the spatial Ricci scalar (the trace of the Ricci tensor of the

spatial metric on the hypersurface of constant t):

$$\frac{\mathcal{R}}{2} = \frac{(kr^2Y)'}{Y^2Y'} = \frac{k}{a_{\perp}^2} + 2\frac{k}{a_{\perp}a_{\parallel}} + \frac{k'r}{a_{\perp}a_{\parallel}}, \quad (6)$$

$$\frac{\mathcal{R}^e}{6} = \frac{1}{6} \frac{\int_0^r \mathcal{R} dV_e}{V_e} = \frac{k(r)}{a_{\perp}^2} \xrightarrow{\text{FLRW}} \frac{k = \text{const}}{a^2}, \quad (7)$$

where the Euclidean volume element – obtained by setting $k = 0$ in Eq. (1) – is used:

$$V_e(t, r) = \int_0^r dV_e = 4\pi \int_0^r Y^2 Y' d\hat{r} = \frac{4\pi}{3} Y^3. \quad (8)$$

The fact that a Euclidean rather than proper average is used leads to backreaction, as discussed in Sect. 2.13. Similarly, Eq. (5) features the Euclidean average of the local matter density, ρ_m :

$$\rho_m(t, r) = \frac{F'(r)}{4\pi Y^2(t, r) Y'(t, r)}, \quad (9)$$

$$F(r) = \int_0^r \rho_m(t, r) dV_e, \quad (10)$$

$$\rho_m^e(t, r) = \frac{F(r)}{V_e} \xrightarrow{\text{FLRW}} \rho_m(t), \quad (11)$$

where the LTB mass function $F(r)$, a constant of integration, is another free function that gives the total gravitating mass up to the shell of coordinate radius r . The local density ρ_m satisfies the continuity equation $\dot{\rho}_m + \theta \rho_m = 0$, where $\theta = H_{\parallel} + 2H_{\perp}$ is the expansion scalar. We note that, as the source is pressureless dust, without pressure gradients, both $F(r)$ and $k(r)$ do not depend on t . The case of the Lemaître metric with pressure is presented in Yamamoto et al. (2016).

Similarly to FLRW, one can interpret the curvature function as related to the total energy per unit of mass of the shell at coordinate radius r :

$$E(r) \equiv -\frac{kr^2}{2} = \frac{1}{2} \dot{Y}^2(t, r) - \frac{GF(r)}{Y(t, r)} - \frac{1}{6} \Lambda Y^2(t, r), \quad (12)$$

where the first term of the energy function E is the kinetic energy per unit of mass of the shell r , the second term is the potential energy per unit of mass due to the total gravitating mass up to the shell r , and the third term is the usual contribution from the cosmological constant (as in the de Sitter-Schwarzschild metric). We note that, thanks to spherical symmetry, one is able to define a potential energy also in cases far away from nearly Newtonian ones and that the potential energy is related to the curvature (Bondi 1947).

Next, similarly to FLRW, one can rewrite Eq. (5) using the equivalent of the density parameters in FLRW:

$$\frac{H_{\perp}^2(t, r)}{H_{\perp 0}^2(r)} = \Omega_{m0}(r) \frac{a_{\perp 0}^3}{a_{\perp}^3} + \Omega_{\Lambda 0}(r) + \Omega_{k0}(r) \frac{a_{\perp 0}^2}{a_{\perp}^2}, \quad (13)$$

where the subscript 0 denotes a quantity evaluated at the present time, t_0 , and

$$\Omega_{m0}(r) = \frac{2GF(r)}{r^3 a_{\perp 0}^3 H_{\perp 0}^2} \quad \Omega_m(t, r) = \Omega_{m0}(r) \frac{H_{\perp 0}^2 a_{\perp 0}^3}{H_{\perp}^2 a_{\perp}^3}, \quad (14)$$

$$\Omega_{\Lambda 0}(r) = \frac{\Lambda}{3H_{\perp 0}^2} \quad \Omega_{\Lambda}(t, r) = \Omega_{\Lambda 0}(r) \frac{H_{\perp 0}^2}{H_{\perp}^2}, \quad (15)$$

$$\Omega_{k0}(r) = -\frac{k(r)}{a_{\perp 0}^2 H_{\perp 0}^2} \quad \Omega_k(t, r) = \Omega_{k0}(r) \frac{H_{\perp 0}^2 a_{\perp 0}^2}{H_{\perp}^2 a_{\perp}^2}, \quad (16)$$

which satisfy $\Omega_m(t, r) + \Omega_{\Lambda}(t, r) + \Omega_k(t, r) = 1$.

2.3. Free functions and gauge fixing

Equation (13) can be used to determine the age of the universe at a radial coordinate r :

$$t - t_{bb}(r) = \frac{1}{H_{\perp 0}(r)} \int_0^{\frac{a_{\perp}(t,r)}{a_{\perp 0}(r)}} \frac{dx}{\sqrt{\Omega_{m0}(r)/x + \Omega_{\Lambda 0}(r)x^2 + \Omega_{k0}(r)}}, \quad (17)$$

where the big bang function $t_{bb}(r)$ is another arbitrary function, which sets the time since the big bang ($a_{\perp} = 0$). If it were $t'_{bb}(r) \neq 0$, the initial singularity would have happened at different times for different shells so that large inhomogeneities would develop in the past, as can be seen from Eq. (9) with $Y \rightarrow 0$. This clearly signals the presence of decaying modes, which would be strongly in contradiction with the inflationary paradigm and are excluded by the choice of a simultaneous big bang (Silk 1977; Biswas et al. 2007; Zibin 2008).

Summarizing, we have seen that the LTB inhomogeneity is specified by three arbitrary functions, $F(r)$, $k(r)$, and $t_{bb}(r)$, which are related, together with $a_{\perp 0}$, by Eq. (17) so that one is not independent. Moreover, one can always make a redefinition of the radial coordinate. Common gauge fixing are $F(r) \propto r^3$ or $a_{\perp 0} = \text{constant}$. It is then clear that one can choose $t_{bb}(r)$ and $k(r)$ as the free functions that specify the model.

Each gauge fixing has pros and cons. For example, $F(r) \propto r^3$ excludes the possibility that there is pure vacuum in some radial interval, and the moment of shell crossing – the time at which $Y' = 0$ so that $g_{rr} = 0$ – clearly depends on the gauge adopted. The numerical codes that we use, VoidDistances2020 (Valkenburg 2012b) and FalconIC (Valkenburg & Hu 2015), adopt the choice $F(r) = 4\pi M_0^4 r^3/3$, where M_0 is an arbitrary mass scale.

2.4. Compensated inhomogeneity profile

As discussed earlier, we consider early-FLRW cosmologies in agreement with the standard scenario of inflation and, therefore, we set

$$t_{bb}(r) = 0. \quad (18)$$

We are then left with the curvature function. Here, we consider the case of an LTB inhomogeneity that matches exactly with the FLRW metric at the finite radius r_b and not only asymptotically. This simplified approach is convenient for the purposes of this work because it allows us to robustly simulate the LTB inhomogeneity inside of a bigger FLRW box. The curvature function is modeled according to the monotonic profile:

$$k(r) = k_b + (k_c - k_b) W_3(r/r_b), \quad (19)$$

where r_b is the coordinate radius of the spherical inhomogeneity, k_b and k_c are the curvature outside and at the center of the inhomogeneity, respectively, and W_3 is the function

$$W_n(x) = \begin{cases} e^{-x^n/(1-x)} & \text{for } 0 \leq x < 1 \\ 0 & \text{for } x \geq 1 \end{cases}. \quad (20)$$

The function $W_n(x)$ interpolates from 1 to 0 when x varies from 0 to 1 while remaining differentiable, which implies that that $k(r)$ is C^∞ everywhere. It is $d^m W_n/dx^m|_0 = 0$ for $0 < m < n$, so that there is no cusp at the center. In the limit $n \rightarrow \infty$, $W_n(x)$ approaches the top-hat function.

For $r \geq r_b$ the curvature profile equals the curvature k_b of the background FLRW such that for $r \geq r_b$ one exactly recovers the

background Λ CDM model: $a_{\perp} = a_{\parallel} = a$. We can then define the local density contrast according to:

$$\delta(t, r) = \frac{\rho_m(t, r)}{\rho_m(t)} - 1, \quad (21)$$

and the (integrated) mass density contrast according to

$$\Delta(t, r) = \frac{\int_0^r \delta(t, \bar{r}) dV_e}{V_e} = \frac{\Omega_m(t, r) H_{\perp}^2(t, r)}{\Omega_m(t) H^2(t)} - 1, \quad (22)$$

where we used the Euclidean average in agreement with Eq. (5). We note that $\Delta(t, r = 0) = \delta(t, r = 0)$. We denote with δ_0 the central contrast today, which is directly related to k_c (see Eq. (35) for the linear relation at early times).

We also note that, because of the matching, it is by construction $\Delta(t, r = r_b) = \delta(t, r = r_b) = 0$. This implies that the central underdensity or overdensity at $0 \leq r < r_t$, determined by the curvature k_c at the center, is automatically compensated by a surrounding overdense or underdense shell at $r_t \leq r < r_b$, where r_t is the transition radius at which $\delta = 0$. A compensating overdense or underdense region is an expected feature of the standard large-scale structure: voids are surrounded by sheets and filaments, and superclusters by voids. We note that it is $r_t = r_t(t)$, as in Eq. (9) the volume element at the denominator is time dependent.

2.5. Physical and light cone distances

The comoving radial coordinate, r , because of the freedom in redefining it, does not possess physical meaning. On the other hand, the proper distance between r_1 and r_2 ($d\ell^2 = d\Omega^2 = 0$ in Eq. (1)) is

$$d_p = \int_{r_1}^{r_2} \frac{Y'(t, r)}{\sqrt{1 - k(r)r^2}} dr \simeq Y(t, r_2) - Y(t, r_1), \quad (23)$$

where the approximation holds for

$$E \sim k(r)r^2 = \frac{Y^2}{a_{\perp}^2/k} = \left(\frac{Y}{\text{curv. radius}} \right)^2 \ll 1. \quad (24)$$

Inside the inhomogeneity ($r < r_b$) the curvature radius is $\approx a_{\perp}/\sqrt{k_c}$, while outside the LTB patch it is $a/\sqrt{k_b}$. We consider models with $k_b = 0$ so that the corrections to Eq. (23) will be due only to the inhomogeneity. We will see that these corrections are also negligible for gigaparsec-scale inhomogeneities ($E \ll 1$; see Fig. 1).

Using Eq. (23) we can then define the corresponding FLRW comoving coordinate as

$$\chi = \frac{d_p}{a(t)} \stackrel{E \ll 1}{=} \frac{Y(t, r)}{a(t)}, \quad (25)$$

so that the FLRW and LTB physical distances coincide (note that $Y' \neq 0$). Thanks to the adopted matching condition, it is $\chi = r$ for $r \geq r_b$. The coordinate χ is the one used in the numerical simulations.

Observationally, the time, t , and radius, r , as a function of the redshift, z , are determined on the past light cone of the central observer by the differential equations for radial null geodesics (see, e.g., Chung & Romano 2006; Enqvist 2008):

$$\frac{dt}{dz} = -\frac{1}{(1+z)H_{\parallel}}, \quad (26)$$

$$\frac{dr}{dz} = \frac{\sqrt{1 - kr^2}}{(1+z)a_{\parallel}H_{\parallel}}, \quad (27)$$




Fig. 1. LTB quantities as a function of the FLRW comoving coordinate, χ , at the present time, t_0 . The two dotted lines mark the positions of the shells relative to r_i and r_b . See Sect. 2.7.

Table 1. Parameters specifying the ALTB model.

FLRW parameters	Value
H_0	$68 \text{ km s}^{-1} \text{ Mpc}^{-1}$
Ω_m	0.3
Ω_k	0
Perturbation parameters	Value
Ω_b	0.048
$\ln(10^{10} A_s)$	3.0
n_s	0.97
τ	0.094
Y_p	0.25
N_{eff}	3.046
$\sum m_\nu$	0
LTB parameters	Value
δ_0	$[-0.6, 0.6]$
r_b	$[500, 4000] \text{ Mpc } h^{-1}$

Notes. The non-LTB parameters define the fiducial BEHOMO cosmology. The amplitude A_s of scalar perturbations and their spectral index n_s are relative to the pivot scale $k_p = 0.05/\text{Mpc}$. This ΛCDM cosmology gives $\sigma_8 = 0.79364$ and $t_0 = 13.862 \text{ Gyr}$. Radiation has been neglected as neither the simulation nor the GR calculations include radiation.

with the initial conditions $t(0) = t_0$ and $r(0) = 0$. The area (d_A) and luminosity (d_L) distances are given by

$$d_A(z) = a_\perp(t(z), r(z)) r(z), \quad (28)$$

$$d_L(z) = (1+z)^2 d_A(z). \quad (29)$$

2.6. Model parameters

The ALTB model is specified by the usual background FLRW parameters, that is, the Hubble constant H_0 , the total matter density parameter Ω_m , and the curvature parameter Ω_k , by the standard perturbation parameters, that is, the baryon density parameter Ω_b , the optical depth τ , the helium fraction Y_p , the effective number of relativistic species N_{eff} , the total neutrino mass $\sum m_\nu$, and the amplitude of the primordial power spectrum A_s and its tilt n_s , and, finally, by the LTB parameters, that is, the central curvature k_c and the inhomogeneity radius r_b . While numerically the profile is specified via k_c , we adopt, in its stead, the derived parameter δ_0 , which is the contrast today at the center of the inhomogeneity and is more intuitive to most cosmologists. Table 1 summarizes all the parameters and their fiducial values.

2.7. Example of inhomogeneity

Figure 1 shows the relevant functions for the case of a central underdensity of present-day contrast $\delta_0 = -0.4$ and comoving radius $r_b = 2000 \text{ Mpc}$, and the fiducial BEHOMO cosmology of Table 1. In particular, one can note that the interior of the inhomogeneity is an open FLRW universe (first panel from the top), that there is a compensating overdensity that surrounds the inner underdensity (second panel), and how the longitudinal Hubble rate deviates from the perpendicular Hubble rate where there is a spatial gradient (fourth panel). Also shown, for later use, are the linear and nonlinear Newtonian potentials together with the energy function (third panel), and the change in redshift induced by the inhomogeneity, together with the peculiar velocity defined in Eq. (38) (last panel). Figure 2 shows the relevant functions on the light cone as compared to their ΛCDM equivalent.




Fig. 2. LTB quantities as a function of redshift. See Sect. 2.7.

From Fig. 1 one can see that an inhomogeneity with a central underdensity of contrast $\delta_0 = -0.4$ could solve the discrepancy between local (Riess et al. 2021) and high-redshift (Aghanim et al. 2020) determinations of the Hubble constant: H_0 goes from the background value of $68 \text{ km s}^{-1} \text{ Mpc}^{-1}$ to the local value of $73 \text{ km s}^{-1} \text{ Mpc}^{-1}$. This is the so-called local void scenario. However, this scenario is ruled out by other observations. Camarena et al. (2021, 2022) constrained the Λ LTB model using the latest available data from the CMB, BAOs, type Ia supernovae, the local H_0 , cosmic chronometers, Compton y -distortion, and the kinetic Sunyaev-Zeldovich (kSZ) effect and showed that an underdensity around the observer as modeled within the Λ LTB model cannot solve the H_0 tension. Appendix A reports the latest constraints by Camarena et al. (2021) using the LTB parameters δ_0 and r_b that we adopt here.

2.8. Newtonianly perturbed FLRW metric

One can regard the LTB inhomogeneity as a perturbation on top of the Λ CDM model. Here, we connect the formalism of the previous sections with that of the Newtonianly perturbed FLRW metric:

$$ds^2 = -d\tilde{t}^2(1 + 2\Phi) + a^2(\tilde{r})(d\tilde{r}^2 + \tilde{r}^2 d\Omega^2)(1 - 2\Phi), \quad (30)$$

where, for simplicity, we assumed a flat background FLRW metric. This will be particularly relevant as N -body simulations are in an FLRW background (see the discussion regarding the N -body gauge in Fidler et al. 2017). This analysis will also be useful to highlight observational effects specific to Λ LTB inhomogeneities. As we will see, in the case of sub-horizon inhomogeneities it is $\Phi \ll 1$.

⁸ One can estimate the change in the expansion rate via linear perturbation theory. An adiabatic perturbation in density causes $\delta H_0/H_0 = -\frac{1}{3}f(\Omega_m)\delta\rho(t_0)/\rho(t_0)$, where $f \approx 0.5$ is the present-day growth rate for the concordance Λ CDM model.

By linearizing the LTB metric and considering a linear gauge transformation, one finds that the Newtonian potential for $r < r_b$ is (Biswas & Notari 2008; Van Acoleyen 2008)

$$\Phi_{\text{lin}}(r) = \frac{3}{5} \int_r^{r_b} \frac{E(\tilde{r})}{\tilde{r}} d\tilde{r} \sim E, \quad (31)$$

and $\Phi_{\text{lin}} = 0$ for $r \geq r_b$, where the potential is written as a function of the LTB coordinate. Hereafter, the subscript “lin” refers to the fact that a linear gauge transformation is used; the potential is always linear, that is, a first-order perturbed quantity. We also note that Φ_{lin} is constant in time, as should be for a linear matter perturbation in a matter-dominated universe. This description should be accurate at $z \gtrsim 10$. The corresponding linear density contrast is

$$\Phi'_{\text{lin}}(r) = -\frac{3}{5} \frac{E(r)}{r}, \quad (32)$$

$$\nabla^2 \Phi_{\text{lin}}(r) = \Phi''_{\text{lin}} + 2 \frac{\Phi'_{\text{lin}}}{r} = -\frac{3}{5} \left[\frac{E'(r)}{r} + \frac{E(r)}{r^2} \right], \quad (33)$$

$$\delta_{\text{lin}}(t, r) = \frac{\nabla^2 \Phi_{\text{lin}}(r)}{4\pi G \rho_m(t) a(t)^2}, \quad (34)$$

where quantities without explicit radial dependence are relative to the FLRW background. We took the derivative with respect to r instead of the Newtonian gauge coordinate \tilde{r} , but the difference is second order. Using Eq. (34) together with Eqs. (12) and (19), one can find the initial evolution of the central density contrast as a function of the central curvature, k_c :

$$\delta_{\text{lin}}(t, 0) = \frac{9k_c}{40\pi G \rho_m(t) a(t)^2}. \quad (35)$$

One could use second-order perturbation theory to improve upon this linear description (Matarrese et al. 1998). However, given that, in general, the LTB inhomogeneity may feature nonlinear contrasts⁹, we now consider the potential as obtained via a nonlinear gauge transformation $\tilde{t} = \tilde{t}(t, r)$ and $\tilde{r} = \tilde{r}(t, r)$, which, following Van Acoleyen (2008), is implicitly defined for $r < r_b$ by

$$Y(r, t) = a(\tilde{t})\tilde{r}(1 - \Phi(\tilde{t}, \tilde{r})), \quad (36)$$

$$t = \tilde{t} + a(\tilde{t}) \int_{\tilde{r}}^{r_b} v(\tilde{t}, \tilde{r}) d\tilde{r}, \quad (37)$$

and by $\tilde{r} = r$ and $\tilde{t} = t$ for $r \geq r_b$, where the peculiar velocity is

$$\begin{aligned} v(\tilde{t}, \tilde{r}) &= \dot{Y}(t, r) - \dot{a}(t)\tilde{r} = \dot{Y}(t, r) - H(t)Y(t, r) \\ &= Y(t, r)[H_{\perp}(t, r) - H(t)]. \end{aligned} \quad (38)$$

This gauge transformation will keep terms up to Φ , $E \sim v^2$ and is valid for sub-horizon inhomogeneities. From Eq. (36) one sees that $\tilde{r} \xrightarrow{\Phi, E \ll 1} \chi$, that is, the coordinate χ defined in Eq. (25) is indeed the one associated with the Newtonian gauge and, therefore, the one adopted by N -body simulations.

One can then use Eqs. (36)–(37) to change the LTB metric of Eq. (1) into the Newtonian gauge of Eq. (30) so as to find the potential Φ . Alternatively, one may proceed by inverting the Poisson equation:

$$\nabla^2 \Phi(\tilde{r}) = \frac{1}{\tilde{r}^2} (\tilde{r}^2 \Phi')' = 4\pi G a^2 \left[\frac{F'}{4\pi Y^2 Y'} - \frac{3}{8\pi G} \left(H^2 - \frac{\Lambda}{3} \right) \right], \quad (39)$$

⁹ See Rigopoulos & Valkenburg (2012) for an alternative approach that uses a gradient series expansion.

where the derivatives are with respect to the variable of the corresponding function. In particular, it is $d\tilde{r} = \frac{Y'}{a}(1 + \mathcal{O}(\Phi))dr$, so that one can integrate on \tilde{r} and obtain

$$\Phi' = a \frac{GF}{Y^2} - \frac{1}{2} a Y \left(H^2 - \frac{\Lambda}{3} \right), \quad (40)$$

where the constant of integration has been chosen in order to have $\Phi'(r_b) = 0$ and the potential is expressed with respect to the LTB coordinate. Integrating again on \tilde{r} , one finally has

$$\begin{aligned} \Phi &= - \int_r^{r_b} \frac{Y' GF}{Y^2} d\tilde{r} + \left(H^2 - \frac{\Lambda}{3} \right) \left(\frac{Y_b^2}{4} - \frac{Y^2}{4} \right) \\ &= \left(H^2 - \frac{\Lambda}{3} \right) \left(\frac{Y_b^2}{4} - \frac{Y^2}{4} \right) + \frac{GF_b}{Y_b} - \frac{GF}{Y} - \int_r^{r_b} \frac{GF'}{Y} d\tilde{r}, \end{aligned} \quad (41)$$

where $Y_b = ar_b$. We note that $\Phi(r_b) = 0$ and that we expressed the potential with respect to the LTB coordinate. It is interesting to note $\nabla^2 \Phi$ gives exactly the LTB contrast in LTB coordinates while the gauge transformation is only valid up to $\mathcal{O}(\Phi)$. Figure 1 (third panel, green solid curve) shows how the potential of Eq. (41) decays during the cosmological-constant dominated phase as compared to the linear-gauge potential of Eq. (31) during matter domination. Also shown (green dot-dashed line) is the linear perturbation result $\Phi(t, r) = \Phi_{\text{lin}}(r)D(t)/a(t)$, where D is the Λ CDM growth function normalized at the matter-dominated epoch. The agreement with Eq. (41) is perfect for linear LTB perturbations but it overestimates the value of the potential in the case of the nonlinear underdensity of Fig. 1.

It is easy to verify that Φ' in Eq. (40) reduces to that of Eq. (32) at early times:

$$\begin{aligned} \tilde{r}\Phi' &\stackrel{(40)}{=} -E + \frac{Y^2}{2} (H_{\perp}^2 - H^2) \simeq -E + Y^2 H^2 \frac{\delta H}{H} \\ &= -E + Y^2 H^2 \frac{\delta E}{5GF/Y} = -E \left(1 - \frac{Y^3 H^2}{5GF} \right) = -\frac{3}{5} E, \end{aligned} \quad (42)$$

where we used Eq. (12) in the first equality, Eq. (31.14, Kaiser 2014) in the third equality and $F \simeq 4\pi Y^3 \rho_m(t)/3$ in the last equality. Moreover, using the result after the second equality, one has

$$v \stackrel{(38)}{=} Y\delta H \stackrel{(42)}{=} \frac{\Phi'}{a} + \frac{E}{\dot{a}r} \stackrel{(32)}{=} -\frac{2}{3} \frac{\Phi'}{a}, \quad (43)$$

in agreement with the expected matter-dominated result (Coles & Lucchin 2002, Eq. (18.1.9)).

2.9. Observables in terms of the Newtonian potential

If one uses the metric functions of the LTB metric of Eq. (1) then the effects of the inhomogeneities are exactly taken into account. However, it is important to discuss and review how the Newtonian potential affects observables. Indeed, the total potential will consist of the sum of the LTB potential and the potential relative to the primordial Gaussian perturbations, and the LTB potential may have observational effects in regimes in which the standard Gaussian potential is inconsequential.

A well-known result is that the redshifts of photons are affected by perturbations according to (see, for example, Bonvin et al. 2006)

$$\frac{\delta z}{1+z} \simeq (\mathbf{v}_O - \mathbf{v}_S) \cdot \mathbf{n} + (\Phi_O - \Phi_S) + 2a \int_{\chi_O}^{\chi_S} \dot{\Phi} d\chi, \quad (44)$$

where the vector \mathbf{n} gives the direction of the source S with velocity \mathbf{v}_S with respect to the observer O with velocity \mathbf{v}_O . In the following, we only consider the contribution from the LTB potential, but there are of course also contributions from standard-model perturbations. The change in redshift can be interpreted as the sum of three effects.

First, there is the differential Doppler shift due to the peculiar motion of source and observer. This contribution is zero if the observer and the source are placed outside the inhomogeneity or (one of them) at its center. Otherwise, one expects a contribution that is proportional to $v \sim Y\Delta H \sim r_b/r_{\text{hor}}$ where $r_{\text{hor}} = H^{-1}$ is the Hubble radius. This contribution is large for gigaparsec-scale inhomogeneities and can significantly alter the luminosity distance–redshift relation; it was indeed used to fit supernova data without dark energy in the void scenario (see Sect. 2.11 for a historical note). Figure 1 (bottom panel) shows this effect for an underdensity of contrast $\delta_0 = -0.4$ and comoving radius $r_b = 2000$ Mpc. Also shown is the peculiar velocity as defined in Eq. (38). One can see that most of the change in redshift can indeed be attributed to a Doppler shift.

The second term gives the so-called Sachs-Wolfe effect, that is, the differential gravitational redshift due to the gravitational potentials at the source's and observer's positions (Sachs & Wolfe 1967). This contribution is zero if the observer and the source are placed outside the inhomogeneity. Otherwise, by comparing Eqs. (12) and (38) it is easy to see that $E \sim Y^2/2 - Y^2 H^2/2 \sim -v^2/2$ so that the potential of Eq. (31) is quadratic in the velocities, $\Phi \propto v^2$. It then follows that $\Phi \propto (r_b/r_{\text{hor}})^2$ and the Sachs-Wolfe effect is subdominant with respect to the Doppler shift. Again from the analysis of Fig. 1 (bottom panel), one can see that $\delta z/1+z \simeq -4 \times 10^{-5}$ at r_b , significantly smaller than the Doppler shift that occurs for source inside the inhomogeneity.

Finally, the last term is the ISW effect, which is present only if the (first-order) gravitational potential evolves with time and is responsible for a nontrivial correlation between CMB anisotropies and the large-scale structure. At high redshift, $10 \lesssim z \lesssim 100$, the standard model is very close to the flat matter-dominated Einstein-de Sitter model. It is then well-known that the (linear) potential is time independent so that the ISW contribution is zero. At later times, however, there are two contributions. First, the universe enters the cosmological constant-dominated phase: this is responsible for the (linear) ISW effect. Second, structures may enter the nonlinear regime so that the so-called Rees-Sciama (RS) effect cannot be neglected. As discussed in Cai et al. (2010), the nonlinear RS correction to the ISW effect acts differently for over and underdensities. Biswas & Notari (2008) explicitly showed that these contributions are suppressed according to $\propto (r_b/r_{\text{hor}})^3$.

For the mildly nonlinear large structures here considered, one expects that RS is subdominant with respect to ISW (Sakai & Inoue 2008). In this case, the potential decays according to the linear ISW modeling:

$$\dot{\Phi} = \frac{3}{2} \Omega_{m0} H_0^2 G(z) P(r), \quad (45)$$

where the (nonlinear) potential is obtained via Eq. (41) and the ISW growth factor, G , is

$$G(z) = (1+z)H(z)[1-f(z)]D(z), \quad (46)$$

where D is the linear growth function, $f \equiv d \ln D / d \ln a \simeq \Omega_m^\gamma(t)$ with $\gamma = 6/11 + 15/11^3(1 - \Omega_m(t))$ is the linear growth rate (Wang & Steinhardt 1998), and $P(r)$ encodes the information on

the inhomogeneity profile. A thorough discussion is available in [Nadathur et al. \(2012\)](#) and [Flender et al. \(2013\)](#).

As thoroughly discussed in [Hui & Greene \(2006\)](#), a perturbation in the redshift affects the luminosity distance. As we have seen, the LTB metric features possibly large contributions from peculiar velocities that are instead negligible in the standard paradigm. This means that proper care has to be adopted when analyzing these models on the light cone. The other important effect to consider is lensing, which modifies the observed flux of an object without changing its redshift. However, in this case, the total effect will be directly computed by binning mass in a suitable number of lens planes. Indeed, the total lensing effect is the sum of the contribution of the LTB potential with the one of the Gaussian perturbations' potential, and these two components make up the various lens planes.

2.10. Scale invariance

As the dynamical equation (Eq. (5)) does not present gradients, the dynamics of the LTB model is scale invariant. This is due to spherical symmetry and the fact that the energy-momentum tensor is dust. The former implies a vanishing magnetic Weyl tensor and consequently no gravitational waves; the latter implies no pressure and so no sound waves. In other words, no direct communication can exist between neighboring world-lines and for this reason such space-times were dubbed “silent” ([Matarrese et al. 1993](#); [Bruni et al. 1995](#)). In particular, pressure gradients would transfer energy between shells and make the energy function E and mass function F time dependent (see [Marra & Paakkonen 2012](#)).

Formally, starting from the solution of Eq. (5) for a given r_b , one can obtain a scaled inhomogeneity with coordinate $\hat{r} = \lambda r$ and size $\hat{r}_b = \lambda r_b$. The Friedmann-like equation is then

$$\frac{\dot{\hat{a}}_{\perp}(t, \hat{r})}{\hat{a}_{\perp}(t, \hat{r})} = \frac{8\pi G}{3} \frac{M_0^4}{\hat{a}_{\perp}^3(t, \hat{r})} + \frac{8\pi G}{3} \rho_{\Lambda} - \frac{\hat{k}(\hat{r})}{\hat{a}_{\perp}^2(t, \hat{r})}, \quad (47)$$

where we adopted the gauge fixing $F(r) = 4\pi M_0^4 r^3/3$ and the functions relative to the scaled inhomogeneity are defined according to

$$\hat{a}_{\perp||}(t, \hat{r}) = a_{\perp||}(t, \hat{r}/\lambda), \quad (48)$$

$$\hat{H}_{\perp||}(t, \hat{r}) = H_{\perp||}(t, \hat{r}/\lambda), \quad (49)$$

$$\hat{k}(t, \hat{r}) = k(t, \hat{r}/\lambda), \quad (50)$$

$$\hat{\rho}_m(t, \hat{r}) = \rho_m(t, \hat{r}/\lambda), \quad (51)$$

$$\hat{Y}(t, \hat{r}) = \lambda Y(t, \hat{r}/\lambda), \quad (52)$$

$$\hat{v}(t, \hat{r}) = \lambda v(t, \hat{r}/\lambda), \quad (53)$$

$$\hat{E}(t, \hat{r}) = \lambda^2 E(t, \hat{r}/\lambda), \quad (54)$$

$$\hat{\Phi}(t, \hat{r}) = \lambda^2 \Phi(t, \hat{r}/\lambda), \quad (55)$$

$$\hat{F}(t, \hat{r}) = \lambda^3 F(t, \hat{r}/\lambda). \quad (56)$$

Starting from one numerical solution, one can then obtain a family of solutions by varying λ .

We note that velocities, and thus Doppler effects, are proportional to λ , explaining why one needs a large inhomogeneity to sizably change the luminosity distance–redshift relation, as in the void scenario discussed in Sect. 2.11. Also, the energy function and the potential scale quadratically with the size so that one expects strong features in the power spectrum of large inhomogeneities.

2.11. A historical note on LTB void models

The LTB model has been studied extensively in the literature as an alternative to dark energy. The relevant case was of an observer sitting near the center of a gigaparsec-scale under-density. It is easy to understand how such an observer would see apparent acceleration: most of our cosmological observables are confined to the light cone and, hence, temporal changes can be associated with spatial changes along photon geodesics. The LTB void model then replaces “faster expansion now than before” with “faster expansion here than there”. Mathematically, the directional derivative on the past light cone follows $d/dt \approx \partial/\partial t - \partial/\partial r$ and the accelerating expansion can be explained by $H'(r) < 0$ ([Enqvist 2008](#)). For 15 years the LTB model was phenomenologically viable, although suffered the extreme fine-tuning of the observer’s position (see [Marra & Notari 2011](#); [Bolejko et al. 2011](#); [Clarkson 2012](#) and references therein). More importantly, it constituted perhaps the only example of a paradigm that departed abruptly from Λ CDM. This allowed cosmologists to ask new questions and develop new methodologies.

However, in 2011, two papers ruled out the LTB model, which already showed problems when confronted with more and more data ([Garcia-Bellido & Haugboelle 2008](#); [Moss et al. 2011](#); [Biswas et al. 2010](#)). [Zhang & Stebbins \(2011\)](#) (see also [Zibin & Moss 2011](#); [Bull et al. 2012](#)) showed that void models without decaying modes produce an excessively large kSZ signal, and [Zibin \(2011\)](#) showed that void models with sizable decaying modes (which could possibly have a small kSZ signal) are ruled out because of y -distortion.

Despite the strong evidence against void models as alternatives to dark energy, one has to point out that those studies considered a homogeneous radiation field. In other words, inhomogeneities were present only in the matter component. [Clarkson & Regis \(2011\)](#) and [Lim et al. \(2013\)](#) considered the more consistent scenario of inhomogeneities also in the radiation and showed that this could alter kSZ and y -distortion predictions.

2.12. The backreaction proposal

Because of the nonlinear nature of general relativity, the average of the solution of Einstein’s equations for an inhomogeneous metric is not the solution of Einstein’s equations for the average of the metric, that is, the operation of smoothing does not commute with solving Einstein’s equations, $\langle G_{\mu\nu}(g_{\alpha\beta}) \rangle \neq G_{\mu\nu}(\langle g_{\alpha\beta} \rangle)$. Consequently, the Friedmann equation – valid for a homogeneous universe – features corrections in the form of extra sources (see [Ellis et al. 1984](#), which is considered the backreaction manifesto).

In the early 2000s, right after the first analyses indicating the acceleration of the universe’s expansion by [Riess et al. \(1998\)](#) and [Perlmutter et al. \(1999\)](#), it was asked if dark energy could actually be explained via the extra sources generated by the nonlinear smoothing, that is, via the backreaction of small-scale inhomogeneities into the large-scale dynamics of the universe. This scenario would elegantly explain the biggest problem of dark energy, of why it appears at $z \approx 1$. The answer would be that structures go nonlinear at $z \approx 1$, transforming a fine tuning into a prediction. It is important to point out that, when the backreaction scenario was proposed, the equation of state w of dark energy was poorly constrained. This is particularly relevant because the effective w that one would measure, if dark energy were caused by backreaction, is not expected to be close to -1 , the value relative to the cosmological constant. In other

words, while the present-day tight constraint $w = -1.03 \pm 0.03$ (Abbott et al. 2022) is, within the backreaction proposal, a coincidence, it is instead a necessary condition for the Λ CDM model.

This proposal started a heated debate on the magnitude of the backreaction effect, which proved difficult to be estimated via (semi) analytical techniques. More information on this is available in Clarkson et al. (2011), Buchert et al. (2015), Green & Wald (2014), and references therein.

In the past few years, the scientific consensus on the relevance of backreaction in cosmology has been sought via the methods of numerical relativity. It seems that backreaction produces a negligible correction to the universe dynamics, although the methodology that has been adopted has some limitations. On one hand, fully GR codes are used, but the implementation of the fluid description of the matter sector raises questions regarding the modeling of the nonlinear structure formation, which is dominated by halo mergers and shell crossing (Giblin et al. 2016; Bentivegna & Bruni 2016; Macpherson et al. 2019). On the other hand, particle-based modeling is adopted at the price of using the weak-field expansion of Einstein’s equations (Adamek et al. 2019). However, codes that adopt a particle description alongside numerical relativity (East et al. 2018; Giblin et al. 2019; Daverio et al. 2019), including the recent code GRAMES (Barrera-Hinojosa & Li 2020a,b), are set to provide important progress toward a definitive answer to the questions raised by the backreaction proposal (Adamek et al. 2020).

2.13. Backreaction in the Λ LTB model

With an LTB perturbation that is exactly matched to a background FLRW metric, one can study in an exact way backreaction, that is, the effect of inhomogeneities on the background dynamics. Indeed, in this very simplified case, the background expansion is set by construction so that one has to simply look at the mismatch between the background energy densities and the averaged ones.

In order to maximize the effect, one could fill the entire universe with an infinite number of spherical patches of different radii and profiles by the Apollonian sphere packing, the three-dimensional extension of the Apollonian gasket (see Fig. 1 of Marra et al. 2007). For this reason, here we are concerned with the background dynamics at $r = r_b$ and not at larger radii.

The effect of backreaction can be read from Eq. (5), which can be rewritten as ($r = r_b$)

$$H^2(t) = \frac{8\pi G}{3}(\langle\rho_m\rangle + \rho_\Lambda) - \left\langle\frac{\mathcal{R}}{6}\right\rangle + P_{\text{inh}}, \quad (57)$$

$$P_{\text{inh}} = \frac{8\pi G}{3}(\rho_m(t) - \langle\rho_m\rangle) - \frac{k}{a^2} + \left\langle\frac{\mathcal{R}}{6}\right\rangle, \quad (58)$$

where $H(t)$ is the background expansion rate, fixed by construction, and P_{inh} represents “the effects of small-scale inhomogeneities in the universe on the dynamic behavior at the smoothed-out scale” (Ellis et al. 1984).

If there were no backreaction, $P_{\text{inh}} = 0$, the Friedmann equation would be sourced by the averages of the energy and curvature content of the inhomogeneity. These are obtained by adopting the actual volume element with curvature, $dV = 4\pi Y^2 Y' / \sqrt{1 + 2E}$. However, Einstein’s equations are nonlinear so that backreaction gives the correction P_{inh} . The correction comes from the fact that, by solving Einstein’s equations for the LTB metric, one finds that it is the Euclidean average of the density and curvature that sources the Friedmann equation.

In the case of the density, for example, the correction is proportional to the difference between the invariant mass and F – also known as the Misner-Sharp mass (see Alfedeel & Hellaby 2010, for an extensive discussion). As the difference is proportional to the energy function $E \sim \Phi \propto (r_b/r_{\text{hor}})^2$, one concludes that the backreaction of small-scale inhomogeneities into the background expansion is negligible (see, however, Lavinto et al. 2013, for a model that does feature large backreaction). A comprehensive discussion of averaging and backreaction in LTB metrics is presented in Sussman (2011).

3. N -body simulations of a perturbed Λ LTB model

We now discuss how we simulate the Λ LTB model. As said in the Introduction, because of spatial gradients, standard primordial perturbations are coupled at first order so that one may expect a different growth of perturbations even on scales at which the evolution is still linear. Besides this, simulations are necessary, just as in Λ CDM, in order to obtain the fully nonlinear structure, which again may be affected by the spatial gradients of an inhomogeneous background. As we are interested in understanding and modeling these effects, each Λ LTB simulation will be coupled with the corresponding Λ CDM one, using the same seed for the initial conditions. This will allow us to study the differential change in quantities as compared to Λ CDM, and reduce possible biases caused by the numerical implementation we adopt.

3.1. Early-FLRW initial conditions

As shown by Alonso et al. (2010, 2012), one can simulate the Λ LTB model by feeding standard Newtonian gravity-only N -body codes with special early-FLRW initial conditions. We give initial conditions at $z_{\text{ini}} = 49$ so that the LTB perturbation is deep into the linear regime and the space-time can be accurately described by a superposition of two kinds of perturbations:

$$\delta(t_{\text{ini}}, \mathbf{x}) = \delta^{\text{LTB}}(t_{\text{ini}}, \mathbf{x}) + \delta^{\text{gau}}(t_{\text{ini}}, \mathbf{x}), \quad (59)$$

where the first term comes from the spherical LTB perturbation and the second one from the statistically isotropic set of primordial Gaussian perturbations. As discussed in the Introduction, the LTB initial conditions are non-Gaussian, with phase coupling induced by the presence of the spherical inhomogeneity.

As we have seen, at early times, the potential Φ^{LTB} induced by the LTB metric is given by Eq. (31), together with Eqs. (12) and (19). Owing to the Poisson equation in Newtonian gravity, the gravitational potential obeys qualitatively exactly the same differential equation as the displacement potential for the matter field, such that initial positions for particles in a simulation can be set by

$$\mathbf{x}(t_{\text{ini}}) = \mathbf{q} + \nabla(\Phi^{\text{LTB}} + \Phi^{\text{gau}}). \quad (60)$$

We generate these initial conditions using FalconIC (2017 version, Valkenburg & Hu 2015), a code that extends Lagrangian perturbation theory to nontrivial theories of gravity.

As the Λ LTB model does not include radiation, we neglect radiation in the N -body simulation, as well as the effect of neutrinos, including massive neutrinos. In other words, the initial transfer function $T_k(z_{\text{ini}})$ is obtained by rescaling the one provided by the Boltzmann solver at $z = 0$ to the initial redshift via the scale-independent radiation-less growth factor $D(z = 49) = 0.0256745$, where $D(z = 0) = 1$. A thorough discussion on initial conditions for simulations is available