

Supplementary information (SI)

Long-term outcomes of canaloplasty and phaco-canaloplasty in the treatment of open angle glaucoma: a single-surgeon experience.

International Ophthalmology

Daniele Tognetto M.D.¹, Gabriella Cirigliano M.D.¹, Stefano Gouigoux M.D.¹, Alberto Grotto M.D.¹, Pier Luigi Guerin M.D.¹, Leandro Inferrera M.D.¹, Dario Marangoni M.D., Ph.D.^{1*}

¹ University Eye Clinic, Department of Medicine, Surgery and Health Sciences, University of Trieste, 34129, Trieste, Italy. *Corresponding author.

Corresponding author:

Dario Marangoni

University Eye Clinic, Department of Medicine, Surgery and Health Sciences

University of Trieste

Piazza dell'Ospitale 1, 34129, Trieste, Italy

dario.marangoni@units.it




Figure S1.

The graphs illustrate individual changes in postoperative IOP and number of glaucoma medications after canaloplasty (blue dots) and phaco-canaloplasty (red dots) over a 48-month period, with data collected at six-month intervals. In each graph the lower right quadrant includes eyes in which both the IOP and number of glaucoma medications were reduced compared to baseline measurements. **CP** = canaloplasty, **PCP** = phaco-canaloplasty.