




Contents lists available at ScienceDirect

Journal of Mathematical Analysis and Applications

journal homepage: www.elsevier.com/locate/jmaa



Regular Articles

An extension of the Poincaré–Birkhoff Theorem coupling twist with lower and upper solutions



Alessandro Fonda ^a, Manuel Garzón ^b, Andrea Sfecci ^{a,*}

^a *Dipartimento di Matematica e Geoscienze, Università degli Studi di Trieste, P.le Europa 1, 34127 Trieste, Italy*

^b *Departamento de Matemática Aplicada, Universidad de Granada, Avenida de la Fuente Nueva S/N, 18071 Granada, Spain*

ARTICLE INFO

Article history:

Received 23 December 2022
Available online 17 July 2023
Submitted by R.O. Popovych

Keywords:

Hamiltonian systems
Periodic boundary value problem
Poincaré–Birkhoff Theorem
Lower and upper solutions

ABSTRACT

In 1983, Conley and Zehnder proved a remarkable theorem on the periodic problem associated with a general Hamiltonian system, giving a partial answer to a conjecture by Arnold. Their pioneering paper has been extended in different directions by several authors. In 2017, Fonda and Ureña established a deeper relation between the results by Conley and Zehnder and the Poincaré–Birkhoff Theorem. In 2020, Fonda and Gidoni pursued along this path in order to treat systems whose Hamiltonian function includes a nonresonant quadratic term. It is the aim of this paper to further extend this fertile theory to Hamiltonian systems which, besides the periodicity-twist conditions always required in the Poincaré–Birkhoff Theorem, also include a term involving a pair of well-ordered lower and upper solutions. Phase-plane analysis techniques are used in order to recover a saddle-type dynamics permitting us to apply the above mentioned results.

© 2023 The Author(s). Published by Elsevier Inc. This is an open access article under the CC BY-NC-ND license (<http://creativecommons.org/licenses/by-nc-nd/4.0/>).

1. Introduction

In 1983, Conley and Zehnder [11] proved a remarkable result on the periodic problem associated with a general Hamiltonian system

$$J\dot{z} = \nabla H(t, z), \tag{1}$$

giving a partial answer to a conjecture by Arnold [1,2]. Here, $J = \begin{pmatrix} 0 & -I \\ I & 0 \end{pmatrix}$ is the standard symplectic matrix, $H: \mathbb{R} \times \mathbb{R}^{2N} \rightarrow \mathbb{R}$ is T -periodic in t , and $\nabla H(t, z)$ denotes its gradient with respect to z . Let us write $z = (q, p)$, with $q = (q_1, \dots, q_N)$ and $p = (p_1, \dots, p_N)$, so that the system reads as

* Corresponding author.

E-mail addresses: a.fonda@units.it (A. Fonda), manuelgarzon@ugr.es (M. Garzón), asfecci@units.it (A. Sfecci).

$$\dot{q} = \partial_p H(t, q, p), \quad \dot{p} = -\partial_q H(t, q, p).$$

Their paper contains two main results.

In [11, Theorem 1], assuming the Hamiltonian function $H(t, z)$ to be periodic in all variables q_1, \dots, q_N and p_1, \dots, p_N , they prove that system (1) has at least $2N + 1$ geometrically distinct T -periodic solutions. They also mention in [11, Theorem 2] that the expected number of solutions is 2^{2N} provided that all the T -periodic solutions are known to be nondegenerate.

In [11, Theorem 3], the function H is assumed to be periodic in q_1, \dots, q_N and to have a quadratic behaviour in p ; namely, that there exist a constant $R > 0$ and a symmetric regular matrix \mathbb{A} such that $H(t, q, p) = \frac{1}{2}\langle \mathbb{A}p, p \rangle +$ “lower order terms”, when $|p| \geq R$. In this setting, they prove that system (1) has at least $N + 1$ geometrically distinct T -periodic solutions.

Conley and Zehnder [11] also mention a possible relation of this second result with the Poincaré–Birkhoff Theorem (the title of the paper is however a bit misleading). This theorem was first conjectured by Poincaré [46] in 1912, shortly before his death, and then proved by Birkhoff in [4,5]. Both of them were interested in proving it because of its consequences in the existence of periodic solutions for some Hamiltonian systems originating from Celestial Mechanics. As mentioned by Zehnder in [50], Arnold called this theorem “the seed of symplectic topology” (cf. also [43]). See [24] for a brief historical account till the centennial anniversary of the theorem.

The pioneering results in [11] have been extended in different directions by several authors (see, e.g., [9, 15, 23, 33, 37, 38, 40, 41, 47, 49]).

More recently, a deeper relation between these results and the Poincaré–Birkhoff Theorem has been established in [30]. Taking, e.g., $\mathcal{D} = \overline{B}(0, R)$, the closed ball with radius R , and assuming H to be periodic in q_1, \dots, q_N , the existence of $N + 1$ geometrically distinct T -periodic solutions was established under the following hypothesis: there exist a constant $R > 0$ and a symmetric regular matrix \mathbb{A} such that the solutions $z(t) = (q(t), p(t))$ of (1) starting with $p(0) \in \mathcal{D}$ are defined on $[0, T]$ and satisfy a “twist condition” like

$$\langle q(T) - q(0), \mathbb{A}p(0) \rangle > 0 \text{ if } p(0) \in \partial\mathcal{D}.$$

It is easily checked that this result generalizes the second Conley–Zehnder Theorem described above. Variants of the twist condition were also proposed in [30,31].

The results in [30] have been extended in [17] in order to include both the above quoted Conley–Zehnder theorems, assuming H to be periodic in q_1, \dots, q_N and possibly also in p_1, \dots, p_L , for some $L \in \{1, \dots, N\}$, together with a very general twist condition, thus finding $N + L + 1$ periodic solutions. The same authors further extended the theory, in a second paper [18], to the case when the Hamiltonian function includes a nonresonant quadratic term. Possible resonance has also been investigated in [10], assuming some Ahmad–Lazer–Paul conditions.

These general existence results have found so far several applications (see [7,16,18,25,26,28,30]), thus generalizing some previously established results for second order equations (cf. [8,13,14,22,27,35,36]). They have even been extended to the study of infinite-dimensional systems, cf. [6,19,21].

It is the aim of this paper to further extend this fertile theory to systems which, besides the periodicity-twist conditions illustrated above, also present a pair of well-ordered lower and upper solutions. In order to better explain this situation, let the considered Hamiltonian system be of the type

$$\begin{cases} \dot{q} = \partial_p H(t, q, p, u, v), & \dot{p} = -\partial_q H(t, q, p, u, v), \\ \dot{u} = \partial_v H(t, q, p, u, v), & \dot{v} = -\partial_u H(t, q, p, u, v). \end{cases} \quad (2)$$

Here $H: \mathbb{R} \times \mathbb{R}^4 \rightarrow \mathbb{R}$ is T -periodic in t , periodic in q , and has a twist involving the (q, p) variables; at the same time, we also assume that there exist some constant lower/upper solutions $\alpha \leq \beta$ involving the

(u, v) variables. In this situation we are able to prove the existence of two geometrically distinct T -periodic solutions. The result will then be extended to higher dimensional systems.

The reader is surely familiar with the method of lower/upper solutions in the case of scalar equations like, e.g.,

$$\ddot{u} = f(t, u).$$

This method has a long history, dating back to the pioneering papers [44,45,48] (see the book [12] for a detailed exposition). We recall that the T -periodic functions $\alpha, \beta: \mathbb{R} \rightarrow \mathbb{R}$ are said to be a lower solution and an upper solution, respectively, if

$$\ddot{\alpha}(t) \geq f(t, \alpha(t)), \quad \ddot{\beta}(t) \leq f(t, \beta(t)),$$

for every $t \in [0, T]$. Recently the theory has been extended to periodic planar systems in [20,29]; this will be the approach adopted here, even if we will only be able to deal with the case of constant lower and upper solutions.

Let us briefly explain why we need to reduce to constant lower/upper solutions. The standard proof procedure in lower/upper solutions theorems is as follows: a) modify the problem below $\alpha(t)$ and above $\beta(t)$, in the u component, by the use of a truncating function; b) show that the modified problem has a solution; c) prove that the u component of this solution stays between $\alpha(t)$ and $\beta(t)$. The technical difficulty encountered in the present paper is that we need to maintain the Hamiltonian structure, hence the modification of the problem has to be made in the Hamiltonian function itself, being careful to preserve the differentiability of the new function. Once the modification has been made, we are allowed to apply the results in [18]. We believe that this technical difficulty could be overcome, but for now the case of nonconstant lower/upper solutions remains an open problem.

Another open problem arises in the case of non-well-ordered lower and upper solutions. Assuming some nonresonance conditions with respect to the higher part of the spectrum, this case is usually treated by topological degree methods. We do not know how to adapt this type of technique to our situation.

In order to maintain a friendly exposition we preferred writing this paper following an increasing order of complexity, first presenting the main ideas in the simplest situation, then extending them to more general systems. The paper is thus organized as follows.

In Section 2 we state our result in the simple case of system (2). The proof is provided in Section 3. Then in Section 4, we provide some variants of the first theorem. In particular, we state a version of the theorem involving a topological annulus, in the spirit of Poincaré's original statement. In Section 5 we illustrate several examples of applications.

In Section 6 we extend the result to higher dimensions, thus generalizing both the Conley–Zehnder theorems presented above. The proof is provided in Section 7. In Section 8 we extend the higher dimensional result to systems whose Hamiltonian function further involves a quadratic term and some examples of possible applications are given in Section 9. In Section 10 we provide a further application to the study of periodic solutions to perturbations of completely integrable systems.

Finally, in Section 11 we establish the most general result of this paper, where the twist condition is stated as an “avoiding cones condition”.

2. Statement of the first result

In this section we consider the Hamiltonian system (2), where the Hamiltonian function $H: \mathbb{R} \times \mathbb{R}^4 \rightarrow \mathbb{R}$ is assumed to be continuous, T -periodic in t , and continuously differentiable in the q, p, u, v variables.

Let us state our assumptions.

Assumption 1 (*Periodicity*). The function $H(t, q, p, u, v)$ is periodic in q .

To fix the ideas, we will assume the period in q to be equal to 2π . Under this setting, T -periodic solutions of (2) appear in equivalence classes made of those solutions whose components $q(t)$ differ by an integer multiple of 2π . We say that two T -periodic solutions are *geometrically distinct* if they do not belong to the same equivalence class. In the sequel we will implicitly use the same notion in different contexts without further mention.

Assumption 2 (*Lower and upper solutions*). There exist some constants $\delta > 0$ and $\alpha \leq \beta$ such that

$$v \partial_v H(t, q, p, u, v) > 0 \quad \text{if } u \in [\alpha - \delta, \alpha] \cup [\beta, \beta + \delta] \text{ and } v \neq 0, \quad (3)$$

and

$$\begin{aligned} \partial_u H(t, q, p, u, 0) &\geq 0 \quad \text{if } u \in [\alpha - \delta, \alpha], \\ \partial_u H(t, q, p, u, 0) &\leq 0 \quad \text{if } u \in [\beta, \beta + \delta]. \end{aligned} \quad (4)$$

The above assumption comes from the definition of lower and upper solutions given in [20,29]. We require here that these lower and upper solutions are constant. More precisely, all constants in $[\alpha - \delta, \alpha]$ are lower solutions, and all constants in $[\beta, \beta + \delta]$ are upper solutions. In the sequel, the constant $\delta > 0$ provided by Assumption 2 will be used without further mention.

Assumption 3 (*Nagumo condition*). There exist $d > 0$ and two continuous functions $f, \varphi: [d, +\infty[\rightarrow]0, +\infty[$ with

$$\int_d^{+\infty} \frac{f(s)}{\varphi(s)} ds = +\infty,$$

satisfying the following property. When $u \in [\alpha - \delta, \beta + \delta]$,

$$\begin{aligned} \partial_v H(t, q, p, u, v) &\geq f(v) && \text{if } v \geq d, \\ \partial_v H(t, q, p, u, v) &\leq -f(-v) && \text{if } v \leq -d, \end{aligned}$$

and

$$|\partial_u H(t, q, p, u, v)| \leq \varphi(|v|) \quad \text{if } |v| \geq d.$$

Assumption 4 (*Linear growth*). For every $K > 0$ there is a constant $C_K > 0$ such that

$$|\partial_q H(t, q, p, u, v)| \leq C_K(|p| + 1) \quad \text{if } u \in [\alpha - \delta, \beta + \delta] \text{ and } |v| \leq K.$$

Remark 1. Notice that, under the above assumption, for any two continuous functions $U, V: [0, T] \rightarrow \mathbb{R}$ with

$$\alpha - \delta \leq U(t) \leq \beta + \delta \quad \text{for every } t \in [0, T], \quad (5)$$

the solutions of the system

$$\dot{q} = \partial_p H(t, q, p, U(t), V(t)), \quad \dot{p} = -\partial_q H(t, q, p, U(t), V(t)) \quad (6)$$

are defined on $[0, T]$. Indeed, let $(q(t), p(t))$ be a solution of system (6) starting at time $t = 0$ from some $(q(0), p(0)) = (q_0, p_0)$. This solution is defined on a maximal interval of future existence $[0, T] \cap [0, \omega[$. Set $K = \|V\|_\infty$. By Assumption 4,

$$|\dot{p}(t)| \leq C_K(|p(t)| + 1) \quad \text{for every } t \in [0, T] \cap [0, \omega[,$$

which, combined with the Gronwall Lemma, yields

$$|p(t)| \leq (|p_0| + 1)e^{C_K T} \quad \text{for every } t \in [0, T] \cap [0, \omega[.$$

Then since H is periodic in q , we have that there is a constant $C > 0$ depending only on U, V and $|p_0|$ such that

$$|\dot{q}(t)| \leq C \quad \text{for every } t \in [0, T] \cap [0, \omega[.$$

Hence

$$|q(t)| \leq |q_0| + CT \quad \text{for every } t \in [0, T] \cap [0, \omega[.$$

This implies that the solution $(q(t), p(t))$ is defined on $[0, T]$, i.e., that $\omega > T$.

We are now ready to state our main result.

Theorem 2. *Let Assumptions 1–4 hold. Assume that there exist $a < b$ and $\rho > 0$ with the following property: For any two continuous functions $U, V: [0, T] \rightarrow \mathbb{R}$ satisfying (5), the solutions of system (6) are such that*

$$\begin{aligned} q(T) - q(0) < 0 & \quad \text{if } p(0) \in [a - \rho, a], \\ q(T) - q(0) > 0 & \quad \text{if } p(0) \in [b, b + \rho]. \end{aligned} \tag{7}$$

Then there exist at least two geometrically distinct T -periodic solutions of system (2) such that $p(0) \in]a, b[$ and

$$\alpha \leq u(t) \leq \beta \quad \text{for every } t \in \mathbb{R}. \tag{8}$$

The same conclusion holds if (7) is replaced by

$$\begin{aligned} q(T) - q(0) > 0 & \quad \text{if } p(0) \in [a - \rho, a], \\ q(T) - q(0) < 0 & \quad \text{if } p(0) \in [b, b + \rho]. \end{aligned}$$

3. The proof of Theorem 2

The proof is based on [18, Corollary 2.4], and will be divided in two steps. In the first step we analyze the dynamics focusing on the (u, v) variables. In the second one, we draw attention to the (q, p) variables.

3.1. Working with the coordinates (u, v)

Let us focus our attention on the couple of variables (u, v) . At first, we are going to modify the original problem (2) outside some suitably chosen set $\mathcal{V} \subseteq \mathbb{R}^4$. We will then prove that all the T -periodic solutions

of the modified system satisfy $z(t) = (q(t), p(t), u(t), v(t)) \in \mathcal{V}$ for every $t \in [0, T]$; hence, such solutions will solve the original problem (2), as well.

Assumption 3 permits us to apply the reasoning in [29, Theorem 3.1] (see also [20, Lemma 15]) in order to find two continuously differentiable functions $\gamma_{\pm}: \mathbb{R} \rightarrow \mathbb{R}$ satisfying

$$\gamma_{-}(s) < -d < d < \gamma_{+}(s) \quad (9)$$

for every $s \in [\alpha - \delta, \beta + \delta]$ and

$$-\partial_u H(t, q, p, u, \gamma_{+}(u)) > \partial_v H(t, q, p, u, \gamma_{+}(u)) \gamma'_{+}(u), \quad (10)$$

$$-\partial_u H(t, q, p, u, \gamma_{-}(u)) < \partial_v H(t, q, p, u, \gamma_{-}(u)) \gamma'_{-}(u) \quad (11)$$

for every $(t, q, p, u) \in [0, T] \times [0, 2\pi] \times \mathbb{R} \times [\alpha - \delta, \beta + \delta]$. Correspondingly, we introduce the set

$$\mathcal{V} = \{z = (q, p, u, v) \mid \alpha \leq u \leq \beta, \gamma_{-}(u) < v < \gamma_{+}(u)\}.$$

Now we can choose a constant $\widehat{d} > d$ satisfying

$$-\widehat{d} < \gamma_{-}(s) < \gamma_{+}(s) < \widehat{d}$$

for every $s \in [\alpha - \delta, \beta + \delta]$.

We consider a continuously differentiable function $\zeta: \mathbb{R} \rightarrow \mathbb{R}$ such that

$$\zeta(u) = \begin{cases} \alpha - \delta & \text{if } u \leq \alpha - 2\delta, \\ u & \text{if } \alpha \leq u \leq \beta, \\ \beta + \delta & \text{if } u \geq \beta + 2\delta \end{cases} \quad (12)$$

and

$$\zeta'(u) > 0 \quad \text{if } u \in]\alpha - 2\delta, \beta + 2\delta[. \quad (13)$$

Notice that

$$\alpha - \delta \leq \zeta(u) \leq \beta + \delta \quad \text{for every } u \in \mathbb{R}.$$

Then we introduce a continuously differentiable function $\chi: [\alpha - \delta, \beta + \delta] \times \mathbb{R} \rightarrow \mathbb{R}$ satisfying the following properties:

$$\chi(u, v) = \begin{cases} -\widehat{d} & \text{if } v < -\widehat{d} - 1, \\ v & \text{if } \gamma_{-}(u) \leq v \leq \gamma_{+}(u), \\ \widehat{d} & \text{if } v > \widehat{d} + 1. \end{cases} \quad (14)$$

Moreover, we assume that

$$\partial_v \chi(u, v) > 0 \quad \text{if } |v| < \widehat{d} + 1. \quad (15)$$

Let $\widehat{H}: \mathbb{R} \times \mathbb{R}^4 \rightarrow \mathbb{R}$ be defined as

$$\widehat{H}(t, q, p, u, v) = H(t, q, p, \zeta(u), \chi(\zeta(u), v)) + \mathcal{H}(u, v),$$

with

$$\mathcal{H}(u, v) = \frac{1}{2} \left[\left[(v - \widehat{d})^+ \right]^2 + \left[(v + \widehat{d})^- \right]^2 - [(u - \beta)^+]^2 - [(u - \alpha)^-]^2 \right],$$

where, as usual, $\xi^+ = \max\{\xi, 0\}$ and $\xi^- = \max\{-\xi, 0\}$. We consider the modified Hamiltonian system

$$\begin{cases} \dot{q} = \partial_p \widehat{H}(t, q, p, u, v), & \dot{p} = -\partial_q \widehat{H}(t, q, p, u, v), \\ \dot{u} = \partial_v \widehat{H}(t, q, p, u, v), & \dot{v} = -\partial_u \widehat{H}(t, q, p, u, v). \end{cases} \tag{16}$$

Notice that $\widehat{H} = H$ in the closure of the set \mathcal{V} . Our aim is to prove the following a priori bound.

Lemma 3. *If $z = (q, p, u, v)$ is a T -periodic solution of (16), then $z(t) \in \mathcal{V}$ for every $t \in [0, T]$ and hence z solves (2).*

The proof of this lemma needs some preparation, hence it will be provided at the end of the section. In what follows, for every $z = (q, p, u, v)$ we introduce the notation

$$\tilde{z} = (q, p, \zeta(u), \chi(\zeta(u), v)).$$

We can explicitly compute

$$\partial_q \widehat{H}(t, z) = \partial_q H(t, \tilde{z}), \quad \partial_p \widehat{H}(t, z) = \partial_p H(t, \tilde{z}), \tag{17}$$

$$\partial_u \widehat{H}(t, z) = [\partial_u H(t, \tilde{z}) + \partial_v H(t, \tilde{z}) \partial_u \chi(\zeta(u), v)] \zeta'(u) + \partial_u \mathcal{H}(u, v), \tag{18}$$

$$\partial_v \widehat{H}(t, z) = \partial_v H(t, \tilde{z}) \partial_v \chi(\zeta(u), v) + \partial_v \mathcal{H}(u, v). \tag{19}$$

Proposition 4. *Any solution of (16) satisfies, for every $t_0 \in \mathbb{R}$,*

$$\begin{aligned} [u(t_0) < \alpha \text{ and } v(t_0) = 0] &\Rightarrow \dot{v}(t_0) < 0, \\ [u(t_0) > \beta \text{ and } v(t_0) = 0] &\Rightarrow \dot{v}(t_0) > 0. \end{aligned}$$

Proof. We first note that, as an immediate consequence of (3), we have

$$\partial_v H(t, q, p, u, 0) = 0 \text{ if } u \in [\alpha - \delta, \alpha] \cup [\beta, \beta + \delta]. \tag{20}$$

We prove the first implication, the second one being similar. Let $t_0 \in \mathbb{R}$ be such that $u(t_0) < \alpha$ and $v(t_0) = 0$. Let $z(t)$ be a solution of (16), with $z(t_0) = (q(t_0), p(t_0), u(t_0), 0)$. Then $\tilde{z}(t_0) = (q(t_0), p(t_0), \zeta(u(t_0)), 0)$. By (18) and (20) we have that

$$\begin{aligned} \dot{v}(t_0) &= -\partial_u \widehat{H}(t_0, z(t_0)) = -\partial_u H(t_0, \tilde{z}(t_0)) \zeta'(u(t_0)) - \partial_u \mathcal{H}(u(t_0), 0) \\ &= -\partial_u H(t_0, \tilde{z}(t_0)) \zeta'(u(t_0)) + u(t_0) - \alpha, \end{aligned} \tag{21}$$

so that

$$\dot{v}(t_0) < -\partial_u H(t_0, \tilde{z}(t_0)) \zeta'(u(t_0)).$$

Then (4) and (13) give the negative sign if $u(t_0) \in]\alpha - 2\delta, \alpha[$. On the other hand, $\zeta'(u)$ vanishes if $u \leq \alpha - 2\delta$ and the conclusion easily follows from (21). \square

Proposition 5. Any solution of (16) satisfies, for every $t_0 \in \mathbb{R}$,

$$\begin{aligned} [u(t_0) \leq \alpha \text{ and } v(t_0) < 0] &\Rightarrow \dot{u}(t_0) < 0, \\ [u(t_0) \leq \alpha \text{ and } v(t_0) > 0] &\Rightarrow \dot{u}(t_0) > 0, \\ [u(t_0) \geq \beta \text{ and } v(t_0) < 0] &\Rightarrow \dot{u}(t_0) < 0, \\ [u(t_0) \geq \beta \text{ and } v(t_0) > 0] &\Rightarrow \dot{u}(t_0) > 0. \end{aligned}$$

Proof. Let us prove the first assertion, the proof of the others being similar. Let $t_0 \in \mathbb{R}$ be such that $u(t_0) \leq \alpha$ and $v(t_0) < 0$. Since $\zeta(u) \in [\alpha - \delta, \alpha]$ when $u \leq \alpha$ and $\chi(u, v) < 0$ when $v < 0$, recalling (3), (15), and (19), if $-\widehat{d} - 1 < v(t_0) < 0$ we get

$$\begin{aligned} \dot{u}(t_0) &= \partial_v H(t_0, \tilde{z}(t_0)) \partial_v \chi(\zeta(u(t_0)), v(t_0)) - (v(t_0) + \widehat{d})^- \\ &\leq \partial_v H(t_0, \tilde{z}(t_0)) \partial_v \chi(\zeta(u(t_0)), v(t_0)) < 0. \end{aligned}$$

On the other hand, if $v(t_0) \leq -\widehat{d} - 1$, then $\partial_v \chi(\zeta(u(t_0)), v(t_0)) = 0$, so that $\dot{u}(t_0) = -(v(t_0) + \widehat{d})^- < 0$. \square

We define the open sets

$$\begin{aligned} A_{NW} &= \{z \in \mathbb{R}^4 \mid u < \alpha, v > 0\}, & A_{NE} &= \{z \in \mathbb{R}^4 \mid u > \beta, v > 0\}, \\ A_{SW} &= \{z \in \mathbb{R}^4 \mid u < \alpha, v < 0\}, & A_{SE} &= \{z \in \mathbb{R}^4 \mid u > \beta, v < 0\}. \end{aligned}$$

As a consequence of the previous propositions, the following can be easily proved.

Proposition 6. The sets A_{NE} and A_{SW} are positively invariant for system (16), while A_{SE} and A_{NW} are negatively invariant.

Proposition 7. Any solution of (16) satisfies, for every $t_0 \in \mathbb{R}$,

$$\begin{aligned} [\alpha \leq u(t_0) \leq \beta \text{ and } v(t_0) > d] &\Rightarrow \dot{u}(t_0) > 0, \\ [\alpha \leq u(t_0) \leq \beta \text{ and } v(t_0) < -d] &\Rightarrow \dot{u}(t_0) < 0. \end{aligned}$$

Proof. Let us prove the first implication. Since $u(t_0) \in [\alpha, \beta]$ and $v(t_0) > d$, we have that $\zeta(u(t_0)) = u(t_0)$ and

$$\dot{u}(t_0) = \partial_v H(t_0, \tilde{z}(t_0)) \partial_v \chi(u(t_0), v(t_0)) + (v(t_0) - \widehat{d})^+.$$

If $v(t_0) \geq \widehat{d} + 1$, then $\partial_v \chi(u(t_0), v(t_0)) = 0$ and $(v(t_0) - \widehat{d})^+ > 0$. On the other hand, if $d < v(t_0) < \widehat{d} + 1$, then $\partial_v \chi(u(t_0), v(t_0)) > 0$ and $(v(t_0) - \widehat{d})^+ \geq 0$; moreover, $\partial_v H(t_0, \tilde{z}(t_0)) > 0$ by Assumption 3, thus proving that $\dot{u}(t_0) > 0$ in both cases. \square

Proof of Lemma 3. Let z be a T -periodic solution of (16). We begin proving that $\alpha \leq u(t) \leq \beta$ for every t . By contradiction, assume that there exists $t_0 \in \mathbb{R}$ such that $u(t_0) > \beta$. If $z(t_0) \in A_{NE}$, by Proposition 6, since the function is periodic, $z(t) \in A_{NE}$ for every $t \in \mathbb{R}$. So, we get a contradiction using Proposition 5. Similarly we cannot have $z(t_0) \in A_{SE}$. Finally, if $v(t_0) = 0$, Proposition 4 takes us to the previous contradicting situations. We have thus proved that $u(t) \leq \beta$ for every $t \in \mathbb{R}$. A similar argument proves that $u(t) \geq \alpha$ for every $t \in \mathbb{R}$.

Now we show that $v(t) < \gamma_+(u(t))$ for every $t \in \mathbb{R}$. Let us define the function $G_+(t) = v(t) - \gamma_+(u(t))$. By Proposition 7, it cannot be that $G_+(t) > 0$ for every $t \in [0, T]$. Assume by contradiction the existence of $t_0 \in \mathbb{R}$ such that $G_+(t_0) = 0$. Then, since $\nabla H(t, z) = \nabla \widehat{H}(t, z)$ for z in the closure of \mathcal{V} , by (10) we have

$$G'_+(t_0) = -\partial_u H(t_0, z(t_0)) - \gamma'_+(u(t_0)) \partial_v H(t_0, z(t_0)) > 0,$$

where $z(t_0) = (q(t_0), p(t_0), u(t_0), \gamma_+(u(t_0)))$. This implies that $G_+(t) > 0$ for every $t > t_0$, which is in contradiction with the periodicity of z . We have thus proved that $G_+(t) < 0$ for every $t \in [0, T]$.

We can similarly prove that $\gamma_-(u(t)) < v(t)$ for every $t \in \mathbb{R}$, using (11). \square

3.2. Working with the coordinates (q, p)

Let us fix $K > 0$ such that

$$|\chi(\zeta(u), v)| < K \text{ for every } (u, v) \in \mathbb{R}^2.$$

Let $z(t) = (q(t), p(t), u(t), v(t))$ be a solution of system (16) starting at time $t = 0$ with $p(0) \in [a - \rho, b + \rho]$. This solution is defined on a maximal interval of future existence $[0, \omega[$. By Assumption 4, recalling (17),

$$|\dot{p}(t)| \leq C_K(|p(t)| + 1) \quad \text{for every } t \in [0, \omega[, \tag{22}$$

hence, setting

$$c = (\max\{|a|, |b|\} + \rho + 1) e^{C_K T}, \tag{23}$$

by the Gronwall Lemma we have that

$$|p(t)| \leq c \quad \text{for every } t \in [0, T] \cap [0, \omega[. \tag{24}$$

Let $\eta: \mathbb{R} \rightarrow \mathbb{R}$ be a C^∞ -smooth nonincreasing function such that

$$\eta(s) = \begin{cases} 1 & \text{if } s \leq c, \\ 0 & \text{if } s \geq c + 1. \end{cases} \tag{25}$$

We can rewrite the Hamiltonian \widehat{H} as

$$\widehat{H}(t, q, p, u, v) = \frac{1}{2}(v^2 - u^2) + \widehat{h}(t, q, p, u, v) \tag{26}$$

and define the new Hamiltonian function

$$\widetilde{H}(t, q, p, u, v) = \frac{1}{2}(v^2 - u^2) + \widetilde{h}(t, q, p, u, v), \tag{27}$$

where

$$\widetilde{h}(t, q, p, u, v) = \eta(|p|) \widehat{h}(t, q, p, u, v).$$

Notice that

$$\text{if } |p| \geq c + 1 \quad \text{then } \widetilde{h}(t, q, p, u, v) = 0 \quad \text{for every } (t, q, u, v) \in \mathbb{R} \times \mathbb{R}^3.$$

The Hamiltonian \tilde{H} induces the system

$$\begin{cases} \dot{q} = \partial_p \tilde{h}(t, q, p, u, v), & \dot{p} = -\partial_q \tilde{h}(t, q, p, u, v), \\ \dot{u} = v + \partial_v \tilde{h}(t, q, p, u, v), & \dot{v} = u - \partial_u \tilde{h}(t, q, p, u, v). \end{cases} \tag{28}$$

Since the Hamiltonian function is periodic in q and the functions $\zeta, \zeta', \chi, \nabla\chi, \eta$ and η' are bounded, there exists a constant $\tilde{C} > 0$ such that

$$|\partial_q \tilde{h}(t, z)| + |\partial_p \tilde{h}(t, z)| + |\partial_u \tilde{h}(t, z)| + |\partial_v \tilde{h}(t, z)| \leq \tilde{C}$$

for every $(t, z) \in \mathbb{R} \times \mathbb{R}^4$. Hence, the Hamiltonian function \tilde{H} is the sum of a nonresonant quadratic term and a function with bounded gradient $\nabla \tilde{h}(t, z)$. In particular, all solutions of (28) are globally defined on $[0, T]$.

We now verify the twist condition for system (28). Let $z = (q, p, u, v)$ be a solution of (28) such that $p(0) \in [a - \rho, b + \rho]$. As long as $|p(t)| \leq c$, we have

$$|\dot{p}(t)| = |\partial_q H(t, q(t), p(t), \zeta(u(t)), \chi(\zeta(u(t))), v(t))| \leq C_K(|p(t)| + 1).$$

Hence, by the Gronwall Lemma, we conclude that $|p(t)| \leq c$ for every $t \in [0, T]$. So, $z = (q, p, u, v)$ is a solution of (16). Then (q, p) is a solution of (6) with $U(t) = \zeta(u(t))$ and $V(t) = \chi(\zeta(u(t)), v(t))$, condition (5) holds and, by (7),

$$\begin{aligned} q(T) &< q(0) && \text{if } p(0) \in [a - \rho, a], \\ q(T) &> q(0) && \text{if } p(0) \in [b, b + \rho]. \end{aligned}$$

Hence, we can apply [18, Corollary 2.4] so to find two T -periodic solutions $z = (q, p, u, v)$ of (28) such that $p(0) \in]a, b[$. By the above estimates, these are indeed solutions of (16) and, recalling Lemma 3, we conclude that they are the T -periodic solutions of the original system (2) we were looking for. \square

4. Some variants of Theorem 2

Let us start with two observations.

Remark 8. Assumption 2 can be generalized by asking that there further exist v_α, v_β such that

$$\begin{aligned} (v - v_\alpha) \partial_v H(t, q, p, u, v) &> 0 && \text{if } u \in [\alpha - \delta, \alpha] \text{ and } v \neq v_\alpha, \\ (v - v_\beta) \partial_v H(t, q, p, u, v) &> 0 && \text{if } u \in [\beta, \beta + \delta] \text{ and } v \neq v_\beta \end{aligned}$$

and

$$\begin{aligned} \partial_u H(t, q, p, u, v_\alpha) &\geq 0 && \text{if } u \in [\alpha - \delta, \alpha], \\ \partial_u H(t, q, p, u, v_\beta) &\leq 0 && \text{if } u \in [\beta, \beta + \delta]. \end{aligned}$$

Remark 9. Instead of a fixed interval $[a, b]$, we could have a varying interval $[a(q), b(q)]$, where $a, b: \mathbb{R} \rightarrow \mathbb{R}$ are continuous 2π -periodic functions. Indeed, if a and b are continuously differentiable, this case can be reduced to the previous one by the symplectic change of variables

$$\Psi(q, p, u, v) = \left(\int_0^q \frac{b(s) - a(s)}{2} ds, \frac{2p - b(q) - a(q)}{b(q) - a(q)}, u, v \right),$$

cf. [43, Exercise 1, p. 132]. If a and b are just continuous, they can be replaced by smooth functions by the use of the Fejer Theorem. Notice that the new Hamiltonian $\tilde{H}(t, \tilde{q}, \tilde{p}, u, v) = H(t, \Psi^{-1}(\tilde{q}, \tilde{p}, u, v))$ is periodic in \tilde{q} with period $\tau := \frac{1}{2} \int_0^{2\pi} (b(s) - a(s)) ds$.

We now propose a variant of our result which is more in the spirit of the Poincaré–Birkhoff Theorem as originally stated by Poincaré [46]. We first recall the definition of rotation number. For $t_1 < t_2$, let $\eta: [t_1, t_2] \rightarrow \mathbb{R}^2$ be a continuous curve such that $\eta(t) \neq (0, 0)$ for every $t \in [t_1, t_2]$. Writing $\eta(t) = \rho(t)(\cos \theta(t), \sin \theta(t))$, with $\rho: \mathbb{R} \rightarrow]0, +\infty[$ and $\theta: \mathbb{R} \rightarrow \mathbb{R}$ continuous, we define

$$\text{Rot}(\eta; [t_1, t_2]) = -\frac{\theta(t_2) - \theta(t_1)}{2\pi}.$$

We need to suitably modify Assumption 4.

Assumption 5 (Energy growth). For every $K > 0$, there is a constant $C_K > 0$ such that

$$\begin{aligned} |q \partial_p H(t, q, p, u, v) - p \partial_q H(t, q, p, u, v)| &\leq C_K(q^2 + p^2 + 1) \\ \text{if } u &\in [\alpha - \delta, \beta + \delta] \text{ and } |v| \leq K. \end{aligned}$$

In the sequel, we denote by $\mathcal{D}(\Gamma)$ the open bounded region delimited by a planar Jordan curve Γ .

Theorem 10. Let Assumptions 2, 3, and 5 hold. Let k be an integer and assume that there exist $\rho > 0$, $\tilde{\rho} > 0$ and two planar Jordan curves Γ_1 and Γ_2 , strictly star-shaped with respect to the origin, with

$$0 \in \mathcal{D}(\Gamma_1) \subseteq \overline{\mathcal{D}(\Gamma_1)} \subseteq \mathcal{D}(\Gamma_2),$$

such that, for any two continuous functions $U, V: [0, T] \rightarrow \mathbb{R}$ satisfying (5), the solutions of system (6) with $\text{dist}((q(0), p(0)), \overline{\mathcal{D}(\Gamma_2)} \setminus \mathcal{D}(\Gamma_1)) \leq \rho$ which are defined on $[0, T]$ satisfy

$$q(t)^2 + p(t)^2 \geq \tilde{\rho} \quad \text{for every } t \in [0, T]$$

and, when $(q(0), p(0)) \notin \overline{\mathcal{D}(\Gamma_2)} \setminus \overline{\mathcal{D}(\Gamma_1)}$,

$$\begin{aligned} \text{Rot}((q, p); [0, T]) &< k \quad \text{if } \text{dist}((q(0), p(0)), \Gamma_1) \leq \rho, \\ \text{Rot}((q, p); [0, T]) &> k \quad \text{if } \text{dist}((q(0), p(0)), \Gamma_2) \leq \rho. \end{aligned} \tag{29}$$

Then the Hamiltonian system (2) has at least two T -periodic solutions $z_i(t)$, $i = 1, 2$, satisfying (8) with

$$(q_i(0), p_i(0)) \in \mathcal{D}(\Gamma_2) \setminus \overline{\mathcal{D}(\Gamma_1)}$$

and

$$\text{Rot}((q_i, p_i); [0, T]) = k.$$

The same is true if (29) is replaced by

$$\begin{aligned} \text{Rot}((q, p); [0, T]) &> k \quad \text{if } \text{dist}((q(0), p(0)), \Gamma_1) \leq \rho, \\ \text{Rot}((q, p); [0, T]) &< k \quad \text{if } \text{dist}((q(0), p(0)), \Gamma_2) \leq \rho. \end{aligned}$$

Proof. At first, let $R_0 > 0$ be such that

$$\text{dist}((q(0), p(0)), \overline{\mathcal{D}(\Gamma_2)}) \leq \rho \quad \Rightarrow \quad q(0)^2 + p(0)^2 \leq R_0. \quad (30)$$

We modify the Hamiltonian H introducing the function \widehat{H} , arguing as in Section 3.1, thus finding the a priori bound in Lemma 3. Let $K > 0$ be such that

$$|\chi(\zeta(u), v)| < K \text{ for every } (u, v) \in \mathbb{R}^2.$$

Let $z(t) = (q(t), p(t), u(t), v(t))$ be any solution of system (16) starting at time $t = 0$ from a point $z(0)$ satisfying $q(0)^2 + p(0)^2 \leq R_0$. This solution is defined on a maximal interval of future existence $[0, \omega[$. Defining $r(t) = q(t)^2 + p(t)^2$, by Assumption 5,

$$|\dot{r}(t)| \leq C_K(r(t) + 1) \quad \text{for every } t \in [0, \omega[,$$

hence, setting

$$c = (R_0 + 1)e^{C_K T},$$

by the Gronwall Lemma we have that

$$r(t) \leq c \quad \text{for every } t \in [0, T] \cap [0, \omega[.$$

Let η be the function introduced in (25). Arguing as above, we can write \widehat{H} as in (26) and define \widetilde{H} as in (27), where now

$$\widetilde{h}(t, q, p, u, v) = \eta(q^2 + p^2) \widehat{h}(t, q, p, u, v).$$

The Hamiltonian function \widetilde{H} induces the system (28), whose solutions are globally defined on $[0, T]$.

Let $z(t) = (q(t), p(t), u(t), v(t))$ be any solution of system (28) with starting point $z(0)$ satisfying $\text{dist}((q(0), p(0)), \overline{\mathcal{D}(\Gamma_2)} \setminus \mathcal{D}(\Gamma_1)) \leq \rho$. Set $U(t) = \zeta(u(t))$ and $V(t) = \chi(\zeta(u(t)), v(t))$. Notice that (5) is satisfied. It can be verified that $r(t) = q(t)^2 + p(t)^2 \leq c$ for every $t \in [0, T]$, with the same constant c as above. Therefore, the couple $(q(t), p(t))$ is a solution of (6). So, by the assumption of the theorem, we get

$$r(t) \geq \tilde{\rho} \quad \text{for every } t \in [0, T]. \quad (31)$$

We now introduce a new smooth cut-off increasing function $\widetilde{\eta}: \mathbb{R} \rightarrow [0, 1]$ such that

$$\widetilde{\eta}(s) = \begin{cases} 0 & \text{if } s \leq \tilde{\rho}/2, \\ 1 & \text{if } s \geq \tilde{\rho} \end{cases}$$

and the new Hamiltonian function

$$\overline{H}(t, q, p, u, v) = \frac{1}{2}(v^2 - u^2) + \widetilde{h}(t, q, p, u, v),$$

where $\widetilde{h}(t, q, p, u, v) = \widetilde{\eta}(q^2 + p^2) \widetilde{h}(t, q, p, u, v)$.

For the new Hamiltonian system $J\dot{z} = \nabla \overline{H}(t, z)$, we introduce the symplectic change of variables

$$q(t) = \sqrt{2r(t)} \cos\left(\theta(t) - \frac{2\pi}{T} kt\right), \quad p(t) = \sqrt{2r(t)} \sin\left(\theta(t) - \frac{2\pi}{T} kt\right),$$

so to get a Hamiltonian system

$$\begin{cases} \dot{\theta} = \partial_r \mathcal{H}(t, \theta, r, u, v), & \dot{r} = -\partial_\theta \mathcal{H}(t, \theta, r, u, v), \\ \dot{u} = \partial_v \mathcal{H}(t, \theta, r, u, v), & \dot{v} = -\partial_u \mathcal{H}(t, \theta, r, u, v), \end{cases} \tag{32}$$

defined for $r > 0$. This system can now be extended also for $r \leq 0$, preserving the regularity. Now, the periodicity Assumption 1 is recovered in the variable θ . A simple computation shows that

$$\dot{r} = q \partial_p H(t, q, p, u, v) - p \partial_q H(t, q, p, u, v),$$

hence Assumption 5 implies that the linear growth Assumption 4 holds for the new variables (θ, r) instead of (q, p) . Then Theorem 2 applies, in view of Remark 9, providing the existence of two geometrically distinct T -periodic solutions of (32), which are then translated, by the inverse change of variables $(\theta, r) \mapsto (q, p)$, into the T -periodic solutions of (2) we are looking for. \square

5. Examples of applications

Both Theorems 2 and 10 open the way to a multitude of applications. We will just sketch here a few. Let us consider for example a system of the type

$$\begin{cases} -\ddot{q} = g(t, q) - e(t) + \partial_q P(t, q, u), \\ -\ddot{u} = -f(u) + \partial_u P(t, q, u), \end{cases} \tag{33}$$

where all functions are continuous and T -periodic in t , with $\int_0^T e(t) dt = 0$. Let $P(t, q, u)$ be 2π -periodic in q and continuously differentiable in (q, u) . Assume moreover $g(t, q)$ to be 2π -periodic in q , with $\int_0^{2\pi} g(t, s) ds = 0$.

Then system (33) is of the type (2), with

$$H(t, q, p, u, v) = \frac{1}{2}(p + E(t))^2 + \frac{1}{2}v^2 + \int_0^q g(t, s) ds - \int_0^u f(\sigma) d\sigma + P(t, q, u),$$

where $E(t) = \int_0^t e(s) ds$. Precisely, we have

$$\begin{cases} \dot{q} = p + E(t), & \dot{p} = -g(t, q) - \partial_q P(t, q, u), \\ \dot{u} = v, & \dot{v} = f(u) - \partial_u P(t, q, u). \end{cases}$$

Corollary 11. *In the above setting, assume in addition that there exist two constants $\alpha < \beta$ such that*

$$f(\alpha) < \partial_u P(t, q, \alpha), \quad \partial_u P(t, q, \beta) < f(\beta), \tag{34}$$

for all $(t, q) \in [0, T] \times [0, 2\pi]$. Then system (33) has at least two geometrically distinct T -periodic solutions (q, u) , with $\alpha \leq u \leq \beta$.

Proof. Notice that $H(t, q, p, u, v)$ is 2π -periodic in q . By continuity, there is a sufficiently small $\delta > 0$ such that Assumption 2 is satisfied. Also Assumption 3 is easily verified, setting $f(s) = s$ and $\varphi(s)$ constant. Moreover, there exists $R > 0$ with the following property: For any continuous function $U: [0, T] \rightarrow \mathbb{R}$ such that $\alpha - \delta \leq U(t) \leq \beta + \delta$ for every $t \in [0, T]$, the solutions of

$$-\ddot{q} = g(t, q) - e(t) + \partial_q P(t, q, U(t))$$

satisfy

$$\begin{aligned} q(T) - q(0) &< 0 & \text{if } \dot{q}(0) < -R, \\ q(T) - q(0) &> 0 & \text{if } \dot{q}(0) > R. \end{aligned}$$

Then taking $b = -a = R + \|E\|_\infty$, Theorem 2 applies. \square

As an immediate consequence, we have the following.

Corollary 12. *The system of coupled pendulums of the form*

$$\begin{cases} \ddot{q} + a \sin q = e(t) - \partial_q P(t, q, u), \\ \ddot{u} + b \sin u = -\partial_u P(t, q, u), \end{cases} \quad (35)$$

where $P(t, q, u)$ and $e(t)$ are as above, has at least two geometrically distinct T -periodic solutions for any $a > 0$ if $\|\partial_u P\|_\infty < b$.

We thus extend two classical results on the pendulum equation (cf. [42]). The first one states that, if $e(t)$ is T -periodic and $\int_0^T e(t) dt = 0$, then the equation

$$\ddot{q} + a \sin q = e(t)$$

has at least two geometrically distinct T -periodic solutions, for any $a > 0$. The second one states that, if $\hat{e}(t)$ is T -periodic and $\|\hat{e}\|_\infty \leq b$, then the equation

$$\ddot{u} + b \sin u = \hat{e}(t)$$

has at least two geometrically distinct T -periodic solutions.

Here are other possible examples of applications.

1. Let $f(u) = |u|^{\ell-1}u$, with $\ell > 0$, and let P be a function as above such that

$$\lim_{u \rightarrow \pm\infty} \frac{\partial_u P(t, q, u)}{|u|^\ell} = 0 \quad \text{uniformly in } (t, q) \in [0, T] \times [0, 2\pi].$$

Then there exist two constants α and β with $-\alpha = \beta > 0$ large enough such that (34) is satisfied. In particular, if $\partial_u P$ is bounded, it is enough to take

$$\alpha < -\|\partial_u P\|_\infty^{1/\ell}, \quad \beta > \|\partial_u P\|_\infty^{1/\ell}.$$

Notice that this includes, e.g., the case of $\partial_u P$ being periodic in u . Hence, in this situation, there exist at least two geometrically distinct T -periodic solutions of system (33).

2. Let $f(u) = |u|^\ell \sin(u)$, with $\ell > 0$, and let P be as in the previous example. Then (34) is satisfied for an infinite number of positive and negative pairs (α_i, β_i) . Hence, there exist infinitely many T -periodic solutions of system (33) with positive u component and infinitely many T -periodic solutions with negative u component.

3. Consider $f(u) = \arctan(u)$. Then, if $\partial_u P$ is bounded with

$$\|\partial_u P\|_\infty < \frac{\pi}{2},$$

there exist at least two geometrically distinct T -periodic solutions of system (33). It is sufficient to take α and β with $-\alpha = \beta > 0$ large enough.

If $P(t, q, u)$ is not periodic in q , we can appeal to Theorem 10. Consider for example the system

$$\begin{cases} -\ddot{q} = g(q) + \partial_q P(t, q, u), \\ -\ddot{u} = -f(u) + \partial_u P(t, q, u), \end{cases} \tag{36}$$

where $P(t, q, u)$ is continuously differentiable in (q, u) . We can prove the following.

Corollary 13. *Assume that*

$$\lim_{|q| \rightarrow \infty} \frac{g(q)}{q} = +\infty \tag{37}$$

and that $\partial_q P$ is bounded. If there exist two constants $\alpha < \beta$ such that (34) holds, then system (36) has infinitely-many T -periodic solutions.

Proof. We will briefly explain the main arguments. At first we modify system (36) by introducing the functions

$$\begin{aligned} \widehat{f}(u) &= (u - \beta)^+ - (u - \alpha)^- + f(\zeta(u))\zeta'(u), \\ \widehat{P}(t, q, u) &= P(t, q, \zeta(u)), \end{aligned}$$

thus obtaining

$$\begin{cases} \dot{q} = p, & \dot{p} = -g(q) - \partial_q \widehat{P}(t, q, u), \\ \dot{u} = v, & \dot{v} = \widehat{f}(u) - \partial_u \widehat{P}(t, q, u) \end{cases} \tag{38}$$

and denoting by $\widehat{H}(t, z) = \frac{1}{2}(v^2 - u^2) + \widehat{h}(t, z)$ its Hamiltonian, for a suitable choice of \widehat{h} . Let $B(0, R) \subseteq \mathbb{R}^2$ be the ball of radius $R > 0$ centred at the origin, and set $\mathcal{B}_R := B(0, R) \times \mathbb{R}^2$. Then, following the arguments in [13, Section 2], we can prove that any solution of system (38) is globally defined in the interval $[0, T]$. Moreover, by the elastic property stated in [13, Lemma 1] we can find $r_3 > r_2 > r_1 > 0$ such that any solution $z(t) = (q(t), p(t), u(t), v(t))$ of (38) starting at $t = 0$ from a point belonging to $\partial\mathcal{B}_{r_2}$ satisfies $z(t) \in \mathcal{B}_{r_3} \setminus \mathcal{B}_{r_1}$ for every $t \in [0, T]$. For such solutions we can find $k > 0$ such that $\text{Rot}((q, p); [0, T]) < k$.

Let us fix any $k' \geq k$. Assumption (37) provides the existence of $R_1 > 0$ such that every solution z of (38), with $z(t) \notin \mathcal{B}_{R_1}$ for any $t \in [0, T]$, satisfies $\text{Rot}((q, p); [0, T]) > k'$. Then, using again [13, Lemma 1], we can find $R_3 > R_2 > R_1$ such that any solution z of (38) starting at $t = 0$ from a point belonging to $\partial\mathcal{B}_{R_2}$ satisfies $z(t) \in \mathcal{B}_{R_3} \setminus \mathcal{B}_{R_1}$ for every $t \in [0, T]$. Hence we have for such solutions $\text{Rot}((q, p); [0, T]) > k'$.

Now, we introduce a smooth cut-off function $\widehat{\eta}: [0, +\infty[\rightarrow [0, 1]$ such that

$$\widehat{\eta}(r) = \begin{cases} 0 & \text{if } r \in [0, r_1/2] \cup [2R_3, +\infty[, \\ 1 & \text{if } r \in [r_1, R_3] \end{cases}$$

and the new Hamiltonian $\widetilde{H}(t, z) = \frac{1}{2}(v^2 - u^2) + \widetilde{h}(t, z)$ where now

$$\tilde{h}(t, z) = \hat{\eta}(\sqrt{q(t)^2 + p(t)^2})\hat{h}(t, z).$$

Finally, we are able to apply Theorem 10 to system (38), thus obtaining two T -periodic solutions satisfying $\text{Rot}((q, p); [0, T]) = k'$ with starting point $z(0) \in \mathcal{B}_{R_2} \setminus \overline{\mathcal{B}_{r_2}}$. Then we can easily see that they are indeed solutions of (36).

Since the above construction can be made for every $k' \geq k$, we have proved the existence of infinitely many T -periodic solutions. \square

We thus extend a result by Ding and Zanolin [13], stating that, if $g(q)$ satisfies (37) and $e(t)$ is T -periodic, the scalar equation

$$\ddot{q} + g(q) = e(t)$$

has infinitely-many T -periodic solutions. A similar result for system (33) remains an open problem, because global existence of the solutions is not guaranteed. However, one could follow the lines of [25,34–36] by assuming that the first equation has 0 as an equilibrium point, and then prove that there are infinitely many T -periodic solutions.

Whenever the function g has a sublinear growth at infinity, the existence of periodic solutions whose minimal period is an arbitrarily large integer multiple of T has been investigated in [14]. These are called *subharmonic* solutions. We could also state such kind of result here, but we avoid the details, for brevity.

Another situation where our results can be applied is when the system in (q, p) has a different rotational behaviour at zero and at infinity. There are many papers dealing with such a problem (see, e.g., [39] and the references therein). This type of situation can be also exploited when dealing with some kind of asymmetric oscillators as, e.g., in [7], where the so-called *jumping nonlinearities* are treated. We do not enter into details, again, to be brief.

6. Going to higher dimensions

We consider system (1), assuming the Hamiltonian function $H: \mathbb{R} \times \mathbb{R}^{2N} \rightarrow \mathbb{R}$ to be continuous, T -periodic in its first variable t , and continuously differentiable with respect to its second variable z , with corresponding gradient $\nabla H(t, z)$.

For $z \in \mathbb{R}^{2N}$, we write $z = (\phi, \psi, q, p, u, v)$, where, for some nonnegative integers L, M and D ,

$$\begin{aligned} \phi &= (\phi_1, \dots, \phi_L) \in \mathbb{R}^L, & \psi &= (\psi_1, \dots, \psi_L) \in \mathbb{R}^L, \\ q &= (q_1, \dots, q_M) \in \mathbb{R}^M, & p &= (p_1, \dots, p_M) \in \mathbb{R}^M, \\ u &= (u_1, \dots, u_D) \in \mathbb{R}^D, & v &= (v_1, \dots, v_D) \in \mathbb{R}^D. \end{aligned}$$

Notice that one or more of these integers could be equal to zero, in which case the corresponding group will not be taken into account; for example, if $L = 0$, then ϕ and ψ will disappear from the list.

Our system (1) then reads as

$$\begin{cases} \dot{\phi} = \partial_{\psi} H(t, z), & \dot{\psi} = -\partial_{\phi} H(t, z), \\ \dot{q} = \partial_p H(t, z), & \dot{p} = -\partial_q H(t, z), \\ \dot{u} = \partial_v H(t, z), & \dot{v} = -\partial_u H(t, z). \end{cases} \tag{39}$$

Let us introduce our assumptions.

Assumption 6 (Periodicity). The function $H(t, z)$ is periodic in each of the variables included in ϕ, ψ, q .

To fix the ideas, we assume that all periods are equal to 2π . The total number of variables in which our Hamiltonian function is 2π -periodic is thus $2L + M$.

In the sequel, inequalities \leq involving vectors are to be interpreted componentwise. Moreover, for any $\sigma \in \mathbb{R}$, we use the notation $\bar{\sigma} = (\sigma, \dots, \sigma) \in \mathbb{R}^D$.

Assumption 7 (*Lower and upper solutions*). There exist some constants $\delta > 0$, $\alpha = (\alpha_1, \dots, \alpha_D)$ and $\beta = (\beta_1, \dots, \beta_D)$ with $\alpha \leq \beta$, having the following property. When $\alpha - \bar{\delta} \leq u \leq \beta + \bar{\delta}$, then, for every $j \in \{1, \dots, D\}$,

$$v_j \partial_{v_j} H(t, z) > 0 \text{ if } u_j \in [\alpha_j - \delta, \alpha_j] \cup [\beta_j, \beta_j + \delta] \text{ and } v_j \neq 0,$$

and

$$\begin{aligned} \partial_{u_j} H(t, z) &\geq 0 && \text{if } u_j \in [\alpha_j - \delta, \alpha_j] \text{ and } v_j = 0, \\ \partial_{u_j} H(t, z) &\leq 0 && \text{if } u_j \in [\beta_j, \beta_j + \delta] \text{ and } v_j = 0. \end{aligned}$$

In the sequel, the constant $\delta > 0$ provided by Assumption 7 will be used without further mention.

Assumption 8 (*Nagumo condition*). For every $j \in \{1, \dots, D\}$ there exist $d_j > 0$ and two continuous functions $f_j, \varphi_j: [d_j, +\infty[\rightarrow]0, +\infty[$ with

$$\int_{d_j}^{+\infty} \frac{f_j(s)}{\varphi_j(s)} ds = +\infty,$$

satisfying the following property. When $\alpha - \bar{\delta} \leq u \leq \beta + \bar{\delta}$, then

$$\begin{aligned} \partial_{v_j} H(t, z) &\geq f_j(v_j) && \text{if } v_j \geq d_j, \\ \partial_{v_j} H(t, z) &\leq -f_j(-v_j) && \text{if } v_j \leq -d_j, \end{aligned}$$

and

$$|\partial_{u_j} H(t, z)| \leq \varphi_j(|v_j|) \text{ if } |v_j| \geq d_j.$$

Assumption 9 (*Linear growth*). For every $K > 0$ there is a constant $C_K > 0$ with the following property. If $\alpha - \bar{\delta} \leq u \leq \beta + \bar{\delta}$ and $|v| \leq K$, then

$$|\partial_q H(t, z)| \leq C_K(|p| + 1).$$

Adapting the argument in Remark 1, under the above assumption, for any two continuous functions $U, V: [0, T] \rightarrow \mathbb{R}^D$ with

$$\alpha - \bar{\delta} \leq U(t) \leq \beta + \bar{\delta} \text{ for every } t \in [0, T], \tag{40}$$

the solutions of the system

$$\begin{cases} \dot{\phi} = \partial_\psi H(t, \phi, \psi, q, p, U(t), V(t)), \\ \dot{\psi} = -\partial_\phi H(t, \phi, \psi, q, p, U(t), V(t)), \\ \dot{q} = \partial_p H(t, \phi, \psi, q, p, U(t), V(t)), \\ \dot{p} = -\partial_q H(t, \phi, \psi, q, p, U(t), V(t)), \end{cases} \tag{41}$$

are defined on $[0, T]$ and, setting $K = \|V\|_\infty$,

$$|p(t)| \leq (|p(0)| + 1) e^{C_K T} \quad \text{for every } t \in [0, T]. \quad (42)$$

In the following, we consider a convex body \mathcal{D} of \mathbb{R}^M , i.e., a closed convex bounded set with nonempty interior. We denote by $\pi_{\mathcal{D}}: \mathbb{R}^M \setminus \mathring{\mathcal{D}} \rightarrow \partial\mathcal{D}$ the projection on the convex set \mathcal{D} and by $\nu_{\mathcal{D}}(\zeta)$ the unit outward normal at $\zeta \in \partial\mathcal{D}$, assuming that \mathcal{D} has a smooth boundary. We say that \mathcal{D} is *strongly convex* if, for any $p \in \partial\mathcal{D}$, the function $\mathcal{F}: \mathcal{D} \rightarrow \mathbb{R}$ defined as $\mathcal{F}(\eta) = \langle \eta - p, \nu_{\mathcal{D}}(p) \rangle$ has a unique maximum point at $\eta = p$.

Here is our first result in this higher dimensional setting, generalizing Theorem 2.

Theorem 14. *Let Assumptions 6, 7, 8, and 9 hold. Assume that there exist $\rho > 0$, a symmetric regular $M \times M$ matrix \mathbb{A} and a strongly convex body \mathcal{D} of \mathbb{R}^M , having a smooth boundary, with the following property: For any two continuous functions $U, V: [0, T] \rightarrow \mathbb{R}^D$ satisfying (40), the solutions of (41) with $p(0) \notin \mathring{\mathcal{D}}$ and $\text{dist}(p(0), \partial\mathcal{D}) \leq \rho$ are such that*

$$\langle q(T) - q(0), \mathbb{A}\nu_{\mathcal{D}}(\pi_{\mathcal{D}}(p(0))) \rangle > 0.$$

Then system (39) has at least $2L + M + 1$ geometrically distinct T -periodic solutions, satisfying

$$p(0) \in \mathring{\mathcal{D}}$$

and

$$\alpha \leq u(t) \leq \beta \quad \text{for every } t \in \mathbb{R}. \quad (43)$$

Remark 15. In analogy with [11, Theorem 2], we expect a larger number of solutions provided that all the T -periodic solutions are assumed to be nondegenerate. We will not pursue this issue for the sake of brevity.

7. Proof of Theorem 14

Since the arguments will be similar to those provided in Section 3, we will try to be brief.

At first we need to suitably modify the Hamiltonian system working componentwise in the (u, v) variables.

For every $j \in \{1, \dots, D\}$, from Assumption 8 we can find some continuously differentiable functions $\gamma_j^\pm: \mathbb{R} \rightarrow \mathbb{R}$ and then $\hat{d}_j > 0$ such that

$$-\hat{d}_j < \gamma_j^-(s) < -d_j, \quad d_j < \gamma_j^+(s) < \hat{d}_j,$$

for every $s \in [\alpha_j - \delta, \beta_j + \delta]$, satisfying

$$\begin{aligned} -\partial_{u_j} H(t, z) &> \partial_{v_j} H(t, z) (\gamma_j^+)'(u_j) & \text{if } v_j = \gamma_j^+(u_j), \\ -\partial_{u_j} H(t, z) &< \partial_{v_j} H(t, z) (\gamma_j^-)'(u_j) & \text{if } v_j = \gamma_j^-(u_j), \end{aligned}$$

for every $(t, z) \in [0, T] \times \mathbb{R}^{2N}$ with $u_j \in [\alpha_j - \delta, \beta_j + \delta]$.

We then define $\gamma^\pm: \mathbb{R}^D \rightarrow \mathbb{R}^D$ as

$$\gamma^\pm(u) = (\gamma_1^\pm(u_1), \dots, \gamma_D^\pm(u_D))$$

and introduce the set

$$\mathcal{V} = \{z = (\phi, \psi, q, p, u, v) \mid \alpha \leq u \leq \beta, \gamma^-(u) < v < \gamma^+(u)\},$$

recalling that we need to check the inequalities componentwise. We can then introduce some functions $\zeta_j: \mathbb{R} \rightarrow \mathbb{R}$ and $\chi_j: [\alpha_j - \delta, \beta_j + \delta] \times \mathbb{R} \rightarrow \mathbb{R}$ similarly as in (12) and (14), define

$$\begin{aligned} \zeta(u) &= (\zeta_1(u_1), \dots, \zeta_D(u_D)), \\ \chi(u, v) &= (\chi_1(u_1, v_1), \dots, \chi_D(u_D, v_D)), \end{aligned}$$

and consider the modified Hamiltonian $\widehat{H}: \mathbb{R} \times \mathbb{R}^{2N} \rightarrow \mathbb{R}$ as

$$\widehat{H}(t, \phi, \psi, q, p, u, v) = H(t, \phi, \psi, q, p, \zeta(u), \chi(\zeta(u), v)) + \mathcal{H}(u, v)$$

with

$$\mathcal{H}(u, v) = \frac{1}{2} \sum_{j=1}^D \left[[(v_j - \widehat{d}_j)^+]^2 + [(v_j + \widehat{d}_j)^-]^2 - [(u_j - \beta_j)^+]^2 - [(u_j - \alpha_j)^-]^2 \right].$$

The modified Hamiltonian system

$$J\dot{z} = \nabla \widehat{H}(t, z) \tag{44}$$

complies the following a priori bound, whose proof can be given adapting the one of Lemma 3, arguing separately on every couple of variables (u_j, v_j) , and verifying the validity of some analogues of Propositions 4, 5, 6, and 7.

Lemma 16. *If $z = (\phi, \psi, q, p, u, v)$ is a T -periodic solution of (44), then $z(t) \in \mathcal{V}$ for every $t \in [0, T]$, hence it solves (39).*

The next step involves the (q, p) variables. The reasoning in Section 3.2 can be adapted to higher dimension. We fix $K > 0$ such that

$$|\chi(\zeta(u), v)| < K \text{ for every } (u, v) \in \mathbb{R}^{2D} \tag{45}$$

and, given the convex body \mathcal{D} and the constant $\rho > 0$ as in the statement of the theorem, we set

$$P_0 = \max\{|p| : p \in \mathcal{D}\} \quad \text{and} \quad c = (P_0 + \rho + 1)e^{C_K T}.$$

Let $z(t)$ be a solution of system (44) having $[0, \omega[$ as its maximal interval of future existence, starting with $p(0)$ satisfying $\text{dist}(p(0), \mathcal{D}) \leq \rho$. Arguing as at the beginning of Section 3.2, from Assumption 9 and (42), we have that

$$|p(t)| \leq c \quad \text{for every } t \in [0, T] \cap [0, \omega[. \tag{46}$$

So, we can introduce the cut-off function η as in (25) in order to change the Hamiltonian function

$$\widehat{H}(t, z) = \frac{1}{2}(|v|^2 - |u|^2) + \widehat{h}(t, z)$$

into the new Hamiltonian function

$$\tilde{H}(t, z) = \frac{1}{2}(|v|^2 - |u|^2) + \tilde{h}(t, z)$$

with $\tilde{h}(t, z) = \eta(|p|)\hat{h}(t, z)$. Notice that $|\nabla\tilde{h}(t, z)| \leq \tilde{C}$ for a certain positive constant \tilde{C} .

Our aim is now to apply [18, Corollary 2.3] to the modified system

$$J\dot{z} = \nabla\tilde{H}(t, z). \tag{47}$$

Any solution $z = (\phi, \psi, q, p, u, v)$ of (47) is defined on $[0, T]$. As in the proof of Theorem 2, we can prove that, if $\text{dist}(p(0), \mathcal{D}) \leq \rho$, then $|p(t)| \leq c$ for every $t \in [0, T]$. In particular, it is a solution of (44). Then, if we set $U(t) = \zeta(u(t))$ and $V(t) = \chi(\zeta(u(t)), v(t))$, we see that (ϕ, ψ, q, p) is a solution of (41). By the assumption of the theorem, if the solution starts with $\text{dist}(p(0), \partial\mathcal{D}) \leq \rho$ and $p(0) \notin \mathring{\mathcal{D}}$, then

$$\langle q(T) - q(0), \mathbb{A}\nu_{\mathcal{D}}(\pi_{\mathcal{D}}(p(0))) \rangle > 0. \tag{48}$$

The application of [18, Corollary 2.3] provides us $2L + M + 1$ geometrically distinct T -periodic solutions of (47) satisfying $p(0) \in \mathring{\mathcal{D}}$. Then, by the above estimates, they are solutions of (44). Finally, from Lemma 16, these are indeed solutions of the original Hamiltonian system (39), and they satisfy $\alpha \leq u(t) \leq \beta$, for every $t \in \mathbb{R}$. \square

8. Variants in higher dimensions

Here are two variants of Theorem 14. In the first one, the twist is formulated as an *avoiding rays condition*.

Theorem 17. *Let Assumptions 6, 7, 8, and 9 hold. Assume that there exist $\rho > 0$ and a convex body \mathcal{D} of \mathbb{R}^M , having a smooth boundary, with the following property: For any two continuous functions $U, V : [0, T] \rightarrow \mathbb{R}^D$ satisfying (40), the solutions of system (41) with $p(0) \notin \mathring{\mathcal{D}}$ and $\text{dist}(p(0), \partial\mathcal{D}) \leq \rho$ are such that*

$$q(T) - q(0) \notin \{\lambda\nu_{\mathcal{D}}(\pi_{\mathcal{D}}(p(0))) : \lambda \geq 0\}. \tag{49}$$

Then the same conclusion of Theorem 14 holds.

Proof. The argument follows the lines of the proof of Theorem 14, with the only difference of applying [18, Corollary 2.1] instead of [18, Corollary 2.3]. \square

Notice that the twist condition (49) may as well be replaced by

$$q(T) - q(0) \notin \{\lambda\nu_{\mathcal{D}}(\pi_{\mathcal{D}}(p(0))) : \lambda \leq 0\}$$

and the same conclusion of Theorem 14 still holds.

We now consider the case when \mathcal{D} is a M -cell, namely

$$\mathcal{D} = [a_1, b_1] \times \cdots \times [a_M, b_M].$$

Theorem 18. *Let Assumptions 6, 7, 8, and 9 hold. Assume that there exist $\rho > 0$ and a M -tuple $\sigma = (\sigma_1, \dots, \sigma_M) \in \{-1, 1\}^M$, with the following property: For any two continuous functions $U, V : [0, T] \rightarrow \mathbb{R}^D$ satisfying (40), the solutions of (41) with $p(0) \notin \mathring{\mathcal{D}}$ and $\text{dist}(p(0), \partial\mathcal{D}) \leq \rho$ are such that, for every $i \in \{1, \dots, M\}$,*

$$\begin{aligned} \sigma_i(q_i(T) - q_i(0)) < 0 & \quad \text{if } p_i(0) \in [a_i - \rho, a_i], \\ \sigma_i(q_i(T) - q_i(0)) > 0 & \quad \text{if } p_i(0) \in [b_i, b_i + \rho]. \end{aligned}$$

Then the same conclusion of Theorem 14 holds.

Proof. Again the proof is similar to the one of Theorem 14, just applying [18, Corollary 2.4] instead of [18, Corollary 2.3]. \square

Remark 19. As noticed in Remark 9, instead of the fixed intervals $[a_i, b_i]$ defining the M -cell \mathcal{D} , we could have varying intervals $[a_i(q_i), b_i(q_i)]$, where $a_i, b_i: \mathbb{R} \rightarrow \mathbb{R}$ are continuous 2π -periodic functions.

We now generalize Theorem 10; we thus drop the periodicity in the q -variables, still maintaining it in the ϕ and ψ variables, as stated below.

Assumption 10 (Periodicity). The function $H(t, z)$ is periodic in each of the variables included in ϕ, ψ .

We also need to suitably modify Assumption 4, in the following way

Assumption 11 (Energy growth). For every $K > 0$ there is a constant $C_K > 0$ such that, for every $i \in \{1, \dots, M\}$,

$$|q_i \partial_{p_i} H(t, q, p, u, v) - p_i \partial_{q_i} H(t, q, p, u, v)| \leq C_K (q_i^2 + p_i^2 + 1)$$

if $u \in [\alpha - \delta, \beta + \delta]$ and $|v| \leq K$.

Here is the generalization of Theorem 10.

Theorem 20. Let Assumptions 7, 8, 10 and 11 hold. Let k_1, \dots, k_M be integers and assume that there exist $\rho > 0, \tilde{\rho} > 0$ and, for each $i \in \{1, \dots, M\}$ there exist two planar Jordan curves Γ_1^i, Γ_2^i , strictly star-shaped with respect to the origin, with

$$0 \in \mathcal{D}(\Gamma_1^i) \subseteq \overline{\mathcal{D}(\Gamma_1^i)} \subseteq \mathcal{D}(\Gamma_2^i),$$

such that, for any two continuous functions $U, V: [0, T] \rightarrow \mathbb{R}$ satisfying (40), the solutions of system (41) with

$$\text{dist}((q_i(0), p_i(0)), \overline{\mathcal{D}(\Gamma_2^i)} \setminus \mathcal{D}(\Gamma_1^i)) \leq \rho \quad \text{for every } i \in \{1, \dots, M\}$$

which are defined on $[0, T]$, satisfy

$$q_i(t)^2 + p_i(t)^2 \geq \tilde{\rho} \quad \text{for every } t \in [0, T]$$

and, when $(q_i(0), p_i(0)) \notin \overline{\mathcal{D}(\Gamma_1^i)} \setminus \mathcal{D}(\Gamma_2^i)$,

$$\begin{aligned} \text{Rot}((q_i, p_i); [0, T]) &< k_i \quad \text{if } \text{dist}((q_i(0), p_i(0)), \Gamma_1^i) \leq \rho, \\ \text{Rot}((q_i, p_i); [0, T]) &> k_i \quad \text{if } \text{dist}((q_i(0), p_i(0)), \Gamma_2^i) \leq \rho. \end{aligned} \tag{50}$$

Then system (39) has at least $2L + M + 1$ distinct T -periodic solutions

$$z^{(1)}(t), \dots, z^{(2L+M+1)}(t),$$

satisfying (43), with

$$(q_i^{(n)}(0), p_i^{(n)}(0)) \in \mathcal{D}(\Gamma_2^i) \setminus \overline{\mathcal{D}(\Gamma_1^i)}$$

and

$$\text{Rot}((q_i^{(n)}, p_i^{(n)}); [0, T]) = k_i,$$

for every $i = 1, \dots, M$ and $n = 1, \dots, 2L + M + 1$. The same is true if, for some $i \in \{1, \dots, M\}$, assumption (50) is replaced by

$$\begin{aligned} \text{Rot}((q_i, p_i); [0, T]) &> k_i \quad \text{if } \text{dist}((q_i(0), p_i(0)), \Gamma_1^i) \leq \rho, \\ \text{Rot}((q_i, p_i); [0, T]) &< k_i \quad \text{if } \text{dist}((q_i(0), p_i(0)), \Gamma_2^i) \leq \rho. \end{aligned}$$

Proof. It is perfectly analogous to the one of Theorem 10, the only difference being the use of Theorem 18 (and Remark 19) instead of Theorem 2. \square

Let us now add a further equation, and consider the more general system

$$\begin{cases} \dot{\phi} = \partial_\psi H(t, z), & \dot{\psi} = -\partial_\phi H(t, z), \\ \dot{q} = \partial_p H(t, z), & \dot{p} = -\partial_q H(t, z), \\ \dot{u} = \partial_v H(t, z), & \dot{v} = -\partial_u H(t, z), \\ J\dot{w} = \partial_w H(t, z). \end{cases} \tag{51}$$

Here, and in the following, the symbol J is always used as the standard symplectic matrix, in different dimensions. Moreover, now $z = (\phi, \psi, q, p, u, v, w)$. We will generalize Theorem 14. Assumptions 6, 7 and 8 will remain the same, while Assumption 9 needs to be modified as follows.

Assumption 12 (Linear growth). Let

$$H(t, z) = \frac{1}{2} \langle \mathbb{B}(t)w, w \rangle + \mathcal{H}(t, z),$$

where the symmetric matrix $\mathbb{B}(t)$ is continuous, T -periodic, and such that

$$w(t) \equiv 0 \quad \text{is the only } T\text{-periodic solution of } J\dot{w} = \mathbb{B}(t)w.$$

The function \mathcal{H} is such that for every $K > 0$ there is a constant $C_K > 0$ with the following property: If $\alpha - \bar{\delta} \leq u \leq \beta + \bar{\delta}$ and $|v| \leq K$, then

$$|\partial_q \mathcal{H}(t, z)| + |\partial_w \mathcal{H}(t, z)| \leq C_K(|p| + 1).$$

Under the above assumption, given any two continuous functions $U, V: [0, T] \rightarrow \mathbb{R}^D$, with

$$\alpha - \bar{\delta} \leq U(t) \leq \beta + \bar{\delta} \quad \text{for every } t \in [0, T], \tag{52}$$

the solutions of the system

$$\begin{cases} \dot{\phi} = \partial_\psi H(t, \phi, \psi, q, p, U(t), V(t), w), \\ \dot{\psi} = -\partial_\phi H(t, \phi, \psi, q, p, U(t), V(t), w), \\ \dot{q} = \partial_p H(t, \phi, \psi, q, p, U(t), V(t), w), \\ \dot{p} = -\partial_q H(t, \phi, \psi, q, p, U(t), V(t), w), \\ J\dot{w} = \partial_w H(t, \phi, \psi, q, p, U(t), V(t), w), \end{cases} \tag{53}$$

are defined on $[0, T]$, cf. Remark 1.

Theorem 21. *Let Assumptions 6, 7, 8 and 12 hold. Assume that there exist $\rho > 0$, a symmetric regular $M \times M$ matrix \mathbb{A} and a strongly convex body \mathcal{D} of \mathbb{R}^M , having a smooth boundary, with the following property: For any two continuous functions $U, V: [0, T] \rightarrow \mathbb{R}^D$ satisfying (52), the solutions of (53) with $p(0) \notin \overset{\circ}{\mathcal{D}}$ and $\text{dist}(p(0), \partial\mathcal{D}) \leq \rho$ are such that*

$$\langle q(T) - q(0), \mathbb{A}\nu_{\mathcal{D}}(\pi_{\mathcal{D}}(p(0))) \rangle > 0. \tag{54}$$

Then system (51) has at least $2L + M + 1$ geometrically distinct T -periodic solutions, such that $p(0) \in \overset{\circ}{\mathcal{D}}$ and $\alpha \leq u(t) \leq \beta$ for every $t \in \mathbb{R}$.

Proof. Following the lines of the proof of Theorem 14, we can introduce a modified system ruled by a Hamiltonian function of the type

$$\widehat{H}(t, z) = \frac{1}{2} \langle \mathbb{B}(t)w, w \rangle + \frac{1}{2} (|v|^2 - |u|^2) + \widehat{h}(t, z).$$

Moreover, \widehat{H} can be introduced so to guarantee that every T -periodic solution of $J\dot{z} = \nabla \widehat{H}(t, z)$ satisfies an a priori bound as in Lemma 16, in particular $\alpha \leq u \leq \beta$. Then, from Assumption 12, we can introduce $K > 0$ as in (45) such that, if $\alpha - \bar{\delta} \leq u \leq \beta + \bar{\delta}$ and $|v| \leq K$, then

$$|\partial_q \widehat{h}(t, z)| + |\partial_w \widehat{h}(t, z)| \leq \widehat{C}_K (|p| + 1).$$

Hence, we can find a constant $c > 0$ such that any solution of $J\dot{z} = \nabla \widehat{H}(t, z)$ starting with $z(0)$ satisfying $\text{dist}(p(0), \mathcal{D}) \leq \rho$ is such that (46) holds.

Again, we can introduce the cut-off function η , as in (25), and the Hamiltonian function

$$\widetilde{H}(t, z) = \frac{1}{2} \langle \mathbb{B}(t)w, w \rangle + \frac{1}{2} (|v|^2 - |u|^2) + \widetilde{h}(t, z),$$

where $\widetilde{h}(t, z) = \eta(|p|) \widehat{h}(t, z)$. Using Assumption 12 we get

$$|\nabla \widetilde{h}(t, z)| \leq \widetilde{C},$$

for a certain positive constant \widetilde{C} . We can rewrite the previous Hamiltonian as

$$\widetilde{H}(t, z) = \frac{1}{2} \left\langle \mathbb{M}(t)(w, (u, v)), (w, (u, v)) \right\rangle + \widetilde{h}(t, z),$$

where

$$\mathbb{M}(t)(w, (u, v)) = (\mathbb{B}(t)w, (-u, v)).$$

Our aim is to apply [18, Corollary 2.3] to the modified system $J\dot{z} = \nabla \widetilde{H}(t, z)$. We easily verify the nonresonance condition

$$\omega(t) \equiv 0 \quad \text{is the only } T\text{-periodic solution of } J\dot{\omega} = \mathbb{M}(t)\omega,$$

where $\omega = (w, (u, v))$.

We finally verify, with the usual argument, that any solution z of the previous system with $0 < \text{dist}(p(0), \mathcal{D}) \leq \rho$ is such that

$$\langle q(T) - q(0), \mathbb{A}p(0) \rangle > 0. \tag{55}$$

The application of [18, Corollary 2.3] provides us $2L + M + 1$ geometrically distinct T -periodic solutions of $J\dot{z} = \nabla \tilde{H}(t, z)$ satisfying $p(0) \in \mathring{D}$. Then, as in the previous proofs, we end showing that they are solutions of (51), too, and they satisfy $\alpha \leq u(t) \leq \beta$, for every $t \in \mathbb{R}$. \square

Clearly enough, the twist condition (54) could be replaced by an avoiding rays condition, as in Theorem 17, or by a sign condition on the edges of an M -cell, like in Theorem 18. Also, we could provide a statement on an annulus, similarly as in Theorem 20. Or even some combination of these could be considered. We avoid the details, for brevity.

9. Examples in higher dimensions

In this section we just briefly mention how the examples given in Section 5 generalize to higher dimensions applying the results of Sections 6 and 8. To this aim, consider a system of the form

$$\begin{cases} \dot{\phi} = \partial_{\psi} P(t, \phi, \psi, q, u), & \dot{\psi} = -\partial_{\phi} P(t, \phi, \psi, q, u), \\ -\ddot{q}_i = g_i(t, q_i) - e_i(t) + \partial_{q_i} P(t, \phi, \psi, q, u), & i = 1, \dots, M, \\ -\ddot{u}_j = -h_j(u_j) + \partial_{u_j} P(t, \phi, \psi, q, u), & j = 1, \dots, D, \end{cases} \tag{56}$$

where all functions are continuous and T -periodic in t . For simplicity, we assume that $P(t, \phi, \psi, q, u)$ is 2π -periodic in each of the variables included in ϕ, ψ and q , so, due to this, we fix an arbitrary cube in \mathbb{R}^{2L+M} of length $[0, 2\pi]$, denoting it by Θ . Moreover, let $P(t, \phi, \psi, q, u)$ be continuously differentiable in (ϕ, ψ, q, u) and assume that, for every $i \in \{1, \dots, M\}$,

$$\int_0^T e_i(t) dt = 0 \quad \text{and} \quad \int_0^{2\pi} g_i(t, s) ds = 0.$$

Then system (56) is of the type (39), with

$$\begin{aligned} H(t, z) = & \frac{1}{2}|p + E(t)|^2 + \frac{1}{2}|v|^2 + \sum_{i=1}^M \int_0^{q_i} g(t, s) ds \\ & - \sum_{j=1}^D \int_0^{u_j} h_j(\sigma) d\sigma + P(t, \phi, \psi, q, u), \end{aligned}$$

where $z = (\phi, \psi, q, p, u, v)$ and $E = (E_1, \dots, E_M): [0, T] \rightarrow \mathbb{R}^M$ is a primitive of the field $e(t) = (e_1(t), \dots, e_M(t))$.

Let us state the analogues of Corollaries 11 and 12.

Corollary 22. *In the above setting, assume moreover that there exist two vectors $\alpha, \beta \in \mathbb{R}^D$, with $\alpha \leq \beta$, such that, for all $j \in \{1, \dots, D\}$,*

$$\begin{aligned} h_j(\alpha_j) &< \partial_{u_j} P(t, \phi, \psi, q, u) && \text{if } u_j = \alpha_j, \\ h_j(\beta_j) &> \partial_{u_j} P(t, \phi, \psi, q, u) && \text{if } u_j = \beta_j, \end{aligned} \tag{57}$$

for all $(t, \phi, \psi, q) \in [0, T] \times \Theta$ and all $u \in \mathbb{R}^D$ such that $\alpha \leq u \leq \beta$. Then system (33) has at least $2L + M + 1$ geometrically distinct T -periodic solutions, with $\alpha \leq u \leq \beta$.

Proof. All the assumptions of Theorem 18 are easily verified, whence the conclusion. \square

As an immediate consequence, we have the following.

Corollary 23. *The system*

$$\begin{cases} \dot{\phi} = \partial_{\psi}P(t, \phi, \psi, q, u), & \dot{\psi} = -\partial_{\phi}P(t, \phi, \psi, q, u), \\ \ddot{q}_i + a_i \sin q_i = e_i(t) - \partial_{q_i}P(t, \phi, \psi, q, u), & i = 1, \dots, M, \\ \ddot{u}_j + b_j \sin u_j = -\partial_{u_j}P(t, \phi, \psi, q, u), & j = 1, \dots, D, \end{cases} \tag{58}$$

where $P(t, \phi, \psi, q, u)$ and $e(t)$ are as above, has at least $2L+M+1$ geometrically distinct T -periodic solutions, for any $a_i > 0$, if $\|\partial_{u_j}P\|_{\infty} < b_j$ for every $j \in \{1, \dots, D\}$.

All the other examples presented in Section 5 can be displayed in this more general setting, also with a mixing of assumptions on each component. In particular, we thus generalize, e.g., the results in [7,8,10,13,14,25,28,32,35,36,40,41]. Also equations with singularities could be considered, as in [22,26]. We will avoid entering further into details, for brevity.

A further example of application is analyzed in the next section.

10. Periodic perturbations of completely integrable systems

In [3,18], perturbations of completely integrable Hamiltonian systems were studied (see also [19], and the references therein). We will now add an extra term to the Hamiltonian function, involving lower and upper solutions.

We consider the system

$$\begin{cases} \dot{\varphi} = \nabla\mathcal{K}(I) + \varepsilon\partial_I P(t, \varphi, I, u), & \dot{I} = -\varepsilon\partial_{\varphi}P(t, \varphi, I, u), \\ -\ddot{u} = \partial_u G(t, u) + \varepsilon\partial_u P(t, \varphi, I, u). \end{cases} \tag{59}$$

Here, we have $(\varphi, I) \in \mathbb{R}^{2M}$ and $u \in \mathbb{R}^D$. We assume that $\mathcal{K}: \mathbb{R}^M \rightarrow \mathbb{R}$ is continuously differentiable and the same for $G: \mathbb{R} \times \mathbb{R}^D \rightarrow \mathbb{R}$ with respect to the second variable. The perturbation function $P: \mathbb{R} \times \mathbb{R}^{2M+D} \rightarrow \mathbb{R}$ is continuous, T -periodic in t , and continuously differentiable with respect to all the other variables, with a bounded gradient. Moreover, the function P is τ_i -periodic in each φ_i , i.e., for every $i \in \{1, \dots, M\}$,

$$P(\dots, \varphi_i + \tau_i, \dots) = P(\dots, \varphi_i, \dots).$$

We also assume that there exist some integers m_1, \dots, m_M for which

$$T\nabla\mathcal{K}(I^0) = (m_1\tau_1, \dots, m_M\tau_M).$$

The expert reader will recognize that we are dealing with a *completely resonant torus*. Here is our result.

Theorem 24. *In the above setting, assume that there exist $I^0 \in \mathbb{R}^M$, a symmetric invertible $M \times M$ matrix \mathbb{A} and $\bar{\rho} > 0$ such that*

$$0 < |I - I^0| \leq \bar{\rho} \implies \langle \nabla\mathcal{K}(I) - \nabla\mathcal{K}(I^0), \mathbb{A}(I - I^0) \rangle > 0. \tag{60}$$

Moreover, let there exist constants $\delta > 0$, $\varsigma > 0$, $\alpha = (\alpha_1, \dots, \alpha_D)$ and $\beta = (\beta_1, \dots, \beta_D)$ with $\alpha \leq \beta$, having the following property: When $\alpha - \delta \leq u \leq \beta + \delta$, for every $j \in \{1, \dots, D\}$ one has

$$\begin{aligned} \partial_{u_j} G(t, u) &\geq \varsigma && \text{if } u_j \in [\alpha_j - \delta, \alpha_j], \\ \partial_{u_j} G(t, u) &\leq -\varsigma && \text{if } u_j \in [\beta_j, \beta_j + \delta]. \end{aligned}$$

Then for every $\sigma > 0$ there exists $\bar{\varepsilon} > 0$ such that, if $|\varepsilon| \leq \bar{\varepsilon}$, there are at least $M + 1$ geometrically distinct solutions of system (59) satisfying

$$\begin{aligned} \varphi(t + T) &= \varphi(t) + T\nabla\mathcal{K}(I^0), & I(t + T) &= I(t), \\ u(t + T) &= u(t), & v(t + T) &= v(t) \end{aligned}$$

and such that

$$|\varphi(t) - \varphi(0) - t\nabla\mathcal{K}(I^0)| + |I(t) - I^0| < \sigma$$

and

$$\alpha \leq u(t) \leq \beta,$$

for every $t \in \mathbb{R}$.

Proof. Since we are looking for solutions with $I(t)$ in the open ball $B(I^0, \sigma)$, we can suitably modify the function \mathcal{K} outside this set and assume that it has a bounded gradient. We perform the change of variables

$$q(t) = \varphi(t) - t\nabla\mathcal{K}(I^0), \quad p(t) = I(t) - I^0,$$

thus obtaining the new Hamiltonian function

$$\mathcal{H}(t, q, p, u, v) = \mathcal{K}(p) + \frac{1}{2}|v|^2 + G(t, u) + \varepsilon Q(t, q, p, u),$$

where the functions

$$\mathcal{K}(p) = \mathcal{K}(I) - \langle \nabla\mathcal{K}(I^0), I - I^0 \rangle, \quad Q(t, q, p, u) = P(t, \varphi, I, u)$$

are implicitly defined. We notice that Q is periodic in q_1, \dots, q_M and both Q and \mathcal{K} have bounded gradient: so, there are $C_{\mathcal{K}}, C_Q > 0$ such that

$$|\nabla\mathcal{K}(p)| \leq C_{\mathcal{K}}, \quad |\nabla Q(t, q, p, u)| \leq C_Q,$$

for every $(t, q, p, u) \in [0, T] \times \mathbb{R}^{2M+D}$. Our goal is now to prove that for every $\bar{\sigma} > 0$ there exists $\bar{\varepsilon} > 0$ such that, if $|\varepsilon| \leq \bar{\varepsilon}$, there are $M + 1$ geometrically distinct T -periodic solutions $z = (q, p, u, v)$ of the system

$$J\dot{z} = \nabla\mathcal{H}(t, z), \tag{61}$$

satisfying $\max\{|q(t) - q(0)|, |p(t)|\} < \bar{\sigma}$ and $\alpha \leq u(t) \leq \beta$, for every $t \in \mathbb{R}$.

Let us fix $\bar{\sigma} \in]0, \bar{\rho}[$. We are in the setting of Theorem 21. Assumptions 6, 7, 8 and 12 can be easily checked. We need to verify the twist condition (54).

Since $\nabla\mathcal{H}(0) = 0$, we can choose $r < \bar{\sigma}/4$ such that

$$|p| \leq 4r \quad \Rightarrow \quad 2T|\nabla\mathcal{K}(p)| < \bar{\sigma} \tag{62}$$

and by (60) there is $\ell > 0$ such that

$$r \leq |p| \leq 4r \quad \Rightarrow \quad \langle \nabla \mathcal{K}(p), \mathbb{A}p \rangle > 4\ell. \tag{63}$$

Let us fix $\delta > 0$ satisfying

$$\delta < \min \left\{ r, \frac{2\ell}{C_{\mathcal{K}}\|\mathbb{A}\|} \right\}.$$

Reducing $\bar{\varepsilon}$ if necessary, every solution of (61), with $|\varepsilon| \leq \bar{\varepsilon}$, is such that $|p(t) - p(0)| < \delta$ for every $t \in [0, T]$.

Let us consider a solution z of (61) with $|\varepsilon| \leq \bar{\varepsilon}$ and initial condition satisfying $2r \leq |p(0)| \leq 3r$. Then $r \leq |p(t)| \leq 4r$ for every $t \in [0, T]$, and so, by (63),

$$\langle \nabla \mathcal{K}(p(t)), \mathbb{A}p(t) \rangle > 4\ell \quad \text{for every } t \in [0, T].$$

Reducing $\bar{\varepsilon}$ if necessary, we have that

$$\begin{aligned} \langle \partial_p H(t, z(t)), \mathbb{A}p(0) \rangle &= \\ &= \langle \nabla \mathcal{K}(p(t)), \mathbb{A}p(t) \rangle - \langle \nabla \mathcal{K}(p(t)), \mathbb{A}(p(t) - p(0)) \rangle + \varepsilon \langle \partial_p Q(t, z(t)), \mathbb{A}p(0) \rangle \\ &> 4\ell - C_{\mathcal{K}}\|\mathbb{A}\|\delta - 3\varepsilon C_Q\|\mathbb{A}\|r > \ell. \end{aligned}$$

Integrating the previous estimate in the interval $[0, T]$, we get

$$\langle q(T) - q(0), \mathbb{A}p(0) \rangle > \ell T > 0.$$

We can thus apply Theorem 21, choosing $\mathcal{D} = \bar{B}(0, 2r)$ and $\rho = r$. We have that

$$|p(t)| \leq |p(0)| + \delta < 2r + \delta < 3r < \bar{\sigma} \quad \text{for every } t \in [0, T].$$

Moreover, by (62), we deduce that $|q(t) - q(0)| < \bar{\sigma}$ for all $t \in [0, T]$.

The T -periodic solutions we have found are then translated, by the inverse change of variables, into the solutions of (59) we are looking for. \square

Remark 25. The twist condition (60) holds true if \mathcal{K} is twice continuously differentiable with $\det \mathcal{K}''(I^0) \neq 0$, by taking $\mathbb{A} = \mathcal{K}''(I^0)$.

Remark 26. A more general twist condition (see [16]) can be considered, e.g.,

$$0 \in \text{cl} \left\{ r \in]0, +\infty[: \min_{|I-I^0|=r} \langle \nabla \mathcal{K}(I) - \nabla \mathcal{K}(I^0), \mathbb{A}(I - I^0) \rangle > 0 \right\}.$$

Remark 27. The same type of result holds, with due changes, for a more general system of the type

$$\begin{cases} \dot{\phi} = \varepsilon \partial_{\psi} P(t, \phi, \psi, \varphi, I, u, w), & \dot{\psi} = -\varepsilon \partial_{\phi} P(t, \phi, \psi, \varphi, I, u, w), \\ \dot{\varphi} = \nabla \mathcal{K}(I) + \varepsilon \partial_I P(t, \phi, \psi, \varphi, I, u, w), & \dot{I} = -\varepsilon \partial_{\varphi} P(t, \phi, \psi, \varphi, I, u, w), \\ -\ddot{u} = \partial_u G(t, u) + \varepsilon \partial_u P(t, \phi, \psi, \varphi, I, u, w), \\ J\dot{w} = \mathbb{B}(t)w + \varepsilon \partial_w P(t, \phi, \psi, \varphi, I, u, w), \end{cases}$$

when P is also periodic in ϕ_1, \dots, ϕ_L and ψ_1, \dots, ψ_L , and $\mathbb{B}(t)$ is a continuous, T -periodic, symmetric matrix such that $w(t) \equiv 0$ is the only T -periodic solution of $J\dot{w} = \mathbb{B}(t)w$.

Remark 28. It would be interesting to see how Theorem 24 could be extended to infinite dimensions in the spirit of [19].

11. The general result

We consider system (1), assuming the Hamiltonian function $H: \mathbb{R} \times \mathbb{R}^{2N} \rightarrow \mathbb{R}$ to be continuous, T -periodic in the variable t , and continuously differentiable with respect to the variable z , with corresponding gradient $\nabla H(t, z)$.

For $z \in \mathbb{R}^{2N}$, we use the notation $z = (x, y)$, with $x = (x_1, \dots, x_N) \in \mathbb{R}^N$ and $y = (y_1, \dots, y_N) \in \mathbb{R}^N$. Moreover, we split each of the variable tuples x and y into five tuples, thus writing

$$x = (x^a, x^b, x^c, x^d, x^e), \quad y = (y^a, y^b, y^c, y^d, y^e),$$

where, for some nonnegative integers N^a, N^b, N^c, N^d, N^e ,

$$\begin{aligned} x^a &= (x_1^a, \dots, x_{N^a}^a) \in \mathbb{R}^{N^a}, & y^a &= (y_1^a, \dots, y_{N^a}^a) \in \mathbb{R}^{N^a}, \\ x^b &= (x_1^b, \dots, x_{N^b}^b) \in \mathbb{R}^{N^b}, & y^b &= (y_1^b, \dots, y_{N^b}^b) \in \mathbb{R}^{N^b}, \\ x^c &= (x_1^c, \dots, x_{N^c}^c) \in \mathbb{R}^{N^c}, & y^c &= (y_1^c, \dots, y_{N^c}^c) \in \mathbb{R}^{N^c}, \\ x^d &= (x_1^d, \dots, x_{N^d}^d) \in \mathbb{R}^{N^d}, & y^d &= (y_1^d, \dots, y_{N^d}^d) \in \mathbb{R}^{N^d}, \\ x^e &= (x_1^e, \dots, x_{N^e}^e) \in \mathbb{R}^{N^e}, & y^e &= (y_1^e, \dots, y_{N^e}^e) \in \mathbb{R}^{N^e}, \end{aligned}$$

and we also introduce the notation

$$z^a = (x^a, y^a), \quad z^b = (x^b, y^b), \quad z^c = (x^c, y^c), \quad z^d = (x^d, y^d), \quad z^e = (x^e, y^e).$$

Notice that one or more of these integers could be equal to zero, in which case the corresponding tuples will not be taken into account; for example, if $N^a = 0$, then x^a, y^a and z^a will disappear from the list. In the following, for simplicity we sometimes write (u, v) instead of (x^e, y^e) , and $D = N^e$.

Let us introduce our assumptions in this general setting.

Assumption 13 (*Periodicity*). The function $H(t, z)$ is periodic in each of the variables included in x^a, x^b, y^a, y^c .

The total number of variables in which our Hamiltonian function is periodic is thus $2N^a + N^b + N^c$.

Assumption 14 (*Lower and upper solutions*). There exist constants $\delta > 0$, $\alpha = (\alpha_1, \dots, \alpha_D)$ and $\beta = (\beta_1, \dots, \beta_D)$ with $\alpha \leq \beta$, having the following property. When $\alpha - \bar{\delta} \leq u \leq \beta + \bar{\delta}$, then, for every $j \in \{1, \dots, D\}$,

$$v_j \partial_{v_j} H(t, z) > 0 \text{ if } u_j \in [\alpha_j - \delta, \alpha_j] \cup [\beta_j, \beta_j + \delta] \text{ and } v_j \neq 0,$$

and

$$\begin{aligned} \partial_{u_j} H(t, z) &\geq 0 && \text{if } u_j \in [\alpha_j - \delta, \alpha_j] \text{ and } v_j = 0, \\ \partial_{u_j} H(t, z) &\leq 0 && \text{if } u_j \in [\beta_j, \beta_j + \delta] \text{ and } v_j = 0. \end{aligned}$$

Assumption 15 (*Nagumo condition*). For every $j \in \{1, \dots, D\}$ there exist $d_j > 0$ and two continuous functions $f_j, \varphi_j: [d_j, +\infty[\rightarrow]0, +\infty[$ with

$$\int_{d_j}^{+\infty} \frac{f_j(s)}{\varphi_j(s)} ds = +\infty,$$

satisfying the following property. When $\alpha - \bar{\delta} \leq u \leq \beta + \bar{\delta}$, then

$$\begin{aligned} \partial_{v_j} H(t, z) &\geq f_j(v_j) && \text{if } v_j \geq d_j, \\ \partial_{v_j} H(t, z) &\leq -f_j(-v_j) && \text{if } v_j \leq -d_j, \end{aligned}$$

and

$$|\partial_{u_j} H(t, z)| \leq \varphi_j(|v_j|) \quad \text{if } |v_j| \geq d_j.$$

Assumption 16 (*Linear growth*). There exists a symmetric $2N^d \times 2N^d$ matrix $\mathbb{B}(t)$, T -periodic and continuous in t , satisfying the nonresonance condition

$$z^d(t) \equiv 0 \quad \text{is the only } T\text{-periodic solution of } J\dot{z}^d(t) = \mathbb{B}(t)z^d(t)$$

and such that, writing

$$H(t, z) = \frac{1}{2} \langle \mathbb{B}(t)z^d, z^d \rangle + \mathcal{H}(t, z),$$

for every $K > 0$ there is a $C_K > 0$ with the following property: If $\alpha - \bar{\delta} \leq u \leq \beta + \bar{\delta}$ and $|v| \leq K$, then

$$|\partial_{x^b} \mathcal{H}(t, z)| + |\partial_{y^c} \mathcal{H}(t, z)| + |\partial_{x^a} \mathcal{H}(t, z)| + |\partial_{y^d} \mathcal{H}(t, z)| \leq C_K(|y^b| + |x^c| + 1).$$

The above assumption guarantees that, for any two continuous functions $U, V: [0, T] \rightarrow \mathbb{R}^D$ with

$$\alpha - \bar{\delta} \leq U(t) \leq \beta + \bar{\delta} \quad \text{for every } t \in [0, T], \tag{64}$$

setting $W(t) = (U(t), V(t))$ and

$$\bar{H}(t, z) = H(t, z^a, z^b, z^c, z^d, W(t)),$$

the solutions of

$$J\dot{z} = \nabla \bar{H}(t, z) \tag{65}$$

are defined on $[0, T]$, cf. Remark 1.

Let us also introduce a C^1 -function $h: \mathbb{R}^{N^b+N^c} \rightarrow \mathbb{R}$ and a regular symmetric $(N^b + N^c) \times (N^b + N^c)$ matrix S such that, for some positive constants C_1, C_2 ,

$$|h(v) - \frac{1}{2} \langle Sv, v \rangle| \leq C_1 \quad \text{and} \quad |\nabla h(v) - Sv| \leq C_2 \quad \text{for every } v \in \mathbb{R}^{N^b+N^c},$$

and let

$$\mathcal{D} = \{v \in \mathbb{R}^{N^b+N^c} : \nabla h(v) = 0\}.$$

We assume that such a set is compact. Our main result is the following.

Theorem 29. *Let Assumptions 13, 14, 15, and 16 hold. Assume in addition that there exists $\rho > 0$ such that, for any two continuous functions $U, V: [0, T] \rightarrow \mathbb{R}^D$ satisfying (64), the solutions of (65) with $(y^b(0), x^c(0)) \notin \bar{\mathcal{D}}$ and $\text{dist}((y^b(0), x^c(0)), \mathcal{D}) \leq \rho$ are such that*

$$(x^b(T) - x^b(0), y^c(T) - y^c(0)) \notin \{\lambda J\nabla h((y^b(0), x^c(0))) : \lambda \geq 0\}.$$

Then system (1) has at least $2N^a + N^b + N^c + 1$ geometrically distinct T -periodic solutions $z(t)$ such that

$$(y^b(0), x^c(0)) \in \mathring{D}$$

and

$$\alpha \leq u(t) \leq \beta \quad \text{for every } t \in \mathbb{R}.$$

Proof. With the same procedure adopted in the proof of Theorem 14 in Section 7, dealing with the $z^e = (u, v)$ coordinates, we can introduce a modified problem ruled by a Hamiltonian of the type

$$\widehat{H}(t, z) = \frac{1}{2} \langle \mathbb{B}(t)z^d, z^d \rangle + \frac{1}{2} (|v|^2 - |u|^2) + \widehat{h}(t, z).$$

Moreover, \widehat{H} can be introduced so to guarantee that every T -periodic solution of $J\dot{z} = \nabla \widehat{H}(t, z)$ satisfies an a priori bound as in Lemma 16, in particular $\alpha \leq u \leq \beta$. Then, from Assumption 16, we can introduce $K > 0$ as in (45) so to obtain

$$|\partial_{x^b} \widehat{h}(t, z)| + |\partial_{y^c} \widehat{h}(t, z)| \leq C_K (|y^b| + |x^c| + 1),$$

when $\alpha - \bar{\delta} \leq u \leq \beta + \bar{\delta}$ and $|v| \leq K$. Hence, as in (46), we can find a constant $c > 0$ such that, if $z(t)$ is a solution of $J\dot{z} = \nabla \widehat{H}(t, z)$ starting with $\text{dist}((y^b(0), x^c(0), \mathcal{D}) \leq \rho$ with maximal interval of future existence $[0, \omega[$, then

$$\max\{|y^b(t)|, |x^c(t)|\} \leq c \quad \text{for every } t \in [0, T] \cap [0, \omega[.$$

Again, we can introduce the cut-off function η as in (25) and the Hamiltonian

$$\widetilde{H}(t, z) = \frac{1}{2} \langle \mathbb{B}(t)z^d, z^d \rangle + \frac{1}{2} (|v|^2 - |u|^2) + \widetilde{h}(t, z),$$

where $\widetilde{h}(t, z) = \eta(|y^b|)\eta(|x^c|)\widehat{h}(t, z)$. Using Assumption 16 we get

$$|\nabla \widetilde{h}(t, z)| \leq \widetilde{C}$$

for a certain positive constant \widetilde{C} .

We can rewrite the previous Hamiltonian as

$$\widetilde{H}(t, z) = \frac{1}{2} \langle \mathbb{M}(t)(z^d, z^e), (z^d, z^e) \rangle + \widetilde{h}(t, z),$$

where

$$\mathbb{M}(t)(z^d, z^e) = \mathbb{M}(t)(z^d, (u, v)) = (\mathbb{B}(t)z^d, (-u, v)).$$

In particular, we can verify the nonresonance condition

$$z^{d,e}(t) \equiv 0 \quad \text{is the only } T\text{-periodic solution of } J\dot{z}^{d,e}(t) = \mathbb{M}(t)z^{d,e}(t),$$

where $z^{d,e} = (z^d, z^e)$. Our aim is to apply [18, Theorem 1.1] to the modified system $J\dot{z} = \nabla \widetilde{H}(t, z)$. Its solutions are globally defined on $[0, T]$.

Let $z = (z^a, z^b, z^c, z^d, z^e)$ be a solution of $J\dot{z} = \nabla\tilde{H}(t, z)$. As in the above proofs we can show that if $\text{dist}((y^b(0), x^c(0)), \mathcal{D}) \leq \rho$, then $\max\{|y^b(t)|, |x^c(t)|\} \leq c$ for every $t \in [0, T]$. Hence, z solves $J\dot{z} = \nabla\hat{H}(t, z)$ and, setting $W(t) = (\zeta(u(t)), \chi(\zeta(u(t)), v(t)))$, we see that (z^a, z^b, z^c, z^d) is a solution of (65).

If $0 < \text{dist}((y^b(0), x^c(0)), \mathcal{D}) \leq \rho$, then

$$(x^b(T) - x^b(0), y^c(T) - y^c(0)) \notin \{\lambda J\nabla h((y^b(0), x^c(0))) : \lambda \geq 0\}, \tag{66}$$

by the hypothesis of the theorem.

The application of [18, Theorem 1.1] provides us $2N^a + N^b + N^c + 1$ geometrically distinct T -periodic solutions of $J\dot{z} = \nabla\hat{H}(t, z)$ satisfying $(y^b(0), x^c(0)) \in \overset{\circ}{\mathcal{D}}$. Then, following the argument of the previous proofs, it can be seen that they are solutions of $J\dot{z} = \nabla H(t, z)$ as well, and $\alpha \leq u \leq \beta$. \square

Remark 30. Theorem 29 generalizes all three Theorems 14, 17 and 18. For example, let the assumptions of Theorem 14 hold. We consider a smooth function $\sigma: \mathbb{R} \rightarrow \mathbb{R}$ such that

$$\sigma(s) = \begin{cases} 0 & \text{if } s \leq 0, \\ 1 & \text{if } s \geq 1 \end{cases} \quad \text{and} \quad \sigma'(s) > 0 \text{ if } s \in]0, 1[$$

and, adapting the notations, we define the function $h: \mathbb{R}^{N^b+N^c} \rightarrow \mathbb{R}$ by

$$h(p) = -\xi(p)\langle \mathbb{A}(p - \pi_{\mathcal{D}}(p)), p - \pi_{\mathcal{D}}(p) \rangle,$$

where

$$\xi(p) = \begin{cases} 0 & \text{if } p \in \mathcal{D}, \\ \frac{1}{2}\sigma(|p - \pi_{\mathcal{D}}(p)|) & \text{if } p \notin \mathcal{D}. \end{cases}$$

Following the proof of [18, Corollary 2.3] one can verify that h satisfies the assumptions in Theorem 29.

Having extended the main theorem in [18] with Theorem 29, we thus have generalized, e.g., the results of [9,11,17,30,33,37,38,40].

Declaration of competing interest

On behalf of all authors, the corresponding author states that there is no conflict of interest.

Data availability

Not applicable.

Acknowledgment

The authors thank the referees for their valuable comments. The second author has been partially supported by the Spanish MINECO (FPI grant PRE2018-083803) and ERDF project MTM2017-82348-C2-1-P. He also wants to thank the Department of Mathematics and Geosciences of the University of Trieste for the kind hospitality.

References

- [1] V.I. Arnold, Sur une propriété topologique des applications globalement canoniques de la mécanique classique, C. R. Acad. Sci. Paris 261 (1965) 3719–3722.
- [2] V.I. Arnold, The stability problem and ergodic properties for classical dynamical systems, in: Proc. Internat. Congr. Math., Moscow, 1966, Mir, Moscow, 1968, pp. 387–392, in Russian, English translation in: AMS Transl. Ser. 2, 70, pp. 5–11, 1968.
- [3] D. Bernstein, A. Katok, Birkhoff periodic orbits for small perturbations of completely integrable Hamiltonian systems with convex Hamiltonians, Invent. Math. 88 (1987) 222–241.
- [4] G.D. Birkhoff, Proof of Poincaré’s geometric theorem, Trans. Am. Math. Soc. 14 (1913) 14–22.
- [5] G.D. Birkhoff, An extension of Poincaré’s last geometric theorem, Acta Math. 47 (1925) 297–311.
- [6] A. Boscaggin, A. Fonda, M. Garrione, An infinite-dimensional version of the Poincaré–Birkhoff theorem on the Hilbert cube, Ann. Sc. Norm. Super. Pisa, Cl. Sci. (5) 20 (2020) 751–770.
- [7] A. Calamai, A. Sfecci, Multiplicity of periodic solutions for systems of weakly coupled parametrized second order differential equations, NoDEA Nonlinear Differ. Equ. Appl. 24 (1) (2017) 4.
- [8] A. Castro, A.C. Lazer, On periodic solutions of weakly coupled systems of differential equations, Boll. Unione Mat. Ital. 18B (1981) 733–742.
- [9] K.C. Chang, On the periodic nonlinearity and the multiplicity of solutions, Nonlinear Anal. 13 (1989) 527–537.
- [10] F. Chen, D. Qian, An extension of the Poincaré–Birkhoff theorem for Hamiltonian systems coupling resonant linear components with twisting components, J. Differ. Equ. 321 (2022) 415–448.
- [11] C.C. Conley, E.J. Zehnder, The Birkhoff – Lewis fixed point theorem and a conjecture of V.I. Arnold, Invent. Math. 73 (1983) 33–49.
- [12] C. De Coster, P. Habets, Two-Point Boundary Value Problems, Lower and Upper Solutions, Elsevier, Amsterdam, 2006.
- [13] T.R. Ding, F. Zanolin, Periodic solutions of Duffing’s equations with superquadratic potential, J. Differ. Equ. 97 (1992) 328–378.
- [14] T.R. Ding, F. Zanolin, Subharmonic solutions of second order nonlinear equations: a time-map approach, Nonlinear Anal. 20 (1993) 509–532.
- [15] P.L. Felmer, Periodic solutions of spatially periodic Hamiltonian systems, J. Differ. Equ. 98 (1992) 143–168.
- [16] A. Fonda, M. Garrione, P. Gidoni, Periodic perturbations of Hamiltonian systems, Adv. Nonlinear Anal. 5 (2016) 367–382.
- [17] A. Fonda, P. Gidoni, An avoiding cones condition for the Poincaré–Birkhoff Theorem, J. Differ. Equ. 262 (2017) 1064–1084.
- [18] A. Fonda, P. Gidoni, Coupling linearity and twist: an extension of the Poincaré–Birkhoff Theorem for Hamiltonian systems, NoDEA Nonlinear Differ. Equ. Appl. 27 (2020) 55.
- [19] A. Fonda, G. Klun, A. Sfecci, Periodic solutions of nearly integrable Hamiltonian systems bifurcating from infinite-dimensional tori, Nonlinear Anal. 201 (2020) 111720.
- [20] A. Fonda, G. Klun, A. Sfecci, Well-ordered and non-well-ordered lower and upper solutions for periodic planar systems, Adv. Nonlinear Stud. 21 (2021) 397–419.
- [21] A. Fonda, G. Klun, A. Sfecci, Periodic solutions of second-order differential equations in Hilbert spaces, Mediterr. J. Math. 18 (5) (2021) 223.
- [22] A. Fonda, R. Manásevich, F. Zanolin, Subharmonic solutions for some second-order differential equations with singularities, SIAM J. Math. Anal. 24 (1993) 1294–1311.
- [23] A. Fonda, J. Mawhin, Multiple periodic solutions of conservative systems with periodic nonlinearity, in: Differential Equations and Applications, Columbus, OH, 1988, Ohio Univ. Press, Athens, 1989, pp. 298–304.
- [24] A. Fonda, M. Sabatini, F. Zanolin, Periodic solutions of perturbed Hamiltonian systems in the plane by the use of the Poincaré – Birkhoff Theorem, Topol. Methods Nonlinear Anal. 40 (2012) 29–52.
- [25] A. Fonda, A. Sfecci, Periodic solutions of weakly coupled superlinear systems, J. Differ. Equ. 260 (2016) 2150–2162.
- [26] A. Fonda, A. Sfecci, Multiple periodic solutions of Hamiltonian systems confined in a box, Discrete Contin. Dyn. Syst. 37 (2017) 1425–1436.
- [27] A. Fonda, R. Toader, Periodic solutions of pendulum-like Hamiltonian systems in the plane, Adv. Nonlinear Stud. 12 (2012) 395–408.
- [28] A. Fonda, R. Toader, Subharmonic solutions of Hamiltonian systems displaying some kind of sublinear growth, Adv. Nonlinear Anal. 8 (2019) 583–602.
- [29] A. Fonda, R. Toader, A dynamical approach to lower and upper solutions for planar systems, Discrete Contin. Dyn. Syst. 41 (2021) 3683–3708.
- [30] A. Fonda, A.J. Ureña, A higher dimensional Poincaré–Birkhoff theorem for Hamiltonian flows, Ann. Inst. Henri Poincaré, Anal. Non Linéaire 34 (2017) 679–698.
- [31] A. Fonda, A.J. Ureña, A Poincaré–Birkhoff theorem for Hamiltonian flows on nonconvex domains, J. Math. Pures Appl. 129 (2019) 131–152.
- [32] A. Fonda, F. Zanolin, Periodic oscillations of forced pendulums with very small length, Proc. R. Soc. Edinb. A 127 (1997) 67–76.
- [33] G. Fournier, D. Lupo, M. Ramos, M. Willem, Limit Relative Category and Critical Point Theory, Dynamics Reported: Expositions in Dynamical Systems, vol. 3, Springer, Berlin, 1994, pp. 1–24.
- [34] P. Gidoni, Existence of a periodic solution for superlinear second order ODEs, J. Differ. Equ. 345 (2023) 401–417.
- [35] P. Hartman, On boundary value problems for superlinear second order differential equations, J. Differ. Equ. 26 (1977) 37–53.
- [36] H. Jacobowitz, Periodic solutions of $x'' + f(x, t) = 0$ via the Poincaré–Birkhoff theorem, J. Differ. Equ. 20 (1976) 37–52.
- [37] F.W. Josellis, Lyusternik–Schnirelman theory for flows and periodic orbits for Hamiltonian systems on $\mathbb{T}^n \times \mathbb{R}^n$, Proc. Lond. Math. Soc. 68 (1994) 641–672.
- [38] J.Q. Liu, A generalized saddle point theorem, J. Differ. Equ. 82 (1989) 372–385.

- [39] A. Margheri, C. Rebelo, F. Zanolin, Maslov index, Poincaré–Birkhoff theorem and periodic solutions of asymptotically linear planar Hamiltonian systems, *J. Differ. Equ.* 183 (2002) 342–367.
- [40] J. Mawhin, Forced second order conservative systems with periodic nonlinearity, in: *Analyse non linéaire*, Perpignan, 1987, *Ann. Inst. Henri Poincaré, Anal. Non Linéaire* 6 (suppl) (1989) 415–434.
- [41] J. Mawhin, Multiplicity of solutions of variational systems involving ϕ -Laplacians with singular ϕ and periodic nonlinearities, *Discrete Contin. Dyn. Syst.* 32 (2012) 4015–4026.
- [42] J. Mawhin, M. Willem, Multiple solutions of the periodic boundary value problem for some forced pendulum-type equations, *J. Differ. Equ.* 52 (1984) 264–287.
- [43] J. Moser, E.J. Zehnder, *Notes on Dynamical Systems*, Amer. Math. Soc., Providence, 2005.
- [44] M. Nagumo, Über die Differentialgleichung $y'' = f(t, y, y')$, *Proc. Phys. Math. Soc. Jpn.* 19 (1937) 861–866.
- [45] E. Picard, Sur l'application des méthodes d'approximations successives à l'étude de certaines équations différentielles ordinaires, *J. Math. Pures Appl.* 9 (1893) 217–271.
- [46] H. Poincaré, Sur un théorème de géométrie, *Rend. Circ. Mat. Palermo* 33 (1912) 375–407.
- [47] P.H. Rabinowitz, On a class of functionals invariant under a \mathbb{Z}^n action, *Trans. Am. Math. Soc.* 310 (1988) 303–311.
- [48] G. Scorza Dragoni, Il problema dei valori ai limiti studiato in grande per le equazioni differenziali del secondo ordine, *Math. Ann.* 105 (1931) 133–143.
- [49] A. Szulkin, A relative category and applications to critical point theory for strongly indefinite functionals, *Nonlinear Anal.* 15 (1990) 725–739.
- [50] E.J. Zehnder, The beginnings of symplectic topology in Bochum in the early eighties, *Jahresber. Dtsch. Math.-Ver.* 121 (2019) 71–90.