



Finite electrodynamics from T-duality

Patricio Gaete^a, Piero Nicolini^{b,c,d,*}

^a Departamento de Física and Centro Científico-Tecnológico de Valparaíso-CCTVal, Universidad Técnica Federico Santa María, Valparaíso, Chile

^b Center for Astro, Particle and Planetary Physics, New York University Abu Dhabi, Abu Dhabi, United Arab Emirates

^c Frankfurt Institute for Advanced Studies (FIAS), Frankfurt am Main, Germany

^d Institut für Theoretische Physik, Goethe-Universität, Frankfurt am Main, Germany

ARTICLE INFO

Article history:

Received 22 February 2022

Received in revised form 5 April 2022

Accepted 8 April 2022

Available online 13 April 2022

Editor: A. Ringwald

ABSTRACT

In this paper, we present the repercussions of Padmanabhan's propagator in electrodynamics. This corresponds to implement T-duality effects in a U(1) gauge theory. By formulating a nonlocal action consistent with the above hypothesis, we derive the profile of static potentials between electric charges via a path integral approach. Interestingly, the Coulomb potential results regularized by a length scale proportional to the parameter $(\alpha')^{1/2}$. Accordingly, fields are vanishing at the origin. We also discuss an array of experimental testbeds to expose the above results. It is interesting to observe that T-duality generates an effect of dimensional fractalization, that resembles similar phenomena in fractional electromagnetism. Finally, our results have also been derived with a gauge-invariant method, as a necessary check of consistency for any non-Maxwellian theory.

© 2022 The Author(s). Published by Elsevier B.V. This is an open access article under the CC BY license (<http://creativecommons.org/licenses/by/4.0/>). Funded by SCOAP³.

1. Introduction

The formulation of a unified description of fundamental interactions governing the Universe from microscopic to macroscopic scales represents a path fraught with many difficulties. For instance, the combination of gravity and quantum mechanics is technically and conceptually challenging, the origin of dark sectors and, more in general, the phenomenology beyond the Standard Model and Einstein gravity are not yet clear. Even more problematic is the inclusion of such a phenomenology within a unique formulation, that works from the Planck scale, $M_P \sim 10^{19}$ GeV, down to the Terascale.

Superstring theory is probably the major contender for a self consistent description of all interactions at quantum level [1]. It enjoys ultraviolet finiteness [2,3] and absence of anomalies [4]. It is, however, not free from limitations. For instance, it is a hard task to identify the string landscape, namely effective field theories that are equipped with genuine stringy effects [5]. In addition, superstring theory is currently in conflict with experimental evidence due to the non observation of supersymmetry signals.¹ It is probably wiser to postpone the analysis of all these problems

and try to better understand the repercussions of some model independent features, i.e., universal characteristics common to any formulation of quantum gravity. For instance, a quantum spacetime is expected to display an intrinsic non-local character and be singularity free. As a result, a lot of efforts has been devoted to improve classical spacetime geometries, by following a variety of quantum gravity paradigms, e.g., noncommutative geometry [6,7], the generalized uncertainty principle [8–11], asymptotically safe gravity [12], loop quantum gravity [13], gravity self-completeness [14–16] and non-local gravity [17,18].

Recently, a regular black hole solution has been derived by implementing T-duality effects [19]. The essence of T-duality is the invariance of string theory under inversion of the compactification radius, $R \longleftrightarrow \alpha'/R$ [20]. This necessarily implies the existence of a minimal compactification radius $l_0 \sim \sqrt{\alpha'}$, namely a scale at which the classical notion of spacetime ceases to exist. Such a feature of spacetime is reflected also in another byproduct of string theory, namely a modification of uncertainty relations

$$\Delta x \simeq \frac{1}{\Delta p} + \alpha' \Delta p \quad (1)$$

known as generalized uncertainty principle [21–23]. Eq. (1) has an additional term that prevents Compton wavelengths to be smaller than $\sqrt{\alpha'}$. The same conclusion descends from noncommutative geometry, a feature of a quantum spacetime to which T-duality is connected both in the case of open and closed strings [24,25].

* Corresponding author.

E-mail addresses: patricio.gaete@usm.cl (P. Gaete), nicolini@fias.uni-frankfurt.de (P. Nicolini).

¹ For a balanced report of the current status of quantum gravity proposals (including string theory) we recommend section 7 of Nicolai's paper [1].

Similarly Padmanabhan's formulation of T-duality is based on a modification of the path integral representation of a field propagator [26]. To accommodate a spacetime "zero-point length", paths smaller or larger than such a length should contribute in the same way to the path integral. As a result, the latter has to be invariant with respect to the duality transformation $s \longleftrightarrow l_0^2/s$, where s is the path proper time.² A propagator of this kind actually reads [26]

$$G(k) = -\frac{l_0}{\sqrt{k^2}} K_1(l_0 \sqrt{k^2}), \quad (2)$$

where $K_1(x)$ is a modified Bessel function of the second kind. Such a function introduces an exponential decaying term for large arguments, that guarantees the ultraviolet finiteness. Interestingly, the exact form of Padmanabhan's propagator (2) can analytically be derived from string theory, as it has been shown in a series of papers [28–30].

It is important to stress that the regularity of the black hole solution [19] discloses another physical meaning of l_0 . From the metric coefficient

$$g_{00} = 1 - \frac{2Mr^2}{(r^2 + l_0^2)^{3/2}} \quad (3)$$

one realizes that the spacetime coincides with Bardeen regular black hole, after switching, the magnetic charge with the zero-point length, $g \rightarrow l_0$. This means that the ultraviolet regulator can be seen as the focal point of another duality, namely that between gravity and gauge theories.³

On these premises, it is now imperative to explore the role of the propagator (2) for the physics of other interactions. Among the motivations for such an investigation, there is the fact that, in general, not only gravity but also customary field theories are subject to non-local corrections at a certain scale, where point-like objects become ill defined. Within such a paradigm, we start in the present paper from the simplest case of a U(1) theory, namely electrodynamics. To this purpose, it is important to clarify the regime we aim to explore. High energy electrodynamics is expected to depart from Maxwell's formulation. Strong fields show non-linearity, a feature that offers a rich phenomenology, e.g. birefringence and photon-photon scattering effects [32], currently under experimental scrutiny [33,34]. On the other hand, we will not consider such non-linear effects but rather we will stay below the so called Schwinger limit, namely

$$E \ll E_c = \frac{m_e^2 c^3}{q_e \hbar} \simeq 1.32 \times 10^{18} \text{ V/m} \quad (4)$$

$$B \ll B_c = \frac{m_e^2 c^2}{q_e \hbar} \simeq 4.41 \times 10^9 \text{ T}. \quad (5)$$

Our approach aims to analyze the short scale behavior of low intensity fields. More importantly, it offers the advantage of understanding the net effect of T-duality without interference from other phenomena of the strong field regime. Finally, we will consider only static conditions throughout the paper. As one can see, this assumption will be enough to capture the essential features of the modified electrodynamics.

The paper is organized as follows: in Sec. 2 we present the derivation of the static forces between two sources by employing

² We incorporate factors of the "zero-point length" by a suitable redefinition of l_0 . For more details see [27].

³ For an expanded interpretation of the dualities emerging from the metric (3), see the analysis in [31] based on the double copy mechanism.

a path integral approach; in Sec. 3 we discuss the significance of our findings and their phenomenological repercussions; in Sec. 4 we propose an alternative derivation of static forces based on a path dependent, gauge invariant formalism; finally in Sec. 5, we summarize our results and we draw the conclusions.

2. Finite Maxwell-like electrodynamics

In this section, we aim to derive the interaction energy between static-point like sources, by assuming that photon exchange is governed by the T-dual propagator (2). The result will turn to be equivalent to that of standard (local) electrodynamics between T-duality modified (i.e. non-local) static sources.

The starting point of our discussion is provided by the following four dimensional spacetime Lagrangian density⁴

$$\mathcal{L} = -\frac{1}{4} F_{\mu\nu} \mathcal{O} F^{\mu\nu}, \quad (6)$$

where \mathcal{O} is the non-local operator

$$\mathcal{O} = \left[l_0 \sqrt{\Delta} K_1(l_0 \sqrt{\Delta}) \right]^{-1}, \quad (7)$$

with $\Delta \equiv \partial_\mu \partial^\mu$. Here K_1 is the modified Bessel function of the second kind and l_0 denotes a minimal length, or zero-point length of spacetime. We stress that the Lagrangian (6) is manifestly gauge invariant, despite the presence of a mass term $\sim 1/l_0$. For further clarifications about this point, see Sec. 4.

The novelty of (6) is connected to the properties of the non-local operator \mathcal{O} . For small momenta, the above action becomes the usual Maxwell action, namely

$$\mathcal{O} \approx 1 \quad \text{for} \quad l_0 \sqrt{\Delta} \ll 1. \quad (8)$$

On the other hand, for large momenta the operator reads:

$$\mathcal{O} \approx \sqrt{\frac{2}{\pi}} e^{l_0 \sqrt{\Delta}} \quad \text{for} \quad l_0 \sqrt{\Delta} \gg 1. \quad (9)$$

The above relation is the key element for having an improvement of the short distance behavior of the field strength $F_{\mu\nu}$.

According to the customary path integral formulation, one can write the functional generator of the Green's functions as

$$Z[J] = \exp\left(-\frac{i}{2} \int d^4x d^4y J^\mu(x) D_{\mu\nu}(x, y) J^\nu(y)\right), \quad (10)$$

where $D_{\mu\nu}(x) = \int \frac{d^4k}{(2\pi)^4} D_{\mu\nu}(k) e^{-ikx}$ is the propagator in the Feynman gauge, which reads

$$D_{\mu\nu}(k) = -\frac{1}{k^2} \left\{ \mathcal{O}^{-1}(k^2) \eta_{\mu\nu} + \left(1 - \mathcal{O}^{-1}(k^2)\right) \frac{k_\mu k_\nu}{k^2} \right\}. \quad (11)$$

With the aid of the expression, $Z = e^{iW[J]}$, and (10), $W[J]$ becomes

$$W[J] = -\frac{1}{2} \int \frac{d^4k}{(2\pi)^4} J_\mu^*(k) \left[-\frac{1}{k^2} \mathcal{O}^{-1}(k^2) \eta_{\mu\nu} \right] J_\nu(k) - \frac{1}{2} \int \frac{d^4k}{(2\pi)^4} J_\mu^*(k) \left[-\frac{1}{k^2} \left(1 - \mathcal{O}^{-1}(k^2)\right) \frac{k_\mu k_\nu}{k^2} \right] J_\nu(k). \quad (12)$$

Now, bearing in mind that the external current $J^\mu(k)$ is conserved, we promptly obtain

⁴ We consider natural units from this point on, namely $\hbar = c = 1$.

$$W[J] = \frac{1}{2} \int \frac{d^4k}{(2\pi)^4} J_\mu^*(k) \left[\frac{l_0 \sqrt{-k^2} K_1(l_0 \sqrt{-k^2})}{k^2} \right] J^\mu(k). \quad (13)$$

For $J_\mu(x) = [Q\delta^{(3)}(\mathbf{x} - \mathbf{x}^{(1)}) + Q'\delta^{(3)}(\mathbf{x} - \mathbf{x}^{(2)})]\delta_\mu^0$, we find that the interaction energy can be brought to the form

$$V = Q Q' \int \frac{d^3k}{(2\pi)^3} \frac{l_0 \sqrt{\mathbf{k}^2}}{\mathbf{k}^2} K_1(l_0 \sqrt{\mathbf{k}^2}) e^{i\mathbf{k}\cdot\mathbf{r}}, \quad (14)$$

where $\mathbf{r} \equiv \mathbf{x}^{(1)} - \mathbf{x}^{(2)}$. By using the integral representation of the Bessel function (see, 3.471(12) of [35])

$$\int_0^\infty x^{\nu-1} \exp\left(-x - \frac{\mu^2}{4x}\right) dx = 2\left(\frac{\mu}{2}\right)^\nu K_\nu(\mu), \quad (15)$$

we express the integral over \mathbf{k} in the form

$$\begin{aligned} & \int \frac{d^3k}{(2\pi)^3} \frac{l_0 \sqrt{\mathbf{k}^2} K_1(l_0 \sqrt{\mathbf{k}^2})}{\mathbf{k}^2} e^{i\mathbf{k}\cdot\mathbf{r}} \\ &= \int_0^\infty dx e^{-x} \int \frac{d^3k}{(2\pi)^3} \frac{e^{-\frac{l_0^2 \mathbf{k}^2}{4x}}}{\mathbf{k}^2} e^{i\mathbf{k}\cdot\mathbf{r}} \\ &= \frac{1}{4(\pi)^{\frac{3}{2}}} \frac{1}{r} \int_0^\infty e^{-x} \gamma\left(1/2, \frac{x r^2}{l_0^2}\right) = \frac{1}{4\pi} \frac{1}{\sqrt{r^2 + l_0^2}} \end{aligned} \quad (16)$$

with $r = |\mathbf{r}|$, $k = |\mathbf{k}|$. Here $\gamma\left(1/2, \frac{x r^2}{l_0^2}\right)$ is the lower incomplete Gamma function defined by the following integral representation

$$\gamma\left(\frac{a}{b}, x\right) \equiv \int_0^x \frac{du}{u} u^{a/b} e^{-u}. \quad (17)$$

Combining (14) and (16), together with $Q' = -Q$, the interaction energy reads

$$V(r) = -\frac{Q^2}{4\pi} \frac{1}{\sqrt{r^2 + l_0^2}}. \quad (18)$$

In contrast to the usual Maxwell's theory, the above energy is regular at the origin. This is the direct consequence of the string T-duality encoded in the propagator (13), in analogy to what found for the Newton's potential in [19].

One can further observe that (16) is nothing but the familiar Green's function, $\tilde{G}(\mathbf{z}, \mathbf{z}')$, for the T-duality modified electrodynamics, namely,

$$\nabla^2 \tilde{G}(\mathbf{z}, \mathbf{z}') = -l_0 \sqrt{-\nabla^2} K_1(l_0 \sqrt{-\nabla^2}) \delta^{(3)}(\mathbf{z} - \mathbf{z}'), \quad (19)$$

where the standard (i.e. Maxwell theory) Green's function, $G_0(\mathbf{z}, \mathbf{z}')$, reads

$$G_0(\mathbf{z}, \mathbf{z}') = \left[l_0 \sqrt{-\nabla^2} K_1(l_0 \sqrt{-\nabla^2}) \right]^{-1} \tilde{G}(\mathbf{z}, \mathbf{z}'). \quad (20)$$

In fact, by transforming $\tilde{G}(\mathbf{z}, \mathbf{z}')$ into Fourier space, we obtain (16). The equation for the Green's function in (19) displays a non-local smearing of the source, that resembles what found in the context of noncommutative geometry [36,37].

Alternatively one can write (19) as

$$\mathcal{D}^2 \tilde{G}(\mathbf{z}, \mathbf{z}') = -\delta^{(3)}(\mathbf{z} - \mathbf{z}'), \quad (21)$$

with

$$\mathcal{D}^2 = \frac{\nabla^2}{l_0 \sqrt{-\nabla^2} K_1(l_0 \sqrt{-\nabla^2})}. \quad (22)$$

Eq. (21) is the equation of a Green's function with a non-local operator and point-like source term. It is the equivalent of (19), that is the equation of a Green's function with a local operator and a non-local source term.

3. Discussion and significance

We present in this section some comments about the interpretation and the phenomenological repercussions of our results.

The simplest Coulomb system is the hydrogen atom, that has recently been proposed as a testbed for exotic quantum field theory effects [38]. In the case of unparticles, the accuracy of measurements of hydrogen energy levels provides limits for the unparticle scale Λ_U , that can compete with those from the muon anomaly [39] and other experiments at the LHC [40]. This is particularly the case for small scaling dimension $d_U \approx 1$, (i.e., in the Maxwell limit), since unparticle mediated electrodynamics is not analytic in this limit and corrections become increasingly large. For the potential (18), the study of effects on the hydrogen atom has been considered in [41]. In this case, however, the constraints are less effective since the theory depends on one parameter only, l_0 , and it is analytic for $l_0 \rightarrow 0$. One can estimate this also by considering

$$\frac{\Delta V(a_0)}{V_0(a_0)} \approx \frac{1}{2} \frac{l_0^2}{a_0^2} < \epsilon \quad (23)$$

where $a_0 \simeq 5.29 \times 10^{-11}$ m is the Bohr radius, $\Delta V(a_0) = V(a_0) - V_0(a_0)$, with $V_0(r)$ the Coulomb potential energy. Here $\epsilon \simeq 7.35 \times 10^{-10}$ is the relative error of standard electrodynamics for the ground state energy, namely the uncertainty on the Rydberg [42, 43]. Accordingly, one finds that $l_0 < 2.03 \times 10^{-15}$ m, in agreement with the full analysis in [41].

Another possible testbed for electrodynamics is offered by the Casimir effect. In case of unparticle modifications, one can find strong bounds on parameters governing the theory, because of the aforementioned singularity of the Maxwell limit [44]. Again, the T-duality electrodynamics is not expected to be strongly constrained in Casimir systems. The only possibility would be to postulate that the action (6), rather than being an extension of standard electrodynamics, describes another sector beyond the Standard Model, e.g., a magnetic monopole as suggested by the role of l_0 in the metric (3). Accordingly, the Maxwell limit would turn to be singular in analogy to the unparticle case.

Interestingly, in the presence of unparticles, one expects the full Casimir plate fractalization. In other words, an un-photon field between the capacitor plates "sees" their dimension as a continuous number, rather than simply two dimensions for a plane. The same phenomenon occurs in the case of the event horizon of a black hole in un-particle mediated gravity [45]. This can be explained by the conformal flatness of two dimensional spacetimes and by the fact that unparticles enjoy scale invariance [46]. Even if this is not the case for T-dual electrodynamics, the fractalization does show up in (18) too, since the spacetime dimension becomes a continuous number depending on l_0 . Indeed one can define the following fractal dimension

$$\mathbb{D} \equiv 3 - \frac{\partial \ln V(r)}{\partial \ln r} \quad (24)$$

corresponding to the power law of a generic static potential $V \sim 1/r^{\mathbb{D}-3}$. For the Coulomb case, the above equation gives $\mathbb{D} = 4$,




Fig. 1. Shape of the potential energy (18) for $\frac{Q}{4\pi} = 1$ and $l_0 = 0.1$. The dashed line represents the Coulomb potential.

signaling the absence of fractalization and the absence of non-local effects, as expected. Conversely for the T-duality case one finds the following scale dependent dimension:

$$\mathbb{D} = 3 + \frac{r^2}{r^2 + l_0^2}. \quad (25)$$

The above equation discloses two regimes: a Coulomb regime corresponding to the standard infrared dimension $\mathbb{D} = 4$ at $r \gg l_0$, and a ultraviolet dimensionally reduced regime, $\mathbb{D} \approx 3.5$ for $r \sim l_0$. The domain of the above dimension is actually $3 \leq \mathbb{D} < 4$. Notice that $\mathbb{D} = 3$ means that the potential is constant at the origin, $V \sim 1/r^0$, namely the signature of finite electrodynamics.

At this point, a remark about the short distance limits, $r \sim l_0$ and $r \ll l_0$, is mandatory. At first sight, it could appear counter-intuitive to consider length scales below the minimal length l_0 . Ideally, one would like to perform a physical limit, $r \rightarrow l_0$, to obtain a finite potential $V \sim 1/l_0$. Such a physical limit would require a “Swiss cheese” spacetime structure. In practice, cutting off a “ball” centered around the charge out of the manifold is mathematically hard or impossible to do. To circumvent such a difficulty, in our formulation we traded the cut off of the manifold ball with an integrand function in (16) that provides negligible contributions to the electrostatic potential for $r < l_0$ (see Fig. 1). As a result, it is formally still possible to consider limits of the fractal dimension for $r \rightarrow 0$. Such a limit tells us that the theory is finite. The other case, namely $r \sim l_0$, tells us that in such a regime the manifold is already subjected to wild quantum mechanically fluctuations. The spacetime is actually a fractal with non-integer dimension.

From (18), one can derive the electric field $\mathbf{E} = -\nabla V/Q$, where V/Q is the electric potential. After taking the gradient one finds:

$$\mathbf{E}(\mathbf{r}) = -\frac{Q r}{4\pi (r^2 + l_0^2)^{3/2}} \hat{\mathbf{r}}. \quad (26)$$

Interestingly, the magnitude of the above field is always smaller than the magnitude of the corresponding field $E_0(r)$ of the Maxwell’s theory, namely $E(r) < E_0(r)$. In addition $E(r)$ vanishes both at large distances and at the origin (see the stationary points of the potential in Fig. 1). In particular the field has a Coulomb like behavior, $E \sim Q/r$ for $r \gg l_0$, and is linearly vanishing at short scales, namely $E \approx Qr/l_0^3$ for $r \ll l_0$. This fact confirms not only the good short distance behavior of fields in the presence of T-duality effects, but it also says that, at an intermediate distance $0 < r_0 < \infty$, there exists a maximum for the magnitude of the electric field, namely $E_{\max} \equiv E(r_0)$. This can be obtained as follows:

$$0 = \frac{dE}{dr} = \frac{Q}{4\pi} \frac{(l_0^2 - 2r^2)}{(r^2 + l_0)^{3/2}} \sim E_{\max} = \frac{Q}{6\sqrt{3}\pi l_0^2}. \quad (27)$$

At this point, we recall that the Schwinger limit for a static particle (4) implies that the amount of energy stored in its electrostatic field cannot exceed its rest mass. For an electron in standard Maxwell electrodynamics, one has⁵

$$(eE_0)^2 \lambda_e^3 < m_e \sim E_0 < E_c \simeq 12m_e^2, \quad (28)$$

where the field energy momentum tensor $T^{00} \sim E^2$ is assumed to be uniformly distributed within the volume $\sim \lambda_e^3$, with the electron Compton wavelength and mass being λ_e and m_e and $\lambda_e \sim 1/m_e$, and $\alpha = e^2 \simeq 1/137$ ($1/e \simeq 12$). Eq. (28) actually coincides with the Schwinger result (4).

If one repeats the above calculation with the field modified by the T-duality in (26), there will be no substantial difference for weak fields. Being $E(r) < E_0(r)$, if the Schwinger limit is satisfied by Maxwell electrodynamics, then T-duality electrodynamics satisfies it too. Conversely, if Maxwell electrodynamics violates the Schwinger limit (namely it requires non-linear corrections), this could not be the case for the action (6). It is therefore instructive to estimate the value of l_0 , at which non-linear effects set in.⁶ To this purpose, one can consider the energy density $\sim e^2 E_{\max}^2$ in a volume $\sim l_0^3$. From (28) one obtains that $l_0 > 2 \times 10^{-17}$ m.

The existence of such a bound mainly says that the scale at which the action (6) requires non-linear corrections is around 10 GeV. This sounds surprising because T-duality softens electrostatic fields. If one considers the general requirement $10 \text{ TeV} < 1/l_0 < M_p$, it descends that any future observation of the T-duality at scales below $(10 \text{ TeV})^{-1} \sim 10^{-22}$ m must be accompanied by non-linearity effects. In other words, there is no energy regime at which T-duality electrodynamics shows up as a linear theory. Such a conclusion confirms early results about an interplay between T-duality and non-linear field theory actions [48].

4. Gauge-invariant calculation

The results up to this point have been obtained by working within a specific gauge, known as Feynman gauge. Electrodynamics is gauge invariant. Therefore there is no issue in following a similar method, if the theory is the Maxwell’s one. In general, one has to make sure that, for a Lagrangian like that in (6), with $O \neq 1$, the resulting potentials (18) do not depend on a specific value of the gauge parameter. For sake of consistency, it is therefore necessary to repeat the calculation with a gauge-invariant, but path-dependent, variables formalism, proposed by one of us in a series of papers [49–51].

To this purpose, we shall first examine the Hamiltonian framework for this theory. As a start, it should be noticed that the theory described by (6) contains higher time derivative terms. In our special case, however, the goal is the study of static potentials and no particular issue arises. For notational convenience, we will keep the customary notation for the differential operator Δ , even if one can practically substitute it with $-\nabla^2$.

The canonical momenta are $\Pi^\mu = -\mathcal{O}F^{0\mu}$. It is easy to see that Π^0 vanishes, we then have the usual constraint equation, which according to Dirac’s theory is written as a weak (\approx) equation, that is, $\Pi^0 \approx 0$. The remaining nonzero momenta must also be written as weak equations, that is, $\Pi^i \approx \mathcal{O}F^{i0}$. The canonical Hamiltonian

⁵ We neglect multiplicative factors that are irrelevant for the estimate of orders of magnitude.

⁶ There exist alternative proposals for the value of the field strength at which non-linearity become important [47]. Accordingly bounds on l_0 can differ.

is now obtained in the usual way via a Legendre transform. It then reads

$$H_C \approx \int d^3x \left\{ -A_0 \partial_i \Pi^i - \frac{1}{2} \Pi_i \mathcal{O}^{-1} \Pi^i + \frac{1}{4} F_{ij} \mathcal{O} F^{ij} \right\}, \quad (29)$$

which must also be written as a weak equation. Next, the primary constraint, $\Pi_0 \approx 0$, must be satisfied for all times. An immediate consequence of this is that, using the equation of motion, $\dot{Z} \approx [Z, H_C]$, we obtain the secondary constraint, $\Gamma_1(x) \equiv \partial_i \Pi^i \approx 0$. It is easy to check that there are no further constraints in the theory, and that the above constraints are first class.

By proceeding in the same way as in [49–51], we obtain the corresponding total (first class) Hamiltonian that generates the time evolution of the dynamical variables by adding all the first class constraints. We thus write $H = H_C + \int d^3x (c_0(x) \Pi_0(x) + c_1(x) \Gamma_1(x))$, where $c_0(x)$ and $c_1(x)$ are arbitrary Lagrange multipliers. Now we recall that, when this new Hamiltonian is employed, the equation of motion of a dynamic variable may be written as a strong equation. With the aid of (29), we find that $\dot{A}_0(x) = [A_0(x), H] = c_0(x)$, which is an arbitrary function. Since $\Pi^0 \approx 0$ always, neither A^0 nor Π^0 are of interest in describing this system and may be discarded from the theory. Actually, the term containing A^0 is redundant, because it can be absorbed by redefining the function $c(x) \equiv c_1(x) - A_0(x)$. As a consequence, the Hamiltonian now reads

$$H = \int d^3x \left\{ c(x) \partial_i \Pi^i - \frac{1}{2} \Pi_i \mathcal{O}^{-1} \Pi^i + \frac{1}{4} F_{ij} \mathcal{O} F^{ij} \right\}. \quad (30)$$

Now the presence of this new arbitrary function, $c(x)$, is undesirable since we have no way of giving it a meaning in a quantum theory. Hence, according to the usual procedure, we impose a gauge condition such that the full set of constraints becomes second class. A particularly appealing and useful choice is given by

$$\Gamma_2(x) \equiv \int_{C_{\xi x}} dz^\nu A_\nu(z) \equiv \int_0^1 d\lambda x^i A_i(\lambda x) = 0, \quad (31)$$

where λ ($0 \leq \lambda \leq 1$) is the parameter describing the space-like straight path $z^i = \xi^i + \lambda(x - \xi)^i$, and ξ is a fixed point (reference point), on a fixed time slice. There is no essential loss of generality if we restrict our considerations to $\xi^i = 0$. As a consequence, the only nontrivial Dirac bracket is given by

$$\begin{aligned} \{A_i(x), \Pi^j(y)\}^* &= \delta_i^j \delta^{(3)}(x-y) \\ &- \partial_i^x \int_0^1 d\lambda x^j \delta^{(3)}(\lambda x - y). \end{aligned} \quad (32)$$

To compute the interaction energy we will calculate the expectation value of the energy operator H in the physical state $|\Phi\rangle$. Let us also mention here that, as was first established by Dirac [52], the physical states $|\Phi\rangle$ correspond to the gauge invariant ones. Thus, the corresponding physical states take the form

$$\begin{aligned} |\Phi\rangle &\equiv |\bar{\Psi}(\mathbf{y}) \Psi(\mathbf{y}')\rangle \\ &= \bar{\psi}(\mathbf{y}) \exp\left(iQ \int_{\mathbf{y}'}^{\mathbf{y}} dz^i A_i(z)\right) \psi(\mathbf{y}') |0\rangle, \end{aligned} \quad (33)$$

where $|0\rangle$ is the physical vacuum state and the line integral appearing in the above expression is along a spacelike path starting at \mathbf{y}' and ending at \mathbf{y} , on a fixed time slice and Q is the external

charge. In this case, each of the states $|\Phi\rangle$ represents a fermion-antifermion pair surrounded by a cloud of gauge fields to maintain gauge invariance.

Moreover, using the Hamiltonian formalism developed in [53], we have

$$\{\Pi_k(\mathbf{x}), \Psi(\mathbf{y})\}^* = iQ \int_0^1 d\lambda y_k \delta^{(3)}(\mathbf{x} - \lambda \mathbf{y}) \Psi(y), \quad (34)$$

and

$$\{\Pi_k(\mathbf{x}), \bar{\Psi}(\mathbf{y})\}^* = -iQ \int_0^1 d\lambda y_k \delta^{(3)}(\mathbf{x} - \lambda \mathbf{y}) \bar{\Psi}(y). \quad (35)$$

Next we will consider the state $\Pi_i(\mathbf{x})|\Phi\rangle$, that is,

$$\begin{aligned} \Pi_i(\mathbf{x})|\Phi\rangle &= \bar{\Psi}(\mathbf{y}) \Psi(\mathbf{y}') \Pi_i(\mathbf{x})|0\rangle \\ &+ ([\Pi_i(\mathbf{x}), \bar{\Psi}(\mathbf{y})] \Psi(\mathbf{y}') + \bar{\Psi}(\mathbf{y}) [\Pi_i(\mathbf{x}), \Psi(\mathbf{y}')]) |0\rangle. \end{aligned} \quad (36)$$

By means of (34) and (35), we can rewrite (36) in the following way

$$\begin{aligned} \Pi_i(x) |\bar{\Psi}(\mathbf{y}) \Psi(\mathbf{y}')\rangle &= \bar{\Psi}(\mathbf{y}) \Psi(\mathbf{y}') \Pi_i(x) |0\rangle \\ &+ Q \int_{\mathbf{y}}^{\mathbf{y}'} dz_i \delta^{(3)}(\mathbf{z} - \mathbf{x}) |\Phi\rangle. \end{aligned} \quad (37)$$

In the case under consideration the expectation value, $\langle H \rangle_\Phi$, reads

$$\langle H \rangle_\Phi = \langle \Phi | \int d^3x -\frac{1}{2} \Pi_i \mathcal{O}^{-1} \Pi^i | \Phi \rangle. \quad (38)$$

Now making use of equation (37), we find that the expectation value can be brought to the form

$$\langle H \rangle_\Phi = \langle H \rangle_0 + \langle H \rangle_\Phi^{(1)}, \quad (39)$$

where $\langle H \rangle_0 = \langle 0 | H | 0 \rangle$. Whereas the $\langle H \rangle_\Phi^{(1)}$ term is given by

$$\begin{aligned} \langle H \rangle_\Phi^{(1)} &= -\frac{Q^2}{2} \int d^3x \int_{\mathbf{y}}^{\mathbf{y}'} dz_i' \delta^{(3)}(\mathbf{x} - \mathbf{z}') \\ &\times \left[-l_0 \sqrt{-\nabla_x^2} K_1 \left(l_0 \sqrt{-\nabla_x^2} \right) \right] \int_{\mathbf{y}}^{\mathbf{y}'} dz_i' \delta^{(3)}(\mathbf{x} - \mathbf{z}), \end{aligned} \quad (40)$$

which can also be expressed solely in terms of the new Green's function, \tilde{G} , namely,

$$\langle H \rangle_\Phi^{(1)} = -\frac{Q^2}{2} \int_{\mathbf{y}}^{\mathbf{y}'} dz^i \int_{\mathbf{y}}^{\mathbf{y}'} dz^j \nabla_z^2 \tilde{G}(\mathbf{z}, \mathbf{z}'). \quad (41)$$

Making use of the foregoing equation and recalling that the integrals over z^i and z'^i are zero except on the contour of integration, we finally obtain the potential for two opposite charges, located at \mathbf{y} and \mathbf{y}' ,

$$V(r) = -\frac{Q^2}{4\pi} \frac{1}{\sqrt{r^2 + l_0^2}}, \quad (42)$$

after subtracting a constant term ($\frac{Q^2}{4\pi} \frac{1}{l_0}$) and $r \equiv |\mathbf{y} - \mathbf{y}'|$

An alternative way of stating the previous result is by considering the expression (see [49]):

$$V \equiv Q (\mathcal{A}_0(\mathbf{y}) - \mathcal{A}_0(\mathbf{y}')), \quad (43)$$

where the physical scalar potential is given by

$$\mathcal{A}_0(x^0, \mathbf{x}) = \int_0^1 d\lambda x^i E_i(\lambda \mathbf{x}), \quad (44)$$

with $i = 1, 2, 3$. This equation follows from the vector gauge-invariant field expression [53]

$$\mathcal{A}_\mu(x) \equiv A_\mu(x) + \partial_\mu \left(- \int_{\xi}^x dz^\mu A_\mu(z) \right), \quad (45)$$

where the line integral is along a space-like path from the point ξ to x , on a fixed slice time. It should be noted that the gauge-invariant variables (45) commute with the sole first constraint (Gauss law), showing in this way that these fields are physical variables [52]. We also recall that Gauss' law for the Maxwell case reads $\partial_i \Pi^i = J^0$, where we have included the external source, J^0 , to represent the presence of "smeared sources". It should be further recalled that the gauge-invariant variables (45) commute with the sole first constraint (Gauss' law), corroborating that these fields are physical variables.

We also note that for

$$J^0(\mathbf{x}) = \frac{3\sqrt{2}}{2} \frac{l_0^2}{\pi^{3/2}} \frac{Q}{(r^2 + l_0^2)^{5/2}}, \quad (46)$$

the electric field is then

$$E^i = Q \partial^i \tilde{G}(x), \quad (47)$$

where \tilde{G} is the Green's function (16).

Finally, replacing this result in (44) and using (43), we readily find that the interaction energy for a pair of opposite "smeared charges" Q , located at $\mathbf{0}$ and \mathbf{r} , is given by

$$V = - \frac{Q^2}{4\pi} \frac{1}{\sqrt{r^2 + l_0^2}}, \quad (48)$$

after subtracting a constant term ($\frac{Q^2}{4\pi} \frac{1}{l_0}$) and $|\mathbf{r}| \equiv r$.

From the above discussion, it has become evident that the understanding of gauge invariance requires a correct identification of physical degrees of freedom of the system. Accordingly, only after such identification has been made, one can legitimately compute the potential by means of the Gauss' law.

5. Final remarks

In this paper we studied the repercussions of Padmanabhan's propagator in electrodynamics. Such a propagator has the property of capturing a feature of string theory known as T-duality. This corresponds to saying that any path integral has to be invariant with respect to the exchange of paths, that result larger and respectively shorter than a fundamental length l_0 . Along this line of reasoning, we computed static potentials due to the exchange of virtual photons, by introducing a non-local gauge field action for the Padmanabhan's propagator.

As a main result, we found that the T-duality actually does efficiently work at short scales. Potentials are finite and fields are vanishing at short distance. We also discussed possible testbeds of

the proposed theory in atomic physics and low energy physics experiments like the Casimir capacitor. In contrast to other non-local field theories, like the unparticles, it is less realistic to obtain competitive constraints with experiments working at energies below $1/l_0$. This is due to the fact that the proposed theory has an analytic limit to Maxwell electrodynamics for $l_0 \rightarrow 0$. We also showed that at higher energies T-duality effects would always show up in combination with non-linear electrodynamics effects. In agreement with unparticles, T-duality introduces a continuous spacetime dimension with consequent fractalization effects for static potentials. This aspect seems to be related to some aspects of the phenomenology of strange metals and, more in general, to the so called fractional electromagnetism [54].

We also presented a double check of the consistency of our approach in relation to the gauge properties of the theory. In particular we proposed an alternative derivation of static potentials based on a gauge-invariant, path-dependent variable formalism.

The topics presented in this work offer one of the rare opportunities for establishing a connection between a quantum gravity modified field theory and a concrete observation of new effects. For this reason, we believe that further investigations have to be done in this direction.

Declaration of competing interest

The authors declare the following financial interests/personal relationships which may be considered as potential competing interests:

Patricio Gaete reports financial support was provided by FONDECYT. Piero Nicolini reports a relationship with Francesco Severi National Institute of Higher Mathematics National Group for Mathematical Physics that includes: non-financial support.

Acknowledgements

One of us (P.G.) was partially supported by FONDECYT (Chile) grant 1180178 and by ANID PIA/APOYO AFB180002. The work of P.N. has partially been supported by GNFM, Italy's National Group for Mathematical Physics.

References

- [1] H. Nicolai, Quantum gravity: the view from particle physics, in: J. Bičák, T. Ledvinka (Eds.), *General Relativity, Cosmology and Astrophysics: Perspectives 100 Years after Einstein's Stay in Prague*, in: *Fundam. Theor. Phys.*, vol. 177, Springer International Publishing, Switzerland, 2014, pp. 369–387, arXiv:1301.5481 [gr-qc].
- [2] D. Amati, M. Ciafaloni, G. Veneziano, *Phys. Lett. B* 197 (1987) 81.
- [3] D. Amati, M. Ciafaloni, G. Veneziano, *Phys. Lett. B* 216 (1989) 41.
- [4] M.B. Green, J.H. Schwarz, *Phys. Lett. B* 149 (1984) 117.
- [5] C. Vafa, The string landscape and the swampland, arXiv:hep-th/0509212, 2005.
- [6] P. Nicolini, A. Smailagic, E. Spallucci, *Phys. Lett. B* 632 (2006) 547, arXiv:gr-qc/0510112 [gr-qc].
- [7] P. Nicolini, *Int. J. Mod. Phys. A* 24 (2009) 1229, arXiv:0807.1939 [hep-th].
- [8] M. Isi, J. Mureika, P. Nicolini, *J. High Energy Phys.* 11 (2013) 139, arXiv:1310.8153 [hep-th].
- [9] B.J. Carr, J. Mureika, P. Nicolini, *J. High Energy Phys.* 07 (2015) 052, arXiv:1504.07637 [gr-qc].
- [10] M. Knipfer, S. Köppel, J. Mureika, P. Nicolini, *J. Cosmol. Astropart. Phys.* 08 (2019) 008, arXiv:1905.03233 [gr-qc].
- [11] B. Carr, H. Mentzer, J. Mureika, P. Nicolini, *Eur. Phys. J. C* 80 (2020) 1166, arXiv:2006.04892 [gr-qc].
- [12] A. Bonanno, M. Reuter, *Phys. Rev. D* 62 (2000) 043008, arXiv:hep-th/0002196.
- [13] L. Modesto, *Phys. Rev. D* 70 (2004) 124009, arXiv:gr-qc/0407097 [gr-qc].
- [14] G. Dvali, C. Gomez, *Fortschr. Phys.* 61 (2013) 742, arXiv:1112.3359 [hep-th].
- [15] P. Nicolini, E. Spallucci, *Adv. High Energy Phys.* 2014 (2014) 805684, arXiv:1210.0015 [hep-th].
- [16] P. Nicolini, *Phys. Lett. B* 778 (2018) 88, arXiv:1712.05062 [gr-qc].
- [17] L. Modesto, J.W. Moffat, P. Nicolini, *Phys. Lett. B* 695 (2011) 397, arXiv:1010.0680 [gr-qc].
- [18] G. Calcagni, L. Modesto, P. Nicolini, *Eur. Phys. J. C* 74 (2014) 2999, arXiv:1306.5332 [gr-qc].

- [19] P. Nicolini, E. Spallucci, M.F. Wondrak, *Phys. Lett. B* 797 (2019) 134888, arXiv:1902.11242 [gr-qc].
- [20] A. Giveon, M. Porrati, E. Rabinovici, *Phys. Rep.* 244 (1994) 77, arXiv:hep-th/9401139.
- [21] M. Maggiore, *Phys. Lett. B* 304 (1993) 65, arXiv:hep-th/9301067.
- [22] L.J. Garay, *Int. J. Mod. Phys. A* 10 (1995) 145, arXiv:gr-qc/9403008.
- [23] A. Kempf, G. Mangano, R.B. Mann, *Phys. Rev. D* 52 (1995) 1108, arXiv:hep-th/9412167.
- [24] N. Seiberg, E. Witten, *J. High Energy Phys.* 09 (1999) 032, arXiv:hep-th/9908142.
- [25] D. Lust, *J. High Energy Phys.* 12 (2010) 084, arXiv:1010.1361 [hep-th].
- [26] T. Padmanabhan, *Phys. Rev. Lett.* 78 (1997) 1854, arXiv:hep-th/9608182.
- [27] T. Padmanabhan, *Phys. Rev. D* 57 (1998) 6206.
- [28] A. Smailagic, E. Spallucci, T. Padmanabhan, String theory T duality and the zero point length of space-time, arXiv:hep-th/0308122, 2003.
- [29] E. Spallucci, M. Fontanini, Zero-point length, extra-dimensions and string T-duality, in: S.A. Grece (Ed.), *New Developments in String Theory Research*, Nova Publishers, 2005.
- [30] M. Fontanini, E. Spallucci, T. Padmanabhan, *Phys. Lett. B* 633 (2006) 627, arXiv:hep-th/0509090.
- [31] D.A. Easson, C. Keeler, T. Manton, *Phys. Rev. D* 102 (2020) 086015, arXiv:2007.16186 [gr-qc].
- [32] S.L. Adler, *Ann. Phys.* 67 (1971) 599.
- [33] S.Z. Akhmadaliev, et al., *Phys. Rev. Lett.* 89 (2002) 061802, arXiv:hep-ex/0111084.
- [34] E. Zavattini, et al., *PVLAS, Phys. Rev. Lett.* 96 (2006) 110406, Erratum: *Phys. Rev. Lett.* 99 (2007) 129901, arXiv:hep-ex/0507107.
- [35] I.S. Gradshteyn, I.M. Ryzhik, *Table of Integrals, Series and Products*, Academic Press, 1980.
- [36] E. Spallucci, A. Smailagic, P. Nicolini, *Phys. Rev. D* 73 (2006) 084004, arXiv:hep-th/0604094.
- [37] M. Kober, P. Nicolini, *Class. Quantum Gravity* 27 (2010) 245024, arXiv:1005.3293 [hep-th].
- [38] M.F. Wondrak, P. Nicolini, M. Bleicher, *Phys. Lett. B* 759 (2016) 589, arXiv:1603.03319 [hep-ph].
- [39] K. Cheung, W.-Y. Keung, T.-C. Yuan, *Phys. Rev. Lett.* 99 (2007) 051803, arXiv:0704.2588 [hep-ph].
- [40] T.M. Aliev, S. Bilmis, M. Solmaz, I. Turan, *Phys. Rev. D* 95 (2017) 095005, arXiv:1701.00498 [hep-ph].
- [41] M.F. Wondrak, M. Bleicher, *Symmetry* 11 (2019) 1478, arXiv:1910.08203 [gr-qc].
- [42] E. Tiesinga, P.J. Mohr, D.B. Newell, B.N. Taylor, *Rev. Mod. Phys.* 93 (2021) 025010.
- [43] A.E. Kramida, *At. Data Nucl. Data Tables* 96 (2010) 586, Erratum: *At. Data Nucl. Data Tables* 126 (2019) 295–298.
- [44] A.M. Frassino, P. Nicolini, O. Panella, *Phys. Lett. B* 772 (2017) 675, arXiv:1311.7173 [hep-ph].
- [45] P. Gaete, J.A. Helayel-Neto, E. Spallucci, *Phys. Lett. B* 693 (2010) 155, arXiv:1005.0234 [hep-ph].
- [46] P. Nicolini, E. Spallucci, *Phys. Lett. B* 695 (2011) 290, arXiv:1005.1509 [hep-th].
- [47] J. Rafelski, G. Soff, W. Greiner, *Phys. Rev. A* 7 (1973) 903.
- [48] A.A. Tseytlin, *Nucl. Phys. B* 469 (1996) 51, arXiv:hep-th/9602064.
- [49] P. Gaete, *Phys. Rev. D* 59 (1999) 127702, arXiv:hep-th/9812245.
- [50] P. Gaete, E. Spallucci, *J. Phys. A* 39 (2006) 6021, arXiv:hep-th/0512178.
- [51] P. Gaete, E. Spallucci, *Phys. Rev. D* 77 (2008) 027702, arXiv:0707.2738 [hep-th].
- [52] P.A.M. Dirac, *Can. J. Phys.* 33 (1955) 650.
- [53] P. Gaete, *Z. Phys. C* 76 (1997) 355.
- [54] G. La Nave, K. Limtragool, P.W. Phillips, *Rev. Mod. Phys.* 91 (2019) 021003, arXiv:1904.01023 [hep-th].