

Table S1. Delivery characteristics and perinatal outcome of 49 pregnancies that received antenatal corticosteroids (ACS) < 14 days before delivery and 49 matched pregnancies that did not

	ACS (n=49)	No ACS (n=49)	P#
Inclusion			
Gestational age	33.4 (32.4-34.7)	33.3 (32.5-34.8)	0.84
EFW	1684 (1439-1923)	1636 (1462-1867)	0.52
UCR	0.69 (0.53-0.83)	0.66 (0.51-0.86)	0.91
Gestational age at corticosteroids	34.4 (33.7-35.8)		
Perinatal outcome			
Gestational age at delivery (weeks)	35.3 (34.2-36.4)	35.6 (34.7-36.4)	0.29
Birth weight (g)	1880 (1720-2090)	1900(1710-2098)	0.90
Male sex	28 (57)	25 (51)	0.69
Abnormal condition at birth	4 (8)	3 (6)	1.00
Major neonatal morbidity*	18 (37)	8 (16)	0.04
Cerebral morbidity	0 (0)	0 (0)	---
Cardiovascular morb.	2 (4)	0 (0)	0.50
Infection/sepsis	4 (8)	2 (4)	0.68
Respiratory morbidity	15 (31)	6 (12)	0.05
Resp. support <1 st wk	12 (25)	5 (10)	0.11
Resp. support after 1 st wk	0	0	---
Mechanical ventilation	1 (2)	0 (0)	1.00
RDS	6 (12)	1 (2)	0.11
Other resp. morbidity	0	0	---
Composite adverse outcome	19 (39)	10 (20)	0.07

Data are given as median (interquartile range) or n (%).

CS: Cesarean section

* Multiple diagnoses possible

Fisher Exact Test or Kruskal-Wallis Test