SUPPLEMENTARY MATERIALS

Table S1. List of diagnosis for disease identification and patients' characterisation

	ICD9-CM
Main inclusion criteria	
Systemic lupus erythematosus	710.0
Cutaneous only lupus	695.4, 373.34
EULAR/ACR criteria	
Leukopenia	288.50, 288.59
Thrombocytopenia	287.3, 287.4, 287.5
Autoimmune haemolysis	283.0
Delirium	297
Psychosis	290-299
Seizure	333.2, 345.0-345.9
Non-scarring alopecia	704.0
Oral ulcers	528.0
Acute pericarditis	420
Pleural and pericardial effusion	510.9, 511, 391.0, 393,423.1, 423.2
Exclusion criteria	
Primary vasculitis	446, 447.5
Myositis	729.1
Polymyositis	710.4
Dermatomyositis	710.3
Psoriatic arthritis	696.0
CREST syndrome or scleroderma	710.1
Comorbidity	
Diabetes	250
Chronic kidney disease	582, 585, 586
Cardiovascular disease (i.e., myocardial	
infarction, coronary artery bypass graft, unstable	402.01, 402.11, 402.91, 410, 411, 413, 414.0 (exc.
angina, angina pectoris, peripheral artery disease,	414.02–07), 414.02–07, V45.81–82, 440.2, 433.9, 444.2, 427.3, 428
heart failure, atrial fibrillation)	444.2, 427.3, 428
Cerebrovascular accident (i.e., stroke and	430–436, V12.54
transient ischemic attack)	,
Hypertension	401-405, 997.91
Dyslipidaemia	272
Dementia/Alzheimer's disease	290, 294.1, 331.2
Parkinson's disease	332
Mood and anviate disandars	300.0, 300.2, 296.0, 296.1, 296.04, 296.14, 296.4, 296.44, 296.5, 296.54, 296.6, 296.7, 296.8, 296.2,
Mood and anxiety disorders	290.44, 290.3, 290.34, 290.0, 290.7, 290.8, 290.2, 296.3, 298.0, 300.4, 311
Chronic hepatic disease	456.0-456.2; 571-573
Autoimmune disease	
Multiple sclerosis	340
Rheumatoid arthritis	714.0, 714.1, 714.2, 714.30, 714.32, 714.33,
Inflammatory bowel disease	564.1
Ankylosing spondylitis	720
Ankylosing sponayillis	120

Idiopathic thrombocytopenic purpura	287.31
Myasthenia gravis	358.0
Sjogren's syndrome	710.2
Osteoporosis	733
Malignancy	140–239
Chronic obstructive pulmonary disease	490, 491, 492, 494, 496

Abbreviations: ICD9-CM, International Classification of Disease, Ninth Revision, Clinical Modification classification; EULAR/ACR, European League Against Rheumatism and American College of Rheumatology.

Variable		2017	2018	2019	2020	2021	2022
			First definition	n			
Overall		4.99	5.46	4.46	5.07	11.19	6.51
		(4.79–5.18)	(5.25-5.66)	(4.27–4.65)	(4.87–5.27)	(10.89 - 11.48)	(6.29–6.74)
Sex	Female	9.08	8.91	7.61	7.54	18.33	8.87
		(5.37-12.80)	(5.27–12.55)	(4.27 - 10.94)	(4.23 - 10.84)	(13.20–23.47)	(5.16–12.57)
	Male	0.91	2.23	1.31	2.61	3.91	4.27
		(0-2.17)	(0.27–4.18)	(0-2.79)	(0.52-4.69)	(1.35–6.46)	(1.48–7.06)
Age	18–29	1.73		1.67	3.35	6.81	3.86
		(0–5.11)	0	(0-4.93)	(0–7.99)	(0.15–13.49)	(0–9.22)
	30–39	1.77	3.56	5.39	1.83	7.52	2.13
		(0–5.23)	(0-8.48)	(0–11.49)	(0–5.43)	(0.15–14.88)	(0-6.32)
	40–49	9.22	9.27	5.89	12.16	11.35	8.66
		(2.83–15.61)	(2.85–15.69)	(0.73–11.06)	(4.63–19.70)	(3.94–18.77)	(1.73–15.60)
	50–59	8.78	5.29	3.09	7.12	18.15	6.54
		(2.70–14.86)	(0.65–9.93)	(0-6.59)	(1.85–12.40)	(9.77–26.54)	(1.31–11.77)
	60–69	2.65	7.77	6.28	3.68	11.89	12.43
		(0-6.32)	(1.55–13.99)	(0.78–11.78)	(0–7.85)	(4.52–19.27)	(4.73–20.13)
	70–79	3.21	4.63	7.48	1.46	14.22	4.45
		(0–7.67)	(0–9.87)	(0.92–14.04)	(0-4.33)	(5.41–23.03)	(0–9.48)
	80+	7.02	8.94	2.09	3.92	5.67	5.85
		(0–14.97)	(0.18–17.7)	(0-6.20)	(0–9.36)	(0-12.08)	(0-12.47)
Geographical	Northern Italy	3.28	5.00	4.62	6.33	12.40	7.55
area		(3.05–3.52)	(4.72–5.29)	(4.35–4.90)	(6.01–6.66)	(11.95–12.86)	(7.19–7.90)
	Central Italy	7.92	7.69	4.86	3.63	10.76	5.03
		(7.36–8.47)	(7.14–8.23)	(4.43–5.30)	(3.26–4.01)	(10.12–11.41)	(4.59–5.47)
	Southern Italy & Islands	4.04	3.28	3.89	4.8	9.34	6.79
		(3.74–4.35)	(3.00–3.55)	(3.59–4.19)	(4.47–5.13)	(8.88–9.80)	(6.40–7.19)
		-	Second definit		-		
Overall		4.58	5.08	4.05	4.82	10.48	6.16
		(4.40–4.77)	(4.88–5.27)	(3.87–4.22)	(4.63–5.02)	(10.19–10.76)	(5.94–6.38)
Sex	Female	8.69	8.52	6.85	7.54	16.84	8.06
		(5.06–12.32)	(4.96–12.08)	(3.68–10.01)	(4.23–10.84)	(11.92–21.76)	(4.53–11.59)

Table S2. Annual (2017–2022) standardised incidence rate (per 100,000 person-years) of SLE, stratified by sex, age, and geographical area.

Variable		2017	2018	2019	2020	2021	2022
	Male	0.46	1.78	1.31	2.17	3.91	4.27
		(0-1.35)	(0.04 - 3.52)	(0-2.79)	(0.27 - 4.07)	(1.35-6.46)	(1.48 - 7.06)
Age	18–29	1.73		1.67	1.68	6.81	3.86
-		(0-5.11)	0	(0-4.93)	(0-4.96)	(0.14 - 13.49)	(0-9.22)
	30–39	1.77	3.56	3.59	1.83	7.52	2.13
		(0-5.23)	(0-8.48)	(0-8.58)	(0-5.43)	(0.15 - 14.88)	(0-6.32)
	40-49	9.22	9.27	4.72	12.16	11.35	8.66
		(2.83 - 15.61)	(2.85 - 15.69)	(0.09 - 9.34)	(4.63–19.7)	(3.94–18.77)	(1.73–15.6)
	50–59	8.78	5.29	3.09	7.12	16.13	6.54
		(2.7 - 14.86)	(0.65–9.93)	(0-6.59)	(1.85 - 12.4)	(8.23–24.04)	(1.31 - 11.77)
	60–69	2.65	7.77	6.28	3.68	11.89	12.43
		(0-6.32)	(1.55 - 13.99)	(0.78 - 11.78)	(0-7.85)	(4.52–19.27)	(4.73–20.13)
	70–79	1.61	3.09	7.48	1.46	14.22	4.45
		(0-4.75)	(0-7.37)	(0.92 - 14.04)	(0-4.33)	(5.41-23.03)	(0-9.48)
	80+	4.68	6.71	2.09	3.92	1.89	1.95
		(0-11.17)	(0-14.29)	(0-6.2)	(0-9.36)	(0-5.59)	(0-5.77)
Geographical	Northern Italy	2.81	5.00	3.65	5.76	11.97	7.55
area		(2.6–3.03)	(4.72–5.29)	(3.4–3.89)	(5.45-6.07)	(11.52–12.41)	(7.19–7.9)
	Central Italy	7.33	7.12	4.86	3.63	9.1	4.47
		(6.8–7.87)	(6.6–7.65)	(4.43–5.3)	(3.26–4.01)	(8.5–9.69)	(4.05 - 4.88)
	Southern Italy & Islands	4.04	2.42	3.89	4.8	9.34	6.07
		(3.74–4.35)	(2.18-2.66)	(3.59–4.19)	(4.47–5.13)	(8.88–9.8)	(5.7–6.45)
			Third definiti	on			
Overall		6.23	6.22	5.02	5.90	14.68	9.67
		(6.02–6.45)	(6.01–6.44)	(4.82–5.22)	(5.69-6.12)	(14.34–15.02)	(9.40–9.95)
Sex	Female	10.66	10.46	8.75	9.05	22.82	13.3
		(6.64–14.69)	(6.51–14.4)	(5.17–12.32)	(5.43-12.66)	(17.1–28.55)	(8.76–17.84)
	Male	1.82	2.23	1.31	2.61	6.08	6.17
		(0.04–3.61)	(0.27-4.18)	(0-2.79)	(0.52-4.69)	(2.89–9.26)	(2.82–9.52)
Age	18–29	3.45		1.67	3.35	8.52	3.86
		(0-8.24)	0	(0-4.93)	(0-7.99)	(1.05 - 15.98)	(0-9.22)
	30–39	1.77	3.56	5.39	5.5	9.4	6.4
		(0-5.23)	(0-8.48)	(0-11.49)	(0-11.72)	(1.16–17.63)	(0-13.64)
	40-49	10.37	11.59	7.07	12.16	22.7	11.55
		(3.6–17.15)	(4.41–18.77)	(1.41 - 12.73)	(4.63–19.7)	(12.21-33.19)	(3.55–19.56)

Variable		2017	2018	2019	2020	2021	2022
	50–59	10.98	7.41	3.09	9.16	20.17	10.90
		(4.17 - 17.78)	(1.92 - 12.9)	(0-6.59)	(3.17–15.14)	(11.33–29.01)	(4.14–17.65)
	60–69	3.97	7.77	7.53	3.68	14.27	17.40
		(0-8.47)	(1.55–13.99)	(1.51–13.56)	(0-7.85)	(6.2–22.35)	(8.29–26.52)
	70–79	4.82	4.63	8.98	1.46	17.06	7.42
		(0-10.27)	(0-9.87)	(1.8–16.17)	(0-4.33)	(7.41–26.72)	(0.92 - 13.92)
	80+	7.02	8.94	2.09	3.92	5.67	7.80
		(0-14.97)	(0.18 - 17.7)	(0-6.2)	(0–9.36)	(0-12.08)	(0.16 - 15.44)
Geographical	Northern Italy	4.83	6.34	5.49	6.33	16.6	11.24
area		(4.54–5.11)	(6.02–6.67)	(5.19-5.80)	(6.01–6.66)	(16.07 - 17.12)	(10.81 - 11.68)
	Central Italy	9.11	8.29	5.42	4.92	14.65	6.98
		(8.51-9.70)	(7.72-8.86)	(4.96–5.87)	(4.49–5.36)	(13.89–15.4)	(6.46–7.50)
	Southern Italy & Islands	4.95	3.28	3.89	6.44	11.12	10.41
		(4.61–5.29)	(3.00 - 3.55)	(3.59 - 4.19)	(6.05–6.82)	(10.61 - 11.62)	(9.92 - 10.90)

Variable		2017	2018	2019	2020	2021	2022		
First definition									
Overall		36.04	39.75	42.73	46.13	55.6	60.57		
		(35.51-36.57)	(39.2–40.3)	(42.15-43.3)	(45.53-46.73)	(54.95–56.26)	(59.89-61.25)		
Sex	Female	60.03	65.84	70.74	75.38	90.54	97.14		
		(50.49-69.58)	(55.94–75.73)	(60.58-80.91)	(64.94-85.82)	(79.14–101.9)	(84.88–109.4)		
	Male	10.93	12.46	13.11	14.33	16.93	18.98		
		(6.55–15.3)	(7.85–17.08)	(8.42–17.8)	(9.44–19.22)	(11.61-22.24)	(13.1–24.86)		
Age	18–29	22.45	21.97	23.32	26.8	34.07	38.65		
-		(10.25 - 34.65)	(10.03-33.91)	(11.11-35.54)	(13.67-39.94)	(19.14–49)	(21.71–55.59)		
	30–39	47.70	51.56	55.72	58.67	67.65	68.28		
		(29.71-65.69)	(32.8–70.32)	(36.11–75.33)	(38.34–78.99)	(45.56-89.74)	(44.63–91.93)		
	40-49	48.40	56.77	63.66	77.85	90.8	105.41		
		(33.77-63.04)	(40.88–72.66)	(46.68-80.63)	(58.78–96.92)	(69.84–111.7)	(81.24–129.6)		
	50–59	48.29	51.85	51.54	54.94	69.58	70.82		
		(34.03-62.56)	(37.34–66.37)	(37.26–65.83)	(40.29–69.59)	(53.17-85.99)	(53.61-88.03)		
	60–69	34.44	41.46	45.2	46.62	57.09	68.36		
		(21.2-47.67)	(27.1–55.83)	(30.44–59.96)	(31.8–61.44)	(40.95–73.24)	(50.3-86.42)		
	70–79	25.70	24.7	29.94	29.24	38.39	38.56		
		(13.11–38.29)	(12.6–36.81)	(16.82–43.06)	(16.43-42.05)	(23.91–52.87)	(23.74–53.38)		
	80+	18.73	22.35	23.03	17.65	17.01	19.5		
		(5.75–31.71)	(8.5–36.21)	(9.42–36.63)	(6.12–29.18)	(5.9–28.12)	(7.41–31.58)		
Geographical area	Northern Italy	42.93	45.05	47.93	52.74	63.38	69.93		
		(42.09–43.78)	(44.19–45.92)	(47.03–48.82)	(51.8–53.67)	(62.35–64.4)	(68.85–71)		
	Central Italy	36.28	42.95	46.49	48.93	58.99	62.75		
		(35.1–37.47)	(41.66–44.24)	(45.15–47.83)	(47.55–50.31)	(57.47–60.5)	(61.19–64.31)		
	Southern Italy & Islands	23.72	26.38	29.00	30.92	37.26	41.87		
		(22.98–24.46)	(25.6–27.16)	(28.19–29.82)	(30.08–31.76)	(36.33–38.18)	(40.89 - 42.85)		
			Second definition						
Overall		31.09	34.96	37.85	41.78	50.55	54.94		
		(30.6–31.58)	(34.44–35.48)	(37.31–38.39)	(41.21–42.35)	(49.93–51.18)	(54.29–55.59)		
Sex	Female	54.50	60.42	64.66	70.1	83.81	89.48		
		(45.41–63.6)	(50.94–69.89)	(54.94–74.37)	(60.03-80.17)	(72.84–94.78)	(77.72–101.2)		
	Male	6.37	8.01	9.18	10.85	13.45	15.19		

Table S3. Annual (2017–2022) standardised prevalence (per 100,000 people) of SLE stratified by sex, age, and geographical area.

Variable		2017	2018	2019	2020	2021	2022
		(3.03–9.71)	(4.31–11.71)	(5.25-13.10)	(6.6–15.11)	(8.72–18.19)	(9.92-20.45)
Age	18–29	22.45	21.97	23.32	25.13	32.37	36.72
C		(10.25 - 34.65)	(10.03-33.91)	(11.11-35.54)	(12.41-37.84)	(17.81-46.92)	(20.21–53.22)
	30–39	42.4	46.22	48.53	51.33	60.13	59.74
		(25.44-59.36)	(28.46-63.99)	(30.23-66.83)	(32.32–70.34)	(39.3-80.96)	(37.62-81.86)
	40–49	47.25	55.61	61.3	75.42	88.28	102.52
		(32.79-61.71)	(39.88–71.34)	(44.64-77.96)	(56.65–94.18)	(67.61–108.9)	(78.69–126.3)
	50–59	45.00	48.68	48.45	52.91	64.54	65.37
		(31.23–58.77)	(34.62-62.74)	(34.6–62.3)	(38.53–67.28)	(48.73-80.35)	(48.84-81.91)
	60–69	29.14	36.28	40.18	41.71	52.34	63.39
		(16.96-41.31)	(22.84-49.72)	(26.26–54.09)	(27.7–55.73)	(36.88–67.8)	(46-80.78)
	70–79	12.85	10.81	17.96	17.54	28.44	29.66
		(3.95 - 21.76)	(2.8 - 18.81)	(7.8–28.13)	(7.62 - 27.47)	(15.98 - 40.9)	(16.66-42.66)
	80+	7.02	13.41	14.65	15.69	11.34	9.75
		(0-14.97)	(2.68 - 24.14)	(3.8–25.51)	(4.82–26.56)	(2.27 - 20.41)	(1.2–18.29)
Geographical area	Northern Italy	37.22	40.41	42.45	47.51	57.63	63.95
		(36.43-38.00)	(39.59–41.23)	(41.61-43.29)	(46.62 - 48.4)	(56.65-58.61)	(62.92–64.98)
	Central Italy	33.74	39.85	43.39	45.83	54.21	57.09
		(32.6–34.88)	(38.61-41.09)	(42.1-44.69)	(44.5–47.17)	(52.77-55.66)	(55.6–58.58)
	Southern Italy & Islands	16.75	18.91	22.65	26.27	32.79	36.64
		(16.13–17.38)	(18.25–19.56)	(21.93-23.38)	(25.5 - 27.05)	(31.92–33.66)	(35.72–37.56)
			Third definition	n			
Overall		42.44	46.43	49.90	53.74	66.55	74.20
		(41.87-43.01)	(45.83-47.03)	(49.28–50.52)	(53.1–54.39)	(65.83-67.27)	(73.44–74.96)
Sex	Female	69.12	75.52	81.39	86.68	105.88	115.68
		(58.88–79.36)	(64.92-86.12)	(70.49–92.29)	(75.49–97.88)	(93.55–118.2)	(102.31–129.0)
	Male	14.57	16.02	16.61	17.80	22.57	26.58
		(9.52-19.61)	(10.79–21.25)	(11.33-21.89)	(12.35-23.25)	(16.43-28.7)	(19.62–33.53)
Age	18–29	27.63	27.04	28.32	31.83	40.88	46.38
		(14.09-41.16)	(13.79–40.28)	(14.86-41.78)	(17.52-46.14)	(24.53-57.24)	(27.83-64.93)
	30–39	56.54	60.44	64.71	71.50	82.68	83.21
		(36.95–76.12)	(40.13-80.76)	(43.58-85.84)	(49.07–93.93)	(58.26–107.1)	(57.11–109.32)
	40-49	56.47	66.04	74.27	87.58	112.24	131.4
		(40.66–72.28)	(48.9–83.18)	(55.94–92.6)	(67.36–107.8)	(88.94–135.5)	(104.42–158.4)
	50–59	54.88	60.32	59.79	64.10	79.67	86.07

Variable		2017	2018	2019	2020	2021	2022
		(39.67–70.08)	(44.67–75.98)	(44.41–75.17)	(48.27–79.92)	(62.11–97.23)	(67.1–105.05)
	60–69	41.06	47.94	52.73	53.98	66.61	82.03
		(26.61–55.51)	(32.5-63.39)	(36.79–68.67)	(38.04–69.93)	(49.17-84.05)	(62.25–101.82)
	70–79	32.13	30.88	37.42	36.55	48.35	48.94
		(18.05–46.21)	(17.35–44.41)	(22.76–52.09)	(22.22-50.87)	(32.1–64.59)	(32.25–65.64)
	80+	21.07	22.35	23.03	17.65	17.01	21.45
		(7.31 - 34.84)	(8.5-36.21)	(9.42-36.63)	(6.12-29.18)	(5.9 - 28.12)	(8.77–34.12)
Geographical area	Northern Italy	50.23	53.1	56.71	61.04	75.94	84.96
		(49.32–51.15)	(52.16–54.04)	(55.74–57.69)	(60.03-62.05)	(74.82–77.06)	(83.78-86.15)
	Central Italy	42.42	49.07	53.18	56.32	69.65	75.11
		(41.14–43.71)	(47.69–50.45)	(51.74–54.61)	(54.85–57.8)	(68–71.29)	(73.41–76.82)
	Southern Italy & Islands	28.76	31.29	33.82	37.40	45.55	54.41
		(27.95–29.57)	(30.44–32.14)	(32.94–34.7)	(36.48–38.33)	(44.53–46.57)	(53.29–55.53)

Figure S1. Graphical algorithm depicting the three different SLE definitions.

