

Table S2. Delivery characteristics and perinatal outcome of 37 pregnancies that received antenatal corticosteroids (ACS) \geq 14 days before delivery and 37 matched pregnancies that did not

	ACS (n=37)	No ACS (n=37)	P#
Inclusion			
Gestational age	33.0 (32.4-34.5)	33.1 (32.4-34.5)	0.83
EFW	1584 (1377-1900)	1626 (1456-1917)	0.77
UCR	0.66 (0.47-0.84)	0.61 (0.50-0.67)	0.12
Gestational age at corticosteroids	31.9 (30.2-33.2)	---	---
Perinatal outcome			
Gestational age at delivery (weeks)	36.9 (34.8-37.8)	36.7 (35.2-37.7)	0.71
Birth weight (g)	2020 (1760-2250)	2020 (1731-2288)	0.93
Male sex	11 (30)	13 (35)	0.80
Abnormal condition at birth	2 (5)	3 (8)	1.00
Major neonatal morbidity*	5 (14)	10 (27)	0.25
Cerebral morbidity	0 (0)	0 (0)	---
Cardiovascular morbidity	0 (0)	4 (11)	0.12
Infection/sepsis	0 (0)	1 (3)	1.00
Respiratory morbidity	5 (14)	5 (14)	1.00
Resp. support <1 st wk	4 (11)	3 (8)	1.00
Resp. support after 1 st wk	0 (0)	0 (0)	---
Mechanical ventilation	1 (3)	0 (0)	1.00
RDS	1 (3)	2 (5)	1.00
Other resp. morbidity	1 (3)	1 (3)	1.00
Composite adverse outcome	5 (14)	11 (30)	0.16

Data are given as median (interquartile range) or n (%).

CS: Cesarean section

* Multiple diagnoses possible

Fisher Exact Test or Kruskal-Wallis Test