

European Journal of Organic Chemistry

Supporting Information

Enhanced Long-Term Stability of a Photosensitizer with a Hydroxamic Acid Anchor in Dye-Sensitized Photocatalytic Hydrogen Generation

Giorgia Salerno, Bianca Cecconi, Ottavia Bettucci, Matteo Monai, Lorenzo Zani,
Daniele Franchi, Massimo Calamante, Alessandro Mordini,* Tiziano Montini,*
Paolo Fornasiero, Norberto Manfredi, and Alessandro Abbotto*

<i>Fig. S1.</i> H_2 production measured using the DYE@ TiO_2 /Pt catalyst in H_2 production from TEOA 10 v/v% solution at pH = 7.0 under irradiation with visible light ($\lambda > 420$ nm).....	2
<i>Fig. S2.</i> 1H NMR Spectrum of 1 in $CDCl_3$	2
<i>Fig. S3.</i> ^{13}C NMR Spectrum of 1 in $CDCl_3$	3
<i>Fig. S4.</i> 1H NMR Spectrum of PTZ1-HA in THF-d8.	3
<i>Fig. S5.</i> ^{13}C NMR Spectrum of PTZ1-HA in THF-d8.	4
<i>Fig. S6.</i> ESI-MS Spectrum of 1.....	5
<i>Fig. S7.</i> ESI-MS Spectrum of PTZ1-HA.	6




Fig. S1. H₂ production measured using the DYE@TiO₂/Pt catalyst in H₂ production from TEOA 10 v/v% solution at pH = 7.0 under irradiation with visible light ($\lambda > 420$ nm).




Fig. S2. ¹H NMR Spectrum of 1 in CDCl₃.




Fig. S3. ^{13}C NMR Spectrum of **1** in CDCl_3




Fig. S4. ^1H NMR Spectrum of **PTZ1-HA** in $\text{THF}-d_8$.




Fig. S5. ^{13}C NMR Spectrum of **PTZ1-HA** in $\text{THF}-d_8$.

BC111neg_150730153335 #1 RT: 0,00 AV: 1 NL: 4,10E3
T: ITMS - c ESI Full ms [150,00-2000,00]




Fig. S6. ESI-MS Spectrum of **1**.




Fig. S7. ESI-MS Spectrum of **PTZ1-HA**.