

Supplementary Material: tables

Supplementary Table 1 Risk factors for Distress according to DT scores at period's illness

Variables	Beta (95% CI)	p value
Age	0.02 (-0.03;0.06)	0.39
Gender (Male vs Female)	-0.10 (-1.18;0.97)	0.84
Number of Symptoms (≥ 7 vs < 7)	0.55 (-0.53;1.64)	0.31
Loss taste (yes vs no)	1.58 (0.15;3.00)	0.02
Loss smell (yes vs no)	-0.34 (-1.84;1.19)	0.66

Supplementary Table 2 Risk factors for anxiety according to HADS-A score at the time of Interview

Variables	Beta (95% CI)	p value
Age	-0.01 (-0.09;0.06)	0.72
Gender (Male vs Female)	-1.64 (-3.39;0.10)	0.07
Number of Symptoms (≥ 7 vs < 7)	1.19 (-0.57;2.96)	0.18
Loss taste (yes vs no)	1.74 (-0.57;4.07)	0.13
Loss smell (yes vs no)	0.24 (-2.23;2.73)	0.84

Supplementary Table 3 Risk factors for Depression according to HADS-D score at the time of Interview

Variables	Beta (95% CI)	p value
Age	0.01 (-0.05;0.07)	0.77
Gender (Male vs Female)	-1.47 (-3.00;0.05)	0.05
Number of Symptoms (≥ 7 vs <7)	0.21 (-1.34;1.75)	0.79
Loss taste (yes vs no)	1.54 (-0.49;3.57)	0.14
Loss smell (yes vs no)	-0.50 (-2.68;1.68)	0.65

Supplementary Table 4. Association between Distress and HADS-Total Scores at time of Interview

Current DT	HADS Total		p value
Scores	Scores	Scores	
	<13	≥13	
<5	52 (71.2%) (13.8%)	4	p<0.001
≥ 5	21 (28.8%) (86.2%)	25	

*for 2 patients no data regarding Current DT