

Fate of microbial metabolites of dietary polyphenols in rats: is the brain their target destination?

Mattia Gasperotti, Sabina Passamonti, Federica Tramer, Domenico Masuero, Graziano Guella, Fulvio Mattivi, and Urska Vrhovsek

ACS Chem. Neurosci., **Just Accepted Manuscript** • DOI: 10.1021/acchemneuro.5b00051 • Publication Date (Web): 18 Apr 2015

Downloaded from <http://pubs.acs.org> on April 20, 2015

Just Accepted

“Just Accepted” manuscripts have been peer-reviewed and accepted for publication. They are posted online prior to technical editing, formatting for publication and author proofing. The American Chemical Society provides “Just Accepted” as a free service to the research community to expedite the dissemination of scientific material as soon as possible after acceptance. “Just Accepted” manuscripts appear in full in PDF format accompanied by an HTML abstract. “Just Accepted” manuscripts have been fully peer reviewed, but should not be considered the official version of record. They are accessible to all readers and citable by the Digital Object Identifier (DOI®). “Just Accepted” is an optional service offered to authors. Therefore, the “Just Accepted” Web site may not include all articles that will be published in the journal. After a manuscript is technically edited and formatted, it will be removed from the “Just Accepted” Web site and published as an ASAP article. Note that technical editing may introduce minor changes to the manuscript text and/or graphics which could affect content, and all legal disclaimers and ethical guidelines that apply to the journal pertain. ACS cannot be held responsible for errors or consequences arising from the use of information contained in these “Just Accepted” manuscripts.

1
2
3
4
5
6
7
8
9
10
11
12
13
14
15
16
17
18
19
20
21
22
23
24
25
26
27
28
29
30
31
32
33
34
35
36
37
38
39
40
41
42
43
44
45
46
47
48
49
50
51
52
53
54
55
56
57
58
59
60

**The fate of microbial metabolites of dietary polyphenols in rats:
is the brain their target destination?**

Mattia Gasperotti[#], Sabina Passamonti^{§*}, Federica Tramer[§], Domenico Masuero[#], Graziano
Guella[‡], Fulvio Mattivi[#], Urska Vrhovsek^{#*}

[#] Department of Food Quality and Nutrition, Research and Innovation Centre, Fondazione
Edmund Mach (FEM), via E. Mach 1, 38010 San Michele all'Adige, Italy

[§] Department of Life Sciences, University of Trieste, via L. Giorgieri 1, 34127 Trieste, Italy

[‡] Department of Physics, University of Trento, via Sommarive 14, 38123, Trento, Italy

Abstract

Different polyphenol compounds are ingested when consuming a serving of fruits rich in polyphenols, spanning from one-phenol hydroxybenzoic acid to more complex polymeric compounds. Only a minor quantity of the polyphenols (5-10%) is absorbed. The remainder reaches the colon and is extensively metabolized by gut microbiota to low-molecular weight metabolites. Their subsequent tissue distribution is still undefined, though these microbial metabolites are currently believed to play a role in human health and disease states.

To fill this knowledge gap, we performed a pharmacokinetics experiment in which a single bolus of 23 polyphenol microbial metabolites (total 2.7 μmol) was administered intravenously to rats to reliably reproduce a physiological post-absorption situation. Tissues and urine were collected shortly thereafter (15 sec to 15 min) and were analyzed by UHPLC-MS/MS to quantitatively track these compounds. Remarkably, the brain was found to be a specific target organ for ten of the 23 polyphenol metabolites injected, which significantly increased in the treated animals. In most cases, their appearance in the brain was biphasic, with an early wave at 2 min (4 compounds) and a second wave starting at 5 min; at 15 min, 9 compounds were still detectable. Most compounds were excreted into the urine. The concentrations in the brain of the treated animals were compared against those of the control group by Student's t-test, with P-values < 0.1 considered statistically significant.

These findings provide new perspectives for understanding the role of diet on brain chemistry. Our experimental approach has enabled us to obtain rich metabolomics information from a single experiment involving a limited number of animals.

Keywords polyphenols, gut microbiota, target metabolomics, mass spectrometry, brain, fruit consumption

Introduction

A diet rich in fruit and vegetables is regarded to be an important factor for protecting the human organism from chronic diseases.¹ However, dietary recommendations require knowledge about food composition, identification of bioactive food components, characterization of bioavailability and tissue distribution, the elimination patterns of metabolites, and components' specific effects on cellular homeostasis.

Polyphenols, which are a wide and heterogeneous group of bioactive compounds found in vegetables and fruits², especially berries³, are regarded to be responsible for protecting the human organism from oxidative stress-related chronic diseases. The bioactivity of polyphenols may serve a much broader purpose, however.

After ingestion, only a minor part of the polyphenols (5-10%) is absorbed in the small intestine. The remainder reaches the colon, where it is extensively metabolized by gut microbiota to low-molecular weight compounds.⁴ Gut microbiota consist of many trillions of microbial cells⁵ whose metabolic activities have been shown to modulate human metabolic phenotypes⁶⁻⁸ and thus play an essential role in human health.

The microbiota break down dietary polyphenols through the actions of glucosidases, esterases, demethylases, dehydroxylases and decarboxylases^{5,9-18} to produce smaller common metabolites, such as phenolic acids and short-chain fatty acids⁹, collectively known as polyphenol microbial metabolites (PMMs). These are absorbed by the colonic epithelium and are found in the blood with patterns that correlate with the diet and individual microbial composition.¹⁹

PMMs are regarded to be responsible for the health effects that correlate with regular berry fruit consumption.^{4,20} Epidemiological studies associate polyphenol consumption with reduced cancer and cardiovascular disease risk.^{21,22} Research on the neuroprotective effects and prevention of brain aging by dietary polyphenols has also increased in recent years.²²⁻²⁸ The

1
2 literature reports that diets rich in polyphenols attenuate neuropathology indicators and cognitive
3 decline.²⁵ Some studies report direct effects of polyphenols on intracellular targets, e.g., three
4 members of the secretase family known to be involved in the amyloidal aggregation have been
5
6 connected with the onset and progression of Alzheimer's disease.²⁹ The question of whether
7
8 these compounds are able to pass the blood-brain barrier and achieve a pharmacologically
9
10 relevant concentration there remains unanswered.
11
12
13
14

15
16 The importance of microbiota for brain function development has been noted by
17
18 numerous studies of experimental animals³⁰⁻³², and these observations can be translated to a
19
20 better understanding of human brain disorders. For instance, autism, a defect of brain
21
22 development leading to impaired social interaction, is now understood to be linked to abnormal
23
24 microbiota population and poor intake of fruits and vegetables.³³ It might be speculated that
25
26 PMMs are involved in normal neural development and function. Indeed, they are chemically
27
28 identical or similar to catabolites of amino acids and neurotransmitters.
29
30

31
32 It seems, however, that PMMs may have some yet uncharacterized bioactivity that has an
33
34 impact on health.²⁰ Current information concerning the absorption and distribution of PMMs in
35
36 mammalian tissues is still poor^{9,34} and not substantial enough to support the hypothesis that
37
38 PMMs are the bioactive agents of our diet. Therefore, it is not possible to make dietary
39
40 recommendations for early prevention of brain dysfunctions.
41

42
43 Given this background, the focus of this study was to examine the ability of selected,
44
45 unconjugated PMMs to enter the brain relative to the main excretory organs. Therefore, we
46
47 performed an experiment to assess the time-dependent tissue distribution of a single intravenous
48
49 injection of a PMM mixture in anaesthetized rats.
50

51
52 The superior selectivity, sensitivity and dynamic range of modern tandem mass-
53
54 spectrometry enabled us to harvest data on a wide range of diverse PMMs in a single *in-vivo*
55
56 experiment. It is noteworthy that our experimental strategy involved the least number of animals,
57
58 in compliance with the 3R principles on animal experimentation.
59
60

1
2 The data obtained show for the first time the simultaneous quantitative profile of several
3
4 PMMs in a mammalian system, proving their distribution in the brain, the main excretory organs
5
6 (liver and kidneys), the heart, the blood and the urine. For some PMMs, specific tropisms were
7
8 also observed, pinpointing target organs for future experimentation. Remarkably, the data
9
10 obtained show that at least 10 PMMs incrementally appear in the brain within 15 min, whereas
11
12 they simultaneously disappear from the blood and/or reach other organs.
13
14
15
16
17
18
19

20 **Results and Discussion**

21
22
23

24 **The tested mixture of polyphenol microbial metabolites: a rational choice**

25

26 The strategy was to simultaneously inject 23 metabolites in a well-defined mixture into a
27
28 single animal (Figure 1).
29

30
31 The selected metabolites are known products of the gut-microbiota metabolism of
32
33 polyphenols found in berries.^{11–13,17,35} Their individual concentrations in the mixture were chosen
34
35 to mimic their average plasma concentration measured after the intake of a berry fruit serving.³⁶
36
37 All individual molecules were commercially available except for urolithin A and B. The latter
38
39 are regarded to be specific biomarkers of dietary ellagitannins metabolized by gut microbiota and
40
41 were synthesized in-house.³⁷
42
43

44 An analytical method based on UHPLC-MS/MS enabled quantitative measurements of
45
46 this set of 23 compounds in biological fluids and tissue extracts from a single rat,³⁸ resulting in a
47
48 remarkable reduction in the number of experimental animals, as recommended by the regulatory
49
50 bodies.³⁹
51
52

53 The *in vivo* injection of a multi-component mixture could give rise to potential problems.
54
55 Some metabolic pathways could transform certain metabolites into other compounds already
56
57 present in the mixture, thus altering the chemical composition of the injected mixture. The
58
59
60

1
2 simultaneous injection of different metabolites could also affect the rate of membrane transport,
3
4 enzyme pathways or cellular regulatory mechanisms, resulting in competitive interactions, rapid
5
6 states of saturation or cooperative responses, to name a few. However, this was a physiological
7
8 situation, closely simulating a nutritionally relevant scenario in the blood. Analyzing a multi-
9
10 component mixture made it possible to simultaneously follow the tissue distribution of each
11
12 compound, especially with respect to the brain, whereas the presence of other congeners or
13
14 further metabolites (e.g., glucuronides or sulfates) could not be followed.
15
16
17
18
19

20 **Dosage, administration route and time-line: a strategy for accurate tissue distribution** 21 **analysis**

22
23
24 The most accurate approach for a study on PMM tissue distribution is to administer the
25
26 PMMs by intravenous injection, thereby minimizing the main factors of inter-individual
27
28 variability in gut metabolism, motility and absorption.⁶ Any other downstream processes, such as
29
30 distribution into tissues and mammalian phase-1 and phase-2 metabolism, are comparatively
31
32 more homogenous among individuals.⁴⁰ As a consequence, a satisfactory variation coefficient
33
34 was obtained for all measured parameters, using the least number of animals (n=4 in this study).
35
36

37
38 The injected dose was 2.7 μmol , which was the sum of 23 individual PMM amounts
39
40 (Figure 1) from 0.9 μmol (gallic acid) to 0.0005 μmol (sinapic acid). Considering that the rats
41
42 (body mass= 288 ± 20 g) had an estimated blood volume of 16 mL⁴¹ and assuming that all of the
43
44 mixture components had free access to the internal volume of blood cells, the initial blood
45
46 concentration was 168 μM , which spanned from 56.25 μM (gallic acid) to 0.31 μM (sinapic
47
48 acid). These values are within the concentration range observed in humans following the
49
50 consumption of a standard serving of berry fruits.⁴²
51
52

53
54 The experiment was designed on the basis of previous data showing the exceptionally
55
56 fast uptake and metabolism of a dietary flavonoid, i.e., cyanidin 3-glucoside, a ubiquitous
57
58
59
60

1
2 pigment in red fruits.⁴⁰ It was assumed that the plasma disappearance kinetics of low-molecular
3 weight flavonoid metabolites and other dietary polyphenols should follow a similar pattern.
4
5

6 The site for the intravenous PMM mix injection was the dorsal penis vein, which is easily
7 accessible through a thin epidermal layer and is afferent to the general circulation.⁴³ Minimally
8 manipulating an anaesthetized animal is seen as a crucial factor because any inflammation
9 mediator that is locally released following surgical trauma might affect the permeability of the
10 blood-brain barrier and/or the basal functions that determine distribution and excretion (i.e.,
11 membrane transporters) or metabolism (i.e., enzymes).
12
13
14
15
16
17
18
19

20 The time points chosen were seemed appropriate for the time course, both in terms of the
21 injected molecules' disappearance from the blood and their appearance in the organs and urine.
22 Under the chosen conditions, the animals were under minimal duration of anesthesia and
23 physical stress. Overall, this experiment is a refinement of previous protocols that we have
24 used.^{40,44}
25
26
27
28
29
30
31
32

33 **The presence of basal levels of the mixture components in control animals**

34 Many of the mixture components arise not only from the microbial metabolism of dietary
35 polyphenols but also from mammalian catabolism of endogenous substrates.³⁶ Thus, some were
36 expected to occur in the rat organs and fluids at basal levels in control animals. Compounds that
37 are exclusively products of microbial metabolism are the urolithins, ferulic acid, gallic acid,
38 coumaric acid and protocatechuic acid.³⁶ However, these might also be expected to occur in
39 control animals as they are markers of normal nutritional status (e.g., ferulic acid). The basal
40 levels of the 23 component mixture were very low in general. Exceptions are mentioned case by
41 case below.
42
43
44
45
46
47
48
49
50
51
52
53
54

55 **Tracing polyphenol metabolites in blood**

1
2 The blood of control rats contained 11 of the 23 injected molecules. Of these endogenous
3
4 compounds (Table 1), six were found at less than 1% of the amount occurring in the injected
5
6 mixture, and another 3 were found at less than 2%. The 2 outliers were *p*-coumaric acid (3.3%)
7
8 and phloroglucinol (17.8%). Thus, the selected mixture represented a valid challenge for a
9
10 pharmacokinetic test because of the low basal levels. This test seemed to reliably simulate what
11
12 happens in the organism after consumption of a standard berry fruit serving after a short “wash
13
14 out” period.
15
16

17
18 Most injected compounds disappeared following apparent pseudo-first order kinetics, as
19
20 shown in Figure 2. The calculated parameters, i.e., the extrapolated amounts at time zero (t_0) and
21
22 at steady state (t_∞), and the disappearance rate constant (k_e), which spanned from 2.73 to 0.38
23
24 min^{-1} ($t/2=0.25$ to 1.84 min) and pointed to a rapid process, are shown in Table 1. The recovery
25
26 in blood extrapolated at t_0 was within 8.5 and 43.5% for 17 of the 23 metabolites (Table 1),
27
28 showing that a major fraction of these compounds was not easily extractable from the whole
29
30 blood. The formation of complexes with the serum proteins therefore still left a significant
31
32 concentration of the unbound form; for the majority of the studied metabolites: this is expected
33
34 to be the active form, which undergoes metabolism and reaches the tissues. However, there were
35
36 a few exceptions.
37
38

39
40 Pyrogallol could never be detected, though its recovery and analysis posed no specific
41
42 issues; the mechanisms of its disappearance were unknown. Two compounds, pyrocatechol and
43
44 3,4-dihydroxyphenyl acetic acid, were detected only at 15 sec. Urolithin B was detected at very
45
46 low levels at both 15 sec and 2 min and rose thereafter by an order of magnitude. Phloroglucinol
47
48 occurred at a similar level during the entire experiment. Gallic and caffeic acid showed a very
49
50 modest recovery, indicating a very modest presence in the free form in the bloodstream.
51
52

53
54 With the few exceptions mentioned above, steady state values were attained even earlier
55
56 than 5 min; therefore, the duration of the experiment was appropriate. In most cases, the
57
58 compounds were essentially disappeared at 15 min.
59
60

1
2 The calculated amounts at t_0 and at any time point were always lower than the sum of the
3 injected dose and any endogenous compounds in the control samples. The calculated t_0 amounts
4 ranged from 43.5% (*m*-coumaric acid) to 1% (gallic acid) and were expressed as a percentage of
5 the injected amounts as shown in Table 1. This result is not surprising given the fast uptake and
6 excretion of one precursor of these metabolites, i.e., the fruit pigment cyanidin 3-glucoside.⁴⁰
7
8
9
10
11
12

13 An explanation for the loss of part of the injected dose and for the extremely rapid blood
14 disappearance may be quick uptake and distribution to the organs. Only some major organs were
15 collected, so a cumulative recovery assessment could not be provided. The compounds also
16 might have been distributed in the large tissues, such as adipose, vascular endothelium or
17 connective tissues. The administered compounds could also have been metabolically
18 transformed, but these were not analyzed in this experiment. This is the first attempt at a
19 simultaneous kinetic characterization of 23 polyphenol metabolites.
20
21
22
23
24
25
26
27
28
29
30

31 **Polyphenol microbial metabolites in the brain**

32
33 The main findings concerned the detection of PMMs in the brain at times when their
34 blood concentrations were vanishingly low (Figure 3). Each plot shows 2 curves representing the
35 total amounts measured in whole brain (dotted line) and those corrected for residual blood
36 (continuous line), as recommended.⁴⁵⁻⁴⁸ Indeed, the brains were not perfused to wash out the
37 blood because this procedure would have altered the chemical equilibrium between the vascular
38 compartment and the whole brain.
39
40
41
42
43
44
45

46 Thirteen of the 23 compounds in the mixture were found in the control brains as
47 endogenous metabolites (Table 2). Disregarding the observation time, 10 of the 23 compounds
48 were found in the brain at a significantly increased amount with respect to the controls (Table 2).
49 In most cases, their appearance in the brain was biphasic, with an early wave at 2 min, as
50 observed with 4 compounds and a second wave starting at 5 min. Nine compounds were detected
51 at 15 min. Gallic acid, which was the most abundant compound in the mixture (Figure 2), best
52
53
54
55
56
57
58
59
60

1
2 showed this bi-phasic accumulation trend. It is noteworthy that the gallic acid accumulated in the
3 brain at approximately 11 times the basal concentration measured in the controls at as early as 15
4 sec. Similarly, caffeic acid had two peaks at 2 and 15 min and accumulated by approximately 20
5 and 34 times, respectively (Table 2).
6
7
8
9

10
11 Importantly, two compounds (*trans*-isoferulic acid and *o*-coumaric acid) were not
12 detected in the brain as endogenous metabolites nor did they ever appear in the brain despite
13 their relatively low plasma elimination rate constant. This proved that the blood-brain barrier was
14 intact during the experiment. Endogenous compounds protocatechuic acid and phloroglucinol
15 never had an increased concentration in the brain.
16
17
18
19
20

21
22 The compounds that accumulated in the brain should be active because, in all cases
23 except urolithin B, they were endogenous metabolites already present in the control brains. Their
24 most predictable effect is unbalancing a pre-existing metabolic equilibrium. Ferulic acid, which
25 accumulated by 15 sec in the experiment, is reportedly able to reduce oxidative damage and
26 amyloid pathology in Alzheimer's disease.⁴⁹ Some PMMs are identical to neurotransmitter
27 metabolites. For instance, vanillic acid and homovanillic acid are catecholamine catabolites that
28 are found in the brain and in cerebrospinal fluid.⁵⁰ Moreover, 3,4-dihydroxyphenyl acetic acid
29 (DOPAC) is a neuronal metabolite of dopamine (via monoamine oxidase) and a direct precursor
30 of homovanillic acid. DOPAC is involved in dopamine catabolism in the pathogenesis of
31 Parkinson's disease.^{51,52} Another mechanism of action involves the enzyme kinetics. Caffeic
32 acid, for example, may bind to tyrosine ammonia lyase, thus altering serotonin homeostasis⁵³, or
33 it may inhibit acetylcholinesterase and butyrylcholinesterase activities, thereby preventing
34 oxidative stress-induced neurodegeneration.⁵⁴ Gallic acid has also been shown to act on key
35 enzyme processes in the brain^{26,55} and can accumulate in the brain after repeated doses.⁵⁶
36
37
38
39
40
41
42
43
44
45
46
47
48
49
50
51
52
53
54

55 **Polyphenol microbial metabolites in other organs**

56
57
58
59
60

1
2 The analyses of PMMs in the heart, the liver, the kidneys, and the urine are shown in
3
4 Figure 5 (and in supplementary Figures 1-4). Eighteen of the 22 compounds attained very high
5
6 concentrations in the urine (Figure 4, supplementary Figure 1). The exceptions were
7
8 pyrocatechol and urolithin B, which were not found in the urine; urolithin A, the concentration of
9
10 which slightly increased; and phloroglucinol, which did not change with respect to the value
11
12 found in control animals. Thus, most PMMs can be defined as nephrotropic. Indeed, 20 of the 22
13
14 compounds dramatically increased their parenchymal concentrations from basal values
15
16 approximately zero up to at least 2 orders of magnitude higher (Figure 4, Supplementary Figure
17
18 2). Due to this extraordinary speed, the apparent rate of accumulation in the kidneys could not be
19
20 calculated. Accumulation lasted no more than 2 min and often less, after which time the
21
22 concentration started to decay. In contrast, the appearance in the urine was slower, with a lag
23
24 phase lasting 15 sec before the onset of urinary excretion in most cases (20 out of 22).
25
26
27

28
29 Only a few compounds were found to be hepato- rather than nephrotropic, including
30
31 phloroglucinol, urolithin A and 4-hydroxybenzoic acid (Figure 4, Supplementary Figure 3). One
32
33 compound (sinapic acid) could not be detected in the liver for 3 consecutive sampling times even
34
35 though it was present in the blood. Therefore, the amount of blood retained in the organ was
36
37 negligible. Similarly, the basal levels of 3-(3-hydroxyphenyl) propanoic acid did not change in
38
39 the liver, despite its presence in the blood at a 2 orders of magnitude higher concentration.
40
41

42
43 Only 3 compounds were found to be cardiotropic, i.e., vanillic acid and both urolithins
44
45 (Figure 4, Supplementary Figure 4). They accumulated to a maximum concentration at 15 sec,
46
47 after which vanillic acid and urolithin A disappeared, whereas urolithin B started to increase
48
49 again at 5 min. Three compounds (phloroglucinol, 3,4-dihydroxyphenyl acetic acid and
50
51 pyrocatechol), though present in the blood, could not be detected in the heart, demonstrating
52
53 negligible blood contamination during the tissue extraction.
54

55
56 The low apparent hepatotropism of most PMMs leads to the conclusion that PMMs
57
58 absorbed from the colon under normal conditions undergo limited first-pass metabolism into the
59
60

1
2 liver so that they can be distributed to the other organs readily after absorption. They are then
3
4 eliminated in the urine after prior accumulation in the kidneys. Only one compound, 4-
5
6 hydroxybenzoic acid, appeared in the urine at 15 sec, likely by glomerular filtration.
7
8
9

10 11 **The case of urolithins**

12
13 Urolithins, with MWs ranging from 212-228 Daltons, are the largest PMM molecules
14
15 (Figure 1) and arise from the microbial metabolism of ellagitannins and ellagic acid conjugates.⁵⁷
16
17 Urolithin A differs from urolithin B by one additional hydroxyl group. No other mammalian
18
19 enzyme pathway is known to produce such end products from more complex precursors. No
20
21 urolithins were detected in the organs of the control animals. Urolithin A and B were
22
23 administered at different doses, i.e., 55 and 25 nmol, respectively. Nevertheless, their
24
25 concentrations in the blood differed by an order of magnitude at 15 sec (332.6 and 20.3
26
27 pmol/mL, respectively). Thus, urolithin B was sequestered (and/or metabolized) more rapidly
28
29 than urolithin A in the tissues.
30
31
32

33
34 Both urolithins displayed a unique blood concentration pattern, tissue distribution and
35
36 urinary excretion (Figure 6). Indeed, though urolithin A had a normal mono-exponential
37
38 disappearance from the blood, urolithin B increased in the blood after 2 min from a very low
39
40 level (see above). Therefore, it seems that urolithin A underwent a very unusual and
41
42 mechanistically unexplained de-hydroxylation reaction. Only bacterial de-hydroxylases are
43
44 known.
45
46

47
48 Both urolithins were negligibly excreted in the urine, which is unique among the PMMs.
49
50 They were presumably excreted as glucuronyl derivatives¹¹, which were not analyzed. Rapid
51
52 uptake of urolithin A into the kidneys was observed, whereas urolithin B appeared in the kidneys
53
54 only after 2 min when it was also present in the blood. The liver slowly absorbed urolithin A up
55
56 to 5 min, but no urolithin B could be detected in the liver at 15 min. This shows that the liver
57
58 could neither convert urolithin A to urolithin B nor take the latter up from the blood.
59
60

1
2 The organ that displayed the highest uptake capacity was the heart, where both urolithin
3
4 A and B were found at 15 sec, at approximately 300 and 2000 pmol/g, respectively. It seems
5
6 likely that the following occurred: i) urolithin A was rapidly (by 15 sec) taken up into the heart
7
8 and de-hydroxylated to urolithin B by an unknown enzyme(s) or in the heart both urolithins
9
10 could be converted to other compounds that were not followed; ii) urolithin B was released from
11
12 the heart into the blood, where it increased from 2 to 5 min; iii) from 5 min on, the urolithin B
13
14 tissue concentrations increased at higher levels than in the blood.
15
16

17
18 Urolithins have a demonstrated activity of reducing protein glycation⁵⁸ in neuronal cells
19
20 and possibly even in the heart. In fact, the advanced glycation end products, derived from the
21
22 glycation reaction in the heart and vascular tissue, are responsible for a reduction in artery and
23
24 heart elasticity and have a role in the progression of cardiovascular complications associated
25
26 with diabetes.⁵⁹
27
28
29
30

31 **Is the brain a target of polyphenol metabolites?**

32
33 The findings would suggest so. To our knowledge, ours is the first report showing that no
34
35 less than 13 PMMs were found at basal levels in control brains. As a consequence, the possibility
36
37 of their passage through the blood-brain barrier must be assumed a priori to ensure exchange
38
39 with the blood compartment.
40
41

42 Ten of the 23 compounds were found to be significantly increased after injecting the
43
44 PMM mixture into the animals. Some of the compounds could have a strictly dietary origin, such
45
46 as gallic acid or *trans*-ferulic acid, whereas others could be catabolites of endogenous
47
48 compounds, such as catecholamine, dopamine, amino acids and others. Figure 6 highlights the
49
50 gallic acid concentration attained in all of the organs considered at various times. The brain
51
52 clearly had the capacity to extract gallic acid from the blood, even higher than that of the liver.
53
54 On the other hand the tightness of the blood-brain barrier is demonstrated by its absolute
55
56 impermeability to *o*-coumaric acid, in spite of its higher hydrophobic index (logP: 2.45, while
57
58
59
60

1
2 gallic acid logP is 0.91). However, both gallic acid and *o*-coumaric acid have similar
3
4 concentrations in the blood.
5

6
7 These data suggest that there are further health implications for the brain-gut microbiota
8
9 axis⁵⁴ and highlight how PMMs could be players in the putative connections/messengers
10
11 between the microbiota and brain. These bidirectional interactions have already been explored
12
13 and highlight the importance of the gut microbiota for brain development and behavior⁵⁵, as well
14
15 as for their influence on anxiety and depression.⁵⁶ Because PMMs are the product of dietary
16
17 polyphenol intake and metabolism by the gut microbiota, it is reasonable to assume that these
18
19 compounds might be able to regulate human health and disease states.⁵⁷ Moreover, the role of
20
21 gut microbiota in the alteration of mammalian blood metabolite levels has already been
22
23 demonstrated through combining data from germ-free and conventional animals.⁵⁸ Their action
24
25 may be related to oxidative stress control or, more likely, to the modulation of biochemical and
26
27 physiological processes by changing the level of endogenous compounds in the brain and other
28
29 organs, as observed in the present study.
30
31
32
33
34
35
36
37

38 **Methods**

39 **Chemicals**

40
41
42 Phloroglucinol, pyrogallol, gallic acid, protocatechuic acid, 3,4-dihydroxyphenylacetic acid, 4-
43
44 hydroxyhippuric acid, 4-hydroxybenzoic acid, pyrocatechol, caffeic acid, vanillic acid, 3-
45
46 hydroxyphenylacetic acid, homovanillic acid, 3-(4-hydroxyphenyl) propanoic acid, 3-(3-
47
48 hydroxyphenyl)propanoic acid, hydroferulic acid, *trans*-ferulic acid, *trans*-isoferulic acid, sinapic
49
50 acid, *m*-coumaric acid, *o*-coumaric acid and *p*-coumaric were purchased from Sigma-Aldrich
51
52 (Saint Luis, Milan, Italy). The isotopically labeled compounds, butyric acid-d7 and cinnamic
53
54 acid-d5, were used as internal standards and purchased from C/D/N Isotopes Inc. (Quebec,
55
56
57
58
59
60

1
2 Canada). Urolithin A and urolithin B were synthesized following a published protocol³⁷ and were
3
4 characterized by NMR for structure confirmation and purity. LC/MS formic acid, Chromasolv
5
6 LC/MS methanol and acetonitrile were purchased from Sigma-Aldrich.
7
8

9 10 11 **Study design**

12
13 The experiment was designed as a multi-component pharmacokinetic study with
14
15 quantitative analysis of organs (liver, kidney, heart and brain) and biofluids (blood and urine).
16
17 The aim was to explore the fate of a polyphenol microbial metabolite mixture in a mammalian
18
19 system after entering the bloodstream. The experiment was divided into four time points: 15 sec,
20
21 2 min, 5 min and 15 min. Each time point was represented by 4 replicates. Four rats received the
22
23 polyphenol microbial metabolite dose, whereas one blank rat served as a control without the
24
25 polyphenol microbial metabolite dose, whereas one blank rat served as a control without the
26
27 polyphenol metabolite mixture but with an identical experimental procedure. The control animals
28
29 that were not injected with the polyphenol microbial metabolites could also be considered as 4
30
31 control replicates to compare at each time point. The short kinetics trial time did not have an
32
33 effect on the concentration and endogenous metabolite pattern in the control animals.
34

35
36 Twenty male rats (*Rattus norvegicus*, Wistar, Harlan Italy S.r.l.) were bred in the animal
37
38 facility at the University of Trieste. The experiment was approved by the bioethical committee of
39
40 the University of Trieste (doc. 865PAS12). Animal experiments were carried out in compliance
41
42 with the provisions of the European Community Council Directive.³⁹ Rats (n=20) at the same
43
44 age (12 weeks) and weight (288 ± 20 g) were maintained in temperature-controlled rooms at 22-
45
46 24 °C, 50-60% humidity and 12 hour light/dark cycles. They were fed until the night before the
47
48 experiment with standard laboratory chow.
49

50
51 During the *in vivo* experiment, the rats were kept under quiet conditions. The cages were
52
53 covered with cloth so that they were either sleeping before the anesthetic injections or were
54
55 awake but without apparent nervous behavior. After the anesthetic, the animals were left alone to
56
57 fall asleep in another covered cage. In all cases, the animals' conditions were monitored.
58
59
60

Polyphenol microbial metabolite mixture for intravenous (i.v.) administration

The polyphenol metabolite mixture comprised 23 low-molecular-weight (110-228 Daltons) metabolites, which were selected from the key polyphenol metabolism products by gut microbiota after the consumption of berry fruits. The metabolites and dosages were selected from reports in the literature. Each metabolite was dissolved in methanol and then mixed together according to the various selected amounts.

Treatment of rats

The rats were divided into two groups for a two-day experiment, with 2 pharmacokinetic time points per day. On the first day, time points 15 min and 2 min were chosen. Time points 5 min and 15 sec were performed on the second day. On the night before the experiment, the rats (10 animals) were starved overnight but were provided with water *ad libitum*. They were divided into 5 different cages for the experiment.

The rats were anaesthetized with intra-peritoneal administration of Tiletamine/Zolazepam (1:1, 25 mg/kg body weight) and Xylazine (10 mg/kg body weight). Each rat received an i.v. administration of 0.3 mL PBS (phosphate buffer solution) containing 2.7 μmol of polyphenol metabolite mixture, dissolved in 30 μL methanol. The blank control animals received an i.v. injection of 0.3 mL PBS with 30 μL methanol.

During anesthesia (10 min) the heart and ventilation rates were controlled. The rats were placed on their backs and the penis was extruded by sliding the prepuce downwards. With the use of a nipper, the glans penis was held at the tip. The dorsal penis vein was then visible and at exactly 10 min after anesthesia, 0.3 mL PBS with (treated) or without (blank) 2.7 μmol polyphenol metabolite mixture was injected using an insulin syringe. The injection site was then pressed for a few seconds and the glans was retracted to prevent bleeding.⁴³ One min before the rat was sacrificed, sodium heparin (0.1 mL, 500 IU) was injected again into the dorsal penis vein

1
2 as before. For time point 15 sec, the procedure was slightly different: after 10 min of anaesthesia,
3
4 0.2 mL PBS and 0.1 mL sodium heparin with (treated) or without (blank) 2.7 μmol polyphenol
5
6 metabolite mixture were simultaneously injected using an insulin syringe.
7

8
9 The rats were sacrificed by decapitation. Blood was drained from the body and urine was
10
11 collected through the urinary bladder with a syringe. The liver, kidney, heart and brain were
12
13 excised from the body, washed with Milli-Q water and immediately frozen in liquid nitrogen and
14
15 stored at $-80\text{ }^{\circ}\text{C}$.
16

17 18 19 20 **Organ collection and extract preparation**

21 22 23 *Extraction from blood and urine*

24
25 After sampling, an aliquot of blood (5 mL) was transferred into aqueous methanol (95:5,
26
27 v/v) at a ratio of 1:9 (v/v). The internal standard cinnamic acid- d_5 was dissolved in aqueous
28
29 methanol at a concentration of 100 mg/L for monitoring the extraction protocol. The sample was
30
31 shaken and extracted for 10 min. The extract was then centrifuged for 5 min at 3600 rpm at 4
32
33 $^{\circ}\text{C}$. The blood extract with the buoyant (non-cellular) portion was transferred to 50 mL dark
34
35 glass vessels and stored at $-80\text{ }^{\circ}\text{C}$.⁴⁴ The choice of extracting blood instead of plasma or serum
36
37 was made to eliminate the need for a centrifugation or blood-clotting step, during which
38
39 metabolic changes might have occurred *ex vivo*.⁶⁰
40
41
42
43

44
45 In the same way, an aliquot of urine was collected with a syringe and extracted with
46
47 aqueous methanol (95:5, v/v). The urine was weighed and the amount of solvent for extraction
48
49 was adjusted to a 1:9 ratio (w/v).
50

51 52 53 *Tissues*

54
55 Organs that were frozen in liquid nitrogen immediately after excision were stored at -80
56
57 $^{\circ}\text{C}$. Frozen liver, kidney, heart and brain were ground under cryogenic conditions using liquid
58
59
60

1
2 nitrogen with a CryoMill (Retsch, Germany) and a single 25-mm i.d. steel ball (30 sec. with a
3 frequency of 25/sec.). Tissue powder (1 g) was transferred (without thawing) into aqueous
4 methanol (5:95 v/v) at a ratio of 1:9 (w/v). The internal standard cinnamic acid-d₅ was dissolved
5 in the aqueous methanol at a concentration of 100 mg/L to monitor the extraction protocol. The
6 samples were extracted on an orbital shaker for 10 min, then centrifuged and decanted as
7 described for blood. The final volume was then adjusted to 10 mL to balance any possible
8 variation in the amount of water among the organs and stored at -80 °C.
9
10
11
12
13
14
15
16
17
18
19

20 **Solid Phase Extraction (SPE) purification and sample preparation**

21
22
23
24 The SPE protocol was applied following a procedure developed by Passamonti et al.⁶¹
25 with modifications as reported by Gasperotti et al.³⁸ The cartridges used for the SPE purification
26 were Biotage Isolute ENV+, 1 g (Uppsala Sweden).
27
28
29

30
31 The internal standard butyric acid-d₇ was dissolved in methanol/water (50:50 v/v) at a
32 concentration of 1 mg/L and added to the sample to monitor the quantitative recovery during
33 sample reconstitution.
34
35
36
37
38
39

40 **Targeted metabolomics analysis by UHPLC-ESI-MS/MS**

41
42 A targeted metabolomics analysis was performed with an ultra-performance liquid
43 chromatographic system (UHPLC) coupled to a tandem mass spectrometer. The system used was
44 an ACQUITY UPLC system coupled to a Xevo TQ triple quadrupole via an electrospray (ESI)
45 interface (Waters, Milford, Massachusetts, USA). The separation of the 23 target metabolites and
46 2 deuterated internal standards was performed with a Waters ACQUITY UPLC column and HSS
47 T3 (100 mm x 2.1 mm, 1.8 μm) equipped with a guard column. The injection volume was 10 μL.
48 Mobile phases of 0.1% formic acid in Milli-Q water (A) and 0.1% formic acid in acetonitrile (B)
49 were used. Chromatographic separation was performed using a gradient as follows: 0 min, 5% B;
50
51
52
53
54
55
56
57
58
59
60

1
2 0-3 min, 5%-20% B; 3-4.30 min, 20% B; 4.30-9 min, 20%-45% B; 9-11 min, 45%-100% B; 11-
3
4 14 min, 100% B; and 14.01-17 min, 5% B as equilibration time. For calibration, a standard
5
6 mixture of polyphenol metabolites was serially diluted in aqueous methanol (50:50) at a
7
8 concentration range of 0.01 µg/L - 100 mg/L. Quantitative data were processed with Targetlynx
9
10 software (Masslynx, Waters).

11
12
13 Details of the UHPLC-ESI-MS/MS method and quantification are described in Gasperotti et al.³⁸
14
15

16 17 18 **Blood residual subtraction for the brain samples**

19
20 The concentrations in the rat brain may be significantly influenced by the quantities of
21
22 metabolites in the residual blood. The correction was made by estimating the amount of
23
24 metabolites in the intravascular blood present in the brain, assuming that the volume of brain
25
26 blood is 47.7 µL/g.⁴⁶ We then subtracted this from the total amount found in the brain. The
27
28 results of this subtraction are presented in Figure 3, which also compares the data of the brain
29
30 before the blood residual subtraction.
31
32

33 34 35 **Calculation of pharmacokinetics parameters**

36
37 Pharmacokinetic parameters were calculated using GraphPad Prism 5 (GraphPad Software, La
38
39 Jolla, CA, USA) with a one-phase decay function suitable for i.v. injection experiments. The
40
41 extrapolated dose at time 0 (t_0) and the dose at steady state (t_∞) were obtained from the plotted
42
43 curves for each metabolite detected and quantified in the blood. The area under the curve (AUC
44
45 $_{0 \rightarrow 15 \text{ min.}}$) and the elimination rate (K_e) were also obtained from the plotted curves.
46
47
48
49
50

51 52 **Statistical analysis**

53
54 The quantitative data for the PMM levels are presented as the mean values \pm standard error ($n =$
55
56 4) for all of the graphs. The concentrations in the brain of the treated animals were compared
57
58 against those of the controls using Student's t-test, with P-values < 0.1 considered statistically
59
60

1
2 significant. Relaxing the significance threshold to 0.1 enabled us to identify the most potentially
3
4 relevant information and minimize type II error in these experiments while involving the least
5
6 number of animals, in compliance with the legislation regulating the use of animals in scientific
7
8 procedures.
9

10 11 12 13 14 15 **AUTHOR INFORMATION**

16 17 **Corresponding authors**

18
19 * To whom correspondence should be addressed:

20
21
22 Tel: +39 0461 615 140. Fax: +39 0461 615 200. E-mail: urska.vrhovsek@fmach.it

23
24
25 Tel: +390405588747. E-mail: spassamonti@units.it

26 27 28 **Funding**

29
30
31 This research was supported by the GMPF International PhD program (to M.G.), the ADP 2012
32
33 Programme funded by the Autonomous Province of Trento, Italy, Agrotur Project (to S.P)
34
35 funded by the Cross-border Cooperation Programme Italy-Slovenia 2007-2013 (European
36
37 Regional Development Fund and National funds).
38

39 40 41 42 **SUPPORTING INFORMATION AVAILABLE**

43
44 A file containing all the supplementary figures and information is presented as supplementary
45
46 online data. This information is available free of charge via the Internet at <http://pubs.acs.org>.
47
48
49
50
51
52
53
54
55
56
57
58
59
60

References

- 1 (1) (2003) Diet, nutrition and the prevention of chronic diseases. World Health Organ Tech Rep
2 Ser 916, i–viii, 1–149, backcover.
- 3 (2) Ovaskainen, M.-L., Törrönen, R., Koponen, J. M., Sinkko, H., Hellström, J., Reinivuo, H.,
4 and Mattila, P. (2008) Dietary Intake and Major Food Sources of Polyphenols in Finnish Adults.
5 J. Nutr. 138, 562–566.
- 6 (3) Del Rio, D., Borges, G., and Crozier, A. (2010) Berry flavonoids and phenolics:
7 bioavailability and evidence of protective effects. British Journal of Nutrition 104, S67–S90.
- 8 (4) Cardona, F., Andrés-Lacueva, C., Tulipani, S., Tinahones, F. J., and Queipo-Ortuño, M. I.
9 (2013) Benefits of polyphenols on gut microbiota and implications in human health. The Journal
10 of Nutritional Biochemistry 24, 1415–1422.
- 11 (5) Moco, S., Martin, F.-P. J., and Rezzi, S. (2012) Metabolomics View on Gut Microbiome
12 Modulation by Polyphenol-rich Foods. J. Proteome Res. 11, 4781–4790.
- 13 (6) Li, M., Wang, B., Zhang, M., Rantalainen, M., Wang, S., Zhou, H., Zhang, Y., Shen, J.,
14 Pang, X., Zhang, M., Wei, H., Chen, Y., Lu, H., Zuo, J., Su, M., Qiu, Y., Jia, W., Xiao, C.,
15 Smith, L. M., Yang, S., Holmes, E., Tang, H., Zhao, G., Nicholson, J. K., Li, L., and Zhao, L.
16 (2008) Symbiotic gut microbes modulate human metabolic phenotypes. PNAS 105, 2117–2122.
- 17 (7) Clayton, T. A., Baker, D., Lindon, J. C., Everett, J. R., and Nicholson, J. K. (2009)
18 Pharmacometabonomic identification of a significant host-microbiome metabolic interaction
19 affecting human drug metabolism. PNAS 106, 14728–14733.
- 20 (8) Clemente, J. C., Ursell, L. K., Parfrey, L. W., and Knight, R. (2012) The Impact of the Gut
21 Microbiota on Human Health: An Integrative View. Cell 148, 1258–1270.
- 22 (9) Selma, M. V., Espín, J. C., and Tomás-Barberán, F. A. (2009) Interaction between Phenolics
23 and Gut Microbiota: Role in Human Health. J. Agric. Food Chem. 57, 6485–6501.
- 24 (10) Rios, L. Y., Gonthier, M.-P., Rémésy, C., Mila, I., Lapiere, C., Lazarus, S. A., Williamson,
25 G., and Scalbert, A. (2003) Chocolate intake increases urinary excretion of polyphenol-derived
26 phenolic acids in healthy human subjects. Am J Clin Nutr 77, 912–918.
- 27 (11) Crozier, A., Del Rio, D., and Clifford, M. N. (2010) Bioavailability of dietary flavonoids
28 and phenolic compounds. Molecular Aspects of Medicine 31, 446–467.
- 29 (12) Monagas, M., Urpi-Sarda, M., Sánchez-Patán, F., Llorach, R., Garrido, I., Gómez-
30 Cordovés, C., Andres-Lacueva, C., and Bartolomé, B. (2010) Insights into the metabolism and
31 microbial biotransformation of dietary flavan-3-ols and the bioactivity of their metabolites. Food
32 Funct 1, 233–253.
- 33 (13) Rechner, A. R., Smith, M. A., Kuhnle, G., Gibson, G. R., Debnam, E. S., Srai, S. K. S.,
34 Moore, K. P., and Rice-Evans, C. A. (2004) Colonic metabolism of dietary polyphenols:
35 influence of structure on microbial fermentation products. Free Radical Biology and Medicine
36 36, 212–225.
- 37 (14) Stalmach, A., Edwards, C. A., Wightman, J. D., and Crozier, A. (2012) Gastrointestinal
38 stability and bioavailability of (poly)phenolic compounds following ingestion of Concord grape
39 juice by humans. Molecular Nutrition & Food Research 56, 497–509.
- 40 (15) Stalmach, A., Edwards, C. A., Wightman, J. D., and Crozier, A. (2013) Colonic catabolism
41 of dietary phenolic and polyphenolic compounds from Concord grape juice. Food Funct. 4, 52–
42 62.
- 43 (16) Dall’Asta, M., Calani, L., Tedeschi, M., Jechiu, L., Brighenti, F., and Del Rio, D. (2012)
44 Identification of microbial metabolites derived from in vitro fecal fermentation of different
45 polyphenolic food sources. Nutrition 28, 197–203.
- 46 (17) Gonthier, M. P., Donovan, J. L., Texier, O., Felgines, C., Remesy, C., and Scalbert, A.
47 (2003) Metabolism of dietary procyanidins in rats. Free Radic. Biol. Med. 35, 837–844.

- 1
2
3 (18) Boto-Ordóñez, M., Rothwell, J. A., Andres-Lacueva, C., Manach, C., Scalbert, A., and
4 Urpi-Sarda, M. (2013) Prediction of the wine polyphenol metabolic space: An application of the
5 Phenol-Explorer database. *Molecular Nutrition & Food Research* n/a–n/a.
- 6 (19) Hanhineva, K., Torronen, R., Bondia-Pons, I., Pekkinen, J., Kolehmainen, M., Mykkanen,
7 H., and Poutanen, K. (2010) Impact of Dietary Polyphenols on Carbohydrate Metabolism. *Int J*
8 *Mol Sci* 11, 1365–1402.
- 9 (20) Williamson, G., and Clifford, M. N. (2010) Colonic metabolites of berry polyphenols: the
10 missing link to biological activity? *Br. J. Nutr.* 104, S48–S66.
- 11 (21) Scalbert, A., Manach, C., Morand, C., Rémésy, C., and Jiménez, L. (2005) Dietary
12 polyphenols and the prevention of diseases. *Crit Rev Food Sci Nutr* 45, 287–306.
- 13 (22) Del Rio, D., Rodriguez-Mateos, A., Spencer, J. P. E., Tognolini, M., Borges, G., and
14 Crozier, A. (2013) Dietary (Poly)phenolics in Human Health: Structures, Bioavailability, and
15 Evidence of Protective Effects Against Chronic Diseases. *Antioxidants & Redox Signaling* 18,
16 1818–1892.
- 17 (23) Jones, Q. R. D., Warford, J., Rupasinghe, H. P. V., and Robertson, G. S. (2012) Target-
18 based selection of flavonoids for neurodegenerative disorders. *Trends in Pharmacological*
19 *Sciences* 33, 602–610.
- 20 (24) Singh, M., Arseneault, M., Sanderson, T., Murthy, V., and Ramassamy, C. (2008)
21 Challenges for research on polyphenols from foods in Alzheimer's disease: bioavailability,
22 metabolism, and cellular and molecular mechanisms. *J. Agric. Food Chem.* 56, 4855–4873.
- 23 (25) Spencer, J. P. E. (2010) The impact of fruit flavonoids on memory and cognition. *Br. J.*
24 *Nutr.* 104 Suppl 3, S40–47.
- 25 (26) Corona, G., Vauzour, D., Hercelin, J., Williams, C. M., and Spencer, J. P. E. (2013)
26 Phenolic Acid Intake, Delivered Via Moderate Champagne Wine Consumption, Improves
27 Spatial Working Memory Via the Modulation of Hippocampal and Cortical Protein
28 Expression/Activation. *Antioxidants & Redox Signaling* 19, 1676–1689.
- 29 (27) Dixon, R. A., and Pasinetti, G. M. (2010) Flavonoids and Isoflavonoids: From Plant
30 Biology to Agriculture and Neuroscience. *Plant Physiol.* 154, 453–457.
- 31 (28) Joseph, J., Cole, G., Head, E., and Ingram, D. (2009) Nutrition, brain aging, and
32 neurodegeneration. *J. Neurosci.* 29, 12795–12801.
- 33 (29) Ho, L., Chen, L. H., Wang, J., Zhao, W., Talcott, S. T., Ono, K., Teplow, D., Humala, N.,
34 Cheng, A., Percival, S. S., Ferruzzi, M., Janle, E., Dickstein, D. L., and Pasinetti, G. M. (2009)
35 Heterogeneity in red wine polyphenolic contents differentially influences Alzheimer's disease-
36 type neuropathology and cognitive deterioration. *J. Alzheimers Dis.* 16, 59–72.
- 37 (30) Diaz Heijtz, R., Wang, S., Anuar, F., Qian, Y., Bjorkholm, B., Samuelsson, A., Hibberd, M.
38 L., Forssberg, H., and Pettersson, S. (2011) Normal gut microbiota modulates brain development
39 and behavior. *Proceedings of the National Academy of Sciences of the United States of America*
40 108.
- 41 (31) Al-Asmakh, M., Anuar, F., Zadjali, F., Rafter, J., and Pettersson, S. (2012) Gut microbial
42 communities modulating brain development and function. *Gut Microbes* 3, 366–373.
- 43 (32) Foster, J. A., and McVey Neufeld, K.-A. (2013) Gut–brain axis: how the microbiome
44 influences anxiety and depression. *Trends in Neurosciences* 36, 305–312.
- 45 (33) Mulle, J. G., Sharp, W. G., and Cubells, J. F. (2013) The gut microbiome: a new frontier in
46 autism research. *Curr Psychiatry Rep* 15, 337.
- 47 (34) Van Duynhoven, J., Vaughan, E. E., Jacobs, D. M., A. Kemperman, R., van Velzen, E. J. J.,
48 Gross, G., Roger, L. C., Possemiers, S., Smilde, A. K., Dore, J., Westerhuis, J. A., and Van de
49 Wiele, T. (2010) Colloquium Paper: Metabolic fate of polyphenols in the human superorganism.
50 *Proceedings of the National Academy of Sciences* 108, 4531–4538.
- 51 (35) Nurmi, T., Mursu, J., Heinonen, M., Nurmi, A., Hiltunen, R., and Voutilainen, S. (2009)
52 Metabolism of Berry Anthocyanins to Phenolic Acids in Humans. *J. Agric. Food Chem.* 57,
53 2274–2281.
54
55
56
57
58
59
60

- 1
2 (36) Wishart, D. S., Jewison, T., Guo, A. C., Wilson, M., Knox, C., Liu, Y., Djoumbou, Y.,
3 Mandal, R., Aziat, F., Dong, E., Bouatra, S., Sinelnikov, I., Arndt, D., Xia, J., Liu, P., Yallou, F.,
4 Bjorn Dahl, T., Perez-Pineiro, R., Eisner, R., Allen, F., Neveu, V., Greiner, R., and Scalbert, A.
5 (2013) HMDB 3.0--The Human Metabolome Database in 2013. *Nucleic Acids Res.* 41, D801–
6 807.
- 7
8 (37) Bialonska, D., Kasimsetty, S. G., Khan, S. I., and Ferreira, D. (2009) Urolithins, Intestinal
9 Microbial Metabolites of Pomegranate Ellagitannins, Exhibit Potent Antioxidant Activity in a
10 Cell-Based Assay. *J. Agric. Food Chem.* 57, 10181–10186.
- 11 (38) Gasperotti, M., Masuero, D., Guella, G., Mattivi, F., and Vrhovsek, U. (2014) Development
12 of a targeted method for twenty-three metabolites related to polyphenol gut microbial
13 metabolism in biological samples, using SPE and UHPLC–ESI-MS/MS. *Talanta* 128, 221–230.
- 14 (39) EUROPEAN PARLIAMENT. (2010) DIRECTIVE 2010/63/EU OF THE EUROPEAN
15 PARLIAMENT AND OF THE COUNCIL of 22 September 2010 on the protection of animals
16 used for scientific purposes.
- 17
18 (40) Vanzo, A., Vrhovsek, U., Tramer, F., Mattivi, F., and Passamonti, S. (2011) Exceptionally
19 Fast Uptake and Metabolism of Cyanidin 3-Glucoside by Rat Kidneys and Liver. *J. Nat. Prod.*
20 74, 1049–1054.
- 21 (41) Lee, H. B., and Blaufox, M. D. (1985) Blood Volume in the Rat. *J Nucl Med* 26, 72–76.
- 22 (42) Manach, C., Williamson, G., Morand, C., Scalbert, A., and Rémésy, C. (2005)
23 Bioavailability and bioefficacy of polyphenols in humans. I. Review of 97 bioavailability
24 studies. *Am J Clin Nutr* 81, 230S–242S.
- 25 (43) Waynforth, H. B., and Flecknell, P. A. (1992) Experimental and surgical technique in the
26 rat. Academic Press, New York.
- 27
28 (44) Vanzo, A., Scholz, M., Gasperotti, M., Tramer, F., Passamonti, S., Vrhovsek, U., and
29 Mattivi, F. (2013) Metabonomic investigation of rat tissues following intravenous
30 administration of cyanidin 3-glucoside at a physiologically relevant dose. *Metabolomics* 9, 88–
31 100.
- 32 (45) Schaffer, S., and Halliwell, B. (2012) Do polyphenols enter the brain and does it matter?
33 Some theoretical and practical considerations. *Genes Nutr* 7, 99–109.
- 34 (46) Friden, M., Ljungqvist, H., Middleton, B., Bredberg, U., and Hammarlund-Udenaes, M.
35 (2010) Improved measurement of drug exposure in the brain using drug-specific correction for
36 residual blood. *J. Cereb. Blood Flow Metab.* 30, 150–161.
- 37 (47) Vauzour, D. (2012) Dietary Polyphenols as Modulators of Brain Functions: Biological
38 Actions and Molecular Mechanisms Underpinning Their Beneficial Effects. *Oxidative Medicine*
39 *and Cellular Longevity* 2012.
- 40 (48) Boer, V. C. J. de, Dihal, A. A., Woude, H. van der, Arts, I. C. W., Wolfram, S., Alink, G.
41 M., Rietjens, I. M. C. M., Keijer, J., and Hollman, P. C. H. (2005) Tissue Distribution of
42 Quercetin in Rats and Pigs. *J. Nutr.* 135, 1718–1725.
- 43 (49) Mancuso, C., Scapagini, G., Currò, D., Giuffrida Stella, A. M., De Marco, C., Butterfield,
44 D. A., and Calabrese, V. (2007) Mitochondrial dysfunction, free radical generation and cellular
45 stress response in neurodegenerative disorders. *Front. Biosci.* 12, 1107–1123.
- 46 (50) De Jong, A. P., Kok, R. M., Cramers, C. A., and Wadman, S. K. (1986) Determination of
47 acidic catecholamine metabolites in plasma and cerebrospinal fluid using gas chromatography-
48 negative-ion mass spectrometry. *J. Chromatogr.* 382, 19–30.
- 49 (51) Florang, V. R., Rees, J. N., Brogden, N. K., Anderson, D. G., Hurley, T. D., and Doorn, J.
50 A. (2007) Inhibition of the oxidative metabolism of 3,4-dihydroxyphenylacetaldehyde, a reactive
51 intermediate of dopamine metabolism, by 4-hydroxy-2-nonenal. *Neurotoxicology* 28, 76–82.
- 52 (52) Fornstedt, B., Pileblad, E., and Carlsson, A. (1990) In Vivo Autoxidation of Dopamine in
53 Guinea Pig Striatum Increases with Age. *Journal of Neurochemistry* 55, 655–659.
- 54 (53) Fernstrom, J. D. (2013) Large neutral amino acids: dietary effects on brain neurochemistry
55 and function. *Amino Acids* 45, 419–430.
- 56
57
58
59
60

- 1
2 (54) Oboh, G., Agunloye, O. M., Akinyemi, A. J., Ademiluyi, A. O., and Adefegha, S. A. (2013)
3 Comparative study on the inhibitory effect of caffeic and chlorogenic acids on key enzymes
4 linked to Alzheimer's disease and some pro-oxidant induced oxidative stress in rats' brain-in
5 vitro. *Neurochem. Res.* 38, 413–419.
6
7 (55) Kade, I. J., and Rocha, J. B. T. (2013) Gallic Acid Modulates Cerebral Oxidative Stress
8 Conditions and Activities of Enzyme-Dependent Signaling Systems in Streptozotocin-Treated
9 Rats. *Neurochem Res* 38, 761–771.
10 (56) Ferruzzi, M. G., Lobo, J. K., Janle, E. M., Cooper, B., Simon, J. E., Wu, Q.-L., Welch, C.,
11 Ho, L., Weaver, C., and Pasinetti, G. M. (2009) Bioavailability of gallic acid and catechins from
12 grape seed polyphenol extract is improved by repeated dosing in rats: implications for treatment
13 in Alzheimer's disease. *J. Alzheimers Dis.* 18, 113–124.
14 (57) Cerdá, B., Espín, J. C., Parra, S., Martínez, P., and Tomás-Barberán, F. A. (2004) The
15 potent in vitro antioxidant ellagitannins from pomegranate juice are metabolised into
16 bioavailable but poor antioxidant hydroxy-6H-dibenzopyran-6- one derivatives by the colonic
17 microflora of healthy humans. *Eur J Nutr* 43, 205–220.
18 (58) Verzelloni, E., Pellacani, C., Tagliazucchi, D., Tagliaferri, S., Calani, L., Costa, L. G.,
19 Brighenti, F., Borges, G., Crozier, A., Conte, A., and Del Rio, D. (2011) Antiglycative and
20 neuroprotective activity of colon-derived polyphenol catabolites. *Mol Nutr Food Res* 55 Suppl 1,
21 S35–43.
22 (59) Orasanu, G., and Plutzky, J. (2009) The Pathologic Continuum of Diabetic Vascular
23 Disease. *Journal of the American College of Cardiology* 53, S35–S42.
24 (60) Gika, H., and Theodoridis, G. (2011) Sample preparation prior to the LC-MS-based
25 metabolomics/metabonomics of blood-derived samples. *Bioanalysis* 3, 1647–1661.
26 (61) Passamonti, S., Vrhovsek, U., Vanzo, A., and Mattivi, F. (2003) The stomach as a site for
27 anthocyanins absorption from food. *FEBS Letters* 544, 210–213.
28
29
30
31
32
33
34
35
36
37
38
39
40
41
42
43
44
45
46
47
48
49
50
51
52
53
54
55
56
57
58
59
60

Figure 1: Mixture of polyphenol microbial metabolites used for the intravenous injection, their chemical formula, molecular weight (MW), dose and relative percentage on the total amount injected, 2690 nmol (dosage:168.13 μ M)





Figure 2: Kinetics of distribution of polyphenol microbial metabolites in blood.

Figure 3: Kinetics of distribution of polyphenol microbial metabolites in brain. Dotted lines are referred to the amount found in the whole brain, while continuous lines are the data after the subtraction of the blood residual in the brain. Data with the asterisk are significantly increased from the control samples ($p < 0.01^{***}$, $p < 0.05^{**}$, $p < 0.1^*$).




Figure 4: Kinetics of distribution of polyphenol microbial metabolites in liver, kidney, urine and heart. Legend of the symbols is reported in the first graph.




Figure 5: Kinetics of distribution of gallic acid and *o*-coumaric acid in brain, liver, kidneys and blood.




Figure 6: Kinetics of distribution of urolithin A and B in all the organs considered for the experiment.




Table 1: Percentage of recovery in blood compared to initial injected dose and pharmacokinetic analysis of polyphenol microbial metabolites (dose at the steady state (t_{∞}), area under the curve (AUC_{0-15}) and elimination rate (K_e)). For the recovery the values given are the percentages of dose found in rats after the i.v injection at four different time points ($t_{0.25}$, t_2 , t_5 , $t_{15 \text{ min}}$) and control, sd is the standard deviation of 4 replicates. For t_0 the percentage reported is an extrapolation based upon the pharmacokinetics curves associate to a first order exponential decay for an i.v injection.

	Time Points (min.)											Pharmacokinetics			
	dose	t_0^*	$t_{0.25}$		t_2		t_5		t_{15}		Control	Steady state t_{∞}	AUC $(0-15)$	k_e	
	nmol	%	average recovery % \pm st. dev.										pmol/m L	pmol* min/m L	min ⁻¹
4-hydroxyhippuric acid	250	13.3	11.7	5.5	13.4	6.2	6.4	6.1	2.4	0.6	0.0	0.0	0.00	15017	0.09
3-hydroxyphenylacetic acid	150	19.9	18.6	6.2	10.5	2.7	6.5	1.5	3.0	1.6	0.6	0.2	304.50	9238	0.38
homovanillic acid	50	18.7	17.5	5.2	8.8	0.8	7.0	1.7	3.0	1.2	0.4	0.3	118.40	3022	0.45
<i>o</i> -coumaric acid	15	34.6	32.0	11.1	15.7	2.3	12.1	2.8	5.2	2.3	0.1	0.1	62.70	1597	0.47
<i>trans</i> -isofेरulic acid	35	25.9	23.2	7.1	9.3	1.8	7.0	2.3	1.9	1.0	0.0	0.0	73.40	2125	0.58
4-hydroxybenzoic acid	100	18.5	16.3	5.3	6.5	0.6	3.8	1.5	1.5	0.4	1.7	0.9	131.50	3864	0.62
vanillic acid	50	18.7	16.3	7.6	6.4	1.8	3.9	1.5	1.8	0.4	1.3	0.5	75.70	1992	0.67
3-(4-hydroxyphenyl)propanoic acid	100	24.8	21.9	6.4	10.1	1.0	8.4	2.2	4.6	1.1	0.4	0.5	371.50	7538	0.69
uroolithin A	55	11.5	9.7	3.3	2.9	0.9	0.8	0.1	0.1	0.1	0.0	0.0	9.80	733	0.71
3-(3-hydroxyphenyl)propanoic acid	450	15.5	13.5	3.6	5.7	0.8	5.1	0.8	2.2	0.6	0.8	0.6	929.30	19569	0.72
<i>p</i> -coumaric acid	15	22.2	19.3	5.0	7.9	1.3	6.6	1.3	3.0	1.1	3.3	5.4	40.50	872	0.74
<i>trans</i> -ferulic acid	60	18.0	14.7	5.9	3.8	0.5	1.9	0.7	0.5	0.5	0.0	0.1	39.30	1387	0.88
caffeic acid	60	1.8	1.4	0.5	0.3	0.0	0.1	0.1	0.0	0.1	0.0	0.0	2.53	108	0.95
sinapic acid	5	14.5	11.6	4.8	2.8	0.3	1.2	0.1	0.6	0.7	0.0	0.0	2.40	84	0.95
hydroferulic acid	75	19.3	15.7	5.7	4.6	0.4	3.6	1.2	1.2	0.7	0.1	0.1	105.30	2535	0.96
<i>m</i> -coumaric acid	15	43.5	35.1	12.6	9.8	1.5	6.8	2.9	2.7	1.8	0.0	0.0	41.60	1044	0.97
protocatechuic acid	65	8.5	6.6	1.2	3.3	0.5	4.0	3.9	2.4	0.7	1.4	0.1	129.70	2101	1.72
gallic acid	900	1.0	0.6	0.5	0.1	0.0	0.1	0.1	0.1	0.0	0.0	0.0	72.40	1296	2.73
phloroglucinol	100	26.3	23.0	5.8	26.2	4.1	22.8	10.8	17.7	9.8	17.0	6.7	§	19951	§
3,4-dihydroxyphenyl acetic acid	75	§	0.3	0.4	0.0	0.0	0.0	0.0	0.0	0.0	0.0	0.0	§	13	§
pyrocatechol	15	§	1.0	0.9	1.2	2.4	0.0	0.0	0.0	0.0	0.0	0.0	§	35	§
uroolithin B	25	§	1.3	1.0	0.4	0.2	9.4	4.2	9.3	12.4	0.0	0.0	§	1721	§

* extrapolated value from the pharmacokinetic curves for the dose at time zero.

§ no available extrapolated data due to the non fitting with the proper pharmacokinetic curves associated to an i.v. injection.

Table 2: Endogenous amounts of polyphenol metabolites in control brain and polyphenol metabolites significantly increased compared to control brain at any time points ($p > 0.1$).

	dose *	control	t _{0.25}	t ₂	t ₅	t ₁₅	fold of change t ₂ /control	fold of change t ₁₅ /control
	<i>pmol</i> <i>/g of</i> <i>anim</i> <i>al</i>	<i>pmol/g</i>						
4-hydroxyhippuric acid	868	5.52				20.06		3.63
3-hydroxyphenylacetic acid	521	n.d.						
homovanillic acid	174	314.81				433.16		1.38
o-coumaric acid	52	n.d.						
isoferulic acid	121	n.d.						
4-hydroxybenzoic acid	347	126.63		206.97		407.92	1.63	3.22
vanillic acid	174	99.71				384.71		3.86
3-(4-hydroxyphenyl)propanoic acid	347	3.88						
urolithin A	191	9.72						
3-(3-hydroxyphenyl)propanoic acid	1562	119.01				192.81		1.62
p-coumaric acid	52	18.34						
<i>trans</i> -ferulic acid	208	6.34		27.01			4.26	
caffeic acid	208	0.11		2.36		3.81	21.45	34.64
sinapic acid	17	n.d.						
hydroferulic acid	260	n.d.						
m-coumaric acid	52	n.d.						
protocatechuic acid	226	183.40						
gallic acid	3125	52.17	573.00	612.72	265.97	610.82	11.74	11.71
phloroglucinol	347	1373.36						
3,4-dihydroxyphenyl acetic acid	260	291.82				501.19		1.72
pyrocatechol	52	n.d.						
urolithin B	87	n.d.			2.09	2.50		> 2.50
compounds increased at any time ($p > 0.1$)			1	4	2	9		

* the dose is expressed as pmol/g of animal considering the rat average weight (288 g) for comparison with the amounts accumulated in rats

For Table of Contents Use Only

Fate of microbial metabolites of dietary polyphenols in rats: is the brain their target destination?

Mattia Gasperotti[#], Sabina Passamonti^{§*}, Federica Tramer[§], Domenico Masuero[#], Graziano Guella[‡], Fulvio Mattivi[#],
Urska Vrhovsek^{**}

[#] Department of Food Quality and Nutrition, Research and Innovation Centre, Fondazione Edmund Mach (FEM), via E. Mach 1, 38010 San Michele all'Adige, Italy

[§] Department of Life Sciences, University of Trieste, via L. Giorgieri 1, 34127 Trieste, Italy

[‡] Department of Physics, University of Trento, via Sommarive 14, 38123, Trento, Italy

