

THE IRREDUCIBLE COMPONENTS OF THE MODULI SPACE OF DIHEDRAL COVERS OF ALGEBRAIC CURVES

FABRIZIO CATANESE, MICHAEL LÖNNE, FABIO PERRONI

ABSTRACT. The main purpose of this paper is to introduce a new invariant for the action of a finite group G on a compact complex curve of genus g . With the aid of this invariant we achieve the classification of the components of the locus (in the moduli space) of curves admitting an effective action by the dihedral group D_n . This invariant has later been used in [CLP13] where the results of Livingston [Liv85] and of Dunfield and Thurston [Du-Th06] have been extended to the ramified case.

1. INTRODUCTION

We study moduli spaces of curves that admit an effective action by a given finite group G . These moduli spaces can be seen as closed algebraic subsets $M_g(G)$ of M_g , the moduli space of smooth curves of genus $g > 1$. We are mainly interested in understanding which are the irreducible components of $M_g(G)$.

To a curve C of genus g with an action by G , we can associate several discrete invariants that are constant under deformation.

On one hand, the *topological type* of the G -action is a homomorphism $\rho: G \rightarrow \text{Map}_g$ well-defined up to inner conjugation induced by different choices of an isomorphism $\text{Map}(C) \cong \text{Map}_g$, see Section 2.

It turns out that the locus $M_{g,\rho}(G)$ of curves admitting a G -action of topological type ρ is a closed irreducible subset of M_g , (see Theorem 2.3).

On the other hand the action of G on C gives rise to a morphism $p: C \rightarrow C/G =: C'$, a G -cover, and the geometry of p encodes several numerical invariants that are constant on $M_{g,\rho}(G)$: the genus g' of C' , the number d of branch points $y_1, \dots, y_d \in C'$ and the orders $m_1 \leq \dots \leq m_d$ of the local monodromies. These numbers $g', d, m_1 \leq \dots \leq m_d$ form the *primary numerical type*.

A second numerical invariant is obtained from the monodromy $\mu: \pi_1(C' \setminus \{y_1, \dots, y_d\}) \rightarrow G$ of the restriction of p to $p^{-1}(C' \setminus \{y_1, \dots, y_d\})$, and is called the *ν -type* or Nielsen function. It is a class function ν

Date: March 10, 2015.

The present work took place in the realm of the DFG Forschergruppe 790 ‘Classification of algebraic surfaces and compact complex manifolds’.

AMS classification: 14H37, 14H30, 14H15, 57M12, 57M60, 20J99 .

which, for each conjugacy class \mathcal{C} in G , counts the number of local monodromies which belong to \mathcal{C} .

Observe that the irreducible closed algebraic sets $M_{g,\rho}(G)$ depend only upon what we call the ‘unmarked topological type’, which is defined as the conjugacy class of the subgroup $\rho(G)$ inside Map_g .

The following is immediate by Riemann’s existence theorem and the irreducibility of the moduli space $M_{g',d}$ of d -pointed curves of genus g' . Given g' and d , the unmarked topological types whose primary numerical type is of the form g', d, m_1, \dots, m_d are in bijection with the quotient of the set of the corresponding monodromies μ modulo the actions by $Aut(G)$ and by $Map(g', d)$.

Here $Map(g', d)$ is the full mapping class group of genus g' and d unordered points. Thus a first step toward the general problem consists in finding a fine invariant that distinguishes these orbits.

In this paper we introduce a new invariant $\hat{\varepsilon}$ for G -actions on smooth curves.

In the case where G is the dihedral group D_n of order $2n$, we show that $\hat{\varepsilon}$ distinguishes the different unmarked topological types, and therefore $\hat{\varepsilon}$ is a fine invariant in the dihedral case.

Our invariant includes and extends two well known invariants that have been studied in the literature: the ν -type (or Nielsen type) of the cover (also called shape in [FV91], cf. Def. 3.9) and the class in the second homology group $H_2(G/H, \mathbb{Z})$ (modulo the action of $Aut(G/H)$) corresponding to the unramified cover $p': C/H \rightarrow C'$, where H is the minimal normal subgroup of G generated by the local monodromies. These invariants, which refine the primary numerical type, provide a fine invariant under some restrictions, for instance when G is abelian or when G acts freely and is the semi-direct product of two finite cyclic groups (as it follows by combining results from [Cat00], [Cat10], [Edm I] and [Edm II]). However, in general, they are not enough to distinguish unmarked topological types, as one can see already for non-free D_n -actions (see Lemma 5.9).

The construction of $\hat{\varepsilon}$ is similar in spirit to the procedure that, using Hopf’s theorem, associates an element in $H_2(G, \mathbb{Z})$ to any free G -action on a smooth curve. So much cannot be achieved in the ‘branched’ case of a non-free action. In this case we are only able to associate to two given actions with the same ν -type, an invariant in a quotient group of $H_2(G, \mathbb{Z})$ which is the ‘difference’ of the respective $\hat{\varepsilon}$ -invariants. Here is the way we do it. For any finite group G , let F be the free group generated by the elements of G and let $R \trianglelefteq F$ be the subgroup of relations, that is $G = F/R$. For any $\Gamma \subset G$, union of non trivial conjugacy classes, let G_Γ be the quotient group of F by the minimal normal subgroup generated by $[F, R]$ and by the elements $\hat{a}\hat{b}\hat{c}^{-1}\hat{b}^{-1} \in F$, for any $a, c \in \Gamma$, $b \in G$, such that $b^{-1}ab = c$. Here we denote by $\hat{g} \in F$ the generator corresponding to $g \in G$. To a given G -cover $p: C \rightarrow C'$

we associate the set Γ of local monodromies, i.e., of elements which i) stabilize some point x of C and ii) act on the tangent space at x by a rotation of angle $\frac{2\pi}{m}$ where m is the order of the stabilizer at x . Upon the choice of a geometric basis for the fundamental group of the complement $C' \setminus \{y_1, \dots, y_d\}$ of the branch set, our cover is given by an element $v = (c_1, \dots, c_d; a_1, b_1, \dots, a_{g'}, b_{g'}) \in G^{d+2g'}$ satisfying certain conditions (a *Hurwitz generating system*), where the first entries correspond to the local monodromies. Thereby $\Gamma = \Gamma_v$ is the union of the conjugacy classes of the c_i 's. The tautological lift \hat{v} of v is $(\hat{c}_1, \dots, \hat{c}_d; \hat{a}_1, \hat{b}_1, \dots, \hat{a}_{g'}, \hat{b}_{g'})$. Finally, define $\varepsilon(v)$ as the class in G_Γ of

$$\prod_1^d \hat{c}_j \cdot \prod_1^{g'} [\hat{a}_i, \hat{b}_i].$$

It turns out that the image of $\varepsilon(v)$ in $(G_\Gamma)/\text{Inn}(G)$ is invariant under the action of $\text{Map}(g', d)$, as shown in Proposition 3.6. Moreover the ν -type of v can be deduced from $\varepsilon(v)$, as it is essentially the image of $\varepsilon(v)$ in the abelianized group G_Γ^{ab} (see the Remark after Def. 3.9).

In order to take into account also the automorphism group $\text{Aut}(G)$, we define

$$G^\cup := \coprod_\Gamma G_\Gamma,$$

the disjoint union of all the G_Γ 's. Now, the group $\text{Aut}(G)$ acts on G^\cup and we get a map

$$\hat{\varepsilon}: (HS(G; g', d)/\text{Aut}(G)) / \text{Map}(g', d) \rightarrow (G^\cup)/\text{Aut}(G)$$

which is induced by $v \mapsto \varepsilon(v)$. Here we denote by $HS(G; g', d)$ the set of all Hurwitz generating systems of length $d + 2g'$.

Finally, we prove that the map $\hat{\varepsilon}$ is injective in the case $G = D_n$ and we determine the image of $\hat{\varepsilon}$, (Theorem 5.1), thus the invariant $\hat{\varepsilon}$ is a fine invariant for D_n -actions. This completes the classification of the unmarked topological types for $G = D_n$, begun in [CLP11].

We finally show how this classification entails the classification of the irreducible components of the loci $M_g(D_n)$.

When $g' = 0$ our G_Γ is related to the group \hat{G} defined in [FV91] (Appendix), where the authors give a proof of a theorem by Conway and Parker. Roughly speaking the theorem says that: if the Schur multiplier $M(G)$ (which is isomorphic to $H_2(G, \mathbb{Z})$) is generated by commutators, then the ν -type is a fine stable invariant, when $g' = 0$. Results of this kind, when $g' > 0$ but for free G -actions and any finite group G , have been proved in [Liv85] and [Du-Th06]. This time the fine stable invariant lives in $H_2(G, \mathbb{Z})/\text{Aut}(G)$.

The natural question whether our $\hat{\varepsilon}$ -invariant is a fine stable invariant for any finite group G and any effective G -action on compact curves has been solved in [CLP13] for genus stabilisation.

The structure of the paper is the following. In Section 2 we introduce the moduli spaces $M_g(G)$ and the subsets $M_{g,\rho}(G)$. Using Riemann's existence theorem, we reduce the problem of the determination of the loci $M_{g,\rho}(G)$ to a combinatorial one. This leads to the concept of topological type and of Hurwitz generating system. In Section 3 we define the map $\hat{\varepsilon}$, the groups $H_{2,\Gamma}(G)$ and we prove some properties. The object of Section 4 is the computation of $H_{2,\Gamma}(D_n)$. These results are all used in Section 5 where we prove the injectivity of $\hat{\varepsilon}$ when $G = D_n$. In Appendix A we collect some results about mapping class groups and their action on fundamental groups. We use these results in the proof of Theorem 5.1.

Appendix B describes the case (see especially theorem 7.8) where two irreducible loci $M_{g,\rho}(D_n)$ coincide.

2. MODULI SPACES OF G -COVERS

Throughout this Section g is an integer, $g > 1$. The moduli space of curves of genus g is denoted by M_g . For any finite group G , $M_g(G)$ is the locus of $[C] \in M_g$ such that there exists an effective action of G on C . For any $[C] \in M_g(G)$, the quotient morphism $p: C \rightarrow C/G = C'$ is a Galois cover with group G , a G -cover, well defined up to isomorphisms.

Riemann's existence theorem allows us to use combinatorial methods to study G -covers, since p determines and is determined by its restriction to $C' \setminus \mathcal{B}$, where $\mathcal{B} = \{y_1, \dots, y_d\} \subset C'$ is the branch locus of p . Fix a base point $y_0 \in C' \setminus \mathcal{B}$ and a point $x_0 \in p^{-1}(y_0)$. Monodromy gives a surjective group-homomorphism

$$(1) \quad \mu: \pi_1(C' \setminus \mathcal{B}, y_0) \longrightarrow G$$

that characterizes p up to isomorphism.

Let us recall that a **geometric basis** of $\pi_1(C' \setminus \mathcal{B}, y_0)$ consists of simple non-intersecting geometric loops based at y_0

$$\gamma_1, \dots, \gamma_d, \alpha_1, \beta_1, \dots, \alpha_{g'}, \beta_{g'}$$

such that we get the presentation

$$\pi_1(C' \setminus \mathcal{B}, y_0) = \langle \gamma_1, \dots, \gamma_d; \alpha_1, \beta_1, \dots, \alpha_{g'}, \beta_{g'} \mid \prod_1^d \gamma_j \cdot \prod_1^{g'} [\alpha_i, \beta_i] = 1 \rangle.$$

Varying a covering in a flat family with connected base, there are some numerical invariants which remain unchanged, the first ones being the respective genera g, g' of the curves C, C' , which are related by the Hurwitz formula:

$$(2) \quad 2(g-1) = |G|[2(g'-1) + \sum_i (1 - \frac{1}{m_i})], \quad m_i := \text{ord}(\mu(\gamma_i)).$$

Observe moreover that a different choice of the geometric basis changes the generators γ_i , but does not change their conjugacy classes (up to permutation), hence another numerical invariant is provided by the number of elements $\mu(\gamma_i)$ which belong to a fixed conjugacy class in the group G .

We formalize these invariants through the following definition.

Definition 2.1. Let G be a finite group, and $g', d \in \mathbb{N}$. A (g', d) -Hurwitz vector in G is an element $v \in G^{d+2g'}$, the Cartesian product of G $(d+2g')$ -times. A (g', d) -Hurwitz vector in G will also be denoted by

$$v = (c_1, \dots, c_d; a_1, b_1, \dots, a_{g'}, b_{g'}).$$

For any $i \in \{1, \dots, d+2g'\}$, the i -th component v_i of v is defined as usual. The *evaluation* of v is the element

$$ev(v) = \prod_1^d c_j \cdot \prod_1^{g'} [a_i, b_i] \in G.$$

A *Hurwitz generating system of length $d+2g'$ in G* is a (g', d) -Hurwitz vector v in G such that the following conditions hold:

- (i) $c_i \neq 1$ for all i ;
- (ii) G is generated by the components v_i of v ;
- (iii) $ev(v) = 1$.

We denote by $HS(G; g', d) \subset G^{d+2g'}$ the set of all Hurwitz generating systems in G of length $d+2g'$.

Definition 2.2. The condition $ev(v) = \prod_1^d c_j \cdot \prod_1^{g'} [a_i, b_i] = 1$ immediately implies that the product $\prod_1^d c_j$ has trivial image in the abelianization G^{ab} of G . Observe that the image of c_j inside G^{ab} only depends on the conjugacy class \mathcal{C}_j of c_j .

Denote by $[\mathcal{C}]$, for each conjugacy class \mathcal{C} in G , its image inside G^{ab} .

One defines the Nielsen class function of v as the function which, on each conjugacy class \mathcal{C} in G , takes the value $\nu(v)(\mathcal{C}) := |\{j | c_j \in \mathcal{C}\}|$.

We shall say that a class function ν is **admissible** if it satisfies:

$$\sum_{\mathcal{C}} \nu(\mathcal{C})[\mathcal{C}] = 0 \in G^{ab}.$$

Notice that, once we fix a base point $y_0 \in C' \setminus \mathcal{B}$ and a geometric basis of $\pi_1(C' \setminus \mathcal{B}, y_0)$, there is a one-to-one correspondence between the set of Hurwitz generating systems of length $d+2g'$ in G and the set of monodromies μ as in (1).

Topological type. We recall a result contained in [Cat00], see also [Cat08].

Define the orbifold fundamental group $\pi_1^{orb}(C' \setminus \mathcal{B}, y_0; m_1, \dots, m_d)$ as the quotient of $\pi_1(C' \setminus \mathcal{B}, y_0)$ by the minimal normal subgroup generated by the elements $(\gamma_i)^{m_i}$. If $p: C \rightarrow C'$ is a G -covering as above, then

its restriction to $C' \setminus \mathcal{B}$ is a regular topological cover with short exact homotopy sequence

$$1 \rightarrow \pi_1(C \setminus p^{-1}\mathcal{B}, x_0) \rightarrow \pi_1(C' \setminus \mathcal{B}, y_0) \rightarrow G \rightarrow 1.$$

The corresponding exact sequence in orbifold covering theory is

$$1 \rightarrow \pi_1(C, x_0) \rightarrow \pi_1^{orb}(C' \setminus \mathcal{B}, y_0; m_1, \dots, m_d) \rightarrow G \rightarrow 1$$

which is completely determined by the monodromy. In turn the exact sequence determines, via conjugation, a homomorphism

$$\rho: G \rightarrow Out^+(\pi_1(C, x_0)) = Map(C) := Diff^+(C)/Diff^0(C)$$

which is fully equivalent to the topological action of G on C .

Here the superscript $+$ denotes orientation-preserving, and the superscript 0 denotes ‘isotopic to the identity’. The image is contained in the index two subgroup of outer automorphisms corresponding to orientation preserving mapping classes under the Dehn-Nielsen-Baer theorem, cf. [FM12, thm. 8.1].

By Lemma 4.12 of [Cat00], all the curves C of a fixed genus g which admit a given topological action ρ of the group G are parametrized by a connected complex manifold; arguing as in Theorem 2.4 of [Cat10] we get

Theorem 2.3. *The triples (C, G, ρ) where C is a complex projective curve of genus $g \geq 2$, and G is a finite group acting effectively on C with a topological action of type ρ are parametrized by a connected complex manifold $\mathcal{T}_{g;G,\rho}$ of dimension $3(g' - 1) + d$, where g' is the genus of $C' = C/G$, and d is the cardinality of the branch locus \mathcal{B} .*

The image $M_{g,\rho}(G)$ of $\mathcal{T}_{g;G,\rho}$ inside the moduli space M_g is an irreducible closed subset of the same dimension $3(g' - 1) + d$.

Obviously, composing ρ with an automorphism $\varphi \in Aut(G)$, i.e. replacing ρ with $\rho \circ \varphi$, does not change the subgroup $\rho(G) \subset Map(C)$. In particular, $M_{g,\rho}(G) = M_{g,\rho \circ \varphi}(G)$, and similarly $\mathcal{T}_{g;G,\rho} = \mathcal{T}_{g;G,\rho \circ \varphi}$.

Notice that $M_g(G) = \bigcup_{\rho} M_{g,\rho}(G)$, hence the components of $M_g(G)$ are in one-to-one correspondence with a subset of the different unmarked topological types. So, the next question which the above result motivates is: when do two monodromies $\mu_1, \mu_2: \pi_1(C' \setminus \mathcal{B}, y_0) \rightarrow G$ have the same unmarked topological type?

The answer is theoretically easy: the two covering spaces have the same unmarked topological type if and only if they are homeomorphic, hence if and only if μ_1 and μ_2 differ by:

- an automorphism of G ;

- and a different choice of a geometric basis. This is realized by the action of a mapping class in

$$\text{Map}(g', d) := \frac{\text{Diff}^+(C', \mathcal{B})}{\text{Diff}^0(C', \mathcal{B})}.$$

To reformulate these conditions in terms of Hurwitz generating systems, notice that $\text{Aut}(G)$ acts on $HS(G; g', d)$ componentwise, and $\text{Map}(g', d)$ acts on $HS(G; g', d)/_{\text{Aut}(G)}$. The latter action is given by the group homomorphism $\text{Map}(g', d) \rightarrow \text{Out}(\pi_1(C' \setminus \mathcal{B}, y_0))$ and the identification between monodromies μ and Hurwitz generating systems. Theorem 2.3 then implies that there is a bijection between the set of unmarked topological types $[\rho]$ with g' and d fixed, and the following orbit space

$$\{[\rho]\} \longleftrightarrow (HS(G; g', d)/_{\text{Aut}(G)}) /_{\text{Map}(g', d)}.$$

In the next sections, we will also use the action of the unpermuted mapping class group

$$\text{Map}^u(g', d+1) := \text{Map}^u(C', \mathcal{B} \cup \{y_0\})$$

on $HS(G; g', d)$, where $\text{Map}^u(g', d+1)$ consists of diffeomorphisms in $\text{Diff}^+(C')$ which are the identity on $\mathcal{B} \cup \{y_0\}$, modulo isotopy. For any $v_1, v_2 \in HS(G; g', d)$, we write $v_1 \sim v_2$ when they are in the same $\text{Map}^u(g', d+1)$ -orbit. While, $v_1 \approx v_2$ means that they represent the same class in $(HS(G; g', d)/_{\text{Aut}(G)}) /_{\text{Map}(g', d)}$. Clearly $v_1 \sim v_2$ implies $v_1 \approx v_2$.

The mapping class group $\text{Map}(g', d)$ acts on $HS(G; g', d)$ only up to conjugation, but, since we are interested in classifying Hurwitz generating systems up to $\text{Aut}(G)$, we will also use the notation $\varphi \cdot v$, meaning $\varphi \cdot [v]$, with $\varphi \in \text{Map}(g', d)$ and $[v] \in HS(G; g', d)/_{\text{Aut}(G)}$.

3. THE TAUTOLOGICAL LIFT

In this section we give the construction of our invariant in several steps. Having defined a suitable group G_Γ , for any $\Gamma \subset G$ union of non-trivial conjugacy classes, we go on to a map ε , which associates to each Hurwitz vector v (with $c_i \in \Gamma$) an element $\varepsilon(v) \in G_\Gamma$. Any automorphism $f \in \text{Aut}(G)$ induces an isomorphism $f_\Gamma: G_\Gamma \rightarrow G_{f(\Gamma)}$, hence $\text{Aut}(G)$ acts on the disjoint union $G^\cup = \coprod_\Gamma G_\Gamma$. We show two key properties of ε :

- it is $\text{Aut}(G)$ -equivariant (Lemma 3.5), hence it descends to a map

$$\tilde{\varepsilon}: HS(G; g', d)/_{\text{Aut}(G)} \rightarrow G^\cup /_{\text{Aut}(G)};$$

- $\tilde{\varepsilon}$ is constant on the orbits of the mapping class group $\text{Map}(g', d)$ (Proposition 3.6).

Therefore ε descends to our invariant $\hat{\varepsilon}$ which is formalized as the map

$$\hat{\varepsilon}: (HS(G; g', d)/Aut(G)) /_{Map(g', d)} \rightarrow G^U /_{Aut(G)}, \quad \forall g', d,$$

induced by ε . We conclude the section with the study of general properties of the invariant that are relevant to this work.

Since our construction is inspired by Hopf's description of the second homology group $H_2(G, \mathbb{Z})$ [Hopf], we begin by recalling this. For a finite group G , fix a presentation of G :

$$1 \rightarrow R \rightarrow F \rightarrow G \rightarrow 1,$$

where F is a free group. Then there is a group isomorphism (cf. [Bro]):

$$(3) \quad H_2(G, \mathbb{Z}) \cong \frac{R \cap [F, F]}{[F, R]}.$$

If $v = (a_1, b_1, \dots, a_{g'}, b_{g'}) \in G^{2g'}$ satisfies $\prod_1^{g'} [a_i, b_i] = 1$, then we can associate a class in $H_2(G, \mathbb{Z})$ in the following way: choose liftings $\hat{a}_i, \hat{b}_i \in F$ of a_i, b_i , then $\prod_1^{g'} [\hat{a}_i, \hat{b}_i] \in R \cap [F, F]$ and its class in $\frac{R \cap [F, F]}{[F, R]}$ gives an element of $H_2(G, \mathbb{Z})$, according to (3). Clearly, this element does not depend on the various choices, moreover it is invariant under the action of the mapping class group, thus giving a topological invariant of v .

The topological meaning of this invariant is the following. If the G -action on C is free, the covering $p: C \rightarrow C'$ is étale, and hence it corresponds to a continuous function $Bp: C' \rightarrow BG$, up to homotopy. Here BG is the classifying space of G . The topological invariant is simply the image $Bp_*([C']) \in H_2(BG, \mathbb{Z}) = H_2(G, \mathbb{Z})$ of the fundamental class $[C'] \in H_2(C', \mathbb{Z})$ of C' under the homomorphism $Bp_*: H_2(C', \mathbb{Z}) \rightarrow H_2(BG, \mathbb{Z})$ induced by Bp . Now, if we view C' as an Eilenberg-Mac Lane space $K(\pi_1(C'), 1)$, then the fundamental class $[C']$ is given by

$$\prod_1^{g'} [\hat{\alpha}_i, \hat{\beta}_i] \in H_2(\pi_1(C'), \mathbb{Z}),$$

where as usual $\alpha_1, \beta_1, \dots, \alpha_{g'}, \beta_{g'}$ is a geometric basis of $\pi_1(C')$ and $\hat{\alpha}_i, \hat{\beta}_i$ are liftings to the free group of a presentation of $\pi_1(C')$. So,

$$Bp_*([C']) = \prod_1^{g'} [\hat{a}_i, \hat{b}_i] \in H_2(G, \mathbb{Z}),$$

where $a_i = \mu(\alpha_i)$, $b_i = \mu(\beta_i)$ and $\mu: \pi_1(C') \rightarrow G$ is the monodromy of $p: C \rightarrow C'$.

From now on, $F = \langle \hat{g} \mid g \in G \rangle$ is the free group generated by the elements of G . Let $R \trianglelefteq F$ be the minimal normal subgroup generated by the relations, that is $G = \frac{F}{R}$.

Definition 3.1. Let G be a finite group and let F, R be as above. For any union of non-trivial conjugacy classes $\Gamma \subset G$, define

$$R_\Gamma = \langle \langle [F, R], \hat{a}\hat{b}\hat{c}^{-1}\hat{b}^{-1} \mid \forall a \in \Gamma, ab = bc \rangle \rangle,$$

$$G_\Gamma = \frac{F}{R_\Gamma}.$$

The map $\hat{a} \mapsto a, \forall a \in G$, induces a group homomorphism $\alpha: G_\Gamma \rightarrow G$. Set $K_\Gamma = \ker(\alpha)$.

Lemma 3.2. *With the notation as before, the following holds. $R_\Gamma \subset R$ and $K_\Gamma = \frac{R}{R_\Gamma}$. In particular K_Γ is contained in the center of G_Γ and the short exact sequence*

$$1 \longrightarrow \frac{R}{R_\Gamma} \longrightarrow G_\Gamma \xrightarrow{\alpha} G \longrightarrow 1$$

is a central extension.

Proof. $[F, R] \subset R$ because R is normal in F and moreover $\hat{a}\hat{b}\hat{c}^{-1}\hat{b}^{-1} \in R$ for any $a, b, c \in G$ with $ab = bc$, therefore $R_\Gamma \subset R$. By the definition of α we have that $K_\Gamma = \frac{R}{R_\Gamma}$. Finally, K_Γ is in the center of G_Γ because $[F, R] \subset R_\Gamma$. \square

The morphism $\alpha: G_\Gamma \rightarrow G$ has a *tautological section* $G \rightarrow G_\Gamma$, $a \mapsto \hat{a}$. This map is not a group homomorphism in general, but every element $\xi \in G_\Gamma$ can be written as $\hat{g}z = z\hat{g}$, with $g = \alpha(\xi) \in G$ and $z \in K_\Gamma$. Here, by abuse of notation, \hat{a} denotes also the class of $\hat{a} \in F$ in $G_\Gamma = F/R_\Gamma$.

Lemma 3.3. *Let $\hat{a}, \xi \in G_\Gamma$. Suppose that \hat{a} is conjugate to ξ in G_Γ and that $a \in \Gamma$. Then $\xi = \widehat{\alpha(\xi)}$.*

Proof. Let $\hat{b}z$ be a conjugating element, that is $\hat{a}\hat{b}z = \hat{b}z\xi$. As $z \in K_\Gamma$, it commutes with any element, hence

$$(4) \quad \hat{a}\hat{b} = \hat{b}\xi.$$

Now apply α and obtain: $ab = b\alpha(\xi)$. By assumption $a \in \Gamma$, hence by definition of G_Γ we have that $\hat{a}\hat{b} = \widehat{b\alpha(\xi)}$. Now using (4) we deduce $\xi = \widehat{\alpha(\xi)}$. \square

Definition 3.4. Given a (g', d) -Hurwitz vector

$$v = (c_1, \dots, c_d; a_1, b_1, \dots, a_{g'}, b_{g'})$$

in G , cf. Definition 2.1, its *tautological lift* \hat{v} is the (g', d) -Hurwitz vector in G_Γ defined by

$$\hat{v} = (\widehat{c}_1, \dots, \widehat{c}_d; \widehat{a}_1, \widehat{b}_1, \dots, \widehat{a}_{g'}, \widehat{b}_{g'})$$

where the factors are the tautological lifts of the factors of v .

Given a (g', d) -Hurwitz vector v in G with $c_i \neq 1, \forall i$, we denote by Γ_v the union of all conjugacy classes of G containing at least one c_i .

For any v as before, let

$$\varepsilon(v) = \prod_1^d \widehat{c}_j \cdot \prod_1^{g'} [\widehat{a}_i \widehat{b}_i] \in G_{\Gamma_v},$$

be the evaluation of the tautological lift \widehat{v} of v in G_{Γ_v} , in analogy to Definition 2.1.

Lemma 3.5. *Let G be any finite group, and let $\Gamma \subset G$ be any union of non trivial conjugacy classes. Then we have:*

- i) any $f \in \text{Aut}(G)$ induces an isomorphism $f_\Gamma: G_\Gamma \rightarrow G_{f(\Gamma)}$;
- ii) $\varepsilon(f(v)) = f_\Gamma(\varepsilon(v))$, $\forall f \in \text{Aut}(G)$ and $\forall v$ a (g', d) -Hurwitz vector with $c_i \neq 1$, $\forall i$, where $\Gamma = \Gamma_v$.

Proof. i) $f \in \text{Aut}(G)$ lifts to an automorphism $\widehat{f} \in \text{Aut}(F)$ defined by

$$\widehat{f}: \widehat{a} \mapsto \widehat{f(a)}.$$

We have: $\widehat{f}(R) \subset R$, and moreover

$$\widehat{f}(\widehat{a\widehat{b}\widehat{c}^{-1}\widehat{b}^{-1}}) = \widehat{f(a)}\widehat{f(b)}\widehat{f(c)}^{-1}\widehat{f(b)}^{-1},$$

for any $a, b, c \in G$. If $a \in \Gamma$, then $f(a) \in f(\Gamma)$ and hence

$$\widehat{f(a)}\widehat{f(b)}\widehat{f(c)}^{-1}\widehat{f(b)}^{-1} \in R_{f(\Gamma)}.$$

$$\begin{aligned} \text{ii)} \quad \varepsilon(f(v)) &= \varepsilon(f(c_1), \dots, f(c_d); f(a_1), \dots, f(b_{g'})) \\ &= \prod_1^d \widehat{f(c_i)} \cdot \prod_1^{g'} [\widehat{f(a_j)}, \widehat{f(b_j)}] \\ &= \prod_1^d \widehat{f(\widehat{c}_i)} \cdot \prod_1^{g'} [\widehat{f(\widehat{a}_j)}, \widehat{f(\widehat{b}_j)}] = f_\Gamma(\varepsilon(v)). \end{aligned}$$

□

Now, we define

$$G^\cup := \prod_\Gamma G_\Gamma,$$

and regard ε as a map $\varepsilon: HS(G; g', d) \rightarrow G^\cup$, $v \mapsto \varepsilon(v) \in G_{\Gamma_v}$. Then the previous lemma means that ε induces a map

$$\tilde{\varepsilon}: HS(G; g', d) / \text{Aut}(G) \rightarrow (G^\cup) / \text{Aut}(G).$$

We have the following

Proposition 3.6. *For any $g', d \in \mathbb{N}$, $\tilde{\varepsilon}$ is $\text{Map}(g', d)$ -invariant, hence it induces a map*

$$\hat{\varepsilon}: (HS(G; g', d) / \text{Aut}(G)) / \text{Map}(g', d) \rightarrow (G^\cup) / \text{Aut}(G).$$

To prove this proposition we need some preliminary results.

Lemma 3.7. *Let Γ_v be associated to a (g', d) -Hurwitz vector v as in Definition 3.4. If the Hurwitz vector v' is related to v by an elementary braid move, then $\varepsilon(v) = \varepsilon(v')$.*

Proof. It suffices to consider the case $g = 0$, $d = 2$ and the elementary braid move σ_1 which maps

$$v = (c_1, c_2) \quad \text{to} \quad v' = (c_2, c_2^{-1}c_1c_2).$$

In G_{Γ_v} we have, thanks to $c_1 \in \Gamma_v$, $c_1c_2 = c_2(c_2^{-1}c_1c_2)$, and the relations of G_{Γ_v} :

$$\varepsilon(v) = \widehat{c_1c_2} = \widehat{c_2c_2^{-1}c_1c_2} = \varepsilon(v').$$

□

Lemma 3.8. *If $\xi, \eta \in G_{\Gamma}$, then*

$$[\xi, \eta] = \left[\widehat{\alpha(\xi)}, \widehat{\alpha(\eta)} \right].$$

Proof. Write $\xi = \widehat{\alpha(\xi)}z$ and $\eta = \widehat{\alpha(\eta)}z'$ with z, z' in K_{Γ} , hence central (Lemma 3.2). Then the conclusion is immediate.

□

Proof. (Of Proposition 3.6.) Let $\varphi \in \text{Map}(g', d)$. Thanks to Lemma 3.7 it suffices to consider the case that φ is a pure mapping class, i.e. that φ does not permute the conjugacy classes associated to the local monodromies. Using Lemma 3.5 ii), we can further ignore the action of G by conjugation and pretend that $\text{Map}(g', d)$ acts on $HS(G; g', d)$.

Since φ is a pure mapping class, the components of v , $\varphi \cdot v$ are conjugate to each other, $v_i \sim (\varphi \cdot v)_i$ for $i = 1, \dots, d$, and where \sim denotes conjugation equivalence. For the components of \hat{v} , $\varphi \cdot \hat{v}$ the same is true, $\hat{v}_i \sim (\varphi \cdot \hat{v})_i$ for $i = 1, \dots, d$. By Lemma 3.3 we thus have:

$$(\hat{v})_i \sim (\varphi \cdot \hat{v})_i \Rightarrow (\varphi \cdot \hat{v})_i = \widehat{\alpha((\varphi \cdot \hat{v})_i)}.$$

Now notice that the homomorphism α of Definition 3.1 induces a map

$$\alpha^{(d+2g')} : HS(G_{\Gamma}; g', d) \longrightarrow HS(G; g', d),$$

which is equivariant under the action of the mapping class group in the following sense: consider the factorizations as a map from the free group on $d + 2g'$ generators to G_{Γ} , resp. G , and the mapping class group as a group of automorphisms of this free group. Then $\alpha^{(d+2g')}$ is equivariant, since such automorphisms act by pre-composition. Hence by the equivariance of $\alpha^{(d+2g')}$

$$\alpha((\varphi \cdot \hat{v})_i) = (\varphi \cdot v)_i$$

and therefore, for $i = 1, \dots, d$,

$$(\varphi \cdot \hat{v})_i = \widehat{\alpha((\varphi \cdot \hat{v})_i)} = \widehat{(\varphi \cdot v)_i} = (\widehat{\varphi \cdot v})_i.$$

By Lemma 3.8 we may change also the entries $(\varphi \cdot \hat{v})_i$, $i > d$ in the commutators to $\widehat{\alpha((\varphi \cdot \hat{v}))} = (\widehat{\varphi \cdot v})_i$ without changing the value of the commutators. Hence

$$ev(\varphi \cdot \hat{v}) = ev(\widehat{\varphi \cdot v}) = \varepsilon(\varphi \cdot v).$$

By the invariance of the evaluation under the mapping class

$$\varepsilon(v) = ev(\hat{v}) = ev(\varphi \cdot \hat{v}) = \varepsilon(\varphi \cdot v)$$

and we have proved our claim. \square

Definition 3.9. Let $v \in HS(G; g', d)$ and let $\nu(v) \in \bigoplus_{\mathcal{C}} \mathbb{Z}\langle \mathcal{C} \rangle$ (\mathcal{C} runs over the set of conjugacy classes of G) be the vector whose \mathcal{C} -component is the number of v_j , $j \leq d$, which belong to \mathcal{C} ($\nu(v)$ is also called the shape of v in [FV91]).

The map

$$\nu: HS(G; g', d) \rightarrow \bigoplus_{\mathcal{C}} \mathbb{Z}\langle \mathcal{C} \rangle$$

obtained in this way induces a map

$$\tilde{\nu}: HS(G; g', d) / Aut(G) \rightarrow (\bigoplus_{\mathcal{C}} \mathbb{Z}\langle \mathcal{C} \rangle) / Aut(G)$$

which is $Map(g', d)$ -invariant, therefore we get a map

$$\hat{\nu}: (HS(G; g', d) / Aut(G)) / Map(g', d) \rightarrow (\bigoplus_{\mathcal{C}} \mathbb{Z}\langle \mathcal{C} \rangle) / Aut(G).$$

For any $v \in HS(G; g', d)$, we call $\hat{\nu}(v)$ the ν -type of v .

Remark 3.10. Let $v \in HS(G; g', d)$ and let $\Gamma_v \subset G$ be the union of the conjugacy classes of v_j , $j \leq d$. The abelianization $G_{\Gamma_v}^{ab}$ of G_{Γ_v} can be described as follows:

$$G_{\Gamma_v}^{ab} \cong \bigoplus_{\mathcal{C} \subset \Gamma_v} \mathbb{Z}\langle \mathcal{C} \rangle \oplus \bigoplus_{g \in G \setminus \Gamma_v} \mathbb{Z}\langle g \rangle,$$

where \mathcal{C} denotes a conjugacy class of G .

Observe that $\nu(v) \in \bigoplus_{\mathcal{C} \subset \Gamma_v} \mathbb{Z}\langle \mathcal{C} \rangle \subset G_{\Gamma_v}^{ab}$ coincides with the vector which is the image in $G_{\Gamma_v}^{ab}$ of $\varepsilon(v) \in G_{\Gamma_v}$ under the natural homomorphism $G_{\Gamma_v} \rightarrow G_{\Gamma_v}^{ab}$.

Definition 3.11. Let $\Gamma \subset G$ be a union of non-trivial conjugacy classes of G . We define

$$H_{2,\Gamma}(G) = \ker(G_{\Gamma} \rightarrow G \times G_{\Gamma}^{ab}),$$

where $G_{\Gamma} \rightarrow G \times G_{\Gamma}^{ab}$ is the morphism with first component α (defined in Definition 3.1) and second component the natural morphism $G_{\Gamma} \rightarrow G_{\Gamma}^{ab}$.

Notice that

$$H_2(G, \mathbb{Z}) \cong \frac{R \cap [F, F]}{[F, R]} \cong \ker \left(\begin{array}{c} F \\ [F, R] \end{array} \rightarrow G \times G_{\emptyset}^{ab} \right).$$

In particular, when $\Gamma = \emptyset$, $H_{2,\Gamma}(G) \cong H_2(G, \mathbb{Z})$.

The next result gives a precise relation between $H_2(G, \mathbb{Z})$ and $H_{2,\Gamma}(G)$.

Lemma 3.12. *Let G be a finite group and let $\Gamma \subset G$ be a union of nontrivial conjugacy classes. Write $G = \frac{F}{R}$ and $G_\Gamma = \frac{F}{R_\Gamma}$. Then, there is a short exact sequence*

$$1 \rightarrow \frac{R_\Gamma \cap [F, F]}{[F, R]} \rightarrow H_2(G, \mathbb{Z}) \rightarrow H_{2, \Gamma}(G) \rightarrow 1.$$

In particular $H_{2, \Gamma}(G)$ is abelian.

Proof. We first define the morphism $H_2(G, \mathbb{Z}) \rightarrow H_{2, \Gamma}(G)$. By Hopf's Theorem we identify $H_2(G, \mathbb{Z})$ with $\frac{R \cap [F, F]}{[F, R]}$ (cf. [Bro]). On the other hand we have:

$$H_{2, \Gamma}(G) = \ker(G_\Gamma \rightarrow G) \cap \ker(G_\Gamma \rightarrow G_\Gamma^{ab}) = \frac{R}{R_\Gamma} \cap [G_\Gamma, G_\Gamma].$$

By Lemma 3.2, $R_\Gamma \subset R$. The homomorphism $R \cap [F, F] \rightarrow \frac{R}{R_\Gamma}$, $r \mapsto rR_\Gamma$, takes values in $H_{2, \Gamma}(G)$. Moreover it descends to a group homomorphism $H_2(G, \mathbb{Z}) \rightarrow H_{2, \Gamma}(G)$ because $[F, R] \subset R_\Gamma$.

To prove the surjectivity, let

$$aR_\Gamma \in \frac{R}{R_\Gamma} \cap [G_\Gamma, G_\Gamma].$$

Since $aR_\Gamma \in [G_\Gamma, G_\Gamma] = \frac{[F, F] \cdot R_\Gamma}{R_\Gamma}$, we may assume $a \in [F, F]$. From $aR_\Gamma \in \frac{R}{R_\Gamma}$, we have $aR_\Gamma = rR_\Gamma$, for some $r \in R$. Since $R_\Gamma \subset R$, we deduce that $a \in R$ and so the surjectivity follows.

The kernel of the morphism so defined is $\frac{R_\Gamma \cap [F, F]}{[F, R]}$.

Since $H_2(G, \mathbb{Z})$ is abelian, so is $H_{2, \Gamma}(G)$. \square

Proposition 3.13. *Let $v_1, v_2 \in HS(G; g', d)$ be two Hurwitz generating systems in G with the same ν -type. Then $\Gamma_{v_1} = \Gamma_{v_2} =: \Gamma$. Moreover, if $ev(v_1) = ev(v_2) \in G$, then the element*

$$\varepsilon(v_1)^{-1} \cdot \varepsilon(v_2) \in H_{2, \Gamma}(G)$$

is invariant under the group $\widetilde{Map}(g', d)$ of isotopy classes of orientation-preserving diffeomorphisms of the pair (C', \mathcal{B}) that fix y_0 . In particular:

- (1) *if v_1 and v_2 are equivalent then the element is trivial;*
- (2) *if the element is non-trivial, then v_1 and v_2 are inequivalent.*

4. COMPUTATION OF $H_{2, \Gamma}(D_n)$

In this section we derive a complete description of $H_{2, \Gamma}(G)$ in the special case that G is equal to the dihedral group

$$\begin{aligned} D_n &= \langle x, y \mid x^n = 1, y^2 = 1, xy = yx^{-1} \rangle \\ &= \{ x^i y^j \mid 0 \leq i < n, 0 \leq j < 2 \} \end{aligned}$$

Proposition 4.1. *Let $n \in \mathbb{N}$, $n \geq 3$. Then we have:*

- (i) *$H_2(D_n, \mathbb{Z})$ is trivial if n is odd and it is isomorphic to $\mathbb{Z}/2\mathbb{Z}$ if n is even;*

(ii) *the natural action of $\text{Aut}(D_n)$ on $H_2(D_n, \mathbb{Z})$ is trivial.*

Proof. (ii) This claim follows directly from (i) and from the fact that the neutral element of $H_2(D_n, \mathbb{Z})$ is fixed by the action of $\text{Aut}(D_n)$.

(i) Identify D_n with the subgroup of $SO(3)$ generated by

$$x := \begin{pmatrix} \cos \frac{2\pi}{n} & -\sin \frac{2\pi}{n} & 0 \\ \sin \frac{2\pi}{n} & \cos \frac{2\pi}{n} & 0 \\ 0 & 0 & 1 \end{pmatrix} \quad \text{and} \quad y := \begin{pmatrix} -1 & 0 & 0 \\ 0 & 1 & 0 \\ 0 & 0 & -1 \end{pmatrix}.$$

Let $u: SU(2) \rightarrow SO(3)$ be the homomorphism $q \mapsto R_q$, where we identify $SU(2)$ with the quaternions $q \in \mathbb{H}$ of norm 1, \mathbb{R}^3 with $\text{Im}\mathbb{H}$, and $R_q(x) = qx\bar{q}$. Consider the binary dihedral group $\tilde{D}_n = u^{-1}(D_n)$. It fits in the following short exact sequence:

$$(5) \quad 1 \rightarrow \mathbb{Z}/2\mathbb{Z} \rightarrow \tilde{D}_n \rightarrow D_n \rightarrow 1,$$

from which we get the 5-term exact sequence (see e.g. [Bro], pg. 47, Exercise 6):

$$(6) \quad H_2(\tilde{D}_n) \rightarrow H_2(D_n) \rightarrow (H_1(\mathbb{Z}/2\mathbb{Z}))_{D_n} \rightarrow H_1(\tilde{D}_n) \rightarrow H_1(D_n) \rightarrow 0,$$

where all the coefficients are in \mathbb{Z} and $(H_1(\mathbb{Z}/2\mathbb{Z}))_{D_n}$ is the group of co-invariants under the D_n -action on $\mathbb{Z}/2\mathbb{Z}$ induced by conjugation by \tilde{D}_n , hence $(H_1(\mathbb{Z}/2\mathbb{Z}))_{D_n} = H_1(\mathbb{Z}/2\mathbb{Z})$ since $\mathbb{Z}/2\mathbb{Z}$ is in the center of \tilde{D}_n .

We have that $H_2(\tilde{D}_n) = \{0\}$, since \tilde{D}_n is a finite subgroup of $SU(2) \cong S^3$ (see [Bro] II pg. 47, Exercise 7). Next, recall that, for any group G , $H_1(G, \mathbb{Z})$ is isomorphic to the abelianization G^{ab} (see [Bro] pg. 36), hence (6) reduces to

$$0 \rightarrow H_2(D_n) \rightarrow \mathbb{Z}/2\mathbb{Z} \rightarrow \tilde{D}_n^{ab} \rightarrow D_n^{ab} \rightarrow 0.$$

To conclude we show that $\ker(\tilde{D}_n^{ab} \rightarrow D_n^{ab}) = \{0\}$ if and only if n is even. With the imaginary units $\underline{i}, \underline{j}, \underline{k} \in \mathbb{H}$ let

$$\xi = \cos\left(\frac{\pi}{n}\right) + \underline{k} \cdot \sin\left(\frac{\pi}{n}\right) \in u^{-1}(x) \quad \text{and} \quad \eta = \underline{j} \in u^{-1}(y).$$

Since $[\xi^\ell, \eta] = \xi^{2\ell}$, $\forall \ell$, we see that, if n is odd, $\xi^n \notin [\tilde{D}_n, \tilde{D}_n]$, but $u(\xi^n) = 1$ and hence $\ker(\tilde{D}_n^{ab} \rightarrow D_n^{ab}) \neq \{0\}$. When n is even, $\tilde{D}_n^{ab} \cong \mathbb{Z}/2\mathbb{Z} \times \mathbb{Z}/2\mathbb{Z}$ and the map $\tilde{D}_n^{ab} \rightarrow D_n^{ab}$ is an isomorphism. \square

Using Lemma 3.2 from [Wie], we deduce the following

Corollary 4.2. *Let $n \in \mathbb{N}$, $n \geq 4$ even. Then, the binary dihedral group \tilde{D}_n is a Schur cover of D_n and the exact sequence (5) identifies $\mathbb{Z}/2\mathbb{Z}$ with $H_2(D_n, \mathbb{Z})$. In particular, for any $(a_1, b_1, \dots, a_{g'}, b_{g'}) \in (D_n)^{2g'}$ with $\prod_1^{g'} [a_i, b_i] = 1$, the image of $\prod_1^{g'} [\hat{a}_i, \hat{b}_i] \in R \cap [F, F]$ in $H_2(D_n, \mathbb{Z}) = \frac{R \cap [F, F]}{[R, F]}$ is given by $\prod_1^{g'} [\tilde{a}_i, \tilde{b}_i]$, where $\tilde{a}_i, \tilde{b}_i \in \tilde{D}_n$ are liftings of a_i, b_i .*

Next the union Γ of non-trivial conjugacy classes comes into play. Recall that the set of reflections $\{x^i y \mid 0 \leq i < n\}$ is a single conjugacy class in case n is odd, and splits into two conjugacy classes in case n even according to the parity of i .

The conjugacy classes in the set of rotations $\{x^i \mid 0 \leq i < n\}$ are of the kind $\{x^i, x^{n-i}\}$ and contain two elements except for $i = 0$ and $i = \frac{n}{2}$ in case n is even.

Corollary 4.3. *Let $\Gamma \subset D_n$ be a union of non-trivial conjugacy classes, $\Gamma \neq \emptyset$. Then $H_{2,\Gamma}(D_n) = \{0\}$ in the following cases: n is odd; n is even and Γ contains some reflection; n is even and Γ contains the non-trivial central element. In the remaining case, $H_{2,\Gamma}(D_n) = \mathbb{Z}/2\mathbb{Z}$.*

Proof. If n is odd, then $H_2(D_n, \mathbb{Z}) = \{0\}$ and hence $H_{2,\Gamma}(D_n) = \{0\}$ for any Γ (Lemma 3.12).

If $n = 2k$ and Γ contains some reflection, say $y \in \Gamma$, then $\widehat{y x^k y^{-1} x^k}^{-1} \in R_\Gamma \cap [F, F]$. But the image of this element in $H_2(D_n, \mathbb{Z})$ is not trivial (Corollary 4.2), hence $H_{2,\Gamma}(D_n) = \{0\}$ (Lemma 3.12). The same argument works if $xy \in \Gamma$.

Assume now $n = 2k$ and $x^k \in \Gamma$. Then $\widehat{x^k y x^k}^{-1} \widehat{y}^{-1} \in R_\Gamma \cap [F, F]$ and its image in $H_2(D_n, \mathbb{Z})$ is not trivial, hence $H_{2,\Gamma}(D_n) = \{0\}$ also in this case.

Finally, if $n = 2k$ and $\Gamma \subset \mathbb{Z}/n\mathbb{Z} \setminus \{x^k\}$, then

$$R_\Gamma = \langle \langle [F, R], \widehat{x^\alpha x^\beta x^\alpha}^{-1} \widehat{x^\beta}^{-1}, \widehat{x^\alpha x^\beta y x^{n-\alpha}}^{-1} \widehat{x^\beta y}^{-1} \mid x^\alpha \in \Gamma \rangle \rangle.$$

First we note that the image of $\widehat{x^\alpha x^\beta x^\alpha}^{-1} \widehat{x^\beta}^{-1}$ in $H_2(D_n, \mathbb{Z})$ is 0. Second, the elements $\widehat{x^\alpha x^\beta y x^{n-\alpha}}^{-1} \widehat{x^\beta y}^{-1}$ generate an abelian group modulo $[F, R]$. Last, the intersection of this subgroup with $[F, F]/[F, R]$ is generated by elements represented by

$$\widehat{x^\alpha x^\beta y x^{n-\alpha}}^{-1} \widehat{x^\beta y}^{-1} \quad . \quad \widehat{x^{n-\alpha} x^\gamma y x^\alpha}^{-1} \widehat{x^\gamma y}^{-1}.$$

It remains to show that these are trivial modulo $[F, R]$, in fact

$$\begin{aligned} &\equiv \widehat{x^\beta y}^{-1} \widehat{x^\alpha x^\beta y x^{n-\alpha}}^{-1} \quad . \quad \widehat{x^{n-\alpha} x^\gamma y x^\alpha}^{-1} \widehat{x^\gamma y}^{-1} \\ &\equiv \widehat{x^\beta y}^{-1} \widehat{x^\alpha x^\beta y} \quad \widehat{x^\gamma y x^\alpha}^{-1} \widehat{x^\gamma y}^{-1} \\ &\equiv \widehat{x^\beta y}^{-1} \widehat{x^\alpha} \quad \underbrace{\widehat{x^\beta y} \quad \widehat{x^\gamma y} \quad \widehat{x^{\gamma-\beta}}}_{\in R} \quad \widehat{x^{\gamma-\beta}}^{-1} \quad \widehat{x^\alpha}^{-1} \widehat{x^\gamma y}^{-1} \\ &\equiv \widehat{x^\beta y}^{-1} \quad \widehat{x^\beta y} \quad \widehat{x^\gamma y} \quad \widehat{x^{\gamma-\beta}} \quad \widehat{x^\alpha} \quad \widehat{x^{\gamma-\beta}}^{-1} \quad \widehat{x^\alpha}^{-1} \widehat{x^\gamma y}^{-1} \\ &\equiv \widehat{x^\gamma y} \quad \widehat{x^{\gamma-\beta}} \quad \widehat{x^\alpha} \quad \widehat{x^{\gamma-\beta}}^{-1} \quad \widehat{x^\alpha}^{-1} \widehat{x^\gamma y}^{-1} \\ &\equiv \widehat{x^{\gamma-\beta}} \quad \widehat{x^\alpha} \quad \widehat{x^{\gamma-\beta}}^{-1} \widehat{x^\alpha}^{-1}. \end{aligned}$$

This last element is trivial modulo $[F, R]$ as noted first. We deduce that $\frac{R_\Gamma \cap [F, F]}{[F, R]} = \{0\}$ and hence $H_{2, \Gamma}(D_n) \cong H_2(D_n, \mathbb{Z}) \cong \mathbb{Z}/2\mathbb{Z}$, by Lemma 3.12. \square

5. THE INJECTIVITY OF $\hat{\varepsilon}$ WHEN $G = D_n$

Recall the following

Notation. For any Hurwitz vector $v = (c_1, \dots, c_d; a_1, b_1, \dots, a_{g'}, b_{g'}) \in G^{d+2g'}$,

$$ev(v) = \prod_{i=1}^d c_i \cdot \prod_{j=1}^{g'} [a_j, b_j] \in G,$$

while, if $c_i \neq 1, \forall i$, $\varepsilon(v) = ev(\hat{v}) \in G_{\Gamma_v}$, where $\hat{v} \in (G_{\Gamma_v})^{d+2g'}$ is the tautological lifting (Definition 3.4).

In this section we prove the following

Theorem 5.1. *Let $G = D_n$, the dihedral group of order $2n$. Then, $\forall g', d$, we have:*

- (i) $\hat{\varepsilon}: (HS(G; g', d)/Aut(G)) /_{Map(g', d)} \rightarrow (G^\cup)/Aut(G) := (\prod_\Gamma G_\Gamma)/Aut(G)$ is injective;
- (ii) The image $Im(\hat{\varepsilon})$ is the inverse image of $Im(\hat{\nu})$ in $(\prod_\Gamma K_\Gamma)/Aut(G)$, where K_Γ is defined in Def. 3.1. In other words, for any $\Gamma \subset G$, union of non trivial conjugacy classes, $Im(\varepsilon) \cap G_\Gamma = H_{2, \Gamma}(G) \cdot (Im(\varepsilon) \cap G_\Gamma)$.
- (iii) For $g' \geq 2$, $Im(\hat{\nu})$ is just the set of admissible class functions ν . For $g' = 1$, $Im(\hat{\nu})$ is the union of the set of admissible class functions ν for which Γ contains some reflection, together with a subset \mathcal{S} of those for which Γ generates a subgroup H of index at most 2 in the subgroup \mathcal{R} of rotations, \mathcal{S} contains the subset for which $H = \mathcal{R}$.

To prove (i), let $[v_1]_\approx, [v_2]_\approx \in (HS)/_\approx$ be equivalence classes up to the combined action of $Aut(G)$ and the mapping class group such that $\hat{\varepsilon}([v_1]_\approx) = \hat{\varepsilon}([v_2]_\approx)$. Then there exists an automorphism $f \in Aut(G)$ such that $f(\Gamma_{v_1}) = \Gamma_{v_2}$ and $f(\varepsilon(v_1)) = \varepsilon(v_2)$. Hence, by Lemma 3.5, we assume without loss of generality $\Gamma_{v_1} = \Gamma_{v_2} = \Gamma$ and $\varepsilon(v_1) = \varepsilon(v_2)$, in particular

$$\varepsilon(v_1) \cdot \varepsilon(v_2)^{-1} = 0 \in H_{2, \Gamma}(G).$$

The outline of the proof is now the following. We address the following mutually exclusive cases: $\Gamma = \emptyset$ (the étale case); $\Gamma \neq \emptyset$ and contains some reflection; $\Gamma \neq \emptyset$ and does not contain reflections. In the first case, for each element of HS , we determine a normal form with respect to \approx , then we show that two different normal forms are distinguished by $H_2(D_n, \mathbb{Z})$ (recall that $Aut(D_n)$ acts trivially on $H_2(D_n, \mathbb{Z})$). In the second case we will show that all Hurwitz generating systems with the

same numerical invariants (n , g' and ν -type) are equivalent with respect to \approx (this agrees with the fact $H_{2,\Gamma}(D_n) = \{0\}$ in this case). In the last case, for every $v \in HS$, we determine a normal form v' with respect to \approx . We see that two different normal forms v'_1 and v'_2 have different invariants, $\varepsilon(v'_1) \neq \varepsilon(v'_2) \in G_\Gamma$. Finally we prove that $v_1 \approx v_2$ if and only if $\exists f \in \text{Aut}(D_n)$ such that $f(\Gamma) = \Gamma$ and $f(\varepsilon(v'_1)) = \varepsilon(v'_2)$. From this (i) follows. We refer to [CLP11] for a useful description of $\text{Aut}(D_n)$.

To prove (ii), we observe that for any $v \in HS(D_n; g', d)$ the orbit $\varepsilon(v) \cdot H_{2,\Gamma_v}(D_n)$ is either $\{\varepsilon(v)\}$ or $\{\varepsilon(v), -\varepsilon(v)\}$ (cf. Cor. 4.3). In the proof of (i) we list all possible normal forms for Hurwitz generating systems and we will see that, in the case where $H_{2,\Gamma_v}(D_n) \cong \mathbb{Z}/2\mathbb{Z} = \{\pm 1\}$, there exists $v' \in HS(D_n; g', d)$ with $\Gamma_{v'} = \Gamma_v$ and $\varepsilon(v') \cdot \varepsilon(v)^{-1} = -1$.

To prove (iii), observe that, given an admissible class function ν , and elements c_1, \dots, c_d which yield the given function ν , the product $c_1 \cdots c_d := c$ is in the commutator subgroup. However, in the dihedral group the commutator subgroup is equal to the set of commutators. Hence we may find a_1, b_1 such that $c^{-1} = [a_1, b_1]$, it suffices, if $c = x^{2\delta}$, to take $a_1 = x^m y, b_1 = x^{m+\delta} y$.

If some c_j is a reflection, without loss of generality, we may assume that $c_j = y$, hence, choosing $m = 1$, we obtain that

$$v := (c_1, \dots, c_d; xy, x^{1+\delta}y, 1, \dots, 1)$$

is a Hurwitz generating system. If $g' \geq 2$, we can just take the Hurwitz generating system

$$v := (c_1, \dots, c_d; y, x^\delta y, x, 1, \dots, 1).$$

In the case where $g' = 1$ and all the c_j are rotations, observe that

$$v := (c_1, \dots, c_d; x^m y, x^{m+\delta} y)$$

is a Hurwitz generating system if the dihedral group is generated by H (the subgroup generated by the c_j 's), together with $x^m y$ and x^δ . This amounts to the condition that H and x^δ generate the subgroup \mathcal{R} of rotations. Since $c = x^{2\delta} \in H$, we see that a necessary condition is that H has index at most two in the group \mathcal{R} of rotations, and a sufficient one is that $H = \mathcal{R}$. In the case where the index is exactly two, so that $n = 2h$ is even, the condition is that δ is an odd number. Observe that in this case if we replace some $c_j = x^{2i}$ by its inverse, the number δ is replaced by $\delta - 2i$, so that condition is indeed a condition on the class function ν .

Remark 5.2. For the case $g' = 0$ we defer to our previous article [CLP11], where all the normal forms were given.

But we can give a direct description as follows.

For $g' = 0$, $Im(\hat{\nu})$ is just the set of admissible class functions ν satisfying one of the following conditions:

(R) there are only two reflection classes, and the subgroup generated by the rotation classes is the whole \mathcal{R}

(O) n is odd and the class function ν takes value at least 4 on the reflection class

(E) n is even, there are at least four reflection classes and either the class function ν takes positive value on both reflection classes, or there exists a rotation class with odd exponent.

In fact, since we want a Hurwitz generating system, it is a necessary condition that there must be at least one reflection (and indeed an even number of reflections by admissibility). Take now an admissible class satisfying this restriction, and take c_1, \dots, c_d realizing the admissible class. Then their product c is in the commutator subgroup, so it is any rotation in case n is odd and a rotation of type x^{2i} in case n is even.

Since there is a reflection, we replace the last reflection, say c_i , by $c_i c^{-1}$, and obtain thus an admissible Hurwitz system without changing the class function.

We must however have an admissible Hurwitz generating system, and we recall that there is an even number of reflections among the c_i 's.

Up to automorphisms of D_n , we have reflections y and $x^m y$ in the components of the Hurwitz vector.

If we have only two reflections $y, x^m y$, we are done if and only if the subgroup generated by the rotations is the whole subgroup \mathcal{R} , since $x^{-m} \in \mathcal{R}$.

Assume that n is odd and there are 4 reflections: assume that the first four are $y, x^{m_1} y, x^{m_2} y, x^{m_3} y$: then we can replace these (without changing the class function) by the following reflections $y, xy, x^{m_2 - m_1 + 1} y, x^{m_3} y$ and we have obtained a Hurwitz generating system.

Assume that n is even and there are at least four reflections:

$$y, x^{m_1} y, x^{m_2} y, x^{m_3} y, \dots$$

We can change the rotations by adding to m_i an even integer $2h_i$, in such a way that the sum $-h_1 + h_2 - h_3 \dots \equiv 0$. Hence if some m_i is odd, we obtain y and xy , and we have a generating system. If instead all m_i 's are even, we obtain y and $x^2 y$, hence we are finally done if and only if there is a rotation with odd exponent.

Case 1: $\Gamma = \emptyset$ (the étale case). In this case $H_{2,\Gamma}(D_n) = H_2(D_n, \mathbb{Z})$, so $v \in HS(D_n)$ implies $\varepsilon(v) \in H_2(D_n, \mathbb{Z})$. In the following, we identify $H_2(D_n, \mathbb{Z}) \cong \mathbb{Z}/2\mathbb{Z} = \{0, 1\}$, when n is even. Then we have:

Proposition 5.3. *Let $n, g' \in \mathbb{N}$ with $n \geq 3$, $g' > 0$. Then, for any $v \in HS(D_n; g')$, we have:*

- (i) $v \approx (y, 1, x, 1, \dots, 1)$, if n is odd or if n is even and $\varepsilon(v) = 0$;

(ii) $v \approx (y, x^{n/2}, x, 1, \dots, 1)$, if n is even and $\varepsilon(v) = 1$.

Proof. Let

$$\bar{v} = v \pmod{\mathbb{Z}/n\mathbb{Z}} \in (\mathbb{Z}/2\mathbb{Z})^{2g'}.$$

Notice that $\bar{v} \in HS(\mathbb{Z}/2\mathbb{Z}; g')$. Since the parameter space for étale $\mathbb{Z}/2\mathbb{Z}$ -coverings of curves of a fixed genus is irreducible (see e.g. [DM] Lemma 5.16, or [BF86], or [Cor87], or [Cat10], Thm. 2.4), there exists $\varphi \in Map_{g'}$ such that $\varphi \cdot \bar{v} = (1, 0, \dots, 0)$. Hence

$$\varphi \cdot v = (x^{\ell_1}y, x^{m_1}, \dots, x^{\ell_{g'}}y, x^{m_{g'}}).$$

The condition $ev(\varphi \cdot v) = 1$ implies that $2m_1 = 0 \pmod{n}$. Hence $m_1 = 0$ or $m_1 = \frac{n}{2} \pmod{n}$.

In the first case, which is the only possible if n is odd,

$$\varphi \cdot v = (x^{\ell_1}y, 1, x^{\ell_2}, \dots, x^{\ell_{g'}}y, x^{m_{g'}}).$$

Consider now $v' := (x^{\ell_2}, x^{m_2}, \dots, x^{\ell_{g'}}y, x^{m_{g'}})$. As $v' \in HS(\mathbb{Z}/n\mathbb{Z}; g'-1)$, from the irreducibility of the parameter space of étale $\mathbb{Z}/n\mathbb{Z}$ -coverings of curves of a fixed genus we deduce that $\exists \varphi' \in Map_{g'-1}$ such that $\varphi' \cdot v' = (x^\lambda, 1, \dots, 1)$, with $(\lambda, n) = 1$ (see e.g. [DM] Lemma 5.16, or [BF86], or [Cor87], or [Cat10], Thm. 2.4). Now, from Proposition 6.3 it follows that $\exists \psi \in Map_{g'}$ such that

$$\psi \cdot v = (x^{\ell_1}y, 1, x^\lambda, 1, \dots, 1).$$

We obtain the normal form (i) after operating with $Aut(D_n)$. The fact that $\varepsilon(v) = 0$ follows from a standard computation (cf. Corollary 4.2). If $m_1 = \frac{n}{2} \pmod{n}$, we have two subcases: $\langle x^{\ell_2}, x^{m_2}, \dots, x^{\ell_{g'}}y, x^{m_{g'}} \rangle = \mathbb{Z}/n\mathbb{Z}$ (which is the case when $n/2$ is even), or $\langle x^{\ell_2}, x^{m_2}, \dots, x^{\ell_{g'}}y, x^{m_{g'}} \rangle = \langle x^2 \rangle$. Proceeding as in the case $m_1 = 0$ we reach the normal form (ii) in the first subcase, otherwise we obtain $(x^{\ell_1}y, x^{\frac{n}{2}}, x^2, 1, \dots, 1)$ in the latter subcase. In the latter case, consider the transformation

$$(7) \quad (a_1, b_1, a_2, b_2) \mapsto (a_2a_1, b_1, b_1a_2b_1^{-1}, a_2b_2a_2b_1^{-1}),$$

which is realized by Map_2 as it preserves the relation $\prod_1^2[\alpha_i, \beta_i] = 1$. Then extend it to $\Pi_{g'}$ using Proposition 6.3 and apply the transformation so obtained to $(x^{\ell_1}y, x^{\frac{n}{2}}, x^2, 1, \dots, 1)$. We obtain:

$$v \approx (x^{\ell_1+2}y, x^{\frac{n}{2}}, x^2, x^{4+\frac{n}{2}}, 1, \dots, 1).$$

Since $\mathbb{Z}/n\mathbb{Z} = \langle x^{\frac{n}{2}}, x^2 \rangle$, there exists $\psi \in Map_{g'}$ such that $\psi \cdot v \approx (x^{\ell_1+2}y, x^{\frac{n}{2}}, x, 1, \dots, 1)$, therefore we obtain the normal form (ii). In both of these subcases we have $\varepsilon(v) = 1$ (cf. Corollary 4.2). \square

Case 2: $\Gamma \neq \emptyset$ and contains some reflection.

Let $v = (c_1, \dots, c_d; a_1, b_1, \dots, a_{g'}, b_{g'})$ be a Hurwitz generating system such that $\{c_1, \dots, c_d\}$ contains some reflection, actually an even number because any product of commutators in D_n is a rotation. If n is odd, all the reflections belong to the same conjugacy class, while

when $n = 2k$ they are divided into two classes. Denote by ν_y (resp. ν_{xy}) the number of c_i 's in the class of y (resp. xy). As the pair (ν_y, ν_{xy}) is not $\text{Aut}(D_n)$ -invariant, we define ν_1, ν_2 by the property that $\{\nu_1, \nu_2\} = \{\nu_y, \nu_{xy}\}$, $\nu_1 \leq \nu_2$ (in [CLP11] we used the notation h for ν_1 , k for ν_2). Recall that, under the above hypotheses, $H_{2,\Gamma}(D_n) = \{0\}$ (Corollary 4.3). Indeed we prove that all the v 's with fixed g', d, n and ν -type are equivalent each other.

Proposition 5.4. *Let $n, g', d \in \mathbb{N}$ with $n \geq 3$, $g', d > 0$. Then, for any $v \in \text{HS}(D_n; g', d)$ such that Γ_v contains some reflection, we have:*

- (i) $v \approx (x^{\underline{r}}, \underbrace{x^{1-|\underline{r}|}y, xy, y, \dots, y}_{\nu_2}; x, 1, \dots, 1, 1)$, if n is odd;
- (ii) $v \approx (x^{\underline{r}}, \underbrace{x^{\varepsilon-|\underline{r}|}y, xy, \dots, xy}_{\nu_2}, \underbrace{y, \dots, y}_{\nu_1}; x, 1, \dots, 1, 1)$, if n is even.

Here $\underline{r} = (r_1, \dots, r_R)$, where $R + \nu_y = d$ in case (i), $R + \nu_y + \nu_{xy} = d$ in case (ii), $0 < r_i \leq r_{i+1} \leq \frac{n}{2}$, $x^{\underline{r}} = (x^{r_1}, \dots, x^{r_R})$, $|\underline{r}| = \sum r_i \pmod{n}$, $\{\nu_1, \nu_2\} = \{\nu_y, \nu_{xy}\}$, $\nu_1 \leq \nu_2$, $\varepsilon \in \{0, 1\}$, $\varepsilon + \nu_2 \equiv 1 \pmod{2}$.

The idea of the proof is the following. Using the action of the unpermuted mapping class group $\text{Map}^u(g', d+1)$ and the fact that at least one c_i is a reflection, we prove that $v \sim (\tilde{c}_1, \dots, \tilde{c}_d; \tilde{a}_1, \tilde{b}_1, \dots, \tilde{a}_{g'}, \tilde{b}_{g'})$, with $\tilde{a}_i, \tilde{b}_i \in \mathbb{Z}/n\mathbb{Z}$, for any i . We collect in the Appendix the relevant facts that will be used about the action of $\text{Map}^u(g', d+1)$ on the fundamental group. Then, using results about étale $\mathbb{Z}/n\mathbb{Z}$ -covers, we deduce that $v \approx v' := (c'_1, \dots, c'_d; x, 1, \dots, 1)$ (Lemma 5.5). At this point we can apply the main theorem of [CLP11] to deduce that, acting with the braid group, it is possible to transform v' to the corresponding normal form. However, we will see that using the entry x in v' , the results in the Appendix and Lemma 2.1 of [CLP11], we can transform directly v' in one of the above forms without using the normal forms for the $g' = 0$ case.

Lemma 5.5. *Let v be as in Proposition 5.4. Then*

$$v \approx v' := (c'_1, \dots, c'_d; x, 1, \dots, 1, 1).$$

Proof. Without loss of generality assume that c_d is a reflection (otherwise act with the braid group). Then, if a_1 is a reflection, by Proposition 6.2 (i), $\exists \varphi \in \text{Map}^u(g', d+1)$ such that

$$\varphi \cdot v = (c_1, \dots, c_{d-1}, (c_d a_1 b_1 a_1^{-1}) c_d (c_d a_1 b_1 a_1^{-1})^{-1}; c_d a_1, b_1, \dots, a_{g'}, b_{g'}).$$

While, if a_1 is a rotation and b_1 is a reflection, by Proposition 6.2 (ii) we have: $\exists \varphi \in \text{Map}^u(g', d+1)$ such that

$$\varphi \cdot v = (c_1, \dots, (c_d [a_1, b_1] a_1^{-1}) c_d (c_d [a_1, b_1] a_1^{-1})^{-1}; a_1, (a_1^{-1} c_d a_1) b_1, \dots, b_{g'}).$$

Notice that in both cases the d -th entry of $\varphi \cdot v$ is a reflection and that $c_d a_1, (a_1^{-1} c_d a_1) b_1 \in \mathbb{Z}/n\mathbb{Z}$. Proceeding in this way we get $\psi \in \text{Map}^u(g', d+1)$ such that $(\psi \cdot v)_i \in \mathbb{Z}/n\mathbb{Z}$, $i = d+1, \dots, 2g'$.

Next, by the main theorem in [Cat10], we conclude that $\psi \cdot v \approx (\tilde{c}_1, \dots, \tilde{c}_d; x^\alpha, 1, \dots, 1)$. We can further assume $(\alpha, n) = 1$. Otherwise, since $D_n = \langle x^\alpha, \tilde{c}_1, \dots, \tilde{c}_d \rangle$, there exists $x^\beta \in \langle \tilde{c}_1, \dots, \tilde{c}_d \rangle$ such that $\mathbb{Z}/n\mathbb{Z} = \langle x^{\alpha+\beta} \rangle$. Using Proposition 6.2 (i) and the braid group, we can multiply x^α by any element of $\langle \tilde{c}_1, \dots, \tilde{c}_d \rangle$. The claim now follows by applying $\text{Aut}(D_n)$. \square

We now complete the proof of Proposition 5.4. Let v' be as in Lemma 5.5 and let $2N$ be the number of reflections in $\{c'_1, \dots, c'_d\}$. Applying Lemma 2.1 of [CLP11] we have

$$(8) \quad v' \approx (x^{\underline{r}}, x^\beta y, x^\alpha y, x^{j_{N-1}} y, x^{j_{N-1}} y, \dots, x^{j_1} y, x^{j_1} y; x, 1, \dots, 1),$$

where $\underline{r} = (r_1, \dots, r_R)$, $0 < r_i \leq r_{i+1} \leq \frac{n}{2}$, $x^{\underline{r}} = (x^{r_1}, \dots, x^{r_R})$.

If $N = 1$ the result is clear. Otherwise we conjugate by x simultaneously the entries of each pair $(x^{j_k} y, x^{j_k} y)$ in (8) without changing the other components, hence we obtain:

$$(9) \quad v' \sim (x^{\underline{r}}, x^\beta y, x^\alpha y, x^{j_{N-1}+2\ell_{N-1}} y, x^{j_{N-1}+2\ell_{N-1}} y, \dots, \dots, x^{j_1+2\ell_1} y, x^{j_1+2\ell_1} y; x, 1, \dots, 1)$$

for any $\ell_1, \dots, \ell_{N-1} \in \mathbb{Z}$.

The equivalence (9) can be proven as follows. We have:

$$\begin{aligned} v' &\sim (x^{\underline{r}}, x^\beta y, \dots, x^{j_1} y, (x^{j_1} y x^{-1}) x^{j_1} y (x^{j_1} y x^{-1})^{-1}; x, x^{-1} x^{j_1} y x, 1, \dots, 1) \\ &\sim (x^{\underline{r}}, x^\beta y, \dots, (x^{-1}) x^{j_1} y(x), x^{j_1} y; x, x^{-1} x^{j_1} y x, 1, \dots, 1) \\ &\sim (x^{\underline{r}}, x^\beta y, \dots, (x^{-1}) x^{j_1} y(x), (x^{-1}) x^{j_1} y(x); x, 1, \dots, 1), \end{aligned}$$

where the first and the third equivalences are given by ξ -twists as in Proposition 6.2 (ii), while the second is a braid twist between the last two components. Iterating these steps we can conjugate by any power of x the entries of $(x^{j_1} y, x^{j_1} y)$ simultaneously. By Lemma 2.3 in [CLP11] we can move $(x^{j_k} y, x^{j_k} y)$ to the right and then conjugate its entries by any power of x as before. This proves (9).

If n is odd, choose ℓ_i in (9) such that $j_i + 2\ell_i = \alpha - 1 \pmod{n}$, then apply the automorphism $x^{\alpha-1} y \mapsto y, x \mapsto x$ to obtain (i).

Assume now that n is even. Without loss of generality we have that $\nu_1 = \nu_y \leq \nu_{xy} = \nu_2$ (otherwise apply $\text{Aut}(D_n)$). Assume further that $x^\beta y$ in (8) is conjugate to xy .

If $x^\alpha y$ is conjugate to xy , choose ℓ_i such that $j_i + 2\ell_i = \alpha$ or $j_i + 2\ell_i = \alpha - 1 \pmod{n}$, so (9) becomes:

$$v' \sim (x^{\underline{r}}, x^\beta y, x^\alpha y, x^\alpha y, \dots, x^\alpha y, x^{\alpha-1} y, \dots, x^{\alpha-1} y; x, 1, \dots, 1).$$

We obtain the normal form (ii) after applying the automorphism $x^\alpha y \mapsto xy, x \mapsto x$.

The remaining case, where $x^\alpha y$ is conjugate to y , is similar. \square

Notice that (9) follows also from Lemma 2.1 in [Kanev05] (see also [GHS02]), which applies to a more general situation. Since we don't

need the whole strength of that result, we preferred to give a complete proof in our case.

Case 3: $\Gamma \neq \emptyset$ and does not contain reflections.

We prove the following

Proposition 5.6. *Let $v, v' \in HS(D_n; g', d)$ with $\Gamma_v, \Gamma_{v'} \subset \mathbb{Z}/n\mathbb{Z}$. Then $v \approx v'$ if and only if there exists $f \in Aut(D_n)$ such that $f(\Gamma_v) = \Gamma_{v'}$ and $f(\varepsilon(v)) = \varepsilon(v')$.*

The “only if” part is clear. So assume $\Gamma_v = \Gamma_{v'} =: \Gamma$ and the existence of f as in the statement. We prove that $v \approx v'$. This is achieved after considering three cases: n is odd; $n = 2k$ and $x^k \in \Gamma$, $n = 2k$ and $x^k \notin \Gamma$. In the first two cases we determine a normal form, with respect to \approx , for each such element of $HS(D_n; g', d)$, and then we show that two such elements are equivalent if and only if they have the same normal form. Notice that in both cases $H_{2,\Gamma}(D_n) = \{0\}$ (Corollary 4.3). In the last case, for any such $v \in HS(D_n; g', d)$, we determine a normal form v' , with respect to the action of $Map(g', d)$ and then we show that $v_1 \approx v_2$ if and only if $\exists f \in Aut(D_n)$ such that $f(\Gamma) = \Gamma$ and $f(\varepsilon(v'_1)) = \varepsilon(v'_2)$. Notice that, in this case $H_{2,\Gamma}(D_n) \cong \mathbb{Z}/2\mathbb{Z}$ (Corollary 4.3).

Notation. Let $v = (c_1, \dots, c_d; a_1, b_1, \dots, a_{g'}, b_{g'}) \in D_n^{d+2g'}$ be a Hurwitz generating system such that $\Gamma_v \subset \mathbb{Z}/n\mathbb{Z}$, i.e. $c_i \in \mathbb{Z}/n\mathbb{Z}$, $\forall i$. We denote by $H = \langle c_1, \dots, c_d \rangle \subset D_n$ the subgroup generated by the c_i 's. Note that, under the above hypotheses, H is normal and it is contained in $\mathbb{Z}/n\mathbb{Z}$. Set $G' := D_n/H$. Then G' is a dihedral group D_m , $m \geq 3$, or is isomorphic to $\mathbb{Z}/2\mathbb{Z} \times \mathbb{Z}/2\mathbb{Z}$, or to $\mathbb{Z}/2\mathbb{Z}$.

Lemma 5.7. *Let $n \in \mathbb{N}$, $n \geq 3$ odd. Let $v \in HS(D_n; g', d)$ with $\Gamma_v \subset \mathbb{Z}/n\mathbb{Z}$. Then*

$$v \approx (x^{\underline{r}}; y, x^h, x, 1, \dots, 1),$$

where $\underline{r} = (r_1, \dots, r_d)$, $x^{\underline{r}} = (x^{r_1}, \dots, x^{r_d})$, $r_1 \leq \dots \leq r_d < \frac{n}{2}$ and $2h = \sum_1^d r_i \pmod{n}$.

Proof. Let us consider

$$\bar{v} := (\bar{a}_1, \bar{b}_1, \dots, \bar{a}_{g'}, \bar{b}_{g'}) \in HS(G'; g', 0),$$

where $\bar{a}_i = a_i \pmod{H}$, $\bar{b}_i = b_i \pmod{H}$. By Proposition 5.3 and by the analogous results for cyclic and $(\mathbb{Z}/2\mathbb{Z} \times \mathbb{Z}/2\mathbb{Z})$ -covers we have:

$$\exists \varphi \in Map_{g'} \quad \text{such that} \quad \varphi \cdot \bar{v} = (\bar{y}, 1, \bar{x}, 1, \dots, 1).$$

By Proposition 6.3, $\exists \tilde{\varphi} \in Map(g', d)$ with

$$\tilde{\varphi} \cdot v = (c_1, \dots, c_d; x^{\ell_1} y, x^{m_1}, x^{\ell_2}, \dots, x^{m_{g'}}),$$

where $x^{m_i} \in H$, $\forall i$, $x^{\ell_2} = x \pmod{H}$ and $x^{\ell_i} \in H$, $\forall i > 2$.

We now apply the ξ -twists as in Proposition 6.2 (i) with $\ell = 2, 3, \dots, g'$ and we deduce that we can multiply all the x^{ℓ_i} , $i > 1$, by any element of H . Hence $\exists \psi \in \text{Map}^u(g', d+1)$ such that

$$\psi \cdot \tilde{\varphi} \cdot v = (c_1, \dots, c_d; x^{\ell_1}y, x^{m_1}, x, x^{m_2}, 1, x^{m_3}, \dots, 1, x^{m_{g'}}).$$

Similarly, using Proposition 6.2 (ii), we get:

$$v \approx (c_1, \dots, c_d; x^{\ell_1}y, x^{m_1}, x, 1, \dots, 1, 1).$$

Now, for any $i = 1, \dots, d$, consider $c_i = x^{s_i}$. If $s_i < \frac{n}{2}$, set $r_i = s_i$, otherwise use the braid group to move c_i to the d -th position and then apply Proposition 6.2 (ii) with $\ell = 1$. After this, c_i becomes $c_i^{-1} = x^{n-s_i}$, then set $r_i = n - s_i$. Finally, using the braid group, we can order the c_i 's such that $r_i \leq r_{i+1}$.

So, we have proved that

$$v \sim (x^{\underline{r}}; x^{\lambda_1}y, x^{\mu_1}, x, 1, \dots, 1),$$

with $r_1 \leq \dots \leq r_d < \frac{n}{2}$. Now the condition $ev(v) = 1$ implies that $2\mu_1 = \sum_1^d r_i \pmod{n}$, therefore set $h := \mu_1 \pmod{n}$.

We reach the normal form after applying the automorphism $x^{\lambda_1}y \mapsto y$, $x \mapsto x$. \square

Lemma 5.8. *Let $n = 2k \in \mathbb{N}$ and let $v \in HS(D_n; g', d)$ with $x^k \in \Gamma_v \subset \mathbb{Z}/n\mathbb{Z}$. Then we have:*

$$v \approx (x^{\underline{r}}; y, x^h, x, 1, \dots, 1),$$

where $\underline{r} = (r_1, \dots, r_d)$, $x^{\underline{r}} = (x^{r_1}, \dots, x^{r_d})$, $r_1 \leq \dots \leq r_d = k$, $2h = \sum_1^d r_i \pmod{n}$ and $h < k$.

Proof. Proceeding as in the proof of the previous lemma, we have:

$$\exists \varphi \in \text{Map}(g', d) \quad \text{such that} \quad \varphi \cdot v = (c_1, \dots, c_d; x^{\ell_1}y, x^{m_1}, x^{\ell_2}, \dots, x^{m_{g'}}),$$

where $x^{m_i} \in H$, $\forall i > 1$, $x^{\ell_2} = x \pmod{H}$ and $x^{\ell_i} \in H$, $\forall i > 2$.

Since we can multiply all the x^{ℓ_i} , $i > 1$, by any element of H (apply Proposition 6.2 (i) with $\ell = 2, 3, \dots, g'$), we have: $\exists \psi \in \text{Map}^u(g', d+1)$ such that

$$\psi \cdot \varphi \cdot v = (c_1, \dots, c_d; x^{\ell_1}y, x^{m_1}, x, x^{m_2}, 1, x^{m_3}, \dots, 1, x^{m_{g'}}).$$

Similarly, using Proposition 6.2 (ii), we get:

$$v \approx (c_1, \dots, c_d; x^{\ell_1}y, x^{m_1}, x, 1, \dots, 1, 1).$$

Now, for any $i = 1, \dots, d$, consider $c_i = x^{s_i}$. If $s_i \leq k$, set $r_i = s_i$, otherwise use the braid group to move c_i to the d -th position and then apply Proposition 6.2 (ii) with $\ell = 1$. In this way c_i becomes c_i^{-1} and so set $r_i = 2k - s_i$.

So, we have proved that

$$v \approx (x^{\underline{r}}; x^{\lambda_1}y, x^{\mu_1}, x, 1, \dots, 1),$$

with $r_i \leq k$, $\forall i$. Now the condition $ev(v) = 1$ implies that $2\mu_1 = \sum_1^d r_i \pmod{n}$. If $\mu_1 < k$, set $h = \mu_1$. Otherwise, apply braid group transformations to achieve the ordering $r_i \leq r_{i+1}$, $\forall i \leq d-1$. By hypotheses $r_d = k$ and we apply Proposition 6.2 with $\ell = 1$. Since x^k is central, this operation does not change r_d , while x^{μ_1} becomes x^{μ_1+k} . Set $h = \mu_1 + k \pmod{n}$.

Finally apply the appropriate element of $Aut(D_n)$ to reach the normal form. \square

We now consider the last case.

Lemma 5.9. *Let $n = 2k \in \mathbb{N}$ and let $v \in HS(D_n; g', d)$ with $\Gamma \subset \mathbb{Z}/n\mathbb{Z} \setminus \{x^k\}$. Then we have:*

- (i) $v \approx v' := (x^{\underline{r}}; y, x^h, x, 1, \dots, 1)$, where $\underline{r} = (r_1, \dots, r_d)$, $x^{\underline{r}} = (x^{r_1}, \dots, x^{r_d})$, $r_1 \leq \dots \leq r_d < k$, $2h = \sum_1^d r_i \pmod{n}$;
- (ii) let $v'_1 = (x^{\underline{r}}; y, x^h, x, 1, \dots, 1)$ and $v'_2 = (x^{\underline{r}}; y, x^{h+k}, x, 1, \dots, 1)$, then $\varepsilon(v'_1) \neq \varepsilon(v'_2) \in (D_n)_\Gamma$;
- (iii) $v'_1 \approx v'_2$ if and only if $\exists f \in Aut(D_n)$ such that $f(\Gamma) = \Gamma$ and $f(\varepsilon(v'_1)) = \varepsilon(v'_2)$.

Proof. The proof of (i) is the same as that of the previous lemma. Since in this case $x^k \notin \Gamma$, we can not achieve $h \leq k$.

To prove (ii) recall that $\varepsilon(v) := ev(\widehat{v}) \in (D_n)_\Gamma$. So, if $ev(\widehat{v}'_1) = ev(\widehat{v}'_2)$, then $ev(\widehat{v}'_2)^{-1} \cdot ev(\widehat{v}'_1) = 0 \in H_{2,\Gamma}(D_n)$. But now a direct computation shows that $ev(\widehat{v}'_2)^{-1} \cdot ev(\widehat{v}'_1) \neq 0$ (Corollary 4.2), a contradiction.

(iii) The ‘‘only if’’ part is clear. So, assume that $\exists f \in Aut(D_n)$ such that $f(\Gamma) = \Gamma$ and $f(\varepsilon(v'_1)) = \varepsilon(v'_2)$. Since $f(\varepsilon(v'_1)) = \varepsilon(f(v'_1))$, we have $\varepsilon(f(v'_1)) = \varepsilon(v'_2)$ and so v'_2 and $f(v'_1)$ have the same ν -type (Remark 3.10). From (i) and (ii) we deduce that

$$f(v'_1) \approx (x^{\underline{r}}; x^{\lambda_1}y, x^{h+k}, x, 1, \dots, 1).$$

Hence, using the automorphism $x^{\lambda_1}y \mapsto y$, $x \mapsto x$, we have that $f(v'_1) \approx v'_2$ and so the claim follows. \square

6. APPENDIX A: AUTOMORPHISMS OF SURFACE-GROUPS

We collect in this Appendix some facts about mapping class groups and their action on fundamental groups. They should be well known to experts, we include them here for completeness.

Let Y be a compact Riemann surface of genus g' and let $\mathcal{B} = \{y_1, \dots, y_d\} \subset Y$ be a finite subset of cardinality d . After the choice of a geometric basis of $Y \setminus \mathcal{B}$, we have the following presentation of the fundamental group:

$$\begin{aligned} \pi_1(Y \setminus \mathcal{B}, y_0) = \\ \langle \gamma_1, \dots, \gamma_d, \alpha_1, \beta_1, \dots, \alpha_{g'}, \beta_{g'} \mid \gamma_1 \cdot \dots \cdot \gamma_d \cdot \prod_{i=1}^{g'} [\alpha_i, \beta_i] = 1 \rangle. \end{aligned}$$

Following [Bir69], there is a short exact sequence
(10)


$$1 \rightarrow \pi_1(Y \setminus \mathcal{B}, y_0) \xrightarrow{\Xi} \text{Map}^u(Y, \{y_0, y_1, \dots, y_d\}) \rightarrow \text{Map}^u(Y, \mathcal{B}) \rightarrow 1$$

which induces an injective group homomorphism

$$\text{Map}^u(Y, \mathcal{B}) \rightarrow \text{Out}(\pi_1(Y \setminus \mathcal{B}, y_0))$$

([Bir69], Thm. 4). The map Ξ is defined as follows. Let $[c] \in \pi_1(Y \setminus \mathcal{B}, y_0)$ be an element of the geometric basis and let $c: [0, 2\pi] \rightarrow Y \setminus \mathcal{B}$ be a simple, smooth loop based at y_0 , representing $[c]$. Let $\Xi([c])$ be the isotopy class of the ξ -twist, ξ_c . Then extend Ξ to the whole group as an homomorphism. Recall that the ξ -twist, ξ_c , can be defined as follows. Let $N \subset Y \setminus \mathcal{B}$ be a tubular neighborhood of c and let $e: A \rightarrow N$ be a diffeomorphism between the annulus $A = \{z = re^{i\theta} \in \mathbb{C} \mid 1 \leq r \leq 2\}$ and N such that $e(\frac{3}{2}, \theta) = c(\theta)$. Define $h: A \rightarrow A$ as follows

$$h(r, \theta) = \begin{cases} (r, \theta + 4\pi(r - 1)), & 1 \leq r \leq \frac{3}{2}; \\ (r, \theta + 4\pi(2 - r)), & \frac{3}{2} \leq r \leq 2. \end{cases}$$



Then h is a diffeomorphism which is the identity when $r = 1, \frac{3}{2}, 2$. Finally, define $\xi_c: Y \rightarrow Y$ as the identity on $Y \setminus N$ and as $e \circ h \circ e^{-1}$ on N .

From the sequence (10), it follows that $\pi_1(Y \setminus \mathcal{B}, y_0)$ is isomorphic through Ξ to a normal subgroup of $\text{Map}^u(Y, \{y_0, y_1, \dots, y_d\})$, hence we get an action by conjugation of $\text{Map}^u(Y, \{y_0, y_1, \dots, y_d\})$ on $\pi_1(Y \setminus \mathcal{B}, y_0)$:

$$[f] \cdot [\xi_c] = [f \circ \xi_c \circ f^{-1}].$$

We have:

Lemma 6.1. *For any $[f] \in \text{Map}^u(Y, \{y_0, y_1, \dots, y_d\})$ and $[c] \in \pi_1(Y \setminus \mathcal{B}, y_0)$, we have:*

$$[f] \cdot [\xi_c] = [\xi_{f\#(c)}],$$

where $f\#(c)(\theta) = (f \circ c)(\theta)$.

Proof. $f \circ \xi_c \circ f^{-1} = (f \circ e) \circ h \circ (f \circ e)^{-1}$ on N , and coincides with the identity on $Y \setminus N$. The result then follows because $f \circ e: A \rightarrow Y$ is a tubular neighborhood of $f_{\#}(c)$. \square

One can define, in the same way, ξ -twists with respect to loops that are not based at y_0 and Lemma 6.1 is still valid. In the following result we give the action of ξ -twists around certain loops in terms of a given geometric basis of $\pi_1(Y \setminus \mathcal{B}, y_0)$.

Proposition 6.2. *Let $\gamma_1, \dots, \gamma_d, \alpha_1, \beta_1, \dots, \alpha_{g'}, \beta_{g'}$ be a fixed geometric basis of $\pi_1(Y \setminus \mathcal{B}, y_0)$.*

- (i) *Let $c \subset Y \setminus \mathcal{B}$ be the loop in Figure 1, image of the two sides of the angle inside the polygon with vertex y_d . Set $u = \prod_{k=1}^{\ell-1} [\alpha_k, \beta_k]$. Then we have:*

$$\begin{aligned} (\xi_c)_*(\alpha_\ell) &= u^{-1} \gamma_d u \alpha_\ell; \\ (\xi_c)_*(\gamma_d) &= (\gamma_d u \alpha_\ell \beta_\ell \alpha_\ell^{-1} u^{-1}) \gamma_d (\gamma_d u \alpha_\ell \beta_\ell \alpha_\ell^{-1} u^{-1})^{-1}; \end{aligned}$$

$$(\xi_c)_*(\alpha_i) = \alpha_i \ (i \neq \ell), \quad (\xi_c)_*(\beta_i) = \beta_i \ (\forall i), \quad (\xi_c)_*(\gamma_j) = \gamma_j \ (j \neq d).$$

- (ii) *Let $c \subset Y \setminus \mathcal{B}$ be the loop in Figure 2, image of the two sides of the angle inside the polygon with vertex y_d . Set $u = \prod_{k=1}^{\ell-1} [\alpha_k, \beta_k]$. Then we have:*

$$\begin{aligned} (\xi_c)_*(\beta_\ell) &= \alpha_\ell^{-1} u^{-1} \gamma_d u \alpha_\ell \beta_\ell; \\ (\xi_c)_*(\gamma_d) &= (\gamma_d u [\alpha_\ell, \beta_\ell] \alpha_\ell^{-1} u^{-1}) \gamma_d (\gamma_d u [\alpha_\ell, \beta_\ell] \alpha_\ell^{-1} u^{-1})^{-1}; \end{aligned}$$

$$(\xi_c)_*(\beta_i) = \beta_i \ (i \neq \ell), \quad (\xi_c)_*(\alpha_i) = \alpha_i \ (\forall i), \quad (\xi_c)_*(\gamma_j) = \gamma_j \ (j \neq d).$$

Proof. (i) The image of α_ℓ under ξ_c is drawn in Figure 3. From this it follows the formula for $(\xi_c)_*(\alpha_\ell)$. Since ξ_c is the identity outside a small tubular neighborhood of c , we have that $(\xi_c)_*(\alpha_i) = \alpha_i$ ($i \neq \ell$), $(\xi_c)_*(\beta_i) = \beta_i$ ($\forall i$) and $(\xi_c)_*(\gamma_j) = \gamma_j$ ($j \neq d$). The formula for $(\xi_c)_*(\gamma_d)$ is now a consequence of $\gamma_1 \cdot \dots \cdot \gamma_d \prod_{i=1}^{g'} [\alpha_i, \beta_i] = 1$, since the product $\gamma_1 \cdot \dots \cdot \gamma_d \prod_{i=1}^{g'} [\alpha_i, \beta_i]$ must be left fixed.

The proof of (ii) is similar. \square




Figure 1.




Figure 2.




Figure 3.

Proposition 6.3. *Let $\Pi_{g'} = \langle \alpha_1, \dots, \beta_{g'} \mid \prod_1^{g'} [\alpha_i, \beta_i] \rangle$ and $\Pi_{g'-1} = \langle \alpha_2, \dots, \beta_{g'} \mid \prod_2^{g'} [\alpha_i, \beta_i] \rangle$. Then, for any $\varphi \in \text{Aut}^0(\Pi_{g'-1})$, there exists $\psi \in \text{Aut}^0(\Pi_{g'})$ and $\delta \in \Pi_{g'}$ such that $\psi(\alpha_1) = \alpha_1$, $\psi(\beta_1) = \beta_1$, $\psi(\alpha_i) = \delta\varphi(\alpha_i)\delta^{-1}$, $\psi(\beta_i) = \delta\varphi(\beta_i)\delta^{-1}$, $i > 1$.*

Proof. We first extend φ to an automorphism

$$\tilde{\varphi} \in \text{Aut} \left(\langle \alpha_2, \dots, \beta_{g'}, \gamma \mid \gamma \cdot \prod_2^{g'} [\alpha_i, \beta_i] \rangle \right)$$

such that $\tilde{\varphi}(\alpha_i) = \varphi(\alpha_i)$, $\tilde{\varphi}(\beta_i) = \varphi(\beta_i)$ and $\tilde{\varphi}(\gamma) = \delta^{-1}\gamma\delta$, $i > 1$. Geometrically this corresponds to representing φ as composition of Dehn twists along curves contained in the complement $Y_{g'-1} \setminus D$ of a closed disk D in a Riemann surface $Y_{g'-1}$ of genus $g' - 1$, where D does not intersect α_i and β_i .

Now simply define $\psi(\alpha_1) = \alpha_1$, $\psi(\beta_1) = \beta_1$, $\psi(\alpha_i) = \delta\tilde{\varphi}(\alpha_i)\delta^{-1}$ and $\psi(\beta_i) = \delta\tilde{\varphi}(\beta_i)\delta^{-1}$, $i > 1$. \square

7. APPENDIX B: LOCI AND TOPOLOGICAL TYPES

In this appendix we show that, up to essentially only one exception, the loci $M_{g,\rho}(D_n)$ are in bijection with the unmarked topological types of D_n -actions.

In general, thanks to the work of Singerman, Ries, and Magaard-Shaska-Shpectorov-Völklein (see [Sing72], [Ries93] and [MSSV01]), this is true more generally for the irreducible components of the loci $M_g(G)$; with a finite number of exceptions, they correspond bijectively to the unmarked topological types of G actions. That there are indeed groups G and different unmarked topological types of G actions which yield the same component was already shown by Ries ([Ries93]).

To have a simple notation, assume in this appendix that H, H' are distinct finite subgroups of the mapping class group Map_g , and denote by $Z := Fix(H)$, $Z' := Fix(H')$ the corresponding irreducible analytic subsets of Teichmüller space \mathcal{T}_g . We already observed that, if we denote by

$$\rho : H \rightarrow Map_g, \rho' : H' \rightarrow Map_g$$

the inclusion homomorphisms, these analytic subsets map to irreducible closed algebraic sets $M_{g,\rho}(H)$, respectively $M_{g,\rho'}(H')$.

We define the generic group of automorphisms of a curve in Z as

$$G := G_H := \bigcap_{C \in Z} Stab_C \quad (Aut(C) \cong Stab_C \subset Map_g).$$

We refer to lemma 4.1 of [MSSV01] for the proof of the following result.

Theorem 7.1. (MSSV) *Suppose $H \subset G$ and Z are as above, with H a proper subgroup of G and $C \in Z$. Then*

$$\delta := \dim(Z) \leq 3.$$

I) if $\delta = 3$, then H has index 2 in G , and $C \rightarrow C/G$ is covering of \mathbb{P}^1 branched on six points, P_1, \dots, P_6 , and with branching indices all equal to 2. Moreover the subgroup H corresponds to the unique genus two double cover of \mathbb{P}^1 branched on the six points, P_1, \dots, P_6 (by Galois theory, intermediate covers correspond to subgroups of G bijectively).

II) If $\delta = 2$, then H has index 2 in G , and $C \rightarrow C/G$ is covering of \mathbb{P}^1 branched on five points, P_1, \dots, P_5 , and with branching indices $2, 2, 2, 2, c_5$. Moreover the subgroup H corresponds to a genus one double cover of \mathbb{P}^1 branched on four of the points P_1, \dots, P_4, P_5 which have branching index 2.

III) If $\delta = 1$, then there are three possibilities.

III-a) H has index 2 in G , and $C \rightarrow C/G$ is covering of \mathbb{P}^1 branched on four points, P_1, \dots, P_4 , with branching indices $2, 2, 2, 2d_4$, where

$d_4 > 1$. Moreover the subgroup H corresponds to the unique genus one double cover of \mathbb{P}^1 branched on the four points, P_1, \dots, P_4 .

III-b) H has index 2 in G , and $C \rightarrow C/G$ is covering of \mathbb{P}^1 branched on four points, P_1, \dots, P_4 , with branching indices $2, 2, c_3, c_4$, where $c_3 \leq c_4 > 2$. Moreover the subgroup H corresponds to a genus zero double cover of \mathbb{P}^1 branched on two points whose branching index equals 2.

III-c) H is normal in G , $G/H \cong (\mathbb{Z}/2)^2$, moreover $C \rightarrow C/G$ is covering of \mathbb{P}^1 branched on four points, P_1, \dots, P_4 , with branching indices $2, 2, 2, c_4$, where $c_4 > 2$. Moreover the subgroup H corresponds to the unique genus zero cover of \mathbb{P}^1 with group $(\mathbb{Z}/2)^2$ branched on the three points P_1, P_2, P_3 whose branching index equals 2.

Corollary 7.2. *Assume that*

$$(*) \quad Z := \text{Fix}(H) = Z' := \text{Fix}(H'),$$

where $H \neq H'$ (hence, in the previous notation, H is a proper subgroup of G).

Then $\delta := \dim(Z) \leq 2$. Moreover, if $\delta = 2$, then necessarily all the five branching indices are equal to 2 ($c_5 = 2$).

Assume further that H, H' have the same cardinality.

Then $G := G_H := \bigcap_{C \in Z} \text{Stab}_C$ induces a sequence of coverings

$$C \rightarrow C/H \rightarrow C/G$$

which is not of type III-c), and, if it is type III-b), then the branching indices $2, 2, c_3, c_4$ must satisfy $c_3 = 2$.

If $C \rightarrow C/H \rightarrow C/G$ is of type III-a), then $C \rightarrow C/H' \rightarrow C/G$ is of type III-b) with branching indices $2, 2, 2, 2d_4$.

If moreover $\delta = 0$, and $H \cong H' \cong D_n$, then necessarily Z and Z' are distinct points.

Proof. We can apply the previous theorem to both H and H' , which are proper subgroups of the same group G .

In the case I) where $\delta = 3$, then both H and H' correspond to the unique genus two cover branched on the six points, contradicting $H \neq H'$.

The same argument applies in case II) when $c_5 > 2$.

In the case where the covering $C \rightarrow C/H \rightarrow C/G$ is of type III-c), then the index of H is four and, since $|H| = |H'|$, the index of H' is also four, hence H' corresponds to the unique genus zero $(\mathbb{Z}/2)^2$ -cover branched on the three points P_1, P_2, P_3 ; again we obtain the contradiction $H = H'$.

If the sequence of coverings $C \rightarrow C/H \rightarrow C/G$ is of type III-b), then the intermediate genus zero covering is unique unless $c_3 = 2$.

If the type is III-a), the type for H' cannot be the same, since we have a unique intermediate genus one cover. Since $\delta = 1$, the index of H' is also equal to two, and the branching indices are $2, 2, 2, 2d_4$ the type of H' must be III-b).

Finally, if $\delta = 0$ and $H \cong H' \cong D_n$, we have a dihedral covering of \mathbb{P}^1 branched in three points, for which there is a unique topological type (see [CLP11]), corresponding to the monodromy factors y, yx, x^{-1} . \square

Remark 7.3. In the case where $H \cong H' \cong D_n$, and $H \neq H'$ we have then $\delta = 1$ or $\delta = 2$. Moreover, both H, H' have index in G equal to 2.

If $\delta = 2$, then H, H' are both of type II), hence they correspond to intermediate genus one covers branched on four of the five points P_1, P_2, \dots, P_5 .

If $\delta = 1$, either H, H' are both of type III-b) (with branching indices $2, 2, 2, c_4$) hence they correspond to intermediate genus zero covers branched on two of the three points P_1, P_2, P_3 ; or, up to exchanging H with H' , H is of type III-a) and H' is of type III-b) with branching indices $2, 2, 2, 2d_4$.

The investigation and detailed classification of these coincidences ($Z = Z'$ and $H \cong H' \cong D_n$, with $H \neq H'$) is interesting, but shall not be fully pursued here. We shall limit ourselves to show that it must be $\delta = 1$, and that there is only one exception, namely, the one where one group is of type III-a) and the other of type III-b).

A common feature of all the cases is however the following situation. Defining $K = H \cap H'$, we have an exact sequence

$$(DD) \quad 1 \rightarrow K \rightarrow G \rightarrow (\mathbb{Z}/2)^2 \rightarrow 1.$$

Proposition 7.4. *Assume that we have an exact sequence (DD) where $H \cong H' \cong D_n$. Then there are elements γ_1, γ_2 with $\gamma_1^2 = \gamma_2^2 = 1$, whose image generate $(\mathbb{Z}/2)^2$, and such that*

$$H = \langle K, \gamma_1 \rangle, \quad H' = \langle K, \gamma_2 \rangle.$$

Moreover (DD) splits if and only if $G \cong D_n \times \mathbb{Z}/2$, H corresponds to the subgroup $D_n \times \{0\}$, and H' is the graph of a homomorphism $\phi : D_n \rightarrow \mathbb{Z}/2$ with Kernel equal to K .

If (DD) does not split, then n is even and either $G \cong D_{2n}$ or $n = 4h$, where h is odd, and G is the semidirect product of $H \cong D_n$ with $\langle \gamma_2 \rangle \cong \mathbb{Z}/2$, such that conjugation by γ_2 acts as follows:

$$y \mapsto yx^2, \quad x \mapsto x^{2h-1}.$$

Remark 7.5. The condition that (DD) splits always holds, except possibly if H, H' are of different types III-a) and III-b).

Indeed if $H, H' \cong D_n$ are both of type II), then the Hurwitz generating systems of $C \rightarrow C/H, C \rightarrow C/H'$ must be of the form

$$v = (c_1, c_2; a, b), \quad v' = (c'_1, c'_2; a', b') \in HS(D_n; 1, 2),$$

where c_2 is conjugate to c_1 (resp. c'_2 is conjugate to c'_1) in G and $c_i^2 = (c'_i)^2 = 1, i = 1, 2$. From Theorem 7.1 it follows that $c_1, c_2, c'_1, c'_2 \notin$

$K = H \cap H'$, therefore c_1, c_2 (resp. c'_1, c'_2) are reflections in H (resp. H').

Since otherwise one of them must be the unique central element of H (resp. of H'); but the unique central element of H is then central in G (H being normal in G), hence it equals the unique central element of H' by the same argument: hence it lies in K and we derive a contradiction.

From this we deduce that the numerical type of v is the same as that of v' , up to automorphisms, and hence using Proposition 5.4 it follows that H and H' have the same unmarked topological type.

Finally, assume that $H, H' \cong D_n$ are both of type III-b). Let $v = (v_1, v_2, v_3, v_4) \in HS(G; 0, 4)$ be the Hurwitz generating system of $C \rightarrow C/G$. By Corollary 7.2 we have:

$$v_1^2 = v_2^2 = v_3^2 = 1 \neq v_4^2.$$

Moreover, by Theorem 7.1, $v_4 \in H \cap H' = K$. Since $v_4^2 \neq 1$, v_4 is a rotation in H and in H' . Now consider the Hurwitz systems of $C \rightarrow C/H$ and of $C \rightarrow C/H'$, which are of the following forms:

$$u = (u_1, u_2, v_4, v'_4), \quad u' = (u'_1, u'_2, v_4, v''_4),$$

where v'_4 is conjugate to v_4 in G . Notice that each of them contains only two reflections, hence by Remark 5.2 (R) v_4 must generate the subgroup of rotations in H (resp. in H'). But $v_4 \in K$ and the subgroup of rotations of K have index 2 in the subgroup of rotations of H (resp. of H') if G is not the direct product of D_n with $\mathbb{Z}/2$.

Proof. (of prop. 7.4).

K is an index two subgroup in D_n . If n is odd, then necessarily $K = \mathcal{R}$, the subgroup of rotations.

1) More generally, if $K = \mathcal{R}$, any element in $H \setminus K$ has order two and splits the exact sequence $1 \rightarrow K \rightarrow H \rightarrow (\mathbb{Z}/2) \rightarrow 1$, and the same holds for H' . Therefore we find elements γ_1, γ_2 as desired, and such that their conjugation action on K is the same, $x^i \mapsto x^{-i}$.

Now, $\gamma := \gamma_1\gamma_2$ centralizes $K = \mathcal{R}$ and $\gamma^2 \in K$. Let x be generator of \mathcal{R} . Hence

$$\gamma_1\gamma_2\gamma_1\gamma_2 = x^r.$$

Replacing γ_2 with γ_2x^a , we replace r by $r + 2a$. If n is odd, then we may assume $r = 0$, whereas if n is even, we also have the case $r = 1$. Observe that $r = 0$ implies the splitting of the above sequence (DD), moreover γ centralizes K .

2) Assume that (DD) splits and $\gamma = \gamma_1\gamma_2$ centralizes K . Then

$$\gamma_1\gamma_2\gamma_1\gamma_2 = 1 \leftrightarrow \gamma\gamma_1\gamma = \gamma_1,$$

which amounts to γ being in the centre of G . Hence

$$G \cong H \times \langle \gamma \rangle \cong H \times \mathbb{Z}/2.$$

Moreover, $H' = \langle K, \gamma_2 \rangle = \langle K, \gamma_1\gamma \rangle$ which proves our assertion.

3) Assume now that $K = \mathcal{R}$ and that (DD) does not split: then we can only achieve $\gamma^2 = x$. Observe now that

$$\gamma_1\gamma\gamma_1 = \gamma_2\gamma_1 = \gamma^{-1}, \gamma^i = 1 \leftrightarrow 2n|i.$$

Since G is generated by γ, γ_1 , it follows that $G \cong D_{4m}$, $n = 2m$.

4) Assume now that $n = 2m$ is even and $K \neq \mathcal{R}$. Then $K \cong D_m$, and K is generated, up to an automorphism, by x^2 and y . The element $z := yx$ has order two and its action by conjugation is

$$x^2 \mapsto x^{-2}, y \mapsto yx^2.$$

In this way we construct also in this case the desired elements γ_1, γ_2 such that $\gamma := \gamma_1\gamma_2$ centralizes K .

Now, γ^2 is in the centre of D_m , hence $\gamma^2 = 1$ if m is odd, or, if $m = 2h$, then $\gamma^2 = x^{2h} = x^m$.

If $\gamma^2 = 1$ we have the splitting and we can apply 2).

Otherwise, if h is even, replace γ_2 by γ_2x^h , which has again order equal to two. Then γ is replaced by $\xi := \gamma_1\gamma_2x^h$. ξ has order two, since

$$\xi^2 = \gamma_1\gamma_2x^h\gamma_1\gamma_2x^h = \gamma_1\gamma_2\gamma_1\gamma_2x^{2h} = \gamma^2x^m = x^{2m} = 1.$$

Observe then that $\gamma_1\gamma_2 = \gamma_2\gamma_1x^m$.

We show that then ξ is in the centre:

$$\gamma_1\xi\gamma_1 := \gamma_1\gamma_1\gamma_2x^h\gamma_1 = \gamma_2\gamma_1x^{-h} = \gamma_1\gamma_2x^{-m-h} = \gamma_1\gamma_2x^h = \xi,$$

$$\gamma_2\xi\gamma_2 := \gamma_2\gamma_1\gamma_2x^h\gamma_2 = \gamma_2\gamma_2\gamma_1x^{m+h}\gamma_2 = \gamma_1x^{-h}\gamma_2 = \gamma_1\gamma_2x^h = \xi.$$

Assume instead that h is odd. Then we observe simply that G is generated by H and by γ_2 , and we set $x := y\gamma_1 \in H$, so that H is generated in the standard way by y, x .

We have $\gamma_2^2 = 1$, moreover conjugation by γ_2 sends

$$x^2 \mapsto x^{-2}, y \mapsto yx^2, \gamma_1 \mapsto \gamma_1x^{2h},$$

since $\gamma_1\gamma_2\gamma_1\gamma_2 = x^{2h}$.

We conclude that

$$x := y\gamma_1 \mapsto yx^2\gamma_1x^{2h} = yx^2yx^{2h} = x^{2h-1}.$$

□

Proposition 7.6. *The case $\delta = 2$ cannot occur for the group D_n .*

Proof.

We have already shown that in case II) the five branching indices must all be equal to 2.

By 7.4 and 7.5 the group G generated by H, H' must be $G = D_n \times C_2$, where $C_2 := \mathbb{Z}/2$, H is the subgroup $D_n \times \{0\}$, while H' is the graph of a homomorphism of H onto C_2 .

We consider first the case where four of the five elements have a reflexion component. To have $(y, 0)$ in the group they generate, the second component of one of these four elements must be trivial. We permute the other three into the first three positions, apply Lemma 2.1

of [CLP11] to make the first two equal. So after a suitable automorphism of D_n the tuple is

$$(y, 1)(y, 1)(yx^\ell, 1)(yx^{\ell+m}, 0)(x^m, 1)$$

where m is an integer in $\{0, \frac{n}{2}\}$.

In case n even and $m = n/2$ we may apply the following automorphism of G , $(x, 0) \mapsto (x, 0)$, $(y, 0) \mapsto (yx^m, 0)$, $(e, 1) \mapsto (x^m, 1)$.

So we are left with the case $m = 0$, where we can apply another automorphism to get

$$(y, 1)(y, 1)(yx, 1)(yx, 0)(e, 1).$$

Since only one of these elements is in H' , and $H' \neq H$, H' is the graph of the homomorphism which sends y to 0 and x to 1, in particular n must be even.

On the two intermediate covers, which are elliptic curves, we get the respective Hurwitz vectors $(yx, yx; x, 1)$ and $(yx', yx'; x', 1)$, where x' is a shorthand for $(x, 1) \in H'$. Obviously they are the same under the automorphism of G which keeps $(y, 0)$ and $(y, 1)$ fixed and sends $(x, 0)$ to $(x, 1)$. This means that $H = H'$, a contradiction.

The analysis in the second case, where two of the five elements have a reflection component, is similar. Up to automorphisms and braid equivalence, there is only one possible tuple and n must equal 2:

$$(x, 1)(0, 1), (x, 1)(y, 1)(y, 0).$$

Here H' must be the graph of the homomorphism which sends y to 1 and x to 0. And on the two intermediate covers we get the respective Hurwitz vectors $(y, y; x, x)$ and $(y', y'; x, x)$, where y' is a shorthand for $(y, 1) \in H'$. They are in one orbit under the automorphism of G which keeps $(x, 0)$ and $(x, 1)$ fix and which exchanges $(y, 0)$ with $(y, 1)$. \square

Theorem 7.7. *Assume that we have two distinct subgroups of Map_g , $H, H' \cong D_n$, and that $Z := \text{Fix}(H) = \text{Fix}(H')$. Then $\delta := \dim(Z) = 1$, and the numerical invariant g' cannot be the same for both actions.*

Proof. We have shown in cor. 7.2 that $\delta \leq 2, \delta \neq 0$, and in proposition 7.6 that $\delta \neq 2$. Hence $\delta = 1$.

By remark 7.3, if we make the assumption that g' is the same, follows then that H and H' are both of type III-b). By prop. 7.4 and remark 7.5, the group is then $G = D_n \times C_2$,

H is $D_n \times \{0\}$ and H' is the graph of some homomorphism from D_n to C_2 . By the last paragraph of Remark 7.5, K coincides with the group of rotations of $H = D_n \times \{0\}$, hence $H' = \ker(f)$, where $f : G = D_n \times C_2 \rightarrow C_2$ is given by $f(y^b x^i, a) = a + b$.

By remark 7.3, the branching indices of $C \rightarrow C/G$ are $2, 2, 2, c_4$, therefore up to a permutation we get

the sequences

$$a = 1, 1, 0, 0, \quad a + b = 0, 1, 1, 0, \quad b = 1, 0, 1, 0.$$

Hence we get that the first coordinates of the second and of the last element are rotations of respective orders d_2, c_4 , where $d_2 = 1$ or $d_2 = 2$.

There is surjection $G \rightarrow D_n$ with the second element in the kernel. Now, if D_n is generated by two reflections σ_1, σ_2 then the rotation $\sigma_1\sigma_2$ generates the group \mathcal{R} of rotations. We conclude that the order $c_4 = n$.

We may assume, up to an automorphism of D_n , that the last element is $(x, 0)$, and the first one is $(y, 1)$.

Then the 4-tuple is:

$$(y, 1)(x^h, 1)(yx^{h-1}, 0)(x, 0),$$

where $h = 0$ or $h = n/2$.

The respective subgroups are generated by $\gamma_3, \gamma_4, \gamma_1\gamma_3\gamma_1 =: \gamma'_3, \gamma_1\gamma_4\gamma_1 =: \gamma'_4$ in one case, by $\gamma_4, \gamma_1^* := \gamma_1^{-1}, \gamma_2\gamma_4\gamma_2, \gamma_2\gamma_1^*\gamma_2$ in the other.

We get the two corresponding length 4 Hurwitz vectors.

$$(yx^{h-1}, x, yx^{h+1}, x^{-1}), \quad (x, y, x, y).$$

The corresponding two Nielsen functions are equal for n odd ($h = 0$), distinct for n even, but in the same orbit under the group $\text{Aut}(D_n)$.

By [CLP11], Theorem 2, H, H' correspond then to the same unmarked topological type.

□

Theorem 7.8. *Assume that we have two distinct subgroups of Map_g , $H, H' \cong D_n$, and that $Z := \text{Fix}(H) = \text{Fix}(H')$. Then $\delta := \dim(Z) = 1$, and case III-a) holds for H , case III-b) holds for H' .*

This case actually occurs.

Proof. That the only possible case is the one described follows by theorem 7.7 and the previous discussion. It suffices thus to give a concrete example.

We have a polygonal group $T(2, 2, 2, 2d)$, generated by elements $\gamma_1, \gamma_2, \gamma_3, \gamma_4$ whose orders are respectively 2, 2, 2, $2d$ and whose product $\gamma_1\gamma_2\gamma_3\gamma_4$ is the identity.

The elliptic induced covering is generated by

$$a := \gamma_1\gamma_2, \quad b := \gamma_2\gamma_3, \quad c := (\gamma_1\gamma_4\gamma_1)^2$$

which satisfy the usual presentation $[a, b] = c$.

Whereas for the genus zero induced covering, since we have a double cover branched on the first two points, we get generators

$$\gamma_3, \quad \gamma_4, \quad \gamma'_3 := \gamma_1\gamma_3\gamma_1, \quad \gamma'_4 := \gamma_1\gamma_4\gamma_1,$$

which satisfy the usual presentation

$$\gamma_3\gamma_4\gamma'_3\gamma'_4 = 1.$$

These elements have respective orders $2, 2d, 2, 2d$, so we guess that γ_4 should map to a generator of the rotation group, say $\gamma_4 \mapsto x$, and γ_3 should map to a reflection, say $\gamma_3 \mapsto y$.

We make the following assumption: since γ_1 normalizes H' , let us just assume that γ_1 centralizes H' .

Then we have a direct product $G = H' \times C_2 \cong D_n \times C_2$, where $n = 2d$, and the cyclic group C_2 of order 2 is generated by γ_1 .

We take as Hurwitz vector for G (images of $\gamma_1, \gamma_2, \gamma_3, \gamma_4$)

$$(0, 1)(yx, 1)(y, 0)(x, 0).$$

Clearly these four elements generate then G .

Now $\gamma_3, \gamma_4, \gamma'_3, \gamma'_4$ are respectively sent to $(y, 0)(x, 0)(y, 0)(x, 0)$, and they generate the group $H' \cong D_n$.

On the other hand

$$a \mapsto (yx, 0), \quad b \mapsto (x^{-1}, 1), \quad c \mapsto (x^2, 0)$$

and clearly $[a, b] \mapsto (x^2, 0)$.

Moreover, the image group H projects, under the first coordinate, onto the dihedral group D_n . □

Acknowledgement:

The authors would like to thank Binru Li and Sascha Weigl for providing the argument which helped us to fix a gap in a previous proof of Prop.7.6.

REFERENCES

- [Artin] Artin, E., Geometric algebra, Interscience Publishers, Inc., New York-London, x+214 (1957).
- [BaCa97] Bauer, I.; Catanese, F. *Generic lemniscates of algebraic functions*. Math. Ann. 307, no. 3, 417–444 (1997).
- [BF86] Biggers, R.; Fried, M. *Irreducibility of moduli spaces of cyclic unramified covers of genus g curves*. Trans. Am. Math. Soc. 295, 59-70 (1986).
- [Bir69] Birman, Joan S. *Mapping class groups and their relationship to braid groups*, Comm. Pure Appl. Math. 22 (1969), 213–238.
- [Bro] Brown, K. S., Cohomology of groups, Graduate Texts in Mathematics, **87**, Springer-Verlag, New York, (1982).
- [Cat88] Catanese, F. *Moduli of algebraic surfaces*. Theory of moduli (Montecatini Terme, 1985), 1–83, Lecture Notes in Math., 1337, Springer, Berlin, (1988).
- [Cat00] Catanese, F., *Fibred Surfaces, varieties isogenous to a product and related moduli spaces*. Amer. J. Math. 122 (2000), no. 1, 1–44.
- [Cat08] Catanese, F. *Differentiable and deformation type of algebraic surfaces, real and symplectic structures*. Symplectic 4-manifolds and algebraic surfaces, 55–167, Lecture Notes in Math., 1938, Springer, Berlin, (2008).

- [Cat10] Catanese, F. *Irreducibility of the space of cyclic covers of algebraic curves of fixed numerical type and the irreducible components of $Sing(\overline{\mathcal{M}}_g)$* . Proceedings volume of "The Conference on Geometry" honoring Shing-Tung Yau's 60th birthday. "Advanced Lectures in Mathematics" series of International Press, in cooperation with the Higher Education Press and the Stefan Banach International Mathematical Centre (Institute of Mathematics Polish Academy of Sciences Publishing House), to appear. arXiv:1011.0316.
- [CLP11] Catanese, F., Lönne, M., Perroni, F. *Irreducibility of the space of dihedral covers of algebraic curves of fixed numerical type*. Atti Accad. Naz. Lincei Cl. Sci. Fis. Mat. Natur. Rend. Lincei (9) Mat. Appl. 22 (2011), 1–19. arXiv:1102.0490 .
- [CLP13] Catanese, F., Lönne, M., Perroni, F., *Genus stabilization for moduli of curves with symmetries*. arXiv:1301.4409
- [Cleb72] Clebsch A., *Zur Theorie der Riemann'schen Flächen*. Math. Ann. 6, 216-230 (1872).
- [Com30] Comessatti, A. *Sulle superficie multiple cicliche*. Rendiconti Seminario Padova 1, 1-45 (1930)
- [Cor87] Cornalba, M. *On the locus of curves with automorphisms*. Ann. Mat. Pura Appl., IV. Ser. 149, 135-151 (1987).
- [Cor08] Cornalba, M. *Erratum: On the locus of curves with automorphisms*. Ann. Mat. Pura Appl. (4) 187, No. 1, 185-186 (2008).
- [DM] Deligne, P., Mumford, D., *The irreducibility of the space of curves of given genus*, Inst. Hautes Études Sci. Publ. Math., no. 36, 75–109 (1969).
- [Du-Th06] Dunfield, N. M.; Thurston, W. P. *Finite covers of random 3-manifolds*. Invent. Math. 166 , no. 3, 457–521 (2006).
- [Edm I] Edmonds, A. L., *Surface symmetry. I*, Michigan Math. J., **29** (1982), n. 2, 171–183.
- [Edm II] Edmonds, A. L., *Surface symmetry. II*, Michigan Math. J., **30** (1983), n. 2, 143–154.
- [FM12] B. Farb, D. Margalit, *A primer on mapping class groups*, volume 49 of *Princeton Mathematical Series*, Princeton University Press, Princeton, NJ, 2012.
- [FV91] Fried, M. D.; Völklein, H., *The inverse Galois problem and rational points on moduli spaces*, Math. Ann., **290**, 1991, n. 4, 771–800.
- [Ful69] W. Fulton: *Hurwitz schemes and irreducibility of moduli of algebraic curves*. Ann. of Math. (2) 90 ,542-575 (1969) .
- [GHS02] Graber, T., Harris, J., Starr, J.: *A note on Hurwitz schemes of covers of a positive genus curve*. arXiv: math. AG/0205056.
- [Hopf] Hopf, H.: *Fundamentalgruppe und zweite Bettische Gruppe*. Comment. Math. Helv. 14 (1942), 257–309.
- [Hur91] Hurwitz, A.: *Ueber Riemann'schen Flächen mit gegebenen Verzweigungspunkten*. Math. Ann. 39, 1–61 (1891).
- [Kanev06] Kanev, V. *Hurwitz spaces of Galois coverings of \mathbf{P}^1 , whose Galois groups are Weyl groups*, J. Algebra 305 (2006) 442–456.
- [Kanev05] Kanev, V. *Irreducibility of Hurwitz spaces*. arXiv: math. AG/0509154.
- [Kluit88] Kluitmann, P.: *Hurwitz action and finite quotients of braid groups*. In: Braids (Santa Cruz, CA 1986). Contemporary Mathematics, vol. 78, pp. 299-325. AMS, Providence (1988).

- [Liv85] Livingston, C.: *Stabilizing surface symmetries*. Mich. Math. J. 32, 249–255 (1985).
- [MSSV01] Magaard, K., Shaska, T., Shpectorov, S., Völklein, H. , *The locus of curves with prescribed automorphism group*. Communications in arithmetic fundamental groups (Kyoto, 1999/2001). Sūrikaiseikikenkyūsho Kōkyūroku No. 1267 (2002), 112–141.
- [Par91] Pardini, R. *Abelian covers of algebraic varieties*. J. Reine Angew. Math. 417 (1991), 191–213.
- [Ries93] Ries, John F. X. , *Subvarieties of moduli space determined by finite groups acting on surfaces*. Trans. Amer. Math. Soc. 335 (1993), no. 1, 385–406.
- [Sia09] Sia, C. *Hurwitz equivalence in tuples of dihedral groups, dicyclic groups, and semidihedral groups*. Electron. J. Combin. 16 , no. 1, Research Paper 95, 17 pp (2009).
- [Sing72] D. Singerman, *Finitely maximal Fuchsian groups*, J. London Math. Soc. 6 (1972), 29–38.
- [Ve06] Vetro, F. : *Irreducibility of Hurwitz spaces of coverings with one special fiber*. Indag. Math. (N.S.) 17 , no. 1, 115–127 (2006).
- [Ve07] Vetro, F. : *Irreducibility of Hurwitz spaces of coverings with monodromy groups Weyl groups of type $W(B_d)$* . Boll. Unione Mat. Ital. Sez. B Artic. Ric. Mat. (8) 10 , no. 2, 405–431 (2007).
- [Ve08] Vetro, F. : *Irreducibility of Hurwitz spaces of coverings with one special fiber and monodromy group a Weyl group of type D_d* . Manuscripta Math. 125 , no. 3, 353–368 (2008).
- [Waj96] Wajnryb, B. *Orbits of Hurwitz action for coverings of a sphere with two special fibers*. Indag. Math. (N.S.) 7, no. 4, 549–558 (1996).
- [Waj99] Wajnryb, B. *An elementary approach to the mapping class group of a surface*. Geom. Topol. 3 , 405–466 (1999).
- [Wie] Wiegold, J., *The Schur multiplier: an elementary approach*, Groups—St. Andrews 1981 (St. Andrews, 1981), London Math. Soc. Lecture Note Ser., **71**, 137–154, Cambridge Univ. Press, (1982).

Authors' Address:

Fabrizio Catanese, Michael Lönne
 Lehrstuhl Mathematik VIII,
 Mathematisches Institut der Universität Bayreuth
 NW II, Universitätsstr. 30
 95447 Bayreuth

Fabio Perroni,
 SISSA - International School for Advanced Studies, School of Mathematics
 via Bonomea, 265 - 34136 Trieste (Italy).

email:
 fabrizio.catanese@uni-bayreuth.de,
 michael.loenne@uni-bayreuth.de,
 perroni@sissa.it.