

# Stable determination of an inclusion in an inhomogeneous elastic body by boundary measurements

ANTONINO MORASSI AND EDI ROSSET

*“Dedicato a Giovanni Alessandrini per il suo sessantesimo compleanno”*

**ABSTRACT.** *In this paper we consider the stability issue for the inverse problem of determining an unknown inclusion contained in an elastic body by all the pairs of measurements of displacement and traction taken at the boundary of the body. Both the body and the inclusion are made by inhomogeneous linearly elastic isotropic material. Under mild a priori assumptions about the smoothness of the inclusion and the regularity of the coefficients, we show that the logarithmic stability estimate proved in [3] in the case of piecewise constant coefficients continues to hold in the inhomogeneous case. We introduce new arguments which allow to simplify some technical aspects of the proof given in [3].*

**Keywords:** Inverse problems, linearized elasticity, inclusions, stability, unique continuation.

**MS Classification 2010:** 35R30, 35Q74, 35R25, 74B05, 74G75.

## 1. Introduction

The inverse problem of determining unknown inclusions in continuous bodies from measurements of physical parameters taken at the boundary of the body has attracted a lot of attention in the last thirty years, see, among other contributions, the reconstruction results obtained in [12, 17, 18]. Inclusions may be due to the presence of inhomogeneities or defects inside the body, and the development of non-invasive testing approaches is of great importance in several practical contexts, ranging from medicine to engineering applications.

Inverse problems of this class are usually ill-posed according to Hadamard’s definition, and one of the main issues is the uniqueness of the solution, that is the determination of the boundary measurements which ensure the unique determination of the defect. Moreover, from the point of view of practical applications, it is crucial to establish how small perturbations on the data may

affect the accuracy of the identification of the inclusion, namely, the study of the stability issue.

The prototype of these inverse problems is the determination of an inclusion inside an electric conductor from boundary measurements of electric potential and current flux. Uniqueness was first proved by Isakov in '88 [14]. The first stability result is due to Alessandrini and Di Cristo [2], who derived a logarithmic stability estimate of the inclusion from all possible boundary measurements, that is from the full Dirichlet-to-Neumann map. More precisely, the authors considered in [2] the case of piecewise-constant coefficients and constructed an ingenious proof which, starting from Alessandrini's identity (first derived in [1]), makes use of fundamental solutions for elliptic equations with discontinuous coefficients, and suitable quantitative forms of unique continuation for solutions to Laplacian equation. An extension of the above result to the case of variable coefficients was derived in [8]. The pioneering work [2] stimulated a subsequent line of research in which methods and results were extended to other frameworks, such as, for example, the stable identification of inclusions in thermal conductors [9, 10], which involves a parabolic equation with discontinuous coefficients.

Concerning the determination of an inclusion in an elastic body from the Dirichlet-to-Neumann map, the uniqueness was proved by Ikehata, Nakamura and Tanuma in [13]. The stability issue has been recently faced in [3]. The statical equilibrium of the defected body is governed by the following system of elliptic equations

$$\operatorname{div}((\mathbb{C} + (\mathbb{C}^D - \mathbb{C})\chi_D)\nabla u) = 0, \quad \text{in } \Omega, \quad (1)$$

where  $u$  is the three-dimensional displacement field inside the elastic body  $\Omega$ ,  $\chi_D$  is the characteristic function of the inclusion  $D$ , and  $\mathbb{C}$ ,  $\mathbb{C}^D$  is the elasticity tensor in the background material and inside the inclusion, respectively. Given inclusions  $D_1$ ,  $D_2$ , let  $\Lambda_{D_i} : H^{1/2}(\partial\Omega) \rightarrow H^{-1/2}(\partial\Omega)$  be the Dirichlet-to-Neumann map which gives the traction at the boundary  $\partial\Omega$  corresponding to a displacement field assigned on  $\partial\Omega$ , when  $D = D_i$ ,  $i = 1, 2$ . Assuming that  $\mathbb{C}$ ,  $\mathbb{C}^{D_1} = \mathbb{C}^{D_2}$  are *constant* and of Lamé type (e.g., isotropic material), and under  $C^{1,\alpha}$ -regularity of the boundary of the inclusion, the authors derived the following stability result. If, for some  $\epsilon$ ,  $0 < \epsilon < 1$ ,

$$\|\Lambda_{D_1} - \Lambda_{D_2}\|_{\mathcal{L}(H^{1/2}(\partial\Omega), H^{-1/2}(\partial\Omega))} \leq \epsilon, \quad (2)$$

then the Hausdorff distance between the two inclusions can be controlled as

$$d_H(\partial D_1, \partial D_2) \leq \frac{C}{|\log \epsilon|^\eta}, \quad (3)$$

where the constants  $C > 0$  and  $\eta$ ,  $0 < \eta \leq 1$ , only depend on the a-priori data.

The piecewise-constant Lamé case can be considered as a simplified mathematical model of real elastic bodies. Therefore, it is of practical interest to extend the stability estimate (3) to *variable* coefficients both in the background,  $\mathbb{C} = \mathbb{C}(x)$ , and in the inclusions,  $\mathbb{C}^{D_i} = \mathbb{C}^{D_i}(x)$ ,  $i = 1, 2$ . More precisely, assuming  $C^{1,1}$  and  $C^\tau$  regularity,  $\tau \in (0, 1)$ , for  $\mathbb{C}$  and  $\mathbb{C}^{D_i}$ , respectively,  $i = 1, 2$ , in this paper we show that (3) continues to hold. Let us emphasize that in order to derive our result the exact knowledge of the elasticity tensor inside the inclusion is not needed. In fact, only the strong convexity conditions (16) and the bounds (17), (20), (22) are required. Moreover, as in [3], the inclusion is allowed to share a portion of its boundary with the boundary of the body  $\Omega$ .

Let us briefly recall the main ideas of our approach and the new mathematical tools we used in the proof of the stability result. Let  $\Gamma^{D_i}$  be the fundamental matrix associated to the elasticity tensor  $(\mathbb{C} + (\mathbb{C}^{D_i} - \mathbb{C})\chi_{D_i})$ ,  $i = 1, 2$ . The main idea is to obtain an upper and a lower bound for  $(\Gamma^{D_2} - \Gamma^{D_1})(y, w)$  for points  $y$  and  $w$  belonging to the connected component of  $\mathbb{R}^3 \setminus (\overline{D_1} \cup \overline{D_2})$  which contains  $\mathbb{R}^3 \setminus \overline{\Omega}$ , and approaching non-tangentially a suitable point  $P \in \partial D_1 \setminus \overline{D_2}$  (or  $\partial D_2 \setminus \overline{D_1}$ ). A first crucial ingredient in determining both upper and lower bounds is the integral representation of  $(\Gamma^{D_2} - \Gamma^{D_1})(y, w)$  given by formula (40). Next, the upper bound follows from an application of Alessandrini's identity (suitably adapted to linear elasticity, see Lemma 6.1 in [3]) and a propagation of smallness argument based on iterated use of the three spheres inequality for solutions to the Lamé system of linear elasticity with smooth variable coefficients.

In proving the lower bound (see Section 4) we introduce new arguments which entail a simplification of the proof given for the piecewise-constant coefficient case. Indeed, a generalization of Theorem 8.1 in [3], which was a key tool in proving the lower bound, should need the derivation of an asymptotic approximation of  $\Gamma^D$  in terms of the fundamental matrix obtained by locally flattening the boundary  $\partial D$  and freezing the coefficients at a point belonging to  $\partial D$ , which does not appear straightforward.

Finally, let us emphasize that the statement of Theorem 8.1 in [3], besides being worth of interest from a theoretical viewpoint, may have relevant interest for its possible applications. In fact, it turned out to be a fundamental ingredient in the proof of Lipschitz stability estimates for the inverse problem of determining the Lamé moduli for a piecewise constant elasticity tensor corresponding to a known partition of the body in a finite number of subdomains having regular interfaces [6], see also [7] for the case of flat interfaces.

The plan of the paper is as follows. Notation and the a priori information are introduced in section 2, together with the statement of the stability result (Theorem 2.2). In section 3 we recall some auxiliary results, we state the upper and lower bounds on  $(\Gamma^{D_2} - \Gamma^{D_1})$ , Theorems 3.4 and 3.5, and we give the proof of the main Theorem 2.2. Section 4 is devoted to the proof of Theorem 3.5.

## 2. The main result

### 2.1. Notation

Let us denote  $\mathbb{R}_+^3 = \{x \in \mathbb{R}^3 \mid x_3 > 0\}$  and  $\mathbb{R}_-^3 = \{x \in \mathbb{R}^3 \mid x_3 < 0\}$ . Given  $x \in \mathbb{R}^3$ , we shall denote  $x = (x', x_3)$ , where  $x' = (x_1, x_2) \in \mathbb{R}^2$ ,  $x_3 \in \mathbb{R}$ . Given  $x \in \mathbb{R}^3$  and  $r > 0$ , we shall use the following notation for balls in three and two dimensions:

$$B_r(x) = \{y \in \mathbb{R}^3 \mid |y - x| < r\}, \quad B_r = B_r(O),$$

$$B'_r(x') = \{y' \in \mathbb{R}^2 \mid |y' - x'| < r\}, \quad B'_r = B'_r(O).$$

DEFINITION 2.1 ( $C^{k,\alpha}$  regularity). *Let  $E$  be a domain in  $\mathbb{R}^3$ . Given  $k, \alpha, k \in \mathbb{N}, 0 < \alpha \leq 1$ , we say that  $E$  is of class  $C^{k,\alpha}$  with constants  $\rho_0, M_0 > 0$ , if, for any  $P \in \partial E$ , there exists a rigid transformation of coordinates under which we have  $P = 0$  and*

$$E \cap B_{\rho_0}(O) = \{x \in B_{\rho_0}(O) \mid x_3 > \varphi(x')\},$$

where  $\varphi$  is a  $C^{k,\alpha}$  function on  $B'_{\rho_0}$  satisfying

$$\varphi(O) = 0,$$

$$|\nabla \varphi(O)| = 0, \quad \text{when } k \geq 1,$$

$$\|\varphi\|_{C^{k,\alpha}(B'_{\rho_0}(O))} \leq M_0 \rho_0.$$

Here and in the sequel all norms are normalized such that their terms are dimensionally homogeneous. For instance

$$\|\varphi\|_{C^{k,\alpha}(B'_{\rho_0}(O))} = \sum_{i=0}^k \rho_0^i \|\nabla^i \varphi\|_{L^\infty(B'_{\rho_0}(O))} + \rho_0^{k+\alpha} |\nabla^k \varphi|_{\alpha, B'_{\rho_0}(O)},$$

where

$$|\nabla^k \varphi|_{\alpha, B'_{\rho_0}(O)} = \sup_{\substack{x', y' \in B'_{\rho_0}(O) \\ x' \neq y'}} \frac{|\nabla^k \varphi(x') - \nabla^k \varphi(y')|}{|x' - y'|^\alpha}.$$

Similarly, for a vector function  $u : \Omega \subset \mathbb{R}^3 \rightarrow \mathbb{R}^3$ , we set

$$\|u\|_{H^1(\Omega, \mathbb{R}^3)} = \left( \int_{\Omega} |u|^2 + \rho_0^2 \int_{\Omega} |\nabla u|^2 \right)^{\frac{1}{2}},$$

and so on for boundary and trace norms such as  $\|\cdot\|_{H^{\frac{1}{2}}(\partial\Omega, \mathbb{R}^3)}, \|\cdot\|_{H^{-\frac{1}{2}}(\partial\Omega, \mathbb{R}^3)}$ .

For any  $U \subset \mathbb{R}^3$  and for any  $r > 0$ , we denote

$$U_r = \{x \in U \mid \text{dist}(x, \partial U) > r\}, \quad (4)$$

$$U^r = \{x \in \mathbb{R}^3 \mid \text{dist}(x, U) < r\}. \quad (5)$$

We denote by  $\mathbb{M}^{m \times n}$  the space of  $m \times n$  real valued matrices and we also use the notation  $\mathbb{M}^n = \mathbb{M}^{n \times n}$ . Let  $\mathcal{L}(X, Y)$  be the space of bounded linear operators between Banach spaces  $X$  and  $Y$ .

For every pair of real  $n$ -vectors  $a$  and  $b$ , we denote by  $a \otimes b$  the  $n \times n$  matrix with entries

$$(a \otimes b)_{ij} = a_i b_j, \quad i, j = 1, \dots, n. \quad (6)$$

For every  $3 \times 3$  matrices  $A, B$  and for every  $\mathbb{C} \in \mathcal{L}(\mathbb{M}^3, \mathbb{M}^3)$ , we use the following notation:

$$(\mathbb{C}A)_{ij} = \sum_{k,l=1}^3 C_{ijkl} A_{kl}, \quad (7)$$

$$A \cdot B = \sum_{i,j=1}^3 A_{ij} B_{ij}, \quad (8)$$

$$|A| = (A \cdot A)^{\frac{1}{2}}, \quad (9)$$

where  $C_{ijkl}, A_{ij}$  and  $B_{ij}$  are the entries of  $\mathbb{C}, A$  and  $B$  respectively.

Finally, let us recall the definition of the Hausdorff distance  $d_H(A, B)$  of two bounded closed sets  $A, B \subset \mathbb{R}^3$

$$d_H(A, B) = \max \left\{ \max_{x \in A} d(x, B), \max_{x \in B} d(x, A) \right\}$$

## 2.2. A-priori information and main result

We make the following a-priori assumptions. The continuous body  $\Omega$  is a bounded domain in  $\mathbb{R}^3$  such that

$$\mathbb{R}^3 \setminus \overline{\Omega} \text{ is connected}, \quad (10)$$

$$|\Omega| \leq M_1 \rho_0^3, \quad (11)$$

$$\Omega \text{ is of class } C^{1,\alpha}, \text{ with constants } \rho_0, M_0, \quad (12)$$

and the inclusion  $D$  is a connected subset of  $\Omega$  satisfying

$$\mathbb{R}^3 \setminus \overline{D} \text{ is connected}, \quad (13)$$

$$D \text{ is of class } C^{1,\alpha}, \text{ with constants } \rho_0, M_0, \quad (14)$$

where  $\rho_0, M_0, M_1$  are given positive constants, and  $0 < \alpha \leq 1$ .

The background material is linearly elastic isotropic, with elasticity tensor  $\mathbb{C} = \mathbb{C}(x)$ , which - without restriction - may be defined in the whole  $\mathbb{R}^3$ . The cartesian components of  $\mathbb{C}(x)$  are

$$C_{ijkl}(x) = \lambda(x)\delta_{ij}\delta_{kl} + \mu(x)(\delta_{ki}\delta_{lj} + \delta_{li}\delta_{kj}), \quad \text{for every } x \in \mathbb{R}^3, \quad (15)$$

where  $\delta_{ij}$  is the Kronecker's delta and the Lamé moduli  $\lambda = \lambda(x)$ ,  $\mu = \mu(x)$  satisfy the strong convexity conditions

$$\mu(x) \geq \alpha_0, \quad 2\mu(x) + 3\lambda(x) \geq \gamma_0, \quad \text{for every } x \in \mathbb{R}^3, \quad (16)$$

for given constants  $\alpha_0 > 0$ ,  $\gamma_0 > 0$ . We shall also assume upper bounds

$$\mu(x) \leq \bar{\mu}, \quad \lambda(x) \leq \bar{\lambda}, \quad \text{for every } x \in \mathbb{R}^3, \quad (17)$$

where  $\bar{\mu} > 0$ ,  $\bar{\lambda} \in \mathbb{R}$  are given constants. Let us notice that (15) clearly implies the major and minor symmetries of  $\mathbb{C}$ , namely

$$C_{ijkl} = C_{klij} = C_{lkij}, \quad i, j, k, l = 1, 2, 3. \quad (18)$$

The inclusion  $D$  is assumed to be made by linearly elastic isotropic material having elasticity tensor  $\mathbb{C}^D = \mathbb{C}^D(x)$  with components

$$C_{ijkl}^D(x) = \lambda^D(x)\delta_{ij}\delta_{kl} + \mu^D(x)(\delta_{ki}\delta_{lj} + \delta_{li}\delta_{kj}), \quad \text{for every } x \in \bar{\Omega}, \quad (19)$$

where the Lamé moduli  $\lambda^D(x)$ ,  $\mu^D(x)$  satisfy the conditions (16)–(17) and, in addition,

$$(\lambda(x) - \lambda^D(x))^2 + (\mu(x) - \mu^D(x))^2 \geq \eta_0^2 > 0, \quad \text{for every } x \in \bar{\Omega}, \quad (20)$$

for a given constant  $\eta_0 > 0$ .

Finally, the elasticity tensors  $\mathbb{C}$  and  $\mathbb{C}^D$  are assumed to be of  $C^{1,1}$  class in  $\mathbb{R}^3$  and of  $C^\tau$  class in  $\bar{\Omega}$ ,  $\tau \in (0, 1)$ , respectively, that is

$$\|\lambda\|_{C^{1,1}(\mathbb{R}^3)} + \|\mu\|_{C^{1,1}(\mathbb{R}^3)} \leq M, \quad (21)$$

$$\|\lambda^D\|_{C^\tau(\bar{\Omega})} + \|\mu^D\|_{C^\tau(\bar{\Omega})} \leq M, \quad (22)$$

for a given constant  $M > 0$ .

For any  $f \in H^{\frac{1}{2}}(\partial\Omega)$ , let  $u \in H^1(\Omega)$  be the weak solution to the Dirichlet problem

$$\begin{cases} \operatorname{div}((\mathbb{C} + (\mathbb{C}^D - \mathbb{C})\chi_D)\nabla u) = 0, & \text{in } \Omega, \\ u = f, & \text{on } \partial\Omega, \end{cases} \quad (23)$$

$$(24)$$

where  $\chi_D$  is the characteristic function of  $D$ . The Dirichlet-to-Neumann map  $\Lambda_D$  associated to (23)–(24),

$$\Lambda_D : H^{1/2}(\partial\Omega) \rightarrow H^{-1/2}(\partial\Omega), \quad (25)$$

is defined in the weak form by

$$\langle \Lambda_D f, v|_{\partial\Omega} \rangle = \int_{\Omega} (\mathbb{C} + (\mathbb{C}^D - \mathbb{C})\chi_D)\nabla u \cdot \nabla v, \quad (26)$$

for every  $v \in H^1(\Omega)$ .

We prove the following logarithmic stability estimate for the inverse problem of recovering the inclusion  $D$  from the knowledge of the map  $\Lambda_D$ .

**THEOREM 2.2.** *Let  $\Omega \subset \mathbb{R}^3$  be a bounded domain satisfying (10)–(12) and let  $D_1, D_2$  be two connected inclusions contained in  $\Omega$  satisfying (13)–(14). Let  $\mathbb{C}(x)$  and  $\mathbb{C}^{D_i}(x)$  be the elasticity tensor of the material of  $\Omega$  and of the inclusion  $D_i$ ,  $i = 1, 2$ , respectively, where  $\mathbb{C}(x)$  given in (15) and  $\mathbb{C}^{D_i}(x)$  given in (19) (for  $D = D_i$ ) satisfy (16), (17), (20), (21) and (22). If, for some  $\epsilon$ ,  $0 < \epsilon < 1$ ,*

$$\|\Lambda_{D_1} - \Lambda_{D_2}\|_{\mathcal{L}(H^{1/2}(\partial\Omega), H^{-1/2}(\partial\Omega))} \leq \frac{\epsilon}{\rho_0}, \quad (27)$$

then

$$d_H(\partial D_1, \partial D_2) \leq C\rho_0 |\log \epsilon|^{-\eta}, \quad (28)$$

where  $C > 0$  and  $\eta$ ,  $0 < \eta \leq 1$ , are constants only depending on  $M_0, \alpha, M_1, \alpha_0, \gamma_0, \bar{\mu}, \bar{\lambda}, \eta_0, \tau, M$ .

**REMARK 2.3.** If in Theorem 2.2 we further assume that the two inclusions are at a prescribed distance from  $\partial\Omega$ , then the result continues to hold even when the local Dirichlet-to-Neumann map is known. The proof can be obtained by adapting the general theory developed by Alessandrini and Kim [4].

### 3. Proof of the main result

In order to state the metric Lemma 3.1 below, we need to introduce some notation.

We denote by  $\mathcal{G}$  the connected component of  $\mathbb{R}^3 \setminus (\overline{D_1 \cup D_2})$  which contains  $\mathbb{R}^3 \setminus \overline{\Omega}$ .

Given  $O = (0, 0, 0)$ , a unit vector  $v$ ,  $h > 0$  and  $\vartheta \in (0, \frac{\pi}{2})$ , we denote by

$$C(O, v, h, \vartheta) = \{x \in \mathbb{R}^3 \mid |x - (x \cdot v)v| \leq \sin \vartheta |x|, 0 \leq x \cdot v \leq h\} \quad (29)$$

the closed truncated cone with vertex at  $O$ , axis along the direction  $v$ , height  $h$  and aperture  $2\vartheta$ . Given  $R, d$ ,  $0 < R < d$  and  $Q = -de_3$ , let us consider the cone

$C\left(O, -e_3, \frac{d^2-R^2}{d}, \arcsin \frac{R}{d}\right)$ , whose lateral boundary is tangent to the sphere  $\partial B_R(Q)$  along the circumference of its base.

Given a point  $P \in \partial D_1 \cap \partial \mathcal{G}$ , let  $\nu$  be the outer unit normal to  $\partial D_1$  at  $P$  and let  $d > 0$  be such that the segment  $[P + d\nu, P]$  is contained in  $\bar{\mathcal{G}}$ . For a point  $P_0 \in \bar{\mathcal{G}}$ , let  $\gamma$  be a path in  $\bar{\mathcal{G}}$  joining  $P_0$  to  $P + d\nu$ . We consider the following neighbourhood of  $\gamma \cup [P + d\nu, P] \setminus \{P\}$  formed by a tubular neighbourhood of  $\gamma$  attached to a cone with vertex at  $P$  and axis along  $\nu$

$$V(\gamma, d, R) = \bigcup_{S \in \gamma} B_R(S) \cup C\left(P, \nu, \frac{d^2 - R^2}{d}, \arcsin \frac{R}{d}\right). \quad (30)$$

Let us also define

$$S_{2\rho_0} = \{x \in \mathbb{R}^3 \mid \rho_0 < \text{dist}(x, \Omega) < 2\rho_0\}. \quad (31)$$

LEMMA 3.1. *Under the assumptions of Theorem 2.2, up to inverting the role of  $D_1$  and  $D_2$ , there exist positive constants  $\bar{d}$ ,  $\bar{c}$ , where  $\frac{\bar{d}}{\rho_0}$  only depends on  $M_0$  and  $\alpha$ , and  $\bar{c} \geq 1$  only depends on  $M_0$ ,  $\alpha$  and  $M_1$ , and there exists a point  $P \in \partial D_1 \cap \partial \mathcal{G}$  such that*

$$d_H(\partial D_1, \partial D_2) \leq \bar{c} \text{dist}(P, D_2), \quad (32)$$

and such that, giving any point  $P_0 \in S_{2\rho_0}$ , there exists a path  $\gamma \subset \Omega^{2\rho_0} \cap \mathcal{G}$  joining  $P_0$  to  $P + \bar{d}\nu$ , where  $\nu$  is the unit outer normal to  $D_1$  at  $P$ , such that, choosing a coordinate system with origin  $O$  at  $P$  and axis  $e_3 = -\nu$ , we have

$$V(\gamma, \bar{d}, \bar{R}) \subset \mathbb{R}^3 \cap \bar{\mathcal{G}}, \quad (33)$$

where  $\frac{\bar{R}}{\rho_0}$  only depends on  $M_0$  and  $\alpha$ .

The thesis of the above lemma is a straightforward consequence of Lemma 4.1 and Lemma 4.2 in [3], and is inspired by results obtained in [5] and [2].

Let  $D$  be a domain of class  $C^{1,\alpha}$  with constants  $\rho_0$ ,  $M_0$  and  $0 < \alpha \leq 1$ . The elasticity tensors  $\mathbb{C}$  and  $\mathbb{C}^D$  given by (15) and (19) respectively, satisfy (16), (17), (21) and (22).

Given  $y \in \mathbb{R}^3$  and a concentrated force  $l\delta(\cdot - y)$  applied at  $y$ , with  $l \in \mathbb{R}^3$ , let us consider the normalized fundamental solution  $u^D \in L^1_{loc}(\mathbb{R}^3, \mathbb{R}^3)$  defined by

$$\begin{cases} \text{div}_x ((\mathbb{C}(x) + (\mathbb{C}^D(x) - \mathbb{C}(x))\chi_D)\nabla_x u^D(x, y; l)) \\ \quad \quad \quad = -l\delta(x - y), & \text{in } \mathbb{R}^3 \setminus \{y\}, \\ \lim_{|x| \rightarrow \infty} u^D(x, y; l) = 0, \end{cases} \quad (34)$$



where  $\delta(\cdot - y)$  is the Dirac distribution supported at  $y$ . It is well-known that

$$u^D(x, y; l) = \Gamma^D(x, y)l, \quad (35)$$

where  $\Gamma^D = \Gamma^D(\cdot, y) \in L^1_{loc}(\mathbb{R}^3, \mathcal{L}(\mathbb{R}^3, \mathbb{R}^3))$  is the *normalized fundamental matrix* for the operator  $\operatorname{div}_x((\mathbb{C}(x) + (\mathbb{C}^D(x) - \mathbb{C}(x))\chi_D)\nabla_x(\cdot))$ . Existence of  $\Gamma^D$  and asymptotic estimates are stated in the following Proposition.

**PROPOSITION 3.2.** *Under the above assumptions, there exists a unique fundamental matrix  $\Gamma^D(\cdot, y) \in C^0(\mathbb{R}^3 \setminus \{y\})$ , such that*

$$\Gamma^D(x, y) = (\Gamma^D(y, x))^T, \quad \text{for every } x \in \mathbb{R}^3, x \neq y, \quad (36)$$

$$|\Gamma^D(x, y)| \leq C|x - y|^{-1}, \quad \text{for every } x \in \mathbb{R}^3, x \neq y, \quad (37)$$

$$|\nabla_x \Gamma^D(x, y)| \leq C|x - y|^{-2}, \quad \text{for every } x \in \mathbb{R}^3, x \neq y, \quad (38)$$

where the constant  $C > 0$  only depends on  $M_0, \alpha, \alpha_0, \gamma_0, \bar{\lambda}, \bar{\mu}, \tau, M$ .

A proof of Proposition 3.2 follows by merging the regularity results by Li and Nirenberg [15] and the analysis by Hofmann and Kim [11], see [3] for details.

Let  $D_i, i = 1, 2$ , be a domain of class  $C^{1,\alpha}$  with constants  $\rho_0, M_0$  and  $0 < \alpha \leq 1$ , and consider the elasticity tensors

$$\mathbb{C}^1 = \mathbb{C}\chi_{\mathbb{R}^3 \setminus D_1} + \mathbb{C}^{D_1}\chi_{D_1}, \quad \mathbb{C}^2 = \mathbb{C}\chi_{\mathbb{R}^3 \setminus D_2} + \mathbb{C}^{D_2}\chi_{D_2}, \quad (39)$$

where  $\mathbb{C}^{D_1}, \mathbb{C}^{D_2}$  given in (19) (with  $D = D_1$  and  $D = D_2$ , respectively) satisfy (16), (17) and (22).

The following Proposition 3.3 states an integral representation involving the normalized fundamental matrices corresponding to inclusions  $D_1$  and  $D_2$ . Similar identities will be introduced in Section 4, in order to prove Theorem 3.5. Since these integral representations are basic ingredients for our approach, we present here a proof of Proposition 3.3, which is more exhaustive with respect to that given in [3, Proof of Lemma 6.2], where some details were implied.

**PROPOSITION 3.3.** *Let  $D_i$  and  $\mathbb{C}^{D_i}, i = 1, 2$ , satisfy the above assumptions. Then, for every  $y, w \in \mathbb{R}^3, y \neq w$ , and for every  $l, m \in \mathbb{R}^3$  we have*

$$\begin{aligned} & (\Gamma^{D_2} - \Gamma^{D_1})(y, w)m \cdot l = \\ & = \int_{\Omega} \mathbb{C}^1 \nabla \Gamma^{D_1}(\cdot, y)l \cdot \nabla \Gamma^{D_2}(\cdot, w)m - \int_{\Omega} \mathbb{C}^2 \nabla \Gamma^{D_1}(\cdot, y)l \cdot \nabla \Gamma^{D_2}(\cdot, w)m. \end{aligned} \quad (40)$$

*Proof.* Formula (40) is obtained by subtracting the two following identities

$$\int_{\mathbb{R}^3} \mathbb{C}^1 \nabla \Gamma^{D_1}(\cdot, y)l \cdot \nabla \Gamma^{D_2}(\cdot, w)m = \Gamma^{D_2}(y, w)m \cdot l, \quad (41)$$

$$\int_{\mathbb{R}^3} \mathbb{C}^2 \nabla \Gamma^{D_1}(\cdot, y) l \cdot \nabla \Gamma^{D_2}(\cdot, w) m = \Gamma^{D_1}(y, w) m \cdot l. \quad (42)$$

To prove (41), let

$$\mathcal{H} = \{f : \mathbb{R}^3 \rightarrow \mathbb{R}^3 \mid f \in C^0(\mathbb{R}^3, \mathbb{R}^3) \cap H^1(\mathbb{R}^3, \mathbb{R}^3), \\ f \text{ with compact support}\}. \quad (43)$$

By the weak formulation of (34) (with  $D = D_1$ ), we have

$$\int_{\mathbb{R}^3} \mathbb{C}^1 \nabla \Gamma^{D_1}(\cdot, y) l \cdot \nabla \varphi = \varphi(y) \cdot l, \quad \text{for every } \varphi \in \mathcal{H}. \quad (44)$$

Let  $\epsilon > 0$ ,  $R > 0$ , with  $\epsilon \leq \frac{|w-y|}{2}$ ,  $R \geq 2 \max\{|y|, |w|\}$ , and choose  $\varphi \in \mathcal{H}$  such that  $\text{supp}(\varphi) \subset B_{2R}(0)$  and  $\varphi|_{B_R(0) \setminus B_\epsilon(w)} \equiv \Gamma^{D_2}(\cdot, w) m$ . Then, (44) can be rewritten as

$$I_{\epsilon, R} + I_\epsilon + I_{R, 2R} = \Gamma^{D_2}(y, w) m \cdot l, \quad (45)$$

where

$$I_{\epsilon, R} = \int_{B_R(0) \setminus B_\epsilon(w)} \mathbb{C}^1 \nabla \Gamma^{D_1}(\cdot, y) l \cdot \nabla \Gamma^{D_2}(\cdot, w) m, \quad (46)$$

$$I_\epsilon = \int_{B_\epsilon(w)} \mathbb{C}^1 \nabla \Gamma^{D_1}(\cdot, y) l \cdot \nabla \varphi, \quad (47)$$

$$I_{R, 2R} = \int_{B_{2R}(0) \setminus B_R(0)} \mathbb{C}^1 \nabla \Gamma^{D_1}(\cdot, y) l \cdot \nabla \varphi. \quad (48)$$

Integrating by parts on  $B_\epsilon(w)$  and recalling that  $y \in \mathbb{R}^3 \setminus \overline{B_\epsilon(w)}$ , we have

$$I_\epsilon = \int_{\partial B_\epsilon(w)} (\mathbb{C}^1 \nabla \Gamma^{D_1}(\cdot, y) l) \nu \cdot \Gamma^{D_2}(\cdot, w) m. \quad (49)$$

For every  $x \in \partial B_\epsilon(w)$  and by our choice of  $\epsilon$ , we have  $|x-y| \geq |y-w| - |w-x| \geq \frac{|y-w|}{2}$ . Therefore, by (37) and (38), we have

$$I_\epsilon \leq C \int_{|x-w|=\epsilon} \frac{1}{|x-y|^2} \frac{1}{|x-w|} \leq \frac{C\epsilon}{|y-w|^2}, \quad (50)$$

where the constant  $C > 0$  only depends on  $M_0$ ,  $\alpha$ ,  $\alpha_0$ ,  $\gamma_0$ ,  $\bar{\lambda}$ ,  $\bar{\mu}$ ,  $\tau$ ,  $M$ .

Analogously, integrating by parts in  $B_{2R}(0) \setminus B_R(0)$  and recalling that  $\varphi = 0$  on  $\partial B_{2R}(0)$  and  $y \in B_{\frac{R}{2}}(0)$ , we have

$$I_{R, 2R} = - \int_{\partial B_R(0)} (\mathbb{C}^1 \nabla \Gamma^{D_1}(\cdot, y) l) \nu \cdot \Gamma^{D_2}(\cdot, w) m. \quad (51)$$

For every  $x \in \partial B_R(0)$  and by our choice of  $R$ , we have  $|x - w| \geq |x| - |w| \geq \frac{R}{2}$  and  $|x - y| \geq \frac{R}{2}$ . Therefore,

$$I_{R,2R} \leq C \int_{|x|=R} \frac{1}{|x-y|^2} \frac{1}{|x-w|} \leq \frac{C}{R}, \quad (52)$$

where the constant  $C > 0$  only depends on  $M_0, \alpha, \alpha_0, \gamma_0, \bar{\lambda}, \bar{\mu}, \tau, M$ .

Using the estimates (50) and (52) in (45), and taking the limit as  $\epsilon \rightarrow 0$  and  $R \rightarrow \infty$ , we obtain (41). Symmetrically, we obtain

$$\int_{\mathbb{R}^3} \mathbb{C}^2 \nabla \Gamma^{D_1}(\cdot, y) l \cdot \nabla \Gamma^{D_2}(\cdot, w) m = \Gamma^{D_1}(w, y) l \cdot m. \quad (53)$$

By using (36), we obtain (42).  $\square$

Let  $P, P \in \partial D_1$ , be the point introduced in Lemma 3.1. In the following two theorems, we use a cartesian coordinate system such that  $P \equiv O = (0, 0, 0)$  and  $\nu = -e_3$ , where  $\nu$  is the unit outer normal to  $D_1$  at  $P$ .

**THEOREM 3.4** (Upper bound on  $(\Gamma^{D_2} - \Gamma^{D_1})$ ). *Under the notation of Lemma 3.1, let*

$$y_h = P - h e_3, \quad (54)$$

$$w_h = P - \lambda_w h e_3, \quad 0 < \lambda_w < 1, \quad (55)$$

with

$$0 < h \leq \bar{h} \rho_0, \quad (56)$$

where  $\bar{h}$  only depends on  $M_0$  and  $\alpha$ .

Then, for every  $l, m \in \mathbb{R}^3$ ,  $|l| = |m| = 1$ , we have

$$|(\Gamma^{D_2} - \Gamma^{D_1})(y_h, w_h) m \cdot l| \leq \frac{C}{\lambda_w h} \epsilon^{C_1 \left(\frac{h}{\rho_0}\right)^{C_2}}, \quad (57)$$

where the positive constants  $C, C_1$  and  $C_2$  only depend on  $M_0, \alpha, M_1, \alpha_0, \gamma_0, \bar{\lambda}, \bar{\mu}, \tau$  and  $M$ .

For the proof of the above result, we refer to [3, Section 7]. To give an idea of the role played by Proposition 3.3 in proving estimate (57), let us recall Alessandrini's identity

$$\int_{\Omega} \mathbb{C}^1 \nabla u_1 \cdot \nabla u_2 - \int_{\Omega} \mathbb{C}^2 \nabla u_1 \cdot \nabla u_2 = \langle (\Lambda_{D_1} - \Lambda_{D_2}) u_2, u_1 \rangle, \quad (58)$$

which holds for every pair of solutions  $u_i \in H^1(\Omega)$  to (1) with  $D = D_i, i = 1, 2$ .

By choosing in the above identity  $u_1(\cdot) = \Gamma^{D_1}(\cdot, y) l, u_2(\cdot) = \Gamma^{D_2}(\cdot, w) m$  with  $y, w \in S_{2\rho_0}$ , the first member of (58) coincides with the second member

of (40), so that, recalling the asymptotic estimate (37) and the hypothesis (27), we obtain the following smallness estimate

$$|(\Gamma^{D_2} - \Gamma^{D_1})(y, w)m \cdot l| \leq C \frac{\epsilon}{\rho_0}, \quad \text{for every } y, w \in S_{2\rho_0}, \quad (59)$$

where  $C > 0$  only depends on  $M_0, \alpha, M_1, \alpha_0, \gamma_0, \bar{\lambda}, \bar{\mu}, \tau, M$ .

This first smallness estimate is then propagated up to the points  $y_h, w_h$ , with a technical construction based on iterated application of the three spheres inequality.

**THEOREM 3.5** (Lower bound on the function  $(\Gamma^{D_2} - \Gamma^{D_1})$ ). *Under the notation of Lemma 3.1, let*

$$y_h = P - he_3. \quad (60)$$

*For every  $i = 1, 2, 3$ , there exists  $\lambda_w \in \{\frac{2}{3}, \frac{3}{4}, \frac{4}{5}\}$  and there exists  $\tilde{h} \in (0, \frac{1}{2})$  only depending on  $M_0, \alpha, \alpha_0, \gamma_0, \bar{\lambda}, \bar{\mu}, \eta_0, \tau, M$ , such that*

$$|(\Gamma^{D_2} - \Gamma^{D_1})(y_h, w_h)e_i \cdot e_i| \geq \frac{C}{h}, \quad \text{for every } h, 0 < h < \tilde{h} \text{ dist}(P, D_2), \quad (61)$$

where

$$w_h = P - \lambda_w h e_3, \quad (62)$$

and  $C > 0$  only depends on  $M_0, \alpha, M_1, \alpha_0, \gamma_0, \bar{\lambda}, \bar{\mu}, \tau, M$  and  $\eta_0$ .

The proof of this key result will be given in Section 4.

We are now in position to prove the main result of this paper.

*Proof of Theorem 2.2.* By the upper bound (57), with  $l = m = e_i$  for  $i \in \{1, 2, 3\}$ , and the lower bound (61), we have

$$C \leq \epsilon^{C_1 \left(\frac{h}{\rho_0}\right)^{C_2}}, \quad \text{for every } h, 0 < h \leq \min\{\bar{h}\rho_0, \tilde{h}d(P, D_2)\} \quad (63)$$

where  $C, C_1, C_2$  only depend on  $M_0, \alpha, M_1, \alpha_0, \gamma_0, \bar{\lambda}, \bar{\mu}, \tau, M$  and  $\eta_0$ . By our regularity assumptions on the domains, there exists  $\tilde{C} > 0$ , only depending on  $M_0, \alpha, M_1$ , such that

$$d(P, D_2) \leq \text{diam}(\Omega) \leq \tilde{C}\rho_0. \quad (64)$$

Set  $h^* = \min\left\{\frac{\bar{h}}{\tilde{C}}, \tilde{h}\right\}$ . Then inequality (63) holds for every  $h$  such that  $h \leq h^*d(P, D_2)$ , with  $h^*$  only depending on  $M_0, \alpha, M_1, \alpha_0, \gamma_0, \bar{\lambda}, \bar{\mu}, \tau, M$  and  $\eta_0$ . Taking the logarithm in (63) and recalling that  $\epsilon \in (0, 1)$ , we obtain

$$h \leq C\rho_0 \left(\frac{1}{|\log \epsilon|}\right)^{\frac{1}{C_2}}, \quad \text{for every } h, 0 < h \leq h^*d(P, D_2), \quad (65)$$

In particular, choosing  $h = h^*d(P, D_2)$ , we have

$$d(P, D_2) \leq C\rho_0 \left( \frac{1}{|\log \epsilon|} \right)^{\frac{1}{c_2}}. \quad (66)$$

The thesis follows from Lemma 3.1.  $\square$

#### 4. Proof of Theorem 3.5

Let us recall that we have chosen a cartesian coordinate system with origin  $P \equiv O$  and  $e_3 = -\nu$ , where  $\nu$  is the unit outer normal to  $D_1$  at  $P$ .

Let  $\mathbb{C}_0 = \mathbb{C}(O)$  be the constant Lamé tensor, having Lamé moduli  $\lambda \equiv \lambda(O)$ ,  $\mu \equiv \mu(O)$ , and let  $\mathbb{C}_0^{D_1} = \mathbb{C}^{D_1}(O)$  be the constant Lamé tensor with Lamé moduli  $\lambda \equiv \lambda^{D_1}(O)$ ,  $\mu \equiv \mu^{D_1}(O)$ . Moreover, let us introduce the elasticity tensors  $\mathbb{C}_0^+ = \mathbb{C}_0\chi_{\mathbb{R}^3} + \mathbb{C}_0^{D_1}\chi_{\mathbb{R}^3_+}$ ,  $\mathbb{C}_0^1 = \mathbb{C}_0\chi_{\mathbb{R}^3 \setminus D_1} + \mathbb{C}_0^{D_1}\chi_{D_1}$ .

Let  $\Gamma, \Gamma_0, \Gamma_0^+, \Gamma_0^{D_1}$  be the fundamental matrices associated to the tensors  $\mathbb{C}, \mathbb{C}_0, \mathbb{C}_0^+, \mathbb{C}_0^1$ , respectively.

In the above notation, we may write, for every  $m, l \in \mathbb{R}^3$ ,  $|l| = |m| = 1$ ,

$$\begin{aligned} |(\Gamma^{D_2} - \Gamma^{D_1})(y_h, w_h)m \cdot l| &\geq |(\Gamma_0^+ - \Gamma_0)(y_h, w_h)m \cdot l| - |(\Gamma^{D_2} - \Gamma)(y_h, w_h)m \cdot l| - \\ &\quad - |(\Gamma - \Gamma_0)(y_h, w_h)m \cdot l| - |(\Gamma_0^+ - \Gamma_0^{D_1})(y_h, w_h)m \cdot l| - \\ &\quad - |(\Gamma_0^{D_1} - \Gamma^{D_1})(y_h, w_h)m \cdot l|. \end{aligned} \quad (67)$$

The following Lemma, which is a straightforward consequence of Proposition 9.3 and formula (9.11), derived in [3], gives a positive lower bound for the term  $|(\Gamma_0^+ - \Gamma_0)(y_h, w_h)e_i \cdot e_i|$ ,  $i = 1, 2, 3$ , for a suitable  $w_h$ .

LEMMA 4.1. *For every  $i = 1, 2, 3$ , there exists  $\lambda_w \in \{\frac{2}{3}, \frac{3}{4}, \frac{4}{5}\}$  such that*

$$|(\Gamma_0^+(y_h, w_h) - \Gamma_0(y_h, w_h))e_i \cdot e_i| \geq \frac{\mathcal{C}}{h}, \quad \text{for every } h > 0, \quad (68)$$

where  $\mathcal{C} > 0$  only depends on  $\alpha_0, \gamma_0, \bar{\lambda}, \bar{\mu}, \eta_0$ .

From now on, let  $\lambda_w$  be chosen accordingly to the above lemma and let  $h \leq \frac{1}{2} \min\{d(P, D_2), \frac{\rho_0}{\sqrt{1+M_0^2}}\}$ .

Term  $\Gamma^{D_2} - \Gamma$ .

Let us consider the vector valued function

$$v(x) = (\Gamma^{D_2} - \Gamma)(x, w_h)m. \quad (69)$$

Let us set  $\rho = d(P, D_2)$ . Since  $d(w_h, P) = \lambda_w h \leq h \leq \frac{\rho}{2}$ , we have that  $d(w_h, D_2) \geq d(P, D_2) - d(w_h, P) \geq \frac{\rho}{2}$ . Therefore  $v(x)$  is a solution to the Lamé system

$$\operatorname{div}_x(\mathbb{C}\nabla_x v(x)) = 0, \quad \text{in } B_{\frac{\rho}{2}}(w_h). \quad (70)$$

By the regularity estimate

$$\sup_{B_{\frac{\rho}{4}}(w_h)} |v(x)| \leq \frac{C}{\rho^{\frac{3}{2}}} \left( \int_{B_{\frac{\rho}{2}}(w_h)} |v(x)|^2 \right)^{\frac{1}{2}}, \quad (71)$$

with  $C$  only depending on  $\alpha_0, \gamma_0, \bar{\lambda}, \bar{\mu}$ , and by applying the asymptotic estimates (37) to  $\Gamma^{D_2}$  and  $\Gamma$ , it follows that

$$\sup_{B_{\frac{\rho}{4}}(w_h)} |v(x)| \leq \frac{C}{\rho}, \quad (72)$$

where  $C > 0$  only depends on  $M_0, \alpha, \alpha_0, \gamma_0, \bar{\lambda}, \bar{\mu}, \tau, M$ .

Since  $d(y_h, w_h) = (1 - \lambda_w)h \leq \frac{h}{3} \leq \frac{\rho}{6}$ ,  $y_h \in B_{\frac{\rho}{4}}(w_h)$  and

$$|(\Gamma^{D_2} - \Gamma)(y_h, w_h)m \cdot l| = |v(y_h) \cdot l| \leq \frac{C}{\rho} = \frac{C}{d(P, D_2)}, \quad (73)$$

for every  $l, m \in \mathbb{R}^3$ ,  $|l| = |m| = 1$ , with  $C$  only depending on  $M_0, \alpha, \alpha_0, \gamma_0, \bar{\lambda}, \bar{\mu}, \tau, M$ .

Term  $\Gamma_0^{D_1} - \Gamma^{D_1}$ .

By the same arguments seen in the proof of Proposition 3.3, we have that, for every  $y, w \in \mathbb{R}^3$ ,  $y \neq w$ , and for every  $l, m \in \mathbb{R}^3$ ,

$$\int_{\mathbb{R}^3} \mathbb{C}^1 \nabla \Gamma^{D_1}(\cdot, y) l \cdot \nabla \Gamma_0^{D_1}(\cdot, w) m = \Gamma_0^{D_1}(y, w) m \cdot l, \quad (74)$$

$$\int_{\mathbb{R}^3} \mathbb{C}_0^1 \nabla \Gamma^{D_1}(\cdot, y) l \cdot \nabla \Gamma_0^{D_1}(\cdot, w) m = \Gamma^{D_1}(y, w) m \cdot l. \quad (75)$$

Choosing  $y = y_h$  and  $w = w_h$ , we have

$$(\Gamma_0^{D_1} - \Gamma^{D_1})(y_h, w_h) m \cdot l = \int_{\mathbb{R}^3} (\mathbb{C}^1 - \mathbb{C}_0^1) \nabla \Gamma^{D_1}(\cdot, y_h) l \cdot \nabla \Gamma_0^{D_1}(\cdot, w_h) m = J + J_0, \quad (76)$$

with

$$J = \int_{D_1} (\mathbb{C}^{D_1} - \mathbb{C}_0^{D_1}) \nabla \Gamma^{D_1}(\cdot, y_h) l \cdot \nabla \Gamma_0^{D_1}(\cdot, w_h) m, \quad (77)$$

$$J_0 = \int_{\mathbb{R}^3 \setminus D_1} (\mathbb{C} - \mathbb{C}_0) \nabla \Gamma^{D_1}(\cdot, y_h) l \cdot \nabla \Gamma_0^{D_1}(\cdot, w_h) m. \quad (78)$$

Let us estimate  $J$ . We have trivially

$$|J| \leq C(I_1 + I_2), \quad (79)$$

where  $C > 0$  only depends on  $M_0, \alpha, \alpha_0, \gamma_0, \bar{\lambda}, \bar{\mu}, \tau, M$  and

$$I_1 = \int_{|x| \geq \rho_0} \frac{|(\mathbb{C}^{D_1} - \mathbb{C}_0^{D_1})(x)|}{|x - y_h|^2 |x - w_h|^2}, \quad (80)$$

$$I_2 = \int_{|x| \leq \rho_0} \frac{|(\mathbb{C}^{D_1} - \mathbb{C}_0^{D_1})(x)|}{|x - y_h|^2 |x - w_h|^2}. \quad (81)$$

Let us first estimate  $I_1$ . Since  $h \leq \frac{\rho_0}{2}$  and  $|x| \geq \rho_0$ , we have that  $|x - y_h| \geq |x| - |y_h| = |x| - h \geq \frac{|x|}{2}$  and similarly  $|x - w_h| \geq \frac{|x|}{2}$ , so that

$$I_1 \leq C \int_{|x| \geq \rho_0} \frac{1}{|x|^4} = \frac{C}{\rho_0}, \quad (82)$$

with  $C$  only depending on  $\bar{\lambda}, \bar{\mu}$ . To estimate  $I_2$ , we use the fact that

$$|(\mathbb{C}^{D_1} - \mathbb{C}_0^{D_1})(x)| = |\mathbb{C}^{D_1}(x) - \mathbb{C}^{D_1}(O)| \leq \frac{C}{\rho_0^\tau} |x|^\tau, \quad (83)$$

with  $C$  only depending on  $M$ , so that

$$I_2 \leq \frac{C}{\rho_0^\tau} (I_2' + I_2''), \quad (84)$$

where

$$I_2' = \int_A \frac{|x|^\tau}{|x - y_h|^2 |x - w_h|^2}, \quad (85)$$

$$I_2'' = \int_B \frac{|x|^\tau}{|x - y_h|^2 |x - w_h|^2}, \quad (86)$$

with  $A = \{|x| \leq \rho_0, |x| < 6|y_h - w_h|\}$ ,  $B = \{6|y_h - w_h| \leq |x| \leq \rho_0\}$ .

We perform the change of variables  $x = |y_h - w_h|z$  in  $I_2'$ , obtaining

$$I_2' \leq 6^\tau |y_h - w_h|^{\tau-1} \int_{|z| \leq 6} \left( z - \frac{y_h}{|y_h - w_h|} \right)^{-2} \left( z - \frac{w_h}{|y_h - w_h|} \right)^{-2}. \quad (87)$$

Since the integral on the right hand side is bounded by an absolute constant, see [16, Chapter 2, Section 11], we have that

$$I_2' \leq C |y_h - w_h|^{\tau-1}, \quad (88)$$

with  $C$  only depending on  $\tau$ .

For every  $x \in B$ , we have

$$|x| \geq 6|y_h - w_h| = 6h(1 - \lambda_w) \geq \frac{6}{5}h, \quad (89)$$

so that

$$|x| \leq |x - y_h| + |y_h| = |x - y_h| + h \leq |x - y_h| + \frac{5}{6}|x|. \quad (90)$$

Hence

$$\frac{1}{6}|x| \leq |x - y_h|, \quad (91)$$

and, similarly,

$$\frac{1}{6}|x| \leq |x - w_h|. \quad (92)$$

By (91)–(92), we have

$$I_2'' \leq 6^4 \int_B |x|^{\tau-4} \leq C \int_{6|y_h-w_h|}^{\rho_0} r^{\tau-2} dr \leq C|y_h - w_h|^{\tau-1}, \quad (93)$$

where  $C$  is an absolute constant.

From (79), (82), (84), (88), (93) and noticing that  $|y_h - w_h| = h(1 - \lambda_w) \geq \frac{h}{5}$ , we have

$$|J| \leq \frac{C}{h} \left( \frac{h}{\rho_0} + \left( \frac{h}{\rho_0} \right)^\tau \right), \quad (94)$$

where  $C$  only depends on  $M_0, \alpha, \alpha_0, \gamma_0, \bar{\lambda}, \bar{\mu}, \tau, M$ .

The term  $J_0$  is estimated analogously with  $\tau$  replaced by 1, and therefore, by (76),

$$|(\Gamma_0^{D_1} - \Gamma^{D_1})(y_h, w_h)m \cdot l| \leq \frac{C}{h} \left( \frac{h}{\rho_0} + \left( \frac{h}{\rho_0} \right)^\tau \right), \quad (95)$$

where  $C$  only depends on  $M_0, \alpha, \alpha_0, \gamma_0, \bar{\lambda}, \bar{\mu}, \tau, M$ .

*Term  $\Gamma_0^+ - \Gamma_0^{D_1}$ .*

Arguing similarly to the proof of Proposition 3.3, we have that, for every  $y, w \in \mathbb{R}^3$ ,  $y \neq w$ , and for every  $l, m \in \mathbb{R}^3$ ,

$$\begin{aligned} (\Gamma_0^+ - \Gamma_0^{D_1})(y, w)m \cdot l &= \int_{\mathbb{R}^3} (\mathbb{C}_0^{D_1} - \mathbb{C}_0)(\chi_{D_1} - \chi_{\mathbb{R}_+^3}) \nabla \Gamma_0^{D_1}(\cdot, y)l \cdot \nabla \Gamma_0^+(\cdot, w)m = \\ &= \int_{D_1 \setminus \mathbb{R}_+^3} (\mathbb{C}_0^{D_1} - \mathbb{C}_0) \nabla \Gamma_0^{D_1}(\cdot, y)l \cdot \nabla \Gamma_0^+(\cdot, w)m - \\ &\quad - \int_{\mathbb{R}_+^3 \setminus D_1} (\mathbb{C}_0^{D_1} - \mathbb{C}_0) \nabla \Gamma_0^{D_1}(\cdot, y)l \cdot \nabla \Gamma_0^+(\cdot, w)m. \end{aligned} \quad (96)$$

Therefore

$$|(\Gamma_0^+ - \Gamma_0^{D_1})(y_h, w_h)m \cdot l| \leq C \int_{A \cup B} \frac{1}{|x - y_h|^2 |x - w_h|^2}, \quad (97)$$



where

$$A = \left\{ x \in (\mathbb{R}_+^3 \setminus D_1) \cup (D_1 \setminus \mathbb{R}_+^3) \mid |x| \geq \frac{\rho_0}{\sqrt{1+M_0^2}} \right\}, \quad (98)$$

$$B = \left\{ x \in (\mathbb{R}_+^3 \setminus D_1) \cup (D_1 \setminus \mathbb{R}_+^3) \mid |x| \leq \frac{\rho_0}{\sqrt{1+M_0^2}} \right\}, \quad (99)$$

and  $C$  only depends on  $M_0, \alpha, \alpha_0, \gamma_0, \bar{\lambda}, \bar{\mu}, \tau, M$ . By our hypotheses,  $h \leq \frac{\rho_0}{2\sqrt{1+M_0^2}}$ . Hence, for every  $x \in A$ ,  $h \leq \frac{|x|}{2}$ ,  $|x - y_h| \geq |x| - h \geq \frac{|x|}{2}$ , and similarly  $|x - w_h| \geq \frac{|x|}{2}$ , so that

$$\int_A \frac{1}{|x - y_h|^2 |x - w_h|^2} \leq 16 \int_{|x| \geq \frac{\rho_0}{\sqrt{1+M_0^2}}} \frac{1}{|x|^4} = \frac{C}{\rho_0}, \quad (100)$$

with  $C$  only depending on  $M_0$ .

By the local representation of the boundary of  $D_1$  as a  $C^{1,\alpha}$  graph, it follows that

$$B \subset \left\{ x \in \mathbb{R}^3 \mid |x'| \leq \frac{\rho_0}{\sqrt{1+M_0^2}}, |x_3| \leq \frac{M_0}{\rho_0^\alpha} |x'|^{1+\alpha} \right\}. \quad (101)$$

By performing the change of variables  $z = \frac{x}{h}$ , we have

$$\begin{aligned} & \int_B \frac{1}{|x - y_h|^2 |x - w_h|^2} \\ & \leq \int_{|x'| \leq \frac{\rho_0}{\sqrt{1+M_0^2}}} dx' \int_{-\frac{M_0}{\rho_0^\alpha} |x'|^{1+\alpha}}^{\frac{M_0}{\rho_0^\alpha} |x'|^{1+\alpha}} \frac{1}{|x - y_h|^2 |x - w_h|^2} dx_3 \\ & = \frac{1}{h} \int_{|z'| \leq \frac{\rho_0}{h\sqrt{1+M_0^2}}} dz' \int_{-M_0 \left(\frac{h}{\rho_0}\right)^\alpha |z'|^{1+\alpha}}^{M_0 \left(\frac{h}{\rho_0}\right)^\alpha |z'|^{1+\alpha}} \frac{1}{|z + e_3|^2 |z + \lambda_w e_3|^2} dz_3 \\ & \leq \frac{1}{h} \int_{\mathbb{R}^2} dz' \int_{-M_0 \left(\frac{h}{\rho_0}\right)^\alpha |z'|^{1+\alpha}}^{M_0 \left(\frac{h}{\rho_0}\right)^\alpha |z'|^{1+\alpha}} \frac{1}{|z + e_3|^2 |z + \lambda_w e_3|^2} dz_3. \end{aligned} \quad (102)$$

Denoting

$$D(z) = (|z'|^2 + (z_3 + 1)^2) (|z'|^2 + (z_3 + \lambda_w)^2), \quad (103)$$

we have

$$\int_B \frac{1}{|x - y_h|^2 |x - w_h|^2} \leq \frac{1}{h} (J_1 + J_2), \quad (104)$$

where

$$J_1 = \int_{|z'| \leq \left(\frac{1}{3M_0}\right)^{\frac{1}{1+\alpha}}} dz' \int_{-M_0 \left(\frac{h}{\rho_0}\right)^\alpha |z'|^{1+\alpha}}^{M_0 \left(\frac{h}{\rho_0}\right)^\alpha |z'|^{1+\alpha}} \frac{1}{D(z)} dz_3, \quad (105)$$

$$J_2 = \int_{|z'| \geq \left(\frac{1}{3M_0}\right)^{\frac{1}{1+\alpha}}} dz' \int_{-M_0 \left(\frac{h}{\rho_0}\right)^\alpha |z'|^{1+\alpha}}^{M_0 \left(\frac{h}{\rho_0}\right)^\alpha |z'|^{1+\alpha}} \frac{1}{D(z)} dz_3. \quad (106)$$

To estimate  $J_1$ , let us notice that, recalling  $h \leq \frac{\rho_0}{2\sqrt{1+M_0^2}}$ ,

$$|z_3 + \lambda_w| \geq \lambda_w - |z_3| \geq \frac{2}{3} - M_0 \left(\frac{h}{\rho_0}\right)^\alpha |z'|^{1+\alpha} \geq \frac{1}{3}, \quad (107)$$

and, a fortiori,  $|z_3 + 1| \geq \frac{1}{3}$ . Hence  $D(z) \geq \frac{1}{3^4}$  and

$$J_1 \leq 3^4 \int_{|z'| \leq \left(\frac{1}{3M_0}\right)^{\frac{1}{1+\alpha}}} 2M_0 \left(\frac{h}{\rho_0}\right)^\alpha |z'|^{1+\alpha} dz' = C \left(\frac{h}{\rho_0}\right)^\alpha, \quad (108)$$

with  $C$  only depending on  $M_0$  and  $\alpha$ .

To estimate  $J_2$  we use the trivial inequality  $D(z) \geq |z'|^4$  when  $\alpha < 1$ , and  $D(z) \geq C(M_0)|z'|^{\frac{7}{2}}$  when  $\alpha = 1$ , so obtaining

$$J_2 \leq C \left(\frac{h}{\rho_0}\right)^\alpha, \quad (109)$$

with  $C$  only depending on  $M_0$  and  $\alpha$ .

By (97), (100), (104), (108), (109), we have

$$|(\Gamma_0^+ - \Gamma_0^{D_1})(y_h, w_h)m \cdot l| \leq \frac{C}{h} \left( \frac{h}{\rho_0} + \left(\frac{h}{\rho_0}\right)^\alpha \right), \quad (110)$$

with  $C$  only depending on  $M_0$ ,  $\alpha$ ,  $\alpha_0$ ,  $\gamma_0$ ,  $\bar{\lambda}$ ,  $\bar{\mu}$ ,  $\tau$ ,  $M$ .

*Term  $\Gamma - \Gamma_0$ .*

Similarly to the proof of Proposition 3.3, we have that, for every  $y, w \in \mathbb{R}^3$ ,  $y \neq w$ ,

$$(\Gamma_0 - \Gamma)(y, w)m \cdot l = \int_{\mathbb{R}^3} (C - C_0) \nabla \Gamma(\cdot, y) l \cdot \nabla \Gamma_0(\cdot, w) m. \quad (111)$$

From this identity, the arguments of the proof are similar to those seen to estimate the addend  $J_0$  in the expression of  $(\Gamma_0^{D_1} - \Gamma^{D_1})(y_h, w_h)l \cdot m$  given by (76), so that

$$|(\Gamma_0 - \Gamma)(y_h, w_h)m \cdot l| \leq \frac{C}{h} \left(\frac{h}{\rho_0}\right), \quad (112)$$

with  $C$  only depending on  $M_0, \alpha, \alpha_0, \gamma_0, \bar{\lambda}, \bar{\mu}, \tau, M$ .

*Conclusion.* Finally, from (67), (68), (73), (95), (110), (112), we have

$$\begin{aligned} & |(\Gamma^{D_2} - \Gamma^{D_1})(y_h, w_h)e_i \cdot e_i| \geq \\ & \geq \frac{C}{h} \left( 1 - C_1 \frac{h}{d(P, D_2)} - C_2 \frac{h}{\rho_0} - C_3 \left( \frac{h}{\rho_0} \right)^\alpha - C_4 \left( \frac{h}{\rho_0} \right)^\tau \right), \end{aligned} \quad (113)$$

with  $C_i, i = 1, \dots, 4$ , only depending on  $M_0, \alpha, \alpha_0, \gamma_0, \bar{\lambda}, \bar{\mu}, \tau, M$  and  $C$  only depending on  $\alpha_0, \gamma_0, \bar{\lambda}, \bar{\mu}$  and  $\eta_0$ . Let  $h_1 = \min\{\frac{1}{2}, \frac{1}{5C_1}\}$ ,  $h_2 = \min\left\{\frac{1}{2\sqrt{1+M_0^2}}, \frac{1}{5C_2}, \frac{1}{(5C_3)^\frac{1}{\alpha}}, \frac{1}{(5C_4)^\frac{1}{\tau}}\right\}$ . If  $h \leq \min\{h_1 d(P, D_2), h_2 \rho_0\}$ , then

$$|(\Gamma^{D_2} - \Gamma^{D_1})(y_h, w_h)e_i \cdot e_i| \geq \frac{C}{5h}, \quad (114)$$

Let  $\tilde{h} = \min\left\{h_1, \frac{h_2}{C}\right\}$ , where  $\tilde{C}$  has been introduced in (64). Then inequality (61) holds for every  $h$  such that  $h \leq \tilde{h} d(P, D_2)$ .

## Acknowledgements

The second author is supported by Università degli Studi di Trieste FRA 2014 “Problemi inversi per PDE, unicità, stabilità, algoritmi” and by GNAMPA of Istituto Nazionale di Alta Matematica.

## REFERENCES

- [1] G. ALESSANDRINI, *Stable determination of conductivity by boundary measurements*, Appl. Anal. **27** (1988), 153–172.
- [2] G. ALESSANDRINI AND M. DI CRISTO, *Stable determination of an inclusion by boundary measurements*, SIAM J. Math. Anal. **37** (2005), 200–217.
- [3] G. ALESSANDRINI, M. DI CRISTO, A. MORASSI, AND E. ROSSET, *Stable determination of an inclusion in an elastic body by boundary measurements*, SIAM J. Math. Anal. **46** (2014), 2692–2729.
- [4] G. ALESSANDRINI AND K. KIM, *Single-logarithmic stability for the Calderón problem with local data*, J. Inverse Ill-Posed Probl. **20** (2012), 389–400.
- [5] G. ALESSANDRINI AND E. SINCICH, *Cracks with impedance, stable determination from boundary data*, Indiana Univ. Math. J. **62** (2013), 947–989.
- [6] E. BERETTA, E. FRANCONI, A. MORASSI, E. ROSSET, AND S. VESSELLA, *Lipschitz continuous dependence of piecewise constant Lamé coefficients from boundary data: the case of non flat interfaces*, Inverse Problems **30** (2014), Paper 125005.

- [7] E. BERETTA, E. FRANCINI, AND S. VESSELLA, *Uniqueness and Lipschitz stability for the identification of Lamé parameters from boundary measurements*, Inverse Problems and Imaging **8** (2014), 611–644.
- [8] M. DI CRISTO, *Stable determination of an inhomogeneous inclusion by local boundary measurements*, J. Comput. Appl. Math. **198** (2007), 414–425.
- [9] M. DI CRISTO AND S. VESSELLA, *Stable determination of the discontinuous conductivity coefficient of a parabolic equation*, SIAM J. Math. Anal. **42** (2010), 183–217.
- [10] M. DI CRISTO AND S. VESSELLA, *Stability analysis of an inverse parabolic problem with discontinuous variable coefficient*, Proc. Roy. Soc. Edinburgh Sect. A-Mathematics **141** (2011), 975–999.
- [11] S. HOFMANN AND S. KIM, *The Green function estimates for strongly elliptic systems of second order*, Manuscripta Math. **124** (2007), 139–172.
- [12] M. IKEHATA, *Reconstruction of inclusion from boundary measurements*, J. Inverse Ill-Posed Probl. **10** (2002), 37–65.
- [13] M. IKEHATA, G. NAKAMURA, AND K. TANUMA, *Identification of the shape of the inclusion in the anisotropic elastic body*, Appl. Anal. **72** (1999), 17–26.
- [14] V. ISAKOV, *On uniqueness of recovery of a discontinuous conductivity coefficient*, Comm. Pure. Appl. Math. **XLI** (1988), 865–877.
- [15] Y. LI AND L. NIRENBERG, *Estimates for elliptic systems from composite material*, Comm. Pure. Appl. Math. **LVI** (2003), 892–925.
- [16] C. MIRANDA, *Partial differential equations of elliptic type*, Springer-Verlag, New York-Berlin, 1970.
- [17] G. UHLMANN AND J.-N. WANG, *Reconstructing discontinuities using complex geometrical optics solutions*, SIAM J. Appl. Math. **68** (2008), 1026–1044.
- [18] G. UHLMANN, J.-N. WANG, AND C.-T. WU, *Reconstruction of inclusions in an elastic body*, J. Math. Pures Appl. **91** (2009), 569–582.

Authors' addresses:

Antonino Morassi  
Dipartimento Politecnico di Ingegneria e Architettura  
Università degli Studi di Udine  
via Cotonificio, 114  
33100 Udine, Italy  
E-mail: [antonino.morassi@uniud.it](mailto:antonino.morassi@uniud.it)

Edi Rosset  
Dipartimento di Matematica e Geoscienze  
Università degli Studi di Trieste  
via Valerio, 12/1  
34127 Trieste, Italy  
E-mail: [rossedi@units.it](mailto:rossedi@units.it)

Received November 23, 2015

Accepted March 24, 2016