




UNIVERSITÀ
DEGLI STUDI DI TRIESTE
Department of Life Sciences
Psychology Unit *Gaetano Kanizsa*

TSPC2016

Proceedings of the
Trieste Symposium on Perception and Cognition
November 4



edited by
Paolo Bernardis
Carlo Fantoni
Walter Gerbino

EUT

Habituation of the alertness response in crayfish is modulated by stimuli with socially relevant information

Andrea Caputi, Walter Gerbino, Piero Giulio Giulianini, Cinzia Chiandetti

Department of Life Sciences, University of Trieste, Trieste, Italy

Keywords: *Procambarus Clarkii*, Looming, Habituation, Social Information, Invertebrate

Habituation consists of a decreased response to repeated stimulation and not involving fatigue or sensory adaptation [1], and is common to many living entities, from single cells to the most complex organisms [2-3]. It has a considerable evolutionary value, since it allows an organism to disregard irrelevant repetitive stimuli.

Here, we tested whether two different stimuli, with possible different social values, may have a different impact on habituation in individuals of crayfish *Procambarus clarkii*, a highly aggressive crustacean often fighting with conspecifics for the control of shelter and resources [4].

The stimuli consisted in a red upright Y-shaped figure, resembling a crayfish displaying its aggressive posture (high intensity stimulus condition), and a red downright Y-shaped figure, identical to the other but rotated of 180° and resembling the animals' relaxed posture (low intensity stimulus condition). A habituation paradigm was used, in which two groups of 24 animals were presented with nine consecutive looming stimulations, thus creating the illusion of an approaching stimulus. Furthermore, a tenth looming presentation of a completely different stimulus (a circular blue moebius) was used to test for a spontaneous recovery of the response. As dependent measure, we recorded the duration of the alert reaction to the approaching shape (measured in video frames, recorded at 29 fps).

Crayfish showed habituation in both conditions ($p=0.002$). The animals tested in the low intensity condition abruptly habituated to the repeating stimulus, and then showed a strong response recovery to the new stimulus ($p=0.001$). By contrast, crayfish in the high intensity condition maintained a consistently higher alertness during the administration of the repeating upright Y-shaped figure ($M=11.61 \pm 1.10$) as compared to those in the low intensity condition ($M=8.72 \pm 1.21$). Moreover, crayfish in the high intensity condition showed no spontaneous recovery to the approaching stimulus when it was changed in shape and colour ($p=0.120$).

Our data confirm previous findings on looming in crayfish [5-6] and indicate that they responded as if they faced real opponents [7]. Indeed, the more sustained pattern of response evoked by the upright stimulus seems to be due to a different social valence between the two schematic stimuli, with one stimulus being perceived as more threatening than the other. Our results show that *P. clarkii* is sensitive to the abstract visual representation of conspecifics.

Acknowledgment: Thanks to Gianmarco Maldarelli for help in testing the animals. The project was supported by a DIAnet FSE fellowship and a UniTs-FRA2015 to CC.

1. Rankin CH, Abrams T, Barry RJ, Bhatnagar S, Clayton D, Colombo J, Coppola G, Geyer MA, Glanzman DL, Marsland S, McSweeney F, Wilson DA, Wu CF, Thompson RF (2009). Habituation revisited: an updated and revised description of the behavioural characteristics of habituation. *Neurobiology of Learning and Memory*, 92: 135-138.
2. Thompson RF (2009). Habituation: a history. *Neurobiology of Learning and Memory*, 92: 127-134.
3. Carew TJ, Sahley CL (1986). Invertebrate learning and memory: from behaviour to molecules. *Annual Reviews of Neuroscience*, 9: 435-487.
4. Issa FA, Adamson DJ, Edwards DH (1999). Dominance hierarchy formation in juvenile crayfish *Procambarus clarkii*. *Journal of experimental biology*, 202: 3497-3506.
5. Beall SP, Langley DJ, Edwards DH (1990). Inhibition of escape tailflip in crayfish during backward walking and the defense posture. *Journal of experimental biology*, 152: 577-582.
6. Glantz RM (1974). Defense reflex and motion detector responsiveness to approaching targets: the motion detector trigger to the defense reflex pathway. *Journal of Comparative Physiology*, 95: 297-314.
7. Sato D, Nagayama T (2012). Development of agonistic encounters in dominance hierarchy formation in juvenile crayfish. *The Journal of Experimental Biology*, 215: 120-1217.