



CERN-EP/2016-293
2017/06/17

CMS-SUS-16-009

Search for supersymmetry in the all-hadronic final state using top quark tagging in pp collisions at $\sqrt{s} = 13$ TeV

—Supplemental Material—

The CMS Collaboration

Submitted to Physical Review D




Figure 1: The purity of the top quark tagger as a function of the reconstructed top quark p_T . The purity is defined as the fraction of reconstructed top quarks that are matched to a generator-level hadronically decaying top quark within a cone of $\Delta R = 0.4$, and was measured in a sample of simulated one-lepton $t\bar{t}$ events. The following event selection requirements were applied: $N_j \geq 4$ for $p_T > 30$ GeV, $|\eta| < 2.4$ and $N_j \geq 2$ for $p_T > 50$ GeV, $|\eta| < 2.4$; $N_b \geq 1$; and $E_T^{\text{miss}} > 200$ GeV.




Figure 2: The event mistag rate of the top quark tagger as a function of E_T^{miss} in the $Z \rightarrow \nu\nu$ simulated sample, with the following event selection requirements applied: $N_j \geq 4$ for $p_T > 30$ GeV, $|\eta| < 2.4$ and $N_j \geq 2$ for $p_T > 50$ GeV, $|\eta| < 2.4$; no electrons, muons, or isolated tracks; $\Delta\phi(E_T^{\text{miss}}, \text{jets})$ matching preselection requirements; and $N_b \geq 1$.




Figure 3: The t -tagged event fraction measured in data and $t\bar{t}$ simulated samples, as a function of the reconstructed top quark candidate p_T . The data are selected from a single muon dataset with the following selection applied: events pass noise filters; $N_j \geq 4$ for $p_T > 30$ GeV, $|\eta| < 2.4$ and $N_j \geq 2$ for $p_T > 50$ GeV, $|\eta| < 2.4$; at least one muon with $p_T > 45$ GeV, $|\eta| < 2.1$; no electrons; muon $p_T + E_T^{\text{miss}} > 150$ GeV; $\Delta\phi(E_T^{\text{miss}}, \text{jets})$ matching preselection requirements; $N_b \geq 1$; and $E_T^{\text{miss}} > 20$ GeV. We also require the presence of at least one candidate from the t -tagger in the event. This candidate can be either: (i) a trijet candidate, composed of three jets with $p_T > 30$ GeV that are within $\Delta R = 1.5$ of the candidate four-momentum, (ii) a dijet candidate, composed of two jets that are within $\Delta R = 1.5$ of the candidate four-momentum, one of which should have a mass between 70 and 110 GeV, or (iii) a monojet candidate, which is simply a single jet with a mass between 110 and 220 GeV. The candidate used to compute the t -tagged event fraction is the candidate whose mass is closest to the top quark mass. The t -tagged event fraction then is defined as the fraction of events for which this top quark candidate satisfies all requirements of the top quark tagging algorithm. The error bar depicts the statistical uncertainty. The ratio of the t -tagged event fraction in data and simulated $t\bar{t}$ is shown in the lower plot, indicating good agreement.