

Reverse sensitivity testing: What does it take to break the model?[☆]

Silvana M. Pesenti^{a,*}, Pietro Millossovich^{a,b}, Andreas Tsanakas^a

^a Cass Business School, City, University of London United Kingdom

^b DEAMS, University of Trieste Italy

ARTICLE INFO

Article history:

Received 23 January 2018

Accepted 4 October 2018

Available online 10 October 2018

Keywords:

Robustness and sensitivity analysis

Risk management

Value-at-Risk

Expected Shortfall

Stress testing

ABSTRACT

Sensitivity analysis is an important component of model building, interpretation and validation. A model comprises a vector of random input factors, an aggregation function mapping input factors to a random output, and a (baseline) probability measure. A risk measure, such as Value-at-Risk and Expected Shortfall, maps the distribution of the output to the real line. As is common in risk management, the value of the risk measure applied to the output is a decision variable. Therefore, it is of interest to associate a critical increase in the risk measure to specific input factors. We propose a global and model-independent framework, termed 'reverse sensitivity testing', comprising three steps: (a) an output stress is specified, corresponding to an increase in the risk measure(s); (b) a (stressed) probability measure is derived, minimising the Kullback–Leibler divergence with respect to the baseline probability, under constraints generated by the output stress; (c) changes in the distributions of input factors are evaluated. We argue that a substantial change in the distribution of an input factor corresponds to high sensitivity to that input and introduce a novel sensitivity measure to formalise this insight. Implementation of reverse sensitivity testing in a Monte Carlo setting can be performed on a single set of input/output scenarios, simulated under the baseline model. Thus the approach circumvents the need for additional computationally expensive evaluations of the aggregation function. We illustrate the proposed approach through numerical examples with a simple insurance portfolio and a model of a London Insurance Market portfolio used in industry.

1. Introduction

1.1. Problem framing and contribution

Risk managers often use complex quantitative models as decision support tools. Of fundamental importance is sensitivity analysis, which is concerned with characterising and providing insight regarding the relation between inputs and outputs. Sensitivity analysis can have different aims, including identifying the most influential inputs (*factor prioritisation*), detecting the direction of input/output relationships, and inferring model structure; see Saltelli

et al. (2008) and Borgonovo and Plischke (2016) for comprehensive reviews. For the specific aim of factor prioritisation, a *sensitivity measure* is typically used, assigning a sensitivity score to each input. When model inputs are subject to uncertainty, *global* sensitivity measures are used, considering the whole possible space of multivariate input scenarios.

In this paper we develop a sensitivity analysis framework appropriate for contexts where the following considerations, typical in several fields, including probabilistic safety assessment, reliability analysis and financial/insurance risk management (Aven & Nøklund, 2010; Gouriéroux, Laurent, & Scaillet, 2000; Saltelli & Tarantola, 2002; Tsanakas & Millossovich, 2016), hold:

- Model inputs are uncertain, hence sensitivity and uncertainty analyses are interlinked and global sensitivity analysis methods are called for.
- A decision criterion is derived by applying a *risk measure* on the distribution of the output. Risk measures are functional mappings random variables to the real line (Artzner, Delbaen, Eber, & Heath, 1999; Szegö, 2005). Risk measures are used in a variety of operations research and risk analysis applications, with Value-at-Risk (VaR) and Expected Shortfall (ES – also

^{*} Earlier versions have been presented at the University of Trieste, the University of Bologna, the *Insurance Data Science Conference* (London), at the *41st AMASES Conference* (Cagliari), the *RSS Applied Probability Section on Probability in Actuarial Science* (London), the *Actuarial and Financial Mathematics Conference* (Brussels), the *21st International Congress on Insurance: Mathematics and Economics* (Vienna) and the workshop on *Risk Measurement and Regulatory Issues in Business* (Montreal). The authors are grateful for discussions with Vali Asimit, Markus Gesmann, Michael Gordy, Andrew Smith and Ruodu Wang. We thank two anonymous referees for their comments, which helped greatly improve the paper.

^{*} Corresponding author.

E-mail address: Silvana.Pesenti@cass.city.ac.uk (S.M. Pesenti).

known as CVaR) particularly popular choices; indicatively see [Rockafellar and Uryasev \(2002\)](#), [Tapiero \(2005\)](#), [Gotoh and Takano \(2007\)](#), [Ahmed, Çakmak, and Shapiro \(2007\)](#), and [Asimit, Bignozzi, Cheung, Hu, and Kim \(2017\)](#).

- The value of the risk measure, applied on the output distribution, gives an indication of criticality for the system whose uncertainty is analysed. For example, in the context of financial risk management, high values of output risk measures may indicate that a portfolio is not admissible, e.g. due to regulatory constraints ([Artzner et al., 1999](#)). In the context of probabilistic safety assessment, legislation postulates acceptable probabilities of failure, e.g. of fatality numbers exceeding a threshold ([Borgonovo & Cillo, 2017](#)).
- The relationship between model inputs and outputs is complex and not necessarily given in analytical form. Furthermore, evaluations of the model are computationally expensive inducing the need to estimate sensitivity measures from a single sample of input and output scenarios ([Beckman & McKay, 1987](#); [Plischke, Borgonovo, & Smith, 2013](#)).

We propose a sensitivity analysis framework, adapted to the above context, termed *reverse sensitivity testing*. We work in the standard setting of sensitivity analysis, where a number of random *input factors* are mapped to a random *output* via an *aggregation function*. The *baseline probability measure* summarises the distribution of inputs and output in the current specification of the model. Our reverse sensitivity testing framework comprises the following steps. First, an output stress is defined, corresponding to an increase in the value of the output risk measure. We focus on the widely used risk measures VaR and ES. The increase in the value of the risk measure is specified so as to produce a stress that is problematic to a decision maker. For example, in a capital management context, a stress on VaR may lead to a situation where insufficient assets are available to satisfy regulatory requirements.

Secondly, a *stressed probability measure* is derived. This is a probability (a) under which the risk measure applied to the model output is at its stressed level and (b) that minimises the Kullback–Leibler (KL) divergence subject to appropriate constraints on the output probability distribution. Thus the stressed probability leads to the most plausible alternative model, under which the output distribution is subjected to the required stress. We derive analytical solutions of the stressed probability measure under an increase of VaR and ES. The form of the solutions allows for numerically efficient implementation via a single set of Monte Carlo simulations.

Finally, the distribution of individual input factors is examined under the baseline and stressed models. Substantial changes in the distribution of a particular input indicate a large sensitivity to that input. A new class of *reverse sensitivity measures* is introduced, quantifying these input changes, and extended to control for statistical dependence between inputs. The sensitivity measures are then used to identify the most influential input factors; in a sense, those factors that may be responsible for ‘breaking the model’.

1.2. Relation to the literature

Prominent sensitivity analysis methods use a (Hoeffding) decomposition of the output variance ([Saltelli, 2002](#); [Saltelli, Chan, & Scott, 2000](#); [Saltelli et al., 2008](#); [Sobol, 1993](#); [Wagner, 1995](#)), as well as moment independent approaches ([Borgonovo, 2007](#); [Borgonovo, Castaings, & Tarantola, 2011](#)). Alternative methods consider partial derivatives of statistical functionals of the output distribution in the direction of parameters of interest, see [Glasserman and Liu \(2010\)](#) for expectation-type and [Hong \(2009\)](#) and [Tsanakas and Millosovich \(2016\)](#) for percentile-based functionals.

The sensitivity measures we propose in the present paper reflect the joint distribution of individual input factors and out-

put; hence our proposed method remains formally within the unifying framework discussed by [Borgonovo, Hazen, and Plischke \(2016\)](#) and thus are (distantly) related to variance-based and moment-independent sensitivity measures. Nonetheless, our proposed sensitivity measures are conceptually different compared to variance-based metrics and other current approaches in the literature. First, our approach involves an assessment of output uncertainty via tail-risk measures rather than the variance. Second, we adopt a reverse approach of stressing the output and then evaluating the impact on the inputs. Our method allows for flexibility in the stress level on the output, giving a nuanced picture of the sensitivity of input factors. Furthermore, the sensitivity measures we propose can take both negative and positive values, indicating the direction in which input factors affect the output. Thus, we view our proposed sensitivity analysis framework as complementary rather than competing with established methods, as it aims to address different questions.

Conceptually, the reverse direction (from output to input) of the proposed method, is related to *regionalised sensitivity analysis* methods ([Osidele & Beck, 2004](#); [Spear, Grieb, & Shang, 1994](#)). However, there is a key difference between regionalised sensitivity analysis and our approach: in the former, states of the output are identified that are ‘out of control’, while in the latter what is ‘out-of-control’ are not individual states but specifications of the output distribution. The numerical tractability of our framework in a Monte Carlo setting is akin to [Beckman and McKay \(1987\)](#).

In the practice of financial risk management and regulation, *reverse stress testing*, starting with a stressed output state and studying the corresponding surface of scenarios that provide the adverse outcome, is frequently used ([BCBS, 2013](#); [EIOPA, 2009](#)). For example, “reverse stresses that result in a depletion of capital...” ([Lloyd’s, 2016](#)) are used in the validation of insurance risk models. The academic literature on reverse stress testing is relatively sparse, with a recent focus towards identifying most likely stress scenarios ([Breuer, Jandačka, Mencía, & Summer, 2012](#); [Glasserman & Xu, 2014](#); [McNeil & Smith, 2012](#)). Our approach differs from reverse stress testing, in that we consider most influential factors in relation to changes in the output distribution and not a particular output state.

The KL-divergence has been widely used in financial risk management, in particular in the context of model uncertainty, where several plausible specifications of the probability measure may co-exist. For example, [Breuer and Csizsár \(2013\)](#), [Glasserman and Xu \(2014\)](#), and [Blanchet, Lam, Tang, and Yuan \(2017\)](#) consider the worst-case probability measure with respect to all probabilities lying within a KL-divergence radius of the baseline probability. In contrast, reflecting our focus on sensitivity rather than model uncertainty, we consider the probability measure with minimal KL-divergence that satisfies given constraints. Our approach is closely related to the work of [Cambou and Filipović \(2017\)](#) with probability set constraints and [Weber \(2007\)](#) with risk measure constraints, see [Section 3](#) for a detailed comparison.

1.3. Structure of the paper

In [Section 2](#), some preliminaries on risk measures and the KL-divergence are given. In [Section 3](#), the optimisation problem yielding stressed probability measures is stated and solved under constraints arising from different risk measures, with emphasis on VaR and ES. Explicit solutions allow easy implementation and inspection of the distributional changes arising. Furthermore, we discuss an extension where the distributions of multiple outputs are stressed. The solutions and their properties are illustrated through an example of a non-linear insurance portfolio evaluated using Monte Carlo simulation.

In [Section 4](#) we propose metrics tailored to the proposed reverse sensitivity testing approach. A comparison study of the proposed *reverse sensitivity measures* with moment independent and variance-based sensitivity measures is conducted, illustrating differences between the concepts but also demonstrating coherent sensitivity rankings of inputs. In addition, a generalisation of reverse sensitivity measures is proposed, with the aim of controlling for dependence between input factors.

[Section 5](#) demonstrates the applicability of the reverse sensitivity testing framework to a commercially used insurance portfolio model.

[Section Appendix A](#) is devoted to a comparison of the stressed and the baseline probability measures through stochastic order relations, in order to establish formal properties of the proposed framework. We find that the distribution of the output under the baseline probability is first-order stochastically dominated by that under the stressed probability. A similar dominance relation is provided for input factors, under the assumption of a non-decreasing aggregation function and positive dependence between input factors. Moreover, stressed probability measures stemming from different stress severities lead to stochastically ordered input factors and output. All proofs are provided in [Appendix B](#).

2. Preliminaries

We consider a measurable space (Ω, \mathcal{A}) and denote by \mathcal{P} the set of all probability measures on (Ω, \mathcal{A}) . For a random variable Z on (Ω, \mathcal{A}) we write $F_Z^Q(\cdot) = Q(Z \leq \cdot)$ for its distribution under $Q \in \mathcal{P}$, and similarly, $E^Q(\cdot)$ for its expectation. Throughout, we use the Kullback–Leibler divergence (*KL-divergence*, [Kullback & Leibler, 1951](#)) as a measure of discrepancy between two probability measures. For $Q^1, Q^2 \in \mathcal{P}$, the KL-divergence, also known as relative entropy, of Q^1 with respect to Q^2 is defined by

$$D_{\text{KL}}(Q^1 \| Q^2) = \begin{cases} \int_{+\infty} \frac{dQ^1}{dQ^2} \log \left(\frac{dQ^1}{dQ^2} \right) dQ^2 & \text{if } Q^1 \ll Q^2 \\ \text{otherwise,} & \end{cases}$$

where we use the convention that $0 \log(0) = 0$. The KL-divergence is non-negative, vanishes if and only if $Q^1 \equiv Q^2$, and is in general not symmetric ([Cover & Thomas, 2012](#); [Kullback, 1959](#)). The KL-divergence is a special case of the class of *f-divergences*, first introduced by [Ali and Silvey \(1966\)](#), for the choice $f(x) = x \log(x)$, $x > 0$. For a given convex function f , the *f-divergence* of Q^1 with respect to Q^2 , for any $Q^1, Q^2 \in \mathcal{P}$, is defined through $D_f(Q^1 \| Q^2) = \int f\left(\frac{dQ^1}{dQ^2}\right) dQ^2$.

Risk measures are tools used in risk management, which associate to every random variable a real number. The application of risk measures leads to a classification of different levels of risk severities, see [Artzner et al. \(1999\)](#) and [Föllmer and Schied \(2011\)](#) for an overview. Moments, such as the mean and standard deviation, can be seen as risk measures. In recent years, percentile-based risk measures ([Acerbi, 2002](#)) have become prominent, with the most commonly used risk measures being Value-at-Risk (VaR) and Expected Shortfall (ES). These risk measures are used extensively in financial regulation for the calculation of capital requirements, specifically VaR for European insurance companies, [EIOPA \(2009\)](#), and ES for banks, [BCBS \(2012, 2013\)](#).

The VaR at level $\alpha \in [0, 1]$ of a random variable Z is defined as the left α -quantile of the distribution of Z , $\text{VaR}_\alpha^Q(Z) = F_Z^{Q,-1}(\alpha) = \inf\{z \in \mathbb{R} \mid F_Z^Q(z) \geq \alpha\}$, where, as usual, $\inf \emptyset = +\infty$. In particular, the essential supremum of Z is $\text{ess sup}^Q Z = F_Z^{Q,-1}(1)$. The ES (also CVaR) of Z at level $\alpha \in [0, 1]$ is defined by

$$\text{ES}_\alpha^Q(Z) = \frac{1}{1-\alpha} \int_\alpha^1 \text{VaR}_u^Q(Z) du = \frac{1}{1-\alpha} E^Q\left(\left(Z - \text{VaR}_\alpha^Q(Z)\right)_+\right)$$

$$+ \text{VaR}_\alpha^Q(Z),$$

where, in the second representation, $\text{VaR}_\alpha^Q(Z)$ can be replaced by any α -quantile of F_Z^Q . Unlike VaR, the ES takes into account the whole tail of the distribution of Z , that is all realisations larger than $\text{VaR}_\alpha^Q(Z)$. See [Föllmer and Schied \(2011\)](#) for a comparison of the two risk measures.

Shortfall risk measures, associated with utility-type arguments, are defined through $\rho^Q(Z) = \inf\{z \in \mathbb{R} \mid E^Q(\ell(Z - z)) \leq z_0\}$ for $Q \in \mathcal{P}$, where ℓ is a non-decreasing, non-constant and convex loss function while z_0 is a point in the interior of the range of ℓ ([Föllmer & Schied, 2002](#)). Examples of shortfall risk measures include entropic risk measures, [Gerber \(1974\)](#), and the class of generalised quantiles called expectiles ([Bellini, Klar, Müller, & Rosazza Gianin, 2014](#); [Newey & Powell, 1987](#)).

3. Deriving the stressed model

3.1. Problem statement

We consider the standard setting of (reverse) sensitivity analysis, involving a (typically complicated) function, mapping model inputs to an output that is used in a decision making process. Mathematically, we define the *input factors* as a random vector $\mathbf{X} = (X_1, \dots, X_n)$ on the measurable space (Ω, \mathcal{A}) . The (measurable) function $g: \mathbb{R}^n \rightarrow \mathbb{R}$, is called the *aggregation function*, which gives, when applied to input factors \mathbf{X} , the one-dimensional random *output* of interest $Y = g(\mathbf{X})$. The variability of the output Y to changes in input factors is of fundamental importance in sensitivity analysis ([Borgonovo & Plischke, 2016](#); [Saltelli et al., 2008](#)) and the focus of this paper. We adopt throughout the convention that large values of the output correspond to adverse states.

We call the triple (\mathbf{X}, g, P) , the *baseline model* with *baseline probability measure* $P \in \mathcal{P}$. The probability P is seen as encoding current beliefs regarding (or software implementation of) the distribution of \mathbf{X} . Under the baseline probability P we suppress the superscript and write, for example, $F_Z(\cdot) = F_Z^P(\cdot)$ and $E(\cdot) = E^P(\cdot)$, and analogously for risk measures, $\text{VaR}_\alpha(\cdot) = \text{VaR}_\alpha^P(\cdot)$ and $\text{ES}_\alpha(\cdot) = \text{ES}_\alpha^P(\cdot)$. We call any $Q \in \mathcal{P}$ an *alternative probability measure* and (\mathbf{X}, g, Q) an *alternative model*. A Radon-Nikodym (RN) density is a non-negative random variable ζ on (Ω, \mathcal{A}) such that $E(\zeta) = 1$. We denote by Q^ζ the probability measure which is absolutely continuous with respect to P with RN-density ζ , that is, $\zeta = \frac{dQ^\zeta}{dP}$.

The starting point of reverse sensitivity analysis is to define a *stress* on the distribution of the output that would be problematic to a decision maker, such as a risk manager or regulator. For example, one may require that the probability of a particular event, representing system failure, increases to an extent that the risk of failure is no longer acceptable. Specific stress definitions using different risk measures are discussed in [Sections 3.2–3.6](#). Subsequently, we call (\mathbf{X}, g, Q) a *stressed model* with *stressed probability measure* $Q \in \mathcal{P}$ if, under Q , the output Y fulfils a set of probabilistic constraints (the stress) and Q has minimal KL-divergence with respect to P . Thus, a stressed probability measure is defined as a solution to

$$\min_{Q \in \mathcal{P}} D_{\text{KL}}(Q \| P),$$

s.t. constraints on the distribution of Y under Q hold. (1)

The optimisation problem (1) is robust in the sense that convergence in the KL-divergence implies weak convergence of the probability measures, [Gibbs and Su \(2002\)](#). This means that an alternative probability which satisfies the constraints of (1) and is close in KL-divergence to the stressed probability, is also close to the stressed probability in the Lévy metric.

Optimisation problem (1) under linear (i.e. moment) constraints was first studied in the seminal paper by [Csiszár \(1975\)](#). In the

context of financial risk management, in particular when risk measures are used, optimisation problem (1) involves non-linear constraints and Csiszár's theory cannot be applied. Relevant research includes Cambou and Filipović (2017) who consider the optimisation problem for general f -divergences and probability set constraints. Weber (2007) works with bounded random variables and considers risk measure constraints such as ES and short-fall risk measures, see Sections 3.3 and 3.4 for a more detailed comparison.

3.2. Probability constraints

Before studying problem (1) with constraints involving the risk measures of Section 2, we consider stresses under which the probabilities of (adverse) outcomes of $Y = g(\mathbf{X})$ are altered. These outcomes are captured by disjoint sets $B_1, \dots, B_I \subseteq \mathbb{R}$, each set B_i associated with an event $\{Y \in B_i\}$ where the system being studied is failing or 'out of control'. In a financial context, where Y is interpreted as a loss, one can identify B_i with a region of extreme losses.

The following result is an immediate consequence of Theorem 3.1 in Csiszár (1975); we also refer to Cambou and Filipović (2017).

Proposition 3.1. *Let $B_1, \dots, B_I \subseteq \mathbb{R}$ be disjoint Borel sets with $P(Y \in B_i) > 0$, $i = 1, \dots, I$, and $\alpha_1, \dots, \alpha_I > 0$ such that $\alpha_1 + \dots + \alpha_I \leq 1$. Then there exists a unique solution to*

$$\min_{Q \ll P} D_{\text{KL}}(Q \| P), \quad \text{s.t. } Q(Y \in B_i) = \alpha_i, \quad i = 1, \dots, I, \quad (2)$$

with RN-density given by $\zeta = \sum_{i=1}^I \frac{\alpha_i}{P(Y \in B_i)} \mathbb{1}_{\{Y \in B_i\}}$, where we write $\alpha_0 = 1 - \sum_{i=1}^I \alpha_i$ and $B_0 = (\bigcup_{i=1}^I B_i)^c$.

The RN-density ζ in Proposition 3.1 is a piecewise constant function of Y . This implies that all outcomes of Y within a set B_i receive the same probability re-weighting by the change to the stressed probability. In particular, if $\alpha_i > P(Y \in B_i)$, under the alternative probability Q the probability of all outcomes in B_i increases. Note that moving from the baseline to the stressed model might induce a new dependence structure in the input factors.

3.3. VaR constraints

We now consider optimisation problem (1) under a constraint on the risk measure VaR, applied to the output Y . A VaR constraint is not equivalent to a probability constraint of optimisation problem (2), when F_Y is not strictly increasing.

Proposition 3.2. *Let $0 < \alpha < 1$ and $q \in \mathbb{R}$ such that $\text{VaR}_\alpha(Y) < q < \text{ess sup } Y$ and consider the optimisation problem*

$$\min_{Q \ll P} D_{\text{KL}}(Q \| P), \quad \text{s.t. } \text{VaR}_\alpha^Q(Y) = q. \quad (3)$$

There exists a unique solution to (3) if and only if $P(q - \varepsilon < Y < q) > 0$ for all $\varepsilon > 0$. The RN-density of the solution is given by

$$\zeta = \frac{\alpha}{P(Y < q)} \mathbb{1}_{\{Y < q\}} + \frac{1 - \alpha}{P(Y \geq q)} \mathbb{1}_{\{Y \geq q\}}.$$

The assumption $P(q - \varepsilon < Y < q) > 0$ for all $\varepsilon > 0$, implies that q cannot be chosen arbitrarily. In particular, problem (3) does not have a solution if the distribution of Y is constant to the left of q (q excluded); this includes the (uncommon in practice) case where Y is a discrete random variable. This complication arises from using the constraint $\text{VaR}_\alpha^Q(Y) = q$ rather than $Q(Y \leq q) = \alpha$. If q cannot be chosen to fulfil the assumptions in Proposition 3.2, the form of ζ in Proposition 3.2 remains meaningful: by Proposition 3.1, it is the solution to an optimisation problem where the constraint $\text{VaR}_\alpha^Q(Y) = q$ is replaced by $Q(Y < q) = \alpha$.

The RN-density ζ of the solution to (3) is a non-decreasing function of Y since $\alpha \leq P(Y \leq \text{VaR}_\alpha(Y)) \leq P(Y < q)$. Hence, under the stressed probability, adverse realisations of the output are given higher probabilities of occurrence.

Remark. Propositions 3.1 and 3.2 hold true for any f -divergence with a strictly convex function f . In particular, the RN-densities ζ of the solutions of (2) and (3) are independent of the choice of f -divergence. We do not provide a proof for this statement, however the steps of the proofs of Propositions 3.1 and 3.2 can be closely retraced if one substitutes the KL-divergence with a general f -divergence. We refer to Ben-Tal, Den Hertog, De Waegenaere, Melenberg, and Rennen (2013) for robust linear optimisation with general f -divergence constraints.

Remark. Let $\text{VaR}_\alpha(Y) < q^* < \text{ess sup } Y$ be a stress for which the distribution function of Y under P is increasing and continuous in a neighbourhood of q^* , so that a solution of problem (3) exists for all q in that neighbourhood. Then, viewed as a function of q , the RN-density $\zeta(q)$ is a.s. continuous under P . Thus, the corresponding probability measure $Q^{\zeta(q)}$, solution of problem (3), converges in total variation distance to $Q^{\zeta(q^*)}$, implying that stressed models are robust with respect to stresses in VaR.

The explicit form of the RN-density in Proposition 3.2 (as well as the subsequent Propositions 3.3–3.4), allows easy implementation of the change of measure in a Monte Carlo simulation context similar to Beckman and McKay (1987). Note that the RN-density is a function of Y , in the sense that $\zeta(\omega) = \eta(Y(\omega))$, $\omega \in \Omega$, for a function η . Then, one can follow the process:

1. Sample M multivariate scenarios $\mathbf{x}^{(1)}, \dots, \mathbf{x}^{(M)}$ from \mathbf{X} under P . Calculate $\mathbf{y}^{(k)} = \mathbf{g}(\mathbf{x}^{(k)})$, $k = 1, \dots, M$.
2. Set $\zeta^{(k)} = \eta(\mathbf{y}^{(k)})$, $k = 1, \dots, M$.
3. The distributions of the output and inputs under the stressed measure Q are estimated by:

$$F_Y^Q(y) = \frac{1}{M} \sum_{k=1}^M \zeta^{(k)} \mathbb{1}_{\{\mathbf{y}^{(k)} \leq y\}}, \quad y \in \mathbb{R},$$

$$F_{X_i}^Q(x) = \frac{1}{M} \sum_{k=1}^M \zeta^{(k)} \mathbb{1}_{\{x_i^{(k)} \leq x\}}, \quad x \in \mathbb{R}, \quad i = 1, \dots, n.$$

Thus, the process of working out the distribution of input factors under the stressed measure is akin to importance sampling, with $\zeta^{(k)}$ playing the role of importance weights. Note that this calculation allows stressing the model without the need to re-simulate scenarios under Q , which can be of practical importance if evaluation of \mathbf{g} is computationally expensive. Straightforward implementation yields a computational cost of $M(n+1)$ for calculating the empirical distribution functions under the stressed model for the output and all input factors. In a simulation environment, convergence can be improved if Quasi Monte Carlo sampling is deployed, which is not a route we pursue here. Note that when simulations are computationally very expensive, meta-modelling techniques are often used in practice.

Example. The following insurance portfolio, similar to Example 1 in Tsanakas and Millossovich (2016), will be used as an illustrative example throughout the paper. An insurance company faces a loss L resulting from two lines of business. The two lines produce losses X_1, X_2 respectively, which are subject to the same multiplicative inflation factor X_3 , such that $L = X_3(X_1 + X_2)$. The insurance company has a reinsurance contract on the loss L with limit l and deductible d . The total portfolio loss for the insurance company is

$$Y = L - (1 - X_4) \min\{(L - d)_+, l\},$$

where X_4 captures the percentage lost due to a default of the reinsurance company.





Fig. 1. Left: simulated RN-density of the solution to (4). Right: simulated empirical distribution functions of the output under the baseline (dashed black) and the stressed (solid grey) model under problem (4).

In this example, the two lines of business X_1, X_2 are truncated Log-Normal(4.98, 0.23^2) and Gamma(100, 2) distributed, with respective means 150, 200 and standard deviations 35, 20. The truncation point for X_1 is chosen to be the 99.9% quantile. The multiplicative factor X_3 follows a truncated Log-Normal(0.05, 0.02^2) distribution with mean 1.05, standard deviation 0.02 and truncation point equal to the 99.9% quantile. The default loss X_4 is modelled through a Beta(0.125, 1.125) distribution, corresponding to mean 0.1 and standard deviation 0.2. To complete the specification of the joint distribution of \mathbf{X} , we further assume that X_1, X_2, X_3 are independent and X_4 is independent of (X_1, X_2, X_3) given L . Additionally, X_4 is taken to be dependent on the aggregated loss L through a Gaussian copula with correlation 0.6. The deductible of the insurance contract is $d = 380$ and the limit $l = 30$.

Consider optimisation problem (3) with a 10% increase in $\text{VaR}_{0.9}$, that is

$$\min_{Q \ll P} D_{\text{KL}}(Q||P), \quad \text{s.t. } \text{VaR}_{0.9}^Q(Y) = 1.1 \text{VaR}_{0.9}(Y). \quad (4)$$

The solution to the problem (4) is estimated from a Monte Carlo sample containing $M = 100,000$ simulated scenarios from (\mathbf{X}, Y) . Simulated values of the RN-density ζ are plotted in the left of Fig. 1, against samples from Y . It is seen that the RN-density is a non-decreasing function of Y and thus gives more weight to adverse outcomes of Y .

The empirical distribution functions of the total loss Y of the insurance company under the baseline probability (dashed black) and the stressed probability (solid grey) are displayed in the right of Fig. 1. The output distribution under the stressed probability lies beneath, and therefore first-order stochastically dominates, the distribution of Y under the baseline probability. We refer to Section Appendix A for a more detailed discussion of stochastic comparisons of stressed and baseline probabilities.

Fig. 2 displays the change in distribution of the input factors when moving from the baseline model to the stressed model. It can be seen that all factors under the stressed probability first-order stochastically dominate the corresponding inputs under the baseline probability. However, not all input factors are impacted the same: the distributions of inputs X_1 and X_4 are stressed more compared to the baseline model. This indicates a higher sensitivity to X_1 and X_4 , compared to X_2 and X_3 . A specific sensitivity measure reflecting the above observations is introduced in Section 4.

Table 1
Distributional characteristics of inputs and output under the baseline and stressed model under problem (4).

Sensitivity	Input factors				Output
	X_1	X_2	X_3	X_4	Y
Mean under P	150	200	1.05	0.10	362
Mean under Q	156	201	1.05	0.14	369
Standard deviation under P	35	20	0.02	0.20	36
Standard deviation under Q	41	21	0.02	0.24	45
Skewness under P	0.6	0.2	0.0	2.5	0.4
Skewness under Q	1.2	0.5	0.1	2.4	1.2
Excess kurtosis under P	0.5	0.1	-0.1	5.6	1.3
Excess kurtosis under Q	0.8	0.2	-0.1	3.9	1.4

Table 1 summarises basic characteristics of the change in the output and the input factors under the two models. Consistently with Fig. 2, it is seen that X_1 and X_4 are the most affected input factors by the change of probability measure. For example, under the stressed probability, X_1, X_4 are subject to a relative increase of the standard deviation of 17%, 20%, respectively.

3.4. VaR and ES constraints

This section addresses optimisation problem (1) with a constraint on both, VaR and ES. Adding to problem (3) a constraint on ES allows to stress the whole tail of the output distribution. Weber (2007) considers optimisation problem (1) with an ES constraint only. In that case there does not exist an analytic solution of the stressed probability and Weber (2007) offers a procedure for a numerical solution.

Proposition 3.3. *Let $0 < \alpha < 1$ and $q, s \in \mathbb{R}$ such that $\text{VaR}_\alpha(Y) < q < s < \text{ess sup } Y$. Assume the cumulant generating function of $Y|Y > q$ under P exists in a neighbourhood of 0 and that $E(Y|Y > q) < s$. Consider the optimisation problem*

$$\min_{Q \in \mathcal{P}} D_{\text{KL}}(Q||P), \quad \text{s.t. } \text{VaR}_\alpha^Q(Y) = q, \text{ES}_\alpha^Q(Y) = s. \quad (5)$$

Define the sets $A_1 = \{Y \geq q\}$ and $A_2 = \{Y > q\}$ and, for $i = 1, 2$, denote by θ_i^* the unique positive solution of the equation

$$E((Y - s)e^{\theta(Y - q)} | A_i) = 0. \quad (6)$$




Fig. 2. Empirical distribution functions of the input factors under the baseline (dashed black) and the stressed model (solid grey) under problem (4). The dark red line displays the difference of the distribution functions according to the axis on the right. (For interpretation of the references to colour in this figure legend, the reader is referred to the web version of this article.)

There exists a unique solution to problem (5) under either

1. $P(q - \varepsilon < Y < q) > 0$ for all $\varepsilon > 0$ and $E(e^{\theta_1^*(Y-q)} | A_1) \leq \frac{P(A_1^c)/P(A_1)}{\alpha/(1-\alpha)}$.
2. $P(Y = q) > 0$ and $P(q - \varepsilon < Y < q) = 0$ for some $\varepsilon > 0$, and $E(e^{\theta_2^*(Y-q)} | A_2) \geq \frac{P(A_2^c)/P(A_2)}{\alpha/(1-\alpha)}$.

The corresponding RN-density of the solution is

$$\zeta_i = \frac{\alpha}{P(A_i^c)} \mathbb{1}_{A_i^c} + \frac{1-\alpha}{E(e^{\theta_i^*(Y-q)} \mathbb{1}_{A_i})} e^{\theta_i^*(Y-q)} \mathbb{1}_{A_i}, \quad i = 1, 2.$$

Note that, compared to stressing solely the VaR, adding an ES constraint may provide a solution even for an output following a discrete distribution. The condition on the moment generating function in cases 1 and 2 restricts the choice of s and q , such that the stressed risk measure values cannot be chosen independently.

The RN-density of Proposition 3.3 under case 1, ζ_1 , is a non-decreasing function of Y . Under Proposition 3.3 case 2, the RN-density ζ_2 is not monotone. However, both RN-densities are exponentially increasing for realisations of Y exceeding q . Thus, under the stressed model, adverse outcomes of Y , such as tail events, admit a higher likelihood compared to the baseline model.

Remark. Let $\text{VaR}_\alpha(Y) < q^* < s^* < \text{ess sup } Y$ be a stress of VaR and ES for which the cumulant generating function of $Y|Y > q^*$ under P exists in a neighbourhood of 0, $E(Y|Y > q^*) < s^*$, the distribution function of Y under P is increasing and continuous in a neighbourhood of q^* and the second inequality in case 1 of

Proposition 3.3 holds strictly. Then, a solution of problem (6) exists in a neighbourhood of (q^*, s^*) . Viewed as a function of (q, s) , the RN-density $\zeta(q, s)$ is a.s. continuous under P . Thus, the corresponding probability measure $Q^{\zeta(q, s)}$, solution of problem (5), converges in total variation distance to $Q^{\zeta(q^*, s^*)}$, implying that stressed models are robust with respect to stresses in VaR and ES.

Example (continued). We consider optimisation problem (5) with a 10% increase in $\text{VaR}_{0.9}$ and a 13% increase in $\text{ES}_{0.9}$. Fig. 3 displays samples of the RN-density of the stressed probability measure, see Proposition 3.3 case 1. For high outcomes of the output Y , the RN-density ζ is exponentially increasing as a function of Y , hence inflates stressed tail probabilities. On the right hand side, the empirical distribution functions of the output under the baseline (dashed black) and the stressed model (solid grey) are shown.

Observe that the stressed distribution of the output appears similar to the stressed distribution of optimisation problem (4), see Fig. 1. This is due to the fact that increasing $\text{VaR}_{0.9}$ by 10% in optimisation problem (4), already leads to an increase of 8.5% in $\text{ES}_{0.9}$ under the stressed model. However, comparing Tables 1 and 2 it is seen that the standard deviation, skewness and kurtosis of Y increase more when stressing VaR and ES, compared to stressing VaR alone.

Similar to optimisation problem (4), the output and the input factors under the baseline probability are first-order stochastically dominated by the stressed probability, as can be seen in Figs. 3 and 4. We refer to Section Appendix A for a formal




Fig. 3. Left: simulated RN-density of the solution. Right: simulated empirical distribution functions of the output under the baseline (dashed black) and the stressed (solid grey) model under problem (5) with a 10% increase in $VaR_{0.9}$ and a 13% increase in $ES_{0.9}$.




Fig. 4. Empirical distribution functions of the input factors under the baseline (dashed black) and the stressed model (solid grey) under problem (5) with a 10% increase in $VaR_{0.9}$ and a 13% increase in $ES_{0.9}$. The dark red line displays the difference of the distribution functions according to the axis on the right. (For interpretation of the references to colour in this figure legend, the reader is referred to the web version of this article.)

Table 2

Distributional characteristics under the baseline and the stressed model under problem (5) with a 10% increase in $\text{VaR}_{0.9}$ and a 13% increase in $\text{ES}_{0.9}$.

Sensitivity	Input factors				Output	
	X_1	X_2	X_3	X_4	Y	
Mean under P	150	200	1.05	0.10	362	
Mean under Q	157	202	1.05	0.14	371	
Standard deviation under P	35	20	0.02	0.20	36	
Standard deviation under Q	43	21	0.02	0.26	50	
Skewness under P	0.6	0.2	0.0	2.5	0.4	
Skewness under Q	1.4	0.5	0.1	2.4	1.7	
Excess kurtosis under P	0.5	0.1	-0.1	5.6	1.3	
Excess kurtosis under Q	1.3	0.2	-0.1	3.7	2.8	

treatment of stochastic comparison of the stressed and baseline probabilities.

3.5. Shortfall risk measure constraints

Optimisation problem (1) with shortfall risk measure constraints is studied in Weber (2007) and is a direct application of Theorem 3.1 in Csiszár (1975). Nonetheless, we present the solution for completeness.

Proposition 3.4. *Let ρ be a shortfall risk measure with loss function ℓ and y_0 , and $q \in \mathbb{R}$ in the support of Y such that $E(\ell(Y - q)) < y_0$. If the moment generating function of $\ell(Y - q)$ exists in a neighbourhood of 0, then the optimisation problem*

$$\min_{Q \in \mathcal{P}} D_{\text{KL}}(Q \| P), \quad \text{s.t. } \rho^Q(Y) = q, \quad (7)$$

has a unique solution whose density is given by $\zeta = \frac{1}{E(e^{\theta^* \ell(Y - q)})} e^{\theta^* \ell(Y - q)}$, where θ^* is the unique positive solution of $E((\ell(Y - q) - y_0) e^{\theta \ell(Y - q)}) = 0$.

3.6. Multivariate output

Problem (1) can be extended to constraints on a multivariate output, that is, to $\mathbf{Y} = (Y_1, \dots, Y_k) = \mathbf{g}(\mathbf{X})$, for $k \in \mathbb{N}$ and aggregation function $\mathbf{g}: \mathbb{R}^n \rightarrow \mathbb{R}^k$. The general case under set constraints can be treated along the lines of Cambou and Filipović (2017), Section 7. We provide below an example based on two outputs.

Example (continued). We revisit the insurance portfolio example of Section 3.3 and view as output both the loss before reinsurance, L , and after reinsurance, Y . To shorten notation let $\nu_L = \text{VaR}_{0.9}(L)$ and $\nu_Y = \text{VaR}_{0.9}(Y)$ and consider the problem of stressing the VaR of L and Y by 10%

$$\min_{Q \in \mathcal{P}} D_{\text{KL}}(Q \| P), \quad \text{s.t. } \text{VaR}_{0.9}^Q(L) = 1.1 \nu_L, \text{VaR}_{0.9}^Q(Y) = 1.1 \nu_Y. \quad (8)$$

The constraints can be written as $Q(L \leq \nu_L) = 0.9$ and $Q(Y \leq \nu_Y) = 0.9$, since the distribution functions of L and Y are increasing and continuous around those stressed VaR. Thus, problem (8) can be solved straightforwardly using the Lagrange multiplier technique. The RN-density of the solution of (8) is constant on the four sets $\{L \leq \nu_L, Y \leq \nu_Y\}$, $\{L > \nu_L, Y \leq \nu_Y\}$, $\{L \leq \nu_L, Y > \nu_Y\}$ and $\{L > \nu_L, Y > \nu_Y\}$.

Fig. 5 displays the simulated empirical distribution of L and Y under the baseline model and the stressed model. Note that stressing both L and Y , in contrast to stressing only Y , do not lead to radical different results, as can be seen comparing Tables 1 and 3. This is because the stressed model, solution of problem (4), already induces an increase of 7% in $\text{VaR}_{0.9}(L)$. Moreover, stressing L , the portfolio loss before reinsurance, in addition to Y , reduces

Table 3

Distributional characteristics under the baseline and the stressed model under problem (8).

Sensitivity	Input factors				Output	
	X_1	X_2	X_3	X_4	L	Y
Mean under P	150	200	1.05	0.10	367	362
Mean under Q	157	202	1.05	0.13	376	370
Standard deviation under P	35	20	0.02	0.2	42	36
Standard deviation under Q	42	21	0.02	0.23	52	46
Skewness under P	0.6	0.2	0.0	2.5	0.4	0.4
Skewness under Q	1.3	0.5	0.1	2.4	1.2	1.3
Excess kurtosis under P	0.5	0.1	-0.1	5.6	0.3	1.3
Excess kurtosis under Q	0.9	0.2	-0.1	4.3	0.7	1.6

the importance of the default of the reinsurance displayed in the change of X_4 under Q .

4. Sensitivity measures for importance ranking

4.1. Definition of sensitivity measures

Plots such as the ones shown in Figs. 2 and 4 provide some insight into the sensitivity of the output risk measure to different input factors. In order to produce a ranking of inputs, it is necessary to introduce a formal sensitivity or importance measure; this is especially the case for models with large numbers of inputs for which succinct sensitivity summaries are needed. Here we develop a sensitivity measure that quantifies changes in input factors under the stressed model, compared to the baseline model.

Before proceeding to the definitions, some preliminaries are due. The random couple (V, W) is comonotonic if it can be written as $(V, W) \stackrel{d}{=} (F_V^{-1}(U), F_W^{-1}(U))$, for a uniformly distributed random variable U on $(0, 1)$. In contrast, (V, W) is countermonotonic if $(V, W) \stackrel{d}{=} (F_V^{-1}(U), F_W^{-1}(1 - U))$. Comonotonicity and countermonotonicity correspond to extremal positive and negative dependence structures respectively, for a random couple with fixed marginals (Müller & Stoyan, 2002). For a random variable V , we denote by $V_{|W}, V_{|W^\dagger}$ the random variables satisfying $V_{|W} \stackrel{d}{=} V_{|W^\dagger} \stackrel{d}{=} V$, such that $(V_{|W}, W)$ is comonotonic and $(V_{|W^\dagger}, W)$ is countermonotonic. Then for any $V' \stackrel{d}{=} V$ it holds that (Rüschendorf, 1983),

$$E(WV_{|W^\dagger}) \leq E(WV') \leq E(WV_{|W}).$$

The subsequent definition introduces a sensitivity measure that captures the extent to which a random variable is affected by a stress on the baseline model, that is, a change in probability measure.

Definition 4.1. Let Q^ξ be an alternative probability with RN-density $\xi = \frac{dQ^\xi}{dP}$. The sensitivity of a random variable Z to the change of measure is given by

$$\mathcal{S}(Z, \xi) = \begin{cases} \frac{E(Z\xi) - E(Z)}{\max_{\psi \stackrel{d}{=} \xi} E(Z\psi) - E(Z)} & E(Z\xi) \geq E(Z), \\ -\frac{E(Z\xi) - E(Z)}{\min_{\psi \stackrel{d}{=} \xi} E(Z\psi) - E(Z)} & \text{otherwise,} \end{cases}$$

where we use the convention $\pm \frac{\infty}{\infty} = \pm 1$ and $\frac{0}{0} = 0$.

In the definition of $\mathcal{S}(Z, \xi)$, the numerator $E(Z\xi) - E(Z)$ reflects the increase in the expectation of Z under the alternative model. The denominator normalises this difference, as it represents the maximal (or minimal) increase of the expectation of Z , under all alternative models with density ψ that are equal in distribution to ξ . This ensures normalisation of the sensitivity measure to $[-1, 1]$. If




Fig. 5. The simulated empirical distribution of the output under the baseline model (dashed black) and under the stressed model (solid grey) under problem (8). The left graph depicts the output before reinsurance, L , whereas the right plot shows the output after reinsurance, Y .

$S(Z, \xi) = 1$ or $S(Z, \xi) = -1$, the alternative model produces a maximal stress on the variable Z , representing a positive or negative impact of the changes in probability measure on Z respectively.

Note that $\arg \max_{\psi \stackrel{d}{=} \xi} E(Z\psi) = \xi_{|Z}$ and $\arg \min_{\psi \stackrel{d}{=} \xi} E(Z\psi) = \xi_{|Z^c}$. This allows for a straightforward calculation of the sensitivity measure. If working within a Monte Carlo simulation context, as is common in risk analysis, $\xi_{|Z}$, resp. $\xi_{|Z^c}$, can be simply obtained by re-arranging samples of ξ to be sorted in the same, resp. opposite, way as samples from Z . This context gives a different perspective on the constraint $\psi \stackrel{d}{=} \xi$: if simulated elements of ξ represent a particular scheme for re-weighting simulated scenarios, then ψ are vectors containing the same weights as ξ , but re-arranged to potentially prioritise different scenarios.

Next we define two sensitivity measures that are specific to the reverse sensitivity analysis framework of this paper.

Definition 4.2. Let Q^ζ be an alternative model with density $\zeta = \frac{dQ^\zeta}{dP} = \eta(Y)$, for a non-decreasing function η . For input X_i and output Y , we define the *reverse* and *forward sensitivity measures* Γ_i and Δ_i by:

$$\Gamma_i = S(X_i, \zeta),$$

$$\Delta_i = S(Y, \zeta_{|X_i}).$$

Here, $\zeta = \eta(Y)$ can be arrived at as the solution of optimisation problems (3), (5) or (7). Γ_i thus reflects the extent to which the reverse sensitivity test affects the expectation of the input factor X_i . Note that for $E(X_i\zeta) \geq E(X_i)$, we can write $\Gamma_i = \frac{\text{Cov}(X_i, \zeta)}{\max_{\psi \stackrel{d}{=} \zeta} \text{Cov}(X_i, \psi)}$, showing that the reverse sensitivity measure can also be understood as a dependence measure between X_i and Y . In this sense it is closely related to the dependence measure introduced by Kachapova and Kachapov (2012). Indeed, sensitivity measures considering the dependence between X_i and Y have a rich history in sensitivity analysis, for an overview see for example Borgonovo et al. (2016). Thus, in contrast to variance-based sensitivity measures, Γ_i and Δ_i can take both negative and positive values, indicating the direction in which input factors affect the output.

A possible criticism of the measure Γ_i and, by extension, the reverse sensitivity testing framework we propose, is as follows. Let Γ_i be high. This implies that stressing the model output Y leads to a substantial change in the distribution of the input factor X_i . However, this is not equivalent to a perturbation in the distribution of X_i leading to a sizeable stress in the distribution of the

output Y . Such a discrepancy, though uncommon, is theoretically possible and has been termed *probabilistic dissonance* (Cooke & van Noortwijk, 1999).

This motivates the introduction of the forward sensitivity measure Δ_i , as a companion measure to Γ_i . The definition of the forward sensitivity measure Δ_i is analogous to that of Γ_i , but with a focus on the change in the expectation of Y when perturbing the distribution of the model input X_i . Recall that $\zeta_{|X_i} = \arg \max_{\psi \stackrel{d}{=} \zeta} E(\psi X_i)$. Therefore, $\zeta_{|X_i}$ is a RN-density with the same distribution as ζ that has the most adverse effect on the input factor X_i . Thus Δ_i captures the impact of a change in the input X_i on the output Y . Reporting Δ_i along with Γ_i can thus produce warning signs of probabilistic dissonance.

Properties of the sensitivity measures Γ_i and Δ_i , reflecting their nature as dependence measures, are summarised below.

Proposition 4.3. Using the above introduced notation, the sensitivity measures Γ_i and Δ_i are well-defined and have the following properties:

1. $-1 \leq \Gamma_i, \Delta_i \leq 1$.
2. $\Gamma_i = \Delta_i = 0$, if X_i, Y are independent.
3. $\Gamma_i = \Delta_i = 1$, if (X_i, Y) is comonotonic.
4. $\Gamma_i = \Delta_i = -1$, if (X_i, Y) is counter-monotonic.
5. $\Gamma_i = \Delta_i \geq 0$, if (X_i, Y) are positively quadrant dependent¹.
6. $\Gamma_i = \Delta_i \leq 0$, if (X_i, Y) are negatively quadrant dependent¹.

Remark. Let ζ be the RN-density of the solution of problem (3) or (5). Then, for an input X_i with continuous distribution function, the corresponding reverse and forward sensitivity measures, Γ_i, Δ_i , are robust in the size of the stress as long as the RN-density is a.s. continuous in a neighbourhood of that stress. We refer to the remarks on robustness in Section 3.3 and 3.4 for details on the conditions required on the RN-density.

The above defined sensitivity measures focus on the difference of expectations under an alternative and the baseline model. If the interest lies in other distributional properties, such as tails, Definition 4.2 can be extended to consider monotone transformations of input factors. Specifically, one can calculate $S(u(X_i), \zeta)$, respectively $S(u(Y), \zeta_{|X_i})$, for an appropriately chosen non-decreasing function u . As the couple $(u(X_i), X_i)$ is comonotonic,

¹ These concepts are reviewed in Section Appendix A.




Fig. 6. Reverse (left plot) and forward (right plot) sensitivity measures $\Gamma_{i,v}, \Delta_{i,v}$ with a 10% increase in VaR and 13% increase in ES.

Table 4

Reverse sensitivity measure Γ_i for a 10% and 20% increase / decrease of $\text{VaR}_{0.9}$.

Input	$\lambda = 0.8$	$\lambda = 0.9$	$\lambda = 1.1$	$\lambda = 1.2$
X_1	-0.83	-0.85	0.88	0.90
X_2	-0.58	-0.51	0.36	0.34
X_3	-0.17	-0.17	0.15	0.14
X_4	-0.93	-0.72	0.60	0.68

the interpretation of the sensitivity measures remains unchanged. One particular example is the choice

$$u_v(X_i) = (X_i - F_{X_i}^{-1}(v))_+ - (F_{X_i}^{-1}(1-v) - X_i)_+, \quad 0.5 \leq v < 1. \quad (9)$$

For $v = 0.5$, the function $u_{0.5}$ is the identity and thus $S(u_{0.5}(X_i), \zeta) = \Gamma_i$, respectively $S(u_{0.5}(X_i), \zeta_{|X_i}) = \Delta_i$. When $v > 0.5$, the function u_v is zero whenever $X_i \in [F_{X_i}^{-1}(1-v), F_{X_i}^{-1}(v)]$ and linearly increasing otherwise. Thus, increasing v places higher emphasis on the tail behaviour of X_i . The random variable $u_v(Y)$ is defined and interpreted in a similar way.

We denote $\Gamma_{i,v} = S(u_v(X_i), \zeta)$ and $\Delta_{i,v} = S(u_v(Y), \zeta_{|X_i})$. It is easily seen that the properties of Proposition 4.3 still apply to $\Gamma_{i,v}, \Delta_{i,v}$. In addition, it holds that

$$S(u_v(aX_i + b), \zeta) = \text{sign}(a)S(u_v(X_i), \zeta),$$

such that the reverse sensitivity measure is invariant under linear transformations of input factors.

Example (continued). Fig. 6 displays the forward and reverse sensitivity measures $\Gamma_{i,v}, \Delta_{i,v}$ for $v \in [0.5, 0.999]$, for the stressed model arising from optimisation problem (5) with a 10% increase in VaR and a 13% increase in ES. Consistently with the example in Section 3.4, the highest sensitivity, for both reverse and forward measures, is displayed by X_1 , followed by X_4, X_2 and X_3 . Furthermore, the ranking is not affected by the level v and is thus not sensitive to emphasising the tails of the distributions. In the next section we present a situation where this no longer holds true.

To illustrate the impact on the sensitivity measure Γ_i of the size of the stress applied to the output risk measure, we fix $\alpha = 0.9$ and let $q = \lambda \text{VaR}_\alpha(Y)$, with λ ranging from 0.8 to 1.2.² Results are given in Table 4. For $\lambda < 1$ it is seen that the sensitivity measure Γ_i takes negative values. This is a result of applying a negative stress on Y , such that the RN-density ζ becomes a decreasing function of Y . The absolute value of the sensitivity measure responds

² Note that in general Proposition 3.2 only applies for $\lambda > 1$. However, continuity of Y in this example implies that the RN-density ζ of Proposition 3.2 is a solution to problem (3) even for $\lambda < 1$.

asymmetrically to positive ($\lambda > 1$) or negative ($\lambda < 1$) stresses on $\text{VaR}_\alpha(Y)$, in extremis even leading to a change in the ranking of inputs. This reflects a different sensitivity to input factors with view to increasing or decreasing the VaR of the output.

4.2. Comparison to other sensitivity measures

By the proposed reverse sensitivity measure Γ_i , we aim to quantify the extent to which a stress in the output distribution impacts different inputs. In the present section we compare Γ_i to the moment independent sensitivity measure introduced in Borgonovo (2007), and to the variance-based sensitivity measures, see Borgonovo and Plischke (2016) for an overview, which are designed to apportion the output variance to individual input factors. Specifically, the first order sensitivity index S_i , the total effects sensitivity index T_i and the moment independent sensitivity measure δ_i are respectively defined as

$$S_i = \frac{\text{Var}(E(Y|X_i))}{\text{Var}(Y)}$$

$$T_i = \frac{E(\text{Var}(Y|\mathbf{X}_{-i}))}{\text{Var}(Y)}$$

$$\delta_i = \frac{1}{2}E\left(\int |f_Y(y) - f_{Y|X_i}(y|X_i)| dy\right),$$

where $\mathbf{X}_{-i} = (X_1, \dots, X_{i-1}, X_{i+1}, \dots, X_n)$, f_Y and $f_{Y|X_i}$ denote the density of the output Y and the density of Y conditional on the input factor X_i , respectively. The measure S_i can be understood as the expected reduction in the variance of Y that would be achieved if input X_i could be fixed, whereas T_i is interpreted as the expected variance that would be left if all inputs but X_i could be fixed (Saltelli et al., 2008; Sobol, 1993; Wagner, 1995). The moment independent sensitivity δ_i can be seen as the expected shift of the output induced by fixing the input factor X_i (Borgonovo, 2007).

Thus, the interpretation of the reverse sensitivity measure Γ_i is quite different to that of S_i, T_i and δ_i , which are designed to answer different questions. Furthermore, Γ_i is designed with reference to a (tail) risk measure like VaR/ES and hence captures distributional impacts differently than the variance-based or the moment independent sensitivities, as is illustrated by the following numerical example.

Example (continued). We return to the simple insurance portfolio example of Section 3.3 (optimisation problem (4)), stressing $\text{VaR}_\alpha(Y)$ by 10%, for $\alpha = 0.5$ and $\alpha = 0.9$. The sensitivity indices S_i, T_i and δ_i are calculated, in addition to Γ_i , where the calculation of the variance-based sensitivities is carried out via estimation of the necessary conditional expectations from the existing Monte

Table 5

Comparison of the variance based sensitivity measures S_i, T_i , the moment independent sensitivity measure δ_i and the reverse sensitivity measure Γ_i with a 10% increase in VaR_α , for $\alpha = 0.5$ and $\alpha = 0.9$, for independent input vector \mathbf{X} .

Input	S_i	T_i	δ_i	Γ_i	
				$\alpha = 0.5$	$\alpha = 0.9$
X_1	0.71	0.73	0.46	0.87	0.89
X_2	0.24	0.26	0.21	0.41	0.36
X_3	0.03	0.03	0.06	0.15	0.15
X_4	0.00	0.01	0.04	0.09	0.07

Carlo sample by local polynomial regression and the estimation of δ_i utilises kernel smoothing as in [Borgonovo et al. \(2011\)](#).

In [Table 5](#), the sensitivity measures are reported for a variation of the model, where the aggregation function g and the marginal distributions of the inputs X_i are unchanged, but the vector \mathbf{X} is independent. Assuming that the input factors are independent facilitates comparison of Γ_i and δ_i with the variance based sensitivity measures, S_i and T_i . It can be seen that all sensitivity measures produce a consistent ranking, with X_4 assigned a very low sensitivity. Note that the first order sensitivities, S_i , sum up to 0.98, implying that the interaction terms are nearly as important as S_3 .

The sensitivity measure Γ_1 is slightly increasing in α , while Γ_2 is decreasing. This reflects the different tail characteristics of X_1 (LogNormal) and X_2 (Gamma); for a high α , the focus is on the right tail of Y , for which the heavier tail of X_1 is more important.

4.3. Controlling for dependence in the sensitivity measure

The literature on sensitivity indices has long been concerned with the implications for sensitivity analyses of statistical dependence between inputs. In particular for variance-based sensitivities, ([Oakley & O'Hagan, 2004](#); [Saltelli & Tarantola, 2002](#)), show that correlation between inputs can impact sensitivity measures in ways that do not reflect the functional dependencies in the model's aggregation function and are thus viewed as spurious; see [Section 4.3 in Borgonovo and Plischke \(2016\)](#) for more discussion of this topic and extensive references. Refinements of variance-based sensitivity indices have been proposed to address dependence between inputs, indicatively see [Xu and Gertner \(2008a\)](#), [Xu and Gertner \(2008b\)](#), and [Mara and Tarantola \(2012\)](#).

Our sensitivity measure Γ_i , similarly to many other sensitivity measures ([Borgonovo et al., 2016](#)), is evaluated on the joint distribution of (X_i, Y) . As a result, it does not control for dependence between inputs and is therefore subject to problems of confounding typical in multivariate analyses. In this section, we put forward a proposal for generalising the sensitivity measure Γ_i , in order to take into account such effects.

To proceed with the definition, denote by $N = \{1, \dots, n\}$ and $\mathbf{X}_T = (X_j)_{j \in T}$ for $T \subseteq N$. Consider now, for $T \subseteq N \setminus \{i\}$, the quantity $\mathcal{S}(X_i - E(X_i | \mathbf{X}_T), \zeta)$, measuring the reverse sensitivity to that part of X_i that is not already explained by the inputs \mathbf{X}_T . With the above in mind, we define the *k*th-order reverse sensitivity measure as

$$\Gamma_i^{(k)} = \frac{1}{c_k} \sum_{T \subseteq N \setminus \{i\}, |T|=k} \mathcal{S}(X_i - E(X_i | \mathbf{X}_T), \zeta), \quad k = 0, \dots, n-1,$$

where $c_k = \binom{n-1}{k}$. Thus $\Gamma_i^{(k)}$ represents the average reverse sensitivity to X_i , after controlling for all subsets of inputs of size k . Note the special cases $\Gamma_i^{(0)} = \Gamma_i$ and $\Gamma_i^{(n-1)} = \mathcal{S}(X_i - E(X_i | \mathbf{X}_{-i}), \zeta)$. If X_i is independent of \mathbf{X}_{-i} , then $\Gamma_i^{(k)} = \Gamma_i$ for all $k = 0, \dots, n-1$.

Example (continued). Continuing with the insurance portfolio example, we work out *k*th-order reverse sensitivity measures for

Table 6

The *k*th-order reverse sensitivity measure $\Gamma_i^{(k)}$, for $k = 0, \dots, 3$, of problem (4) for a 10% increase of $\text{VaR}_{0.9}(Y)$.

Input	$k = 0$	$k = 1$	$k = 2$	$k = 3$
X_1	0.88	0.81	0.76	0.71
X_2	0.36	0.30	0.29	0.31
X_3	0.15	0.13	0.12	0.13
X_4	0.60	0.48	0.34	0.18

$k = 0, \dots, 3$, with the RN-density ζ derived from problem (4) for an 10% increase of $\text{VaR}_{0.9}(Y)$. The results are summarised in [Table 6](#) and show that, as the order of the sensitivity measure increases, the sensitivity of some input factors is impacted more than that of others. This is particularly noticeable for X_4 : for $k = 3$, where all other input factors are controlled for, the sensitivity drops substantially.

Comparing the first column in [Table 6](#), that is Γ_i , with [Table 5](#), the sensitivity for the independent input vector, we see that the rank of X_2 and X_4 are reversed. The observed impact of dependence between inputs on the importance ranking of X_4 can be understood as follows. X_4 represents the percentage of reinsurance recovery lost due to default. In our baseline model, X_4 is dependent on $L = X_3(X_1 + X_2)$, with Gaussian copula correlation of 0.6. The correlation reflects the notion that such recovery losses are more likely under those scenarios when they are most needed (i.e. L is large), leading to a high sensitivity to X_4 .

5. Financial application: a London Insurance Market portfolio

In this section we demonstrate the use of the sensitivity measures $\Gamma_{i,v}$ and $\Delta_{i,v}$, in a more realistic insurance risk model with a higher number of inputs. This is a proprietary model of a London Insurance Market portfolio, currently in use by a participant in that market. We have been supplied by the model owner with a Monte Carlo sample of size $M = 500,000$, containing simulated observations from input factors $\mathbf{X} = (X_1, \dots, X_{72})$ and output Y . Each of the X_i 's represents a normalised loss for a particular part of the portfolio and is measured on the same scale. The output Y stands for the portfolio loss.

The aggregation function g is linear, specifically

$$Y = g(\mathbf{X}) = \sum_{j=1}^{72} w_j X_j,$$

for a vector of weights $\mathbf{w} = (w_1, \dots, w_{72})$. The linearity of g is not used for sensitivity calculations, since the reverse sensitivity testing framework makes no assumptions on the form of g . We do not have access to the joint probability distribution that was used to generate samples from the random vector \mathbf{X} ; in fact the distribution of \mathbf{X} is not given in closed form, as samples from \mathbf{X} are themselves outputs of a different model, which remains a completely black box to us.

We consider optimisation problem (5) with risk measure constraints on VaR and ES given by $q = \text{VaR}_{0.95}^Q(Y) = 1.08 \text{VaR}_{0.95}(Y)$ and $s = \text{ES}_{0.95}^Q(Y) = 1.1 \text{ES}_{0.95}(Y)$. In [Fig. 7](#), the reverse and forward sensitivity measures $\Gamma_{i,v}, \Delta_{i,v}$, for $v = 0.5$ and $v = 0.95$, are presented for all 72 inputs. The input factors are ordered according to $\Gamma_{i,0.5}$ and the sizes of the markers reflect the weights w_i attached to the individual input factors X_i .

Observations on the plot of [Fig. 7](#):

- The ranking of input factors according to $\Gamma_{i,0.5}$ and $\Gamma_{i,0.95}$ is not fully consistent; moving focus to the tails of input factors changes the order of the sensitivity measures. Hence, under the




Fig. 7. Reverse and forward sensitivity measures $\Gamma_{i,v}$, $\Delta_{i,v}$ for the London Insurance Market portfolio, for $v = 0.5$ and $v = 0.95$.

stressed model, for some input factors the expectation is affected more, while for others the impact is higher in the tail.

- For $v = 0.5$, the ranking produced by the reverse and forward sensitivity metrics is not equivalent. However, once the focus is moved towards the tails of risk factor distributions (e.g. $v = 0.95$), the discrepancy of the two sensitivity measures diminishes.
- There is no clear relation between the sizes of the markers and the ranking of input factors. This means that the sensitivity measure $\Gamma_{i,v}$ does not solely reproduce the size of the weight w_i .

To elaborate on the last of those points, in Fig. 8 (left), the reverse sensitivities $\Gamma_{i,0.95}$ are plotted against the weights w_i . There is a broadly increasing relation, which is not unreasonable. Given the linearity of the aggregation function, a higher weight w_i implies a higher local sensitivity $\frac{\partial g}{\partial x_i}$ (Borogovno & Plischke, 2016). But the relation is by no means deterministic: weight is a weak predictor of the reverse sensitivity measure $\Gamma_{i,v}$.

Furthermore, the reverse sensitivity measure does not only reflect the shape of the input risk factor distributions. In Fig. 8 (right), $\Gamma_{i,0.95}$ is displayed against the scaled percentiles $\frac{\text{VaR}_{0.95}(X_i)}{E(X_i)} - 1$, not showing a clear pattern. Hence the two plots in Fig. 8 demonstrate that the proposed reverse sensitivity measure does not reproduce easily observed characteristics of the aggregation function g or of the distributions of the inputs X_i .

6. Conclusions

We proposed a reverse sensitivity testing framework that is appropriate for contexts where model inputs are uncertain and the relationship between model inputs and outputs is complex and

not necessarily given in analytical form. At the core of the reverse sensitivity framework is a stress on the output distribution, corresponding to an increase in the value of a risk measure applied on the output and representing a plausible but adverse model change. This leads to stressed probabilities under which the distribution of the input factors (marginals and dependence structure) is altered such that the output distribution is subjected to the required stress.

We provided analytical solutions of the stressed probability measure under an increase of the VaR and ES risk measures. These explicit solutions facilitate straightforward implementation in a Monte Carlo simulation context and inspection of changes in the distributions of inputs. A new class of reverse sensitivity measures is introduced, quantifying the extent that the distribution of an input factor is distorted by the transition to a stressed probability. Analysis of stochastic order relations induced by the change of measure provides assurance that the proposed method has desirable properties.

The reverse sensitivity framework can be easily deployed by a risk analyst with access only to a set of input / output scenarios, simulated under the baseline model. Thus there is no need for a detailed consideration of the model structure or of simulating additional scenarios, involving computationally expensive model evaluations. Thus the proposed framework is immediately applicable to industry applications.

Appendix A. Stochastic comparisons

The proposed reverse sensitivity testing framework is based on the change from a baseline probability measure P to a stressed probability Q . The optimisation problems of Section 3 ensure that under Q the value of particular risk measures applied on Y increases. But the broader changes in the distributions of input




Fig. 8. Reverse sensitivity measure $\Gamma_{i, 0.95}$ for the London Insurance Market portfolio, against weights w_i (left) and scaled input percentiles $\frac{\text{VaR}_{0.95}(X_i)}{E(X_i)} - 1$ (right).

factors \mathbf{X} and output Y arising from the change of measure are also of interest in a risk management context. For Q to be meaningfully called a ‘stressed measure’, we argue that three properties should be fulfilled. First, under Q the distribution of the output should dominate (in a suitable stochastic order relation) the output distribution under the baseline model. Second, under the assumptions of a non-decreasing aggregation function and positive dependence between input factors, the distribution of the input vector \mathbf{X} under Q should stochastically dominate the distribution of \mathbf{X} under P . Third, an increase in the extent to which risk measures are stressed should be reflected in the distributions of output and inputs under the corresponding stressed probabilities. In this section we aim to give precise conditions under which the above properties are fulfilled. Note that most of the discussion is not contingent on Q being a solution of one of the optimisation problems of [Section 3](#).

We adopt the standard definitions of stochastic order relations. For distribution functions F, G we write $F \leq_{st} G$ if G is larger than F in first-order stochastic dominance, that is $F(\mathbf{x}) \geq G(\mathbf{x})$ for all $\mathbf{x} \in \mathbb{R}^n$. For univariate F, G , we denote $F \leq_{icx} G$ if G is larger than F in increasing convex (or stop-loss) order, that is $\int_u^1 F^{-1}(s) ds \leq \int_u^1 G^{-1}(s) ds$ for all $u \in (0, 1)$. The following dependence concepts are of importance, see [Denuit, Dhaene, Goovaerts, and Kaas \(2006\)](#):

- An m -dimensional random vector \mathbf{Z} is *stochastically increasing* (or *positively regression dependent*) in a random variable W , denoted by $\mathbf{Z} \uparrow_{si} W$, if $P(\mathbf{Z} > \mathbf{z} | W = w)$ is non-decreasing in w , for all $\mathbf{z} \in \mathbb{R}^m$.
- An m -dimensional random vector \mathbf{Z} is *associated* if $\text{Cov}(h_1(\mathbf{Z}), h_2(\mathbf{Z})) \geq 0$, for all component-wise non-decreasing functions $h_1, h_2 : \mathbb{R}^m \rightarrow \mathbb{R}$ for which the covariance exists.
- The random couple (W, Z) is *positively quadrant dependent* (PQD) if $P(W \leq w, Z \leq z) \geq P(W \leq w)P(Z \leq z)$ for all $w, z \in \mathbb{R}$.
- The random couple (W, Z) is *negatively quadrant dependent* (NQD) if $P(W \leq w, Z \leq z) \leq P(W \leq w)P(Z \leq z)$ for all $w, z \in \mathbb{R}$.

For a pair of random variables (W, Z) the above definitions are successively weaker: $\mathbf{Z} \uparrow_{si} W$ implies that (Z, W) is associated, which implies PQD, see [Esary, Proschan, and Walkup \(1967\)](#). We write $\mathbf{Z}_{-k} = (Z_1, \dots, Z_{k-1}, Z_{k+1}, \dots, Z_m)$, $1 \leq k \leq m$ for the $(m-1)$ -dimensional sub-vector of \mathbf{Z} deprived of its k th component.

The next two propositions characterise the stochastic ordering of inputs and output under two different probabilities Q^1, Q^2 , making alternative assumptions on distributions under P , on g and on the form of the corresponding two RN-densities.

Proposition A.1. *Let $Q^1, Q^2 \in \mathcal{P}$ be two probability measures with $\frac{dQ^1}{dP} = \eta_1(Y)$, $\frac{dQ^2}{dP} = \eta_2(Y)$, for some non-negative functions η_j , $j = 1, 2$. If the RN-densities cross once, such that for some $d \in \mathbb{R}$*

$$\eta_2(y) \begin{cases} \leq \eta_1(y) & y < d \\ \geq \eta_1(y) & y \geq d, \end{cases} \quad (\text{A.1})$$

then the following hold:

1. $F_Y^{Q^1} \leq_{st} F_Y^{Q^2}$
2. For given $i \in \{1, \dots, n\}$, if $E((X_i - t)_+ | Y = y)$ is non-decreasing in y for all $t \in \mathbb{R}$, then $F_{X_i}^{Q^1} \leq_{icx} F_{X_i}^{Q^2}$.
3. For given $i \in \{1, \dots, n\}$, if $X_i \uparrow_{si} Y$, then $F_{X_i}^{Q^1} \leq_{st} F_{X_i}^{Q^2}$.

Proposition A.2. *Let $Q^1, Q^2 \in \mathcal{P}$ be two probability measures with $\frac{dQ^1}{dP} = \eta_1(Y)$, $\frac{dQ^2}{dP} = \eta_2(Y)$ for some non-negative functions η_j , $j = 1, 2$. Assume that $\eta_2 - \eta_1$ is non-decreasing. Then the following hold:*

1. $F_Y^{Q^1} \leq_{st} F_Y^{Q^2}$.
2. If the aggregation function g is non-decreasing in coordinate i and X_i is independent of \mathbf{X}_{-i} , then $F_{X_i}^{Q^1} \leq_{st} F_{X_i}^{Q^2}$.
3. Assume that the aggregation function g is non-decreasing.
 - (a) For given $i \in \{1, \dots, n\}$, if (X_i, Y) is PQD, then $F_{X_i}^{Q^1} \leq_{st} F_{X_i}^{Q^2}$.
 - (b) If \mathbf{X} is associated, then $F_{\mathbf{X}}^{Q^1} \leq_{st} F_{\mathbf{X}}^{Q^2}$.

Part 1. of both [Propositions Appendix A.1](#) and [Appendix A.2](#) reflects the comparative impact of the stress on the output Y , while parts 2. and 3. characterise the impact of the stress on the inputs. An example where the assumption of [Proposition Appendix A.1](#), part 3., is satisfied is the following. Suppose the input vector \mathbf{X} is multivariate normal and $Y = h(\sum_{i=1}^n w_i X_i)$ for an increasing function h and $w_i \in \mathbb{R}$ for all i . If $\text{Cov}(X_i, h^{-1}(Y)) = \sum_{j=1}^n w_j \text{Cov}(X_i, X_j) \geq 0$, then $X_i \uparrow_{si} Y$ holds. The assumption in [Proposition Appendix A.2](#) part 3.(a) holds for example if $\mathbf{X}_{-i} \uparrow_{st} X_i$ and g is non-decreasing.




Fig. A.1. Left: simulated RN-densities of the solution to (4) with a 10% (black) and 15% (grey) increase in VaR. Right: simulated RN-densities of the solution to (5) case 1, with a 10% increase in VaR and 9% (black) and 13% (grey) increase in ES.

Propositions Appendix A.1 and Appendix A.2 allow for a stochastic comparison of the output and the input factors under the stressed and the baseline model. In particular, Proposition Appendix A.1 applies to the solutions of problems (3), (5) and (7) with $Q^2 = Q$ and $Q^1 = P$. Proposition Appendix A.2 applies to optimisation problem (1), with $Q^2 = Q$ and $Q^1 = P$, if the RN-density of the solution is a non-decreasing function of Y . Recall that the RN-density of the solutions to (3), (5) case 1, and (7) are non-decreasing. Moreover, for a stressed model under which the input X_i stochastically dominates, in first-order or increasing convex order, the distribution of the input under the baseline model, the introduced sensitivity measure Γ_i is positive.

Proposition Appendix A.1 also enables to contrast stressed probabilities corresponding to different stress levels. For example, when solving optimisation problem (3) with two different VaR constraints, the output under the stressed model corresponding to a higher VaR should stochastically dominate the output under the other stressed model. The next lemma associates Proposition Appendix A.1 with solutions of the optimisation problems (3) and (5).

Lemma A.3. *The crossing condition of Proposition Appendix A.1 is satisfied for:*

1. Two solutions Q^1, Q^2 of optimisation problem (3) with constraints $\text{VaR}_\alpha^{Q^1}(Y) = q_1$ respectively $\text{VaR}_\alpha^{Q^2}(Y) = q_2$, and $q_1 < q_2$.
2. Two solutions Q^1, Q^2 of optimisation problem (5) with constraints $\text{VaR}_\alpha^{Q^1}(Y) = \text{VaR}_\alpha^{Q^2}(Y) = q$ and $\text{ES}_\alpha^{Q^1}(Y) = s_1$, respectively $\text{ES}_\alpha^{Q^2}(Y) = s_2$, and $s_1 < s_2$.

The second part of Lemma Appendix A.3 holds true for both types of solutions of (5).

Example (continued). Applying Proposition Appendix A.1 to the two optimisation problems in this example, we immediately verify that the output under the stressed probabilities first-order stochastically dominates the output under the baseline probability, see Figs. 1 and 3. Moreover, the aggregation function g is non-decreasing and it can be verified that, for instance, (X_4, Y) is PQD. Hence, following Proposition Appendix A.2 part 3.(a), the distribution of X_4 under the stressed probability first-order stochastically dominates that under the baseline probability. This can be seen in Figs. 2 and 4.

An illustration of Lemma Appendix A.3 is given in Fig. A.1. The left plot shows the RN-densities of solutions to (3) with two different stress levels. The black line corresponds to an increase of VaR of 10%, the same as in Fig. 1, and the grey line to an increase of VaR of 15%. The plot to the right displays the RN-densities of solutions to (5) for an increase of 10% in VaR and 9% in ES (black) and an increase of 10% in VaR and 13% in ES (grey), see Fig. 3. It is seen how in both cases, the two RN-densities satisfy the crossing condition of Proposition Appendix A.1.

Appendix B. Proofs

Proposition 3.1. A similar result to Proposition 3.1 can be found in Cambou and Filipović (2017), and we also refer to Csiszár (1975) for a general form of the solution. It is immediately verified that ζ is a RN-density for which $Q^\zeta(Y \in B_i) = \alpha_i$, $i = 1, \dots, I$. Let ξ be any RN-density that satisfies $Q^\xi(Y \in B_i) = \alpha_i$, $i = 1, \dots, I$. Using Jensen inequality, the KL-divergence of Q^ξ with respect to P fulfils

$$\begin{aligned} D_{\text{KL}}(Q^\xi \| P) &= \sum_{i=0}^I E(\xi \log(\xi) | Y \in B_i) P(Y \in B_i) \\ &\geq \sum_{i=0}^I E(\xi | Y \in B_i) \log(E(\xi | Y \in B_i)) P(Y \in B_i) \\ &= \sum_{i=0}^I \alpha_i \log\left(\frac{\alpha_i}{P(Y \in B_i)}\right) \\ &= D_{\text{KL}}(Q^\zeta \| P). \end{aligned}$$

Therefore Q^ζ is a solution of (2). Uniqueness follows by strict convexity of the KL-divergence, see Csiszár (1975). \square

Proposition 3.2. Assume that $P(q - \varepsilon < Y < q) > 0$ for all $\varepsilon > 0$. Then, it is immediate to verify that ζ is a RN-density such that $\text{VaR}_\alpha^{Q^\zeta}(Y) = q$. Let $\xi = \frac{dQ^\xi}{dP}$ be a RN-density for which $\text{VaR}_\alpha^{Q^\xi}(Y) = q$. By Jensen inequality, the KL-divergence of Q^ξ with respect to P is

$$\begin{aligned} D_{\text{KL}}(Q^\xi \| P) &= E(\xi \log(\xi) | Y < q) P(Y < q) + E(\xi \log(\xi) | Y \geq q) P(Y \geq q) \end{aligned}$$

$$\begin{aligned} &\geq Q^\xi(Y < q) \log\left(\frac{Q^\xi(Y < q)}{P(Y < q)}\right) + Q^\xi(Y \geq q) \log\left(\frac{Q^\xi(Y \geq q)}{P(Y \geq q)}\right) \\ &= k(Q^\xi(Y < q), P(Y < q)), \end{aligned}$$

where we define $k(x, y) = x \log(\frac{x}{y}) + (1-x) \log(\frac{1-x}{1-y})$, for $0 < x < 1, 0 < y < 1$. Inspection shows that, for fixed $0 < y < 1$, $x \rightarrow k(x, y)$ is non-increasing on $(0, y]$. Moreover it holds

$$Q^\xi(Y < q) \leq \alpha \leq P(Y \leq \text{VaR}_\alpha(Y)) \leq P(Y < q).$$

The KL-divergence of Q^ξ is thus larger than the KL-divergence of Q^ζ ,

$$\begin{aligned} D_{\text{KL}}(Q^\xi \| P) &\geq k(Q^\xi(Y < q), P(Y < q)) \\ &\geq k(\alpha, P(Y < q)) \\ &= \alpha \log\left(\frac{\alpha}{P(Y < q)}\right) + (1-\alpha) \log\left(\frac{1-\alpha}{P(Y \geq q)}\right) \\ &= D_{\text{KL}}(Q^\zeta \| P), \end{aligned}$$

and Q^ζ is a solution of (3). Uniqueness follows by strict convexity of the KL-divergence.

Assume now that there exists $\varepsilon > 0$ such that $P(q - \varepsilon < Y < q) = 0$. If $P(Y = q) = 0$, by the absolute continuity of the probability measures, the optimisation problem (3) does not admit a solution. Hence, we assume that $P(Y = q) > 0$. Let Q^ξ be a RN-density for which $\text{VaR}_{\alpha}^{Q^\xi}(Y) = q$. Denote $r = Q^\xi(Y \leq q)$ and $p = P(Y \leq q)$. The KL-divergence of Q^ξ with respect to P is

$$\begin{aligned} D_{\text{KL}}(Q^\xi \| P) &= E(\xi \log(\xi) | Y \leq q)p + E(\xi \log(\xi) | Y > q)(1-p) \\ &\geq r \log\left(\frac{r}{p}\right) + (1-r) \log\left(\frac{1-r}{1-p}\right) \\ &= D_{\text{KL}}(Q^{\xi^r} \| P), \end{aligned}$$

where we define $\xi^u = \frac{dQ^{\xi^u}}{dP} = \frac{u}{p} \mathbb{1}_{\{Y \leq q\}} + \frac{1-u}{1-p} \mathbb{1}_{\{Y > q\}}$, $0 \leq u \leq 1$. The family of RN-densities ξ^u fulfil $\text{VaR}_{\alpha}^{Q^{\xi^u}}(Y) = q$ if and only if $\alpha \leq u < \alpha \frac{p}{P(Y < q)}$. In particular this holds for the RN-density ξ^r . Hence the optimisation problem (3) is reduced to minimise $D_{\text{KL}}(Q^{\xi^u} \| P)$ subject to $\alpha \leq u < \alpha \frac{p}{P(Y < q)}$. As a function of u the KL-divergence $D_{\text{KL}}(Q^{\xi^u} \| P)$ is non-increasing on $(0, p]$, hence the optimisation problem does not admit a solution as $\alpha \frac{p}{P(Y < q)} < p$. \square

Proposition 3.3. For $i = 1, 2$, Eq. (6) can be rewritten as

$$\frac{\partial}{\partial \theta} E(e^{\theta(Y-q)} | A_i) = \frac{E((Y-q)e^{\theta(Y-q)} | A_i)}{E(e^{\theta(Y-q)} | A_i)} = s - q.$$

The left hand side is increasing for positive θ , negative for $\theta = 0$ and diverges for $\theta \uparrow \theta_{\max}$, where $\theta_{\max} = \sup\{\theta > 0 | E(e^{\theta Y} | A_i) < \infty\}$, by properties of the moment generating function. Thus, for $i = 1, 2$, there exists a unique positive solution θ_i^* of (6).

Case 1. The RN-density ζ_1 fulfils the constraints in (5) since $Q^{\zeta_1}(Y < q) = \alpha$, $Q^{\zeta_1}(Y \leq q) \geq \alpha$ and the ES constraint is equivalent to $(1-\alpha)(s-q) = E^{Q^{\zeta_1}}((Y-q)_+)$. Let $\xi = \frac{dQ^\xi}{dP}$ be a RN-density satisfying the constraints of problem (5) and denote $r = Q^\xi(A_1^c)$ and $p = P(A_1^c)$. Using Jensen's inequality, the KL-divergence of Q^ξ with respect to P fulfils

$$\begin{aligned} D_{\text{KL}}(Q^\xi \| P) &= E(\xi \log(\xi) \mathbb{1}_{A_1^c}) + E(\xi \log(\xi) \mathbb{1}_{A_1}) \\ &\quad + \theta_1^*(1-\alpha)(s-q) - E(\xi \log(e^{\theta_1^*(Y-q)}) \mathbb{1}_{A_1}) \\ &\geq r \log\left(\frac{r}{p}\right) + \theta_1^*(1-\alpha)(s-q) \\ &\quad + E\left(\xi \log\left(\frac{\xi}{e^{\theta_1^*(Y-q)}}\right) \middle| A_1\right) (1-p). \end{aligned}$$

Recall that the perspective of a convex function f , defined by $h(x, y) = yf(x/y)$ is itself convex, see [Boyd and Vandenberghe \(2004\)](#). Applying then Jensen's inequality to $h(x, y) = y \log(\frac{x}{y})$, the third term becomes

$$\begin{aligned} &E\left(\xi \log\left(\frac{\xi}{e^{\theta_1^*(Y-q)}}\right) \middle| A_1\right) (1-p) \\ &\geq E(\xi | A_1) \log\left(\frac{E(\xi | A_1)}{E(e^{\theta_1^*(Y-q)} | A_1)}\right) (1-p) \\ &= (1-r) \log\left(\frac{(1-r)}{E(e^{\theta_1^*(Y-q)} \mathbb{1}_{A_1})}\right). \end{aligned}$$

Collecting all terms,

$$\begin{aligned} D_{\text{KL}}(Q^\xi \| P) &\geq r \log\left(\frac{r}{p}\right) + \theta_1^*(1-\alpha)(s-q) \\ &\quad + (1-r) \log\left(\frac{(1-r)}{E(e^{\theta_1^*(Y-q)} \mathbb{1}_{A_1})}\right) \\ &= k(r, p, E(e^{\theta_1^*(Y-q)} \mathbb{1}_{A_1})), \end{aligned}$$

where we define $k(x, y, z) = x \log(\frac{x}{y}) + \theta_1^*(1-\alpha)(s-q) + (1-x) \log(\frac{1-x}{z})$, for $0 < x < 1$ and $y, z > 0$. For fixed $y, z > 0$, the function $x \rightarrow k(x, y, z)$ is decreasing on $(0, \frac{y}{y+z}]$. The condition on θ_1^* in 1. is equivalent to

$$\alpha \leq \frac{p}{p + E(e^{\theta_1^*(Y-q)} \mathbb{1}_{A_1})}.$$

Therefore, noting that $r \leq \alpha$, we obtain

$$\begin{aligned} D_{\text{KL}}(Q^\xi \| P) &\geq k(r, p, E(e^{\theta_1^*(Y-q)} \mathbb{1}_{A_1})) \geq k(\alpha, p, E(e^{\theta_1^*(Y-q)} \mathbb{1}_{A_1})) \\ &= D_{\text{KL}}(Q^{\zeta_1} \| P). \end{aligned}$$

The last equality follows since

$$\begin{aligned} D_{\text{KL}}(Q^{\zeta_1} \| P) &= \alpha \log\left(\frac{\alpha}{p}\right) + \frac{1-\alpha}{E(e^{\theta_1^*(Y-q)} \mathbb{1}_{A_1})} \\ &\quad \times E\left(e^{\theta_1^*(Y-q)} \mathbb{1}_{A_1} \log\left(\frac{1-\alpha}{E(e^{\theta_1^*(Y-q)} \mathbb{1}_{A_1})} e^{\theta_1^*(Y-q)}\right)\right) \\ &= \alpha \log\left(\frac{\alpha}{p}\right) + (1-\alpha) \log\left(\frac{1-\alpha}{E(e^{\theta_1^*(Y-q)} \mathbb{1}_{A_1})}\right) \\ &\quad + \theta_1^* \frac{1-\alpha}{E(e^{\theta_1^*(Y-q)} \mathbb{1}_{A_1})} E(e^{\theta_1^*(Y-q)} (Y-q)_+) \\ &= \alpha \log\left(\frac{\alpha}{p}\right) + (1-\alpha) \log\left(\frac{1-\alpha}{E(e^{\theta_1^*(Y-q)} \mathbb{1}_{A_1})}\right) \\ &\quad + \theta_1^*(1-\alpha)(s-q) \\ &= k(\alpha, p, E(e^{\theta_1^*(Y-q)} \mathbb{1}_{A_1})). \end{aligned}$$

Therefore Q^{ζ_1} is a solution of (5). Uniqueness follows by strict convexity of the KL-divergence.

Case 2. The proof of case 2 is similar to that of case 1, replacing the set A_1 with A_2 and ζ_1 with ζ_2 . The RN-density ζ_2 fulfils the constraints (5). Letting $\xi = \frac{dQ^\xi}{dP}$ be a RN-density satisfying the constraints of problem (5), then the KL-divergence of Q^ξ with respect to P can be bounded by

$$D_{\text{KL}}(Q^\xi \| P) \geq k(Q^\xi(A_2^c), P(A_2^c), E(e^{\theta_2^*(Y-q)} \mathbb{1}_{A_2})),$$

where the function $k(x, y, z)$ has been defined above. For fixed $y, z > 0$, the function $x \rightarrow k(x, y, z)$ is increasing on $[\frac{y}{y+z}, 1)$. Moreover,

the condition on θ_2^* in 2. is equivalent to

$$\frac{P(A_2^c)}{P(A_2^c) + E(e^{\theta_2^*(Y-q)} \mathbb{1}_{A_2})} \leq \alpha.$$

Since $\alpha \leq Q^\xi(A_2^c)$ we obtain

$$D_{\text{KL}}(Q^\xi \| P) \geq k(\alpha, P(A_2^c), E(e^{\theta_2^*(Y-q)} \mathbb{1}_{A_2})) = D_{\text{KL}}(Q^{\xi_2} \| P),$$

which is the KL-divergence of Q^{ξ_2} . \square

Proposition A1. Let $y \leq d$, then $Q^2(Y \leq y) = E(\eta_2(Y) \mathbb{1}_{\{Y \leq y\}}) \leq E(\eta_1(Y) \mathbb{1}_{\{Y \leq y\}}) = Q^1(Y \leq y)$. For $y > d$, it holds $Q^2(Y \leq y) = 1 - Q^2(Y > y) = 1 - E(\eta_2(Y) \mathbb{1}_{\{Y > y\}}) \leq 1 - E(\eta_1(Y) \mathbb{1}_{\{Y > y\}}) = Q^1(Y \leq y)$. For the second part we have, for all $t \in \mathbb{R}$, using the tower property under P ,

$$\begin{aligned} E^{Q^2}((X_i - t)_+) &= E^{Q^2}(E((X_i - t)_+ | Y)) \geq E^{Q^1}(E((X_i - t)_+ | Y)) \\ &= E^{Q^1}((X_i - t)_+), \end{aligned}$$

by first-order stochastic dominance of Y with respect to the measures Q^1, Q^2 . The last claim follows using a similar argument. \square

Proposition A2. The RN-densities have to cross once due to normalisation, therefore part 1. applies. In the rest of the proof, let $h = \eta_2 - \eta_1$.

To prove part 2., let g be non-decreasing in coordinate i and X_i independent of \mathbf{X}_{-i} . For any $t \in \mathbb{R}$, using the Fortuin–Kasteleyn–Ginibre inequality (Wüthrich & Merz, 2013), we have

$$\begin{aligned} Q^2(X_i > t) - Q^1(X_i > t) &= E(h(Y) \mathbb{1}_{\{X_i > t\}}) = E(E(h(Y) \mathbb{1}_{\{X_i > t\}} | \mathbf{X}_{-i})) \\ &\geq E(E(h(Y) | \mathbf{X}_{-i}))P(X_i > t) = 0, \end{aligned}$$

proving first-order stochastic dominance.

To show part 3.(a), assume that g is non-decreasing and (X_i, Y) are PQD. Hence, for all $t \in \mathbb{R}$,

$$Q^2(X_i > t) - Q^1(X_i > t) = E(\mathbb{1}_{\{X_i > t\}} h(Y)) \geq 0,$$

where the last inequality follows from Lemma 3 in Lehmann (1966). Part 3.(b) follows by association of the vector $(h(Y), \mathbf{X})$, using a similar argument. \square

Lemma A3. The first claim follows since $\alpha \leq P(Y < q_1) \leq P(Y < q_2)$. For part 2., consider first the case where $P(q - \varepsilon < Y < q) > 0$ for all $\varepsilon > 0$. Denote by θ_1^*, θ_2^* the solutions to (6) with q and s_1 , respectively s_2 . Hence, $\theta_1^* \leq \theta_2^*$, and there exists a $d > q$ such that for all $\omega \in \Omega$ with $Y(\omega) > d$ we have

$$e^{(\theta_2^* - \theta_1^*)(Y(\omega) - q)} \geq \frac{E(e^{\theta_2^*(Y-q)} \mathbb{1}_{A_1})}{E(e^{\theta_1^*(Y-q)} \mathbb{1}_{A_1})},$$

which implies $\eta_2 \geq \eta_1$ for all ω with $Y(\omega) > d$. Since on A_1^c , $\eta_1 = \eta_2$ P -a.s. the RN-densities admit a (unique) crossing point. The argument also holds if A_1 is replaced with A_2 . \square

Proposition 4.3. We also refer to Theorem 6 in Kachapova and Kachapov (2012). The first two properties are immediate. For 3. if X_i and Y are comonotonic, ζ and $\zeta|_{X_i}$ are also comonotonic since ζ is a non-decreasing function of Y and $\zeta|_{X_i}$ a non-decreasing function of X_i . Part 4. follows by a similar argument. Properties 5. and 6. are consequences of the invariance of PQD (NQD) under non-decreasing (non-increasing) transformations, see Lemma 1 in Lehmann (1966). \square

References

Acerbi, C. (2002). Spectral measures of risk: A coherent representation of subjective risk aversion. *Journal of Banking and Finance*, 26(7), 1505–1518.
 Ahmed, S., Çakmak, U., & Shapiro, A. (2007). Coherent risk measures in inventory problems. *European Journal of Operational Research*, 182(1), 226–238.

Ali, S. M., & Silvey, S. D. (1966). A general class of coefficients of divergence of one distribution from another. *Journal of the Royal Statistical Society. Series B Methodological*, 131–142.
 Artzner, P., Delbaen, F., Eber, J.-M., & Heath, D. (1999). Coherent measures of risk. *Mathematical Finance*, 9(3), 203–228. doi:10.1111/1467-9965.00068.
 Asimit, A. V., Bignozzi, V., Cheung, K. C., Hu, J., & Kim, E.-S. (2017). Robust and pareto optimality of insurance contracts. *European Journal of Operational Research*, 262(2), 720–732.
 Aven, T., & Nøklund, T. (2010). On the use of uncertainty importance measures in reliability and risk analysis. *Reliability Engineering & System Safety*, 95(2), 127–133.
 BCBS (2012). Consultative Document May 2012. Fundamental review of the trading book. *Technical Report*. Basel Committee on Banking Supervision, Bank for International Settlements.
 BCBS (2013). Consultative Document October 2013. Fundamental review of the trading book: A revised market risk framework. *Technical Report*. Basel Committee on Banking Supervision, Bank for International Settlements.
 Beckman, R. J., & McKay, M. D. (1987). Monte Carlo estimation under different distributions using the same simulation. *Technometrics*, 29(2), 153–160.
 Bellini, F., Klar, B., Müller, A., & Rosazza Gianin, E. (2014). Generalized quantiles as risk measures. *Insurance: Mathematics and Economics*, 54, 41–48. doi:10.1016/j.insmatheco.2013.10.015.
 Ben-Tal, A., Den Hertog, D., De Waegenaere, A., Melenberg, B., & Rennen, G. (2013). Robust solutions of optimization problems affected by uncertain probabilities. *Management Science*, 59(2), 341–357.
 Blanchet, J., Lam, H., Tang, Q., & Yuan, Z. (2017). Applied robust performance analysis for actuarial applications. *Technical Report*. The Society of Actuaries (SOA).
 Borgonovo, E. (2007). A new uncertainty importance measure. *Reliability Engineering & System Safety*, 92(6), 771–784.
 Borgonovo, E., Castaings, W., & Tarantola, S. (2011). Moment independent importance measures: New results and analytical test cases. *Risk Analysis*, 31(3), 404–428.
 Borgonovo, E., & Cillo, A. (2017). Deciding with thresholds: Importance measures and value of information. *Risk Analysis*, 37(10), 1828–1848.
 Borgonovo, E., Hazen, G. B., & Plischke, E. (2016). A common rationale for global sensitivity measures and their estimation. *Risk Analysis*, 36(10), 1871–1895.
 Borgonovo, E., & Plischke, E. (2016). Sensitivity analysis: A review of recent advances. *European Journal of Operational Research*, 248(3), 869–887.
 Boyd, S., & Vandenberghe, L. (2004). *Convex optimization*. Cambridge University Press.
 Breuer, T., & Csiszár, I. (2013). Systematic stress tests with entropic plausibility constraints. *Journal of Banking & Finance*, 37(5), 1552–1559.
 Breuer, T., Jandačka, M., Mencía, J., & Summer, M. (2012). A systematic approach to multi-period stress testing of portfolio credit risk. *Journal of Banking & Finance*, 36(2), 332–340.
 Cambou, M., & Filipović, D. (2017). Model uncertainty and scenario aggregation. *Mathematical Finance*, 27(2), 534–567.
 Cooke, R. M., & van Noortwijk, J. M. (1999). Local probabilistic sensitivity measures for comparing FORM and Monte Carlo calculations illustrated with dike ring reliability calculations. *Computer Physics Communications*, 117(1–2), 86–98.
 Cover, T. M., & Thomas, J. A. (2012). *Elements of information theory*. John Wiley & Sons.
 Csiszár, I. (1975). I-divergence geometry of probability distributions and minimization problems. *The Annals of Probability*, 3(1), 146–158.
 Denuit, M., Dhaene, J., Goovaerts, M., & Kaas, R. (2006). *Actuarial theory for dependent risks: Measures, orders and models*. John Wiley & Sons.
 EIOPA (2009). Directive 2009/138/EC of the European parliament and of the council. *Technical Report*. European Insurance and Occupational Pensions Authority.
 Esary, J. D., Prochan, F., & Walkup, D. W. (1967). Association of random variables, with applications. *The Annals of Mathematical Statistics*, 38(5), 1466–1474.
 Föllmer, H., & Schied, A. (2002). Convex measures of risk and trading constraints. *Finance and Stochastics*, 6(4), 429–447. doi:10.1007/s007800200072.
 Föllmer, H., & Schied, A. (2011). *Stochastic finance: An introduction in discrete time* (3rd). de Gruyter.
 Gerber, H. U. (1974). On additive premium calculation principles. *ASTIN Bulletin: The Journal of the IAA*, 7(3), 215–222.
 Gibbs, A. L., & Su, F. E. (2002). On choosing and bounding probability metrics. *International Statistical Review*, 70(3), 419–435.
 Glasserman, P., & Liu, Z. (2010). Sensitivity estimates from characteristic functions. *Operations Research*, 58(6), 1611–1623.
 Glasserman, P., & Xu, X. (2014). Robust risk measurement and model risk. *Quantitative Finance*, 14(1), 29–58.
 Gotoh, J.-Y., & Takano, Y. (2007). Newsvendor solutions via conditional value-at-risk minimization. *European Journal of Operational Research*, 179(1), 80–96.
 Gouriéroux, C., Laurent, J. P., & Scaillet, O. (2000). Sensitivity analysis of values at risk. *Journal of Empirical Finance*, 7(3), 225–245.
 Hong, L. J. (2009). Estimating quantile sensitivities. *Operations Research*, 57(1), 118–130.
 Kachapova, F., & Kachapov, I. (2012). A measure of monotonicity of two random variables. *Journal of Mathematics and Statistics*, 8(2), 221–228.
 Kullback, S. (1959). *Information theory and statistics*. John Wiley & Sons.
 Kullback, S., & Leibler, R. A. (1951). On information and sufficiency. *The Annals of Mathematical Statistics*, 22(1), 79–86.
 Lehmann, E. L. (1966). Some concepts of dependence. *The Annals of Mathematical Statistics*, 37(5), 1137–1153.
 Lloyd's (2016). Internal model validation: Supplementary notes to the Solvency II model validation guidance (April 2014). *Technical Report*. Lloyd's.

- Mara, T. A., & Tarantola, S. (2012). Variance-based sensitivity indices for models with dependent inputs. *Reliability Engineering & System Safety*, 107, 115–121.
- McNeil, A. J., & Smith, A. D. (2012). Multivariate stress scenarios and solvency. *Insurance: Mathematics and Economics*, 50(3), 299–308.
- Müller, A., & Stoyan, D. (2002). *Comparison methods for stochastic models and risks*. John Wiley & Sons.
- Newey, W. K., & Powell, J. L. (1987). Asymmetric least squares estimation and testing. *Econometrica: Journal of the Econometric Society*, 819–847.
- Oakley, J. E., & O'Hagan, A. (2004). Probabilistic sensitivity analysis of complex models: A Bayesian approach. *Journal of the Royal Statistical Society: Series B Statistical Methodology*, 66(3), 751–769.
- Osidele, O. O., & Beck, M. B. (2004). An inverse approach to the analysis of uncertainty in models of environmental systems. *Integrated Assessment*, 4(4), 265–282.
- Plischke, E., Borgonovo, E., & Smith, C. L. (2013). Global sensitivity measures from given data. *European Journal of Operational Research*, 226(3), 536–550.
- Rockafellar, R. T., & Uryasev, S. (2002). Conditional value-at-risk for general loss distributions. *Journal of Banking & Finance*, 26(7), 1443–1471.
- Rüschendorf, L. (1983). Solution of a statistical optimization problem by rearrangement methods. *Metrika*, 30(1), 55–61.
- Saltelli, A. (2002). Sensitivity analysis for importance assessment. *Risk Analysis*, 22(3), 579–590.
- Saltelli, A., Chan, K., & Scott, E. M. (2000). *Sensitivity analysis*. John Wiley & Sons.
- Saltelli, A., Ratto, M., Andres, T., Campolongo, F., Cariboni, J., Gatelli, D., et al. (2008). *Global sensitivity analysis: The primer*. John Wiley & Sons.
- Saltelli, A., & Tarantola, S. (2002). On the relative importance of input factors in mathematical models. *Journal of the American Statistical Association*, 97(459), 702–709. doi:10.1198/016214502388618447.
- Sobol, I. M. (1993). Sensitivity estimates for nonlinear mathematical models. *Mathematical Modelling and Computational Experiments*, 1(4), 407–414.
- Spear, R. C., Grieb, T. M., & Shang, N. (1994). Parameter uncertainty and interaction in complex environmental models. *Water Resources Research*, 30(11), 3159–3169.
- Szegö, G. (2005). Measures of risk. *European Journal of Operational Research*, 163(1), 5–19.
- Tapiero, C. S. (2005). Value at risk and inventory control. *European Journal of Operational Research*, 163(3), 769–775.
- Tsanakas, A., & Millosovich, P. (2016). Sensitivity analysis using risk measures. *Risk Analysis*, 36(1), 30–48.
- Wagner, H. M. (1995). Global sensitivity analysis. *Operations Research*, 43(6), 948–969.
- Weber, S. (2007). Distribution-invariant risk measures, entropy, and large deviations. *Journal of Applied Probability*, 44(1), 16–40.
- Wüthrich, M. V., & Merz, M. (2013). *Financial modeling, actuarial valuation and solvency in insurance*. Springer.
- Xu, C., & Gertner, G. Z. (2008a). A general first-order global sensitivity analysis method. *Reliability Engineering & System Safety*, 93(7), 1060–1071.
- Xu, C., & Gertner, G. Z. (2008b). Uncertainty and sensitivity analysis for models with correlated parameters. *Reliability Engineering & System Safety*, 93(10), 1563–1573.