




CIF Angina







Supplementary Table 1: Cumulative incidence rates of Angina/MI, taking into account death as a competing risk

Follow-up	12 months	24 months	48 months	96 months	120 months
Groups of patients	Angina/MI (%)				
DM & HypovitD	11.7	15.6	20.1	21.9	24.6
nonDM & HypovitD	8.7	13.8	16	19.9	22.7
DM & non HypovitD	10.3	11.6	13	18.2	23.6
Non DM & non HypovitD	6.7	11.1	14.6	16.7	16.7
	Death before Angina/MI event (%)				
DM & HypovitD	13.2	15.6	19.2	27.5	34.8
nonDM & HypovitD	6	8.9	13.9	21.2	24.8
DM & non HypovitD	5.1	7.5	9.7	24.8	30.4
Non DM & non HypovitD	5	6.4	7.6	17.4	21.6

Supplementary Table 2: Cumulative incidence rates of HF, taking into account death as a competing risk

Follow-up	12 months	24 months	48 months	96 months	120 months
Groups of patients	HF (%)				
DM & HypovitD	8.1	10.9	16.3	22	22.8
nonDM & HypovitD	6	7.8	9.8	12	12.6
DM & non HypovitD	6	12.1	13.6	16.2	16.2
Non DM & non HypovitD	4.4	5.3	7.1	8.9	9.9
	Death before HF event (%)				
DM & HypovitD	13,3	15,2	20	26,5	33,2
nonDM & HypovitD	5,2	9,1	13,2	22,7	25,6
DM & non HypovitD	6,2	7,4	7,4	19,7	28,8
Non DM & non HypovitD	5,4	6,4	7	15,5	24,3