

PAPER • OPEN ACCESS

## Stability for the Calderón's problem for a class of anisotropic conductivities via an ad hoc misfit functional

To cite this article: Sonia Foschiatti *et al* 2021 *Inverse Problems* **37** 125007

View the [article online](#) for updates and enhancements.

You may also like

- [Electrical impedance tomography and Calderón's problem](#)  
G Uhlmann
- [Lipschitz stability for the inverse conductivity problem for a conformal class of anisotropic conductivities](#)  
Romina Gaburro and Eva Sincich
- [Imaging of isotropic and anisotropic conductivities from power densities in three dimensions](#)  
François Monard and Donsub Rim



**IOP | ebooks™**

Bringing together innovative digital publishing with leading authors from the global scientific community.

Start exploring the collection—download the first chapter of every title for free.

# Stability for the Calderón's problem for a class of anisotropic conductivities via an ad hoc misfit functional

Sonia Foschiatti<sup>1</sup> , Romina Gaburro<sup>2,\*</sup>  and Eva Sincich<sup>1</sup> 

<sup>1</sup> Dipartimento di Matematica e Geoscienze, Università degli Studi di Trieste, Italy

<sup>2</sup> Department of Mathematics and Statistics, University of Limerick, Castletroy, Limerick, Ireland

E-mail: [sonia.foschiatti@phd.units.it](mailto:sonia.foschiatti@phd.units.it), [romina.gaburro@ul.ie](mailto:romina.gaburro@ul.ie) and [esincich@units.it](mailto:esincich@units.it)

Received 10 July 2021, revised 3 October 2021

Accepted for publication 29 October 2021

Published 17 November 2021



CrossMark

## Abstract

We address the stability issue in Calderón's problem for a special class of anisotropic conductivities of the form  $\sigma = \gamma A$  in a Lipschitz domain  $\Omega \subset \mathbb{R}^n$ ,  $n \geq 3$ , where  $A$  is a known Lipschitz continuous matrix-valued function and  $\gamma$  is the unknown piecewise affine scalar function on a given partition of  $\Omega$ . We define an ad hoc misfit functional encoding our data and establish stability estimates for this class of anisotropic conductivity in terms of both the misfit functional and the more commonly used local Dirichlet-to-Neumann map.

Keywords: Calderón's problem, anisotropic conductivity, stability, misfit functional

## 1. Introduction

The paper addresses the so-called Calderón's inverse conductivity problem of recovering the conductivity  $\sigma$  of a body  $\Omega \subset \mathbb{R}^n$  by taking measurements of voltage and electric current on its surface  $\partial\Omega$ . More specifically, the case when the conductivity is anisotropic and it is *a priori* known to be of type  $\sigma = \gamma A$ , where  $A$  is a known Lipschitz continuous matrix valued function on  $\Omega$  and  $\gamma$  is a piecewise-affine unknown function on a given partition of  $\Omega$ , is considered. It is well known that in absence of internal sources or sinks, the electrostatic potential  $u$  in a conducting body, described by a domain  $\Omega \subset \mathbb{R}^n$ , is governed by the elliptic equation

\*Author to whom any correspondence should be addressed.



Original content from this work may be used under the terms of the [Creative Commons Attribution 4.0 licence](https://creativecommons.org/licenses/by/4.0/). Any further distribution of this work must maintain attribution to the author(s) and the title of the work, journal citation and DOI.

$$\operatorname{div}(\sigma \nabla u) = 0 \text{ in } \Omega, \quad (1.1)$$

where the symmetric, positive definite matrix  $\sigma(x) = (\sigma_{ij}(x))_{i,j=1}^n$ ,  $x \in \Omega$  represents the (possibly anisotropic) electric conductivity. The inverse conductivity problem consists of finding  $\sigma$  when the so called Dirichlet-to-Neumann (D–N) map

$$\Lambda_\sigma : H^{\frac{1}{2}}(\partial\Omega) \ni u|_{\partial\Omega} \rightarrow \sigma \nabla u \cdot \nu|_{\partial\Omega} \in H^{-\frac{1}{2}}(\partial\Omega)$$

is given for any  $u \in H^1(\Omega)$  solution to (1.1). Here,  $\nu$  denotes the unit outer normal to  $\partial\Omega$ . If measurements can be taken only on one portion  $\Sigma$  of  $\partial\Omega$ , then the relevant map is called the local D–N map ( $\Lambda_\sigma^\Sigma$ ).

This problem arises in many different fields such as geophysics, known as DC method, medicine, known as electrical impedance tomography (EIT) and non-destructive testing of materials. The first mathematical formulation of the inverse conductivity problem is due to Calderón [23], where he addressed the problem of whether it is possible to determine the (isotropic) conductivity  $\sigma = \gamma I$  by the D–N map. This seminal paper opened the way to the solution to the uniqueness issue where one is asking whether  $\sigma$  can be determined by the knowledge of  $\Lambda_\sigma$  or its local version when measurements are available on a portion of  $\partial\Omega$  only.

The case when measurements can be taken over the full boundary has been studied extensively in the past and the fundamental papers [2, 43, 44, 54, 61] had led the way of solving the problem of uniqueness in the isotropic case. We also recall the uniqueness results of Druskin who, independently from Calderón, dealt directly with the geophysical setting of the problem in [28–30]. His uniqueness result obtained in [29] was for conductivities described by piecewise constant functions (see also [11]). The problem of recovering the conductivity  $\sigma$  by local measurements has been treated more recently (see [45, 46]). We also refer to the fundamental papers [12, 25] and the result in [50] for the two dimensional case. In the present paper, we consider the issue of stability in the inverse conductivity problem, therefore we refer to [22, 24, 63] for an overview regarding the issues of uniqueness and reconstruction of the conductivity.

Regarding the stability issue, Alessandrini proved in [1] that, in the isotropic case and dimension  $n \geq 3$ , assuming *a priori* bounds on  $\sigma$  of the form  $\|\sigma\|_{H^s(\Omega)} \leq E$ ,  $s > \frac{n}{2} + 2$ , leads to a continuous dependence of  $\sigma$  in  $\Omega$  upon  $\Lambda_\sigma$  of logarithmic type. We also refer to [14, 15, 50] for subsequent results in this direction and to [3] for an overview of the stability issue. Even though stability at the boundary  $\partial\Omega$  is of Lipschitz type (see [5, 6]), Mandache [53] showed that in the interior of  $\Omega$ , the inconvenient logarithmic type of stability is the best possible, in any dimension  $n \geq 2$ , under *a priori* smoothness assumptions on  $\sigma$ . It seems therefore reasonable to think that, in order to restore stability in a really (Lipschitz) stable fashion, one needs to replace in some way the *a priori* assumptions expressed in terms of regularity bounds with *a priori* pieces of information of a different type that suit the underlying physical problem. Alessandrini and Vessella showed in [11] that when  $\sigma$  is isotropic and piecewise constant on a given partition of  $\Omega$ , then Lipschitz stability can be restored in terms of the local D–N map (conditional stability). Rondi [56] proved that the Lipschitz constant has an exponential behaviour with respect to the number of subdomain of the partition. From a medical imaging point of view, the partition of  $\Omega$  may represent different volumes occupied by different tissues or organs and one can think that their geometrical configuration is given by means of other imaging modalities such as MRI. We also recall [7, 8, 17–21, 57, 58] where similar Lipschitz stability results have been obtained for the classical and fractional Calderón’s problem, the Lamé parameters and for a Schrödinger type of equation.

In this paper we address the issue of stability in Calderón's problem in presence of anisotropy. This choice is motivated by the fact that anisotropy appears quite often in nature. Most tissues in the human body are anisotropic. In the theory of homogenization, anisotropy results as a limit in layered or fibrous structures such as rock stratum or muscle, as a result of crystalline structure or of deformation of an isotropic material. In the geophysical context, in 1920, Schlumberger [59] recognized that anisotropy may affect geological formations' electrical properties and anisotropic effects when measuring electromagnetic fields in geophysical applications have been studied ever since. Individual minerals are typically anisotropic but rocks composed of them can appear to be isotropic.

From a mathematical point of view, the inverse problem with anisotropic conductivities is an open problem. Since Tartar's observation [42] that any diffeomorphism of  $\Omega$  which keeps the boundary points fixed has the property of leaving the D–N map unchanged, whereas  $\sigma$  is modified, different lines of research have been pursued. One direction has been to find the conductivity up to a diffeomorphism which keeps the boundary fixed (see [13, 16, 45–47, 54, 60]). Another direction has been the one to formulate suitable *a priori* assumptions (possibly fitting some physical context) which constrain the structure of the unknown anisotropic conductivity. For instance, one can formulate the hypothesis that the directions of anisotropy are known while some scalar space dependent parameter is not. Along this line of reasoning, we mention the results in [1, 5, 6, 36, 37, 43, 48]. We also refer to [4, 13, 16, 26, 27, 34, 45] and for related results in the anisotropic case and to [4, 40, 41] for examples of non-uniqueness.

Here, we follow this second direction by *a priori* assuming that the conductivity is of type

$$\sigma(x) = \sum_{m=1}^N \gamma_m(x) \chi_{D_m}(x) A(x), \quad \text{for any } x \in \Omega, \quad (1.2)$$

where  $\gamma_m(x)$  is an unknown affine scalar function on  $D_m$ ,  $A$  is a known Lipschitz continuous matrix-valued function on  $\Omega$  and  $\{D_m\}_{m=1}^N$  is a given partition of  $\Omega$  (the precise assumptions on  $\sigma$ ,  $A$  and  $\{D_m\}_{m=1}^N$  are given in subsections 2.1 and 2.2). Allowable partitions for our machinery to work include, in the geophysical setting, models of layered media and bodies with multiple inclusions. The ill-posed nature of the EIT inversion is aggravated the deeper one tries to image inside a body  $\Omega$  [55], where EIT image resolution becomes quite poor (see [39]), leading to blurry images. Thus, in a geophysical context for example, it becomes difficult to recognise individual thin sediments and rock layers or fractures in the deep subsurface, but the 'average' effect at large scale of fine layering and fracturing are still shown as equivalent anisotropic media. It seems therefore reasonable to model the conductivity  $\sigma$  within each layer  $D_m$  by an anisotropic conductivity  $\sigma_m$  to make it up for the finer layering structure within  $D_m$  that otherwise might have been neglected the deeper one goes inside  $\Omega$  due to poor resolution.

In order to introduce the misfit functional, consider two anisotropic conductivities  $\sigma^{(1)}$  and  $\sigma^{(2)}$  of type (1.2). If measurements are locally taken on an open portion  $\Sigma \subset \partial\Omega$ , we conveniently enlarge the physical domain  $\Omega$  to an augmented domain  $\tilde{\Omega}$  and consider Green's functions  $G_i$  for  $\operatorname{div}(\sigma^{(i)} \nabla \cdot)$  in  $\tilde{\Omega}$ , for  $i = 1, 2$ , with poles  $y, z \in \tilde{\Omega} \setminus \bar{\Omega}$  respectively. Hence we express the error in the measurements corresponding to  $\sigma^{(1)}$  and  $\sigma^{(2)}$  by means of the misfit functional

$$\mathcal{J}(\sigma^{(1)}, \sigma^{(2)}) = \int_{D_y \times D_z} |S_{U_0}(y, z)|^2 dy dz, \quad (1.3)$$

where  $D_y, D_z$  are suitably chosen sets compactly contained in  $\tilde{\Omega} \setminus \bar{\Omega}$  and  $S_{U_0}(y, z)$  is defined by the surface integral

$$S_{\mathcal{L}_0}(y, z) = \int_{\Sigma} [G_2(\cdot, z)\sigma^{(1)}(\cdot)\nabla G_1(\cdot, y) \cdot \nu - G_1(\cdot, y)\sigma^{(2)}(\cdot)\nabla G_2(\cdot, z) \cdot \nu] dS. \quad (1.4)$$

We have obtained the following stability estimate of Hölder type:

$$\|\sigma^{(1)} - \sigma^{(2)}\|_{L^\infty(\Omega)} \leq C(\mathcal{J}(\sigma^{(1)}, \sigma^{(2)}))^{1/2}, \quad (1.5)$$

where  $C > 0$  is a constant that depends on the *a priori* information only. The augmented domain  $\tilde{\Omega}$  is chosen in such a way that  $G_1(\cdot, y)|_{\partial\Omega}$ ,  $G_2(\cdot, z)|_{\partial\Omega}$  are supported in  $\Sigma$  in the trace sense, hence belonging to the domain of the local D–N maps  $\Lambda_{\sigma_i}^\Sigma$ ,  $i = 1, 2$  (see section 2.3 for the formal definitions of the local D–N map and the appropriate spaces). Therefore, not only (1.5), together with the well-known Alessandrini’s identity [2], implies a Lipschitz stability estimate of  $\sigma$  in terms of the more commonly used local D–N map in the mathematical literature, but it also indicates that the set of measurements  $\{G(\cdot, y)|_{\partial\Omega}\}$ , with  $y, z \in \tilde{\Omega} \setminus \tilde{\Omega}$  is enough to stably determine  $\sigma$ . A Lipschitz stability estimate in terms of  $\Lambda_\sigma^\Sigma$  was obtained in [37] for the case  $\sigma = \gamma A$ , with  $\gamma$  piecewise constant instead.

The piecewise affine parametrizations considered in the present work tie in well with the finite elements method for computations. Compared to the previous work in [37], this accounts to iteratively determine boundary values and normal derivatives of the conductivity at the various interfaces of the domain partition. In turn, this involves on one hand an asymptotic analysis of Green’s functions (for the conductivity operator on  $\tilde{\Omega}$ ) and their derivatives up to order two at the interfaces between contiguous domains of the partition of  $\Omega$ , where the conductivity changes (possibly) discontinuously (see proposition 3.1). On the other hand, our machinery also relies on an argument of quantitative estimates of propagation of smallness of singular solutions and their derivatives up to order two as well (see proposition 3.2). The argument in [37] was based on an iterative determination of boundary values of the conductivity at the various interfaces of the domain partition only. This, in turn, required an asymptotic analysis of Green’s functions and their first derivatives only at such interfaces together with quantitative estimates of propagation of smallness of the singular solutions and their first derivatives only.

With the stability estimate (2.13) at hand, one can apply certain iterative methods for reconstruction within a subspace of piecewise affine functions with a starting model at a distance less than the radius of convergence to the unique solution [9, 31–33]. This radius is known to be roughly inversely proportional to the stability constant appearing in the estimate. More importantly, we can iteratively construct the best piecewise affine approximation for a given domain partition. Since the stability constant will grow at least exponentially with the number of subdomains in the partition [56], the radius of convergence shrinks accordingly. One can expect accurate piecewise affine approximations with relatively less subdomains (compared to the piecewise constant case of [37]) to describe the subsurface, noting that the domain partition need not be uniform and may show a local refinement, and hence our result provides the necessary insight for developing a practical approach with relatively minor prior information.

To the best of our knowledge a first stability estimate in terms of an ad hoc misfit functional was achieved in the mathematical literature in [9] in the context of the full waveform inversion. Such an estimate proved to be key for the implementation and reliability of a reconstruction procedure (see [9, 33]) based on the use of Cauchy data only, being the latter independent on the availability of the D–N map. In the more recent result in [35] an ad hoc misfit functional has been introduced in the context of imaging elastic media.

We also observe that another advantage of choosing the misfit functional over the local D–N map (even if available) to model the measurements error in EIT is motivated by its potentially simpler numerical implementation, compared to the computation of the norm of bounded linear operators between  $H^{\frac{1}{2}}$  spaces and their duals. Moreover, the misfit functional

could also provide, again in the context of a possible numerical reconstruction of  $\sigma$ , additional features compared to the more traditional least-squares approach, allowing, in particular, for a distinction between the computational and the observational measurements. This is due to the introduction of the possibly distinct sets  $D_y$  and  $D_z$  that can almost be arbitrarily chosen outside the physical domain  $\Omega$ . For example,  $D_y$  could be an arbitrarily chosen set for the numerical data acquisition for the sake of the simulations, where  $D_z$  could model a more realistic set that fits the geometric disposition of the electrodes in the actual measurements acquisition. Hence, in the discrete setting, such distinction can potentially require minimal information about the observational acquisition geometry of the electrodes employed for the observational measurements. This is due to the definition of the misfit functional that does not compare simulations and observations directly, but it rather compares products of observed and simulated measurements. Note also that with a slight modification, our arguments can apply when the local Neumann-to-Dirichlet (N–D) map is available instead, see for instance the discussion in [6].

The paper is organized as follows. In section 2 we introduce the main assumptions on the domain  $\Omega$  and the anisotropic conductivity  $\sigma$ . Section 2 contains the formal definitions of the local D–N map (subsection 2.3), the misfit functional (subsection 2.4) and the statement of our main result (theorem 2.1). A Lipschitz stability estimate in terms of the local D–N map follows as a straightforward consequence (corollary 2.2). Section 3 is devoted to the introduction of some technical tools of asymptotic estimates for the Green function (proposition 3.1) and propagation of smallness (proposition 3.2) needed for the machinery of the proof of theorem 2.1. The proof of theorem 2.1 and corollary 2.2 are also contained in this section. Section 4 contains the proofs of proposition 3.1 and proposition 3.2.

## 2. Misfit functional and the main result

### 2.1. Assumptions about the domain $\Omega$

For  $n \geq 3$ , a point  $x \in \mathbb{R}^n$  will be denoted by  $x = (x', x_n)$ , where  $x' \in \mathbb{R}^{n-1}$  and  $x_n \in \mathbb{R}$ . Moreover, given a point  $x \in \mathbb{R}^n$ , we will denote with  $B_r(x)$ ,  $B'_r(x')$  the open balls in  $\mathbb{R}^n$ ,  $\mathbb{R}^{n-1}$  respectively centred at  $x$  and  $x'$  with radius  $r$  and by  $Q_r(x)$  the cylinder

$$Q_r(x) = B'_r(x') \times (x_n - r, x_n + r).$$

Set  $B_r = B_r(0)$ ,  $Q_r = Q_r(0)$ , the positive real half space  $\mathbb{R}_+^n = \{(x', x_n) \in \mathbb{R}^n : x_n > 0\}$ , the positive semisphere centred at the origin  $B_r^+ = B_r \cap \mathbb{R}_+^n$ , the positive semicylinder  $Q_r^+ = Q_r \cap \mathbb{R}_+^n$ . Similar definitions for  $\mathbb{R}_-^n$ ,  $B_r^-$  and  $Q_r^-$ .

Let us recall a couple of definitions concerning the regularity of the boundary of the domain.

**Definition 2.1.** Let  $\Omega$  be a bounded domain in  $\mathbb{R}^n$ . A portion  $\Sigma$  of  $\partial\Omega$  is of Lipschitz class with constants  $r_0, L > 0$  if for each point  $P \in \Sigma$  there exists a rigid transformation of coordinates under which  $P$  coincides with the origin and

$$\Omega \cap Q_{r_0} = \{x \in Q_{r_0} : x_n > \varphi(x')\},$$

where  $\varphi$  is a Lipschitz function on  $B'_{r_0}$  such that  $\varphi(0) = 0$  and  $\|\varphi\|_{C^{0,1}(B'_{r_0})} \leq Lr_0$ .

**Definition 2.2.** Let  $\Omega$  be a domain in  $\mathbb{R}^n$ . A subset  $\Sigma$  of  $\partial\Omega$  is a flat portion of size  $r_0$  if for each point  $P \in \Sigma$  there exists a rigid transformation of coordinates under which  $P$  coincides with the origin and

$$\Sigma \cap Q_{r_0} = \{x \in Q_{r_0} : x_n = 0\}, \quad \Omega \cap Q_{r_0} = \{x \in Q_{r_0} : x_n > 0\}.$$

From now on, we will consider  $\Omega \subset \mathbb{R}^n$ ,  $n \geq 3$  as a bounded, measurable domain with boundary  $\partial\Omega$  of Lipschitz class with positive constants  $r_0, L$  as in definition 2.1 and satisfying

$$|\Omega| \leq Nr_0^n, \quad (2.1)$$

where  $|\Omega|$  denotes the Lebesgue measure of  $\Omega$ . Moreover, we assume that there exists a partition of bounded subdomains  $D = \{D_m\}_{m=1}^N$  contained in  $\Omega$  such that the following conditions hold:

- (a)  $D_m$  for  $m = 1, \dots, N$  are connected, pairwise non-overlapping subdomains with boundaries  $\partial D_m$  which are of Lipschitz class with constants  $r_0, L$
- (b)  $\bar{\Omega} = \bigcup_{m=1}^N \bar{D}_m$ ;
- (c) (Chain of subdomains.) First, we assume that there exists one region, let us call it  $D_1$ , such that the intersection  $\partial D_1 \cap \Sigma$  contains a *flat* portion  $\Sigma_1$  of size  $r_0/3$  (see definition 2.2) and that for every  $i \in \{2, \dots, N\}$  there exists a collection of indices  $m_1, \dots, m_K \in \{1, \dots, N\}$  such that  $D_{m_1} = D_1$  and  $D_{m_K} = D_i$  and the subdomains are pairwise disjoint. Secondly, we assume that, for every fixed sub-index  $k = 1, \dots, K$  of the chain, the intersection  $\partial D_{m_k} \cap \partial D_{m_{k+1}}$  contains a *flat* portion  $\Sigma_{m_{k+1}}$  of size  $r_0/3$  such that  $\Sigma_{m_{k+1}} \subset \Omega$  for  $k = 1, \dots, K-1$ . Finally, for each of these flat sub-portions  $\Sigma_{m_{k+1}}$ ,  $k = 1, \dots, K-1$ , there exist a point  $P_{k+1} \in \Sigma_{m_{k+1}}$  and a rigid transformation of coordinates under which  $P_{k+1}$  coincides with the origin and

$$\begin{aligned} \Sigma_{m_{k+1}} \cap Q_{r_0/3} &= \{x \in Q_{r_0/3} : x_n = 0\}, \\ D_{m_k} \cap Q_{r_0/3} &= \{x \in Q_{r_0/3} : x_n < 0\}, \\ D_{m_{k+1}} \cap Q_{r_0/3} &= \{x \in Q_{r_0/3} : x_n > 0\}. \end{aligned}$$

Later, we will add a domain  $D_0 \subset \mathbb{R}^n \setminus \bar{\Omega}$  so that, when indexing the chain of subdomains, we agree that  $D_{m_0} = D_0$ .

## 2.2. A priori information on the anisotropic conductivity $\sigma$

Our stability result for the Calderón inverse problem concerns a special family of anisotropic conductivities  $\sigma$ . Let us describe in details their form. The conductivities  $\sigma(x) = \{\sigma_{ij}(x)\}$  are real-valued, symmetric  $n \times n$  matrices such that  $\sigma \in L^\infty(\Omega, \text{Sym}_n)$  and have the form

$$\sigma(x) = \gamma(x)A(x) \quad (2.2a)$$

$$\gamma(x) = \sum_{m=1}^N \gamma_m(x)\chi_{D_m}(x), \quad \gamma_m(x) = s_m + S_m \cdot x, \quad \text{for any } x \in \Omega, \quad (2.2b)$$

where the scalars  $s_m \in \mathbb{R}$  and the vectors  $S_m \in \mathbb{R}^n$ ,  $m = 1, \dots, N$  are the unknowns,  $A(x)$  is a known fixed matrix and  $D = \{D_m\}_{m=1}^N$  is the known partition of  $\Omega$  introduced in section 2.1. Furthermore,

- (a) The scalar functions  $\gamma_m$  are bounded, piecewise linear and there is a positive constant  $\bar{\gamma} > 1$  such that

$$\bar{\gamma}^{-1} \leq \gamma_m(x) \leq \bar{\gamma}, \quad \text{for any } m = 1, \dots, N, \text{ for any } x \in \Omega; \quad (2.3)$$

- (b) The matrix  $A(x)$  satisfies the following Lipschitz continuity condition: there exists a constant  $\bar{A} > 0$  such that  $\|A\|_{C^{0,1}(\Omega)} \leq \bar{A}$ ;

(c) There exists a constant  $\lambda > 1$  such that

$$\lambda^{-1}|\xi|^2 \leq A(x)\xi \cdot \xi \leq \lambda|\xi|^2, \quad \text{for a.e. } x \in \Omega, \text{ for every } \xi \in \mathbb{R}^n. \quad (2.4)$$

**Definition 2.3.** The set of positive constants  $\{N, r_0, L, \lambda, \bar{\gamma}, \bar{A}, n\}$  with  $N \in \mathbb{N}$  and the space dimension  $n \geq 3$ , is called the *a priori data*.

In the paper several constants depending on the *a priori data* will appear. In order to simplify our notation, we will denote them by  $C, C_1, C_2, \dots$ , avoiding in most cases to point out their specific dependence on the *a priori data* which may vary from case to case.

### 2.3. The local D–N map

By now, assume simply that  $\Omega$  is a bounded domain with  $\partial\Omega$  of Lipschitz class. Since Dirichlet data are different from zero on a small portion  $\Sigma \subset \partial\Omega$ , we introduce a suitable trace space for the formulation of the local D–N map.

**Definition 2.4.** Let  $\Sigma$  be a non-empty (flat) open portion of  $\partial\Omega$ . The subspace of  $H^{1/2}(\partial\Omega)$  of trace functions which are compactly supported in  $\Sigma$  is defined as

$$H_{\text{co}}^{1/2}(\Sigma) = \left\{ f \in H^{1/2}(\partial\Omega) : \text{supp } f \subset \Sigma \right\}. \quad (2.5)$$

The trace space  $H_{00}^{1/2}(\Sigma)$  is the closure of  $H_{\text{co}}^{1/2}(\Sigma)$  with respect to the  $H^{1/2}(\partial\Omega)$ -norm. We denote by  $H_{00}^{-1/2}(\Sigma)$  the dual of the trace space  $H_{00}^{1/2}(\Sigma)$ .

**Definition 2.5.** The local D–N map associated with  $\sigma$  and  $\Sigma$  is the operator

$$\begin{aligned} \Lambda_{\sigma}^{\Sigma} : H_{00}^{1/2}(\Sigma) &\rightarrow H_{00}^{-1/2}(\Sigma) \\ g &\mapsto \sigma \nabla u \cdot \nu|_{\Sigma}, \end{aligned} \quad (2.6)$$

where  $\nu$  is the unit outward normal of  $\partial\Omega$  and  $u \in H^1(\Omega)$  is the weak solution to the boundary value problem

$$\begin{cases} \operatorname{div}(\sigma(\cdot)\nabla u) = 0, & \text{in } \Omega, \\ u = g, & \text{on } \partial\Omega. \end{cases}$$

The map (2.6) can be identified with the bilinear form  $H_{00}^{1/2}(\Sigma) \times H_{00}^{1/2}(\Sigma) \rightarrow \mathbb{R}$  defined by

$$\langle \Lambda_{\sigma}^{\Sigma} g, \eta \rangle = \int_{\Omega} \sigma(x) \nabla u(x) \cdot \nabla \varphi(x) dx, \quad (2.7)$$

where  $\eta \in H_{00}^{1/2}(\Sigma)$  and  $\varphi \in H^1(\Omega)$  is any function such that  $\varphi|_{\Sigma} = \eta$ . In (2.7) the bracket  $\langle \cdot, \cdot \rangle$  denotes the  $L^2(\partial\Omega)$ -pairing between  $H_{00}^{1/2}(\Sigma)$  and its dual  $H_{00}^{-1/2}(\Sigma)$ .

For simplicity, we will denote by  $\|\cdot\|_*$  the  $\mathcal{L}(H_{00}^{1/2}(\Sigma), H_{00}^{-1/2}(\Sigma))$ -norm of the Banach space of bounded linear operators from  $H_{00}^{1/2}(\Sigma)$  to  $H_{00}^{-1/2}(\Sigma)$ .

### 2.4. Misfit functional

To begin with, we introduce the Green function  $G$  in an augmented domain  $\tilde{\Omega}$  as follows. From the assumptions on the domain  $\Omega$  (section 2.1) there is a point  $P_1 \in \Sigma$  that coincides with the



origin, up to a rigid transformation of coordinates. For simplicity, let us assume that the locally flat portion  $\Sigma_1$  coincides with the entire portion  $\Sigma$ . Let us define the domain  $D_0 \subset \mathbb{R}^n \setminus \overline{\Omega}$  as

$$D_0 = \left\{ x \in (\mathbb{R}^n \setminus \overline{\Omega}) \cap B_{r_0} \mid |x_i| < \frac{r_0}{3}, \quad i = 1, \dots, n-1, \quad -\frac{r_0}{3} < x_n < 0 \right\}, \quad (2.8)$$

and such that

$$\partial D_0 \cap \partial \Omega \subset \subset \Sigma.$$

We define the augmented domain  $\tilde{\Omega}$  as the set

$$\tilde{\Omega} = \overline{\Omega \cup D_0}. \quad (2.9)$$

It turns out that  $\tilde{\Omega}$  is of Lipschitz class with constants  $\frac{r_0}{3}$  and  $\tilde{L}$ , where  $\tilde{L}$  depends on  $L$  only.

Denote

$$(D_0)_r = \{x \in D_0 : \text{dist}(x, \partial D_0) > r\}, \quad r \in \left(0, \frac{r_0}{6}\right).$$

Finally, we introduce two sets contained in  $D_0$ : the sets  $D_y$  and  $D_z$  which are compactly supported in  $D_0$ , i.e.  $D_y, D_z \subset \subset D_0$ . In the following sections, we might identify these sets with the set  $(D_0)_r$ , but in general, they can be freely chosen in  $D_0$ .

Consider two anisotropic conductivities  $\sigma^{(i)}$ ,  $i = 1, 2$  as in section 2.2. Without loss of generality, we can extend them to the augmented domain  $\tilde{\Omega}$  by setting their value equal to the identity matrix on  $D_0$ , so that they are of the form

$$\begin{aligned} \sigma^{(i)}(x) &= \gamma^{(i)}(x)A(x), \quad \text{for any } x \in \Omega, \\ \sigma^{(i)}|_{D_0} &= I, \quad \gamma^{(i)}|_{D_0} = 1. \end{aligned}$$

We denote with the same symbol  $\sigma$  the extended conductivity.

Denote with  $G_i$  the Green function associated with the operator  $\text{div}(\sigma^{(i)}(\cdot)\nabla\cdot)$  and  $\tilde{\Omega}$ , for  $i = 1, 2$ . For every  $y \in D_0$ ,  $G_i(\cdot, y)$  is the weak solution to the Dirichlet problem

$$\begin{cases} \text{div}(\sigma^{(i)}(\cdot)\nabla G_i(\cdot, y)) = -\delta(\cdot - y) & \text{in } \tilde{\Omega}, \\ G_i(\cdot, y) = 0 & \text{on } \partial\tilde{\Omega}, \end{cases} \quad (2.10)$$

where  $\delta(\cdot - y)$  is the Dirac distribution centred at  $y$ .

We recall the following properties for the Green's functions (see [51]):

$$G(x, y) = G(y, x), \quad \forall x \neq y,$$

and

$$0 < G(x, y) < C|x - y|^{2-n}, \quad \forall x \neq y, \quad (2.11)$$

where  $C$  is a positive constant depending on  $\lambda$  and  $n$ . For  $(y, z) \in D_y \times D_z$ , define the following surface integral

$$S_{\mathcal{U}_0}(y, z) = \int_{\Sigma} [G_2(x, z) \sigma^{(1)}(x) \nabla G_1(x, y) \cdot \nu - G_1(x, y) \sigma^{(2)}(x) \nabla G_2(x, z) \cdot \nu] \, dS(x). \quad (2.12)$$

We define the *misfit functional* as the quantity

$$\mathcal{J}(\sigma^{(1)}, \sigma^{(2)}) = \int_{D_y \times D_z} |S_{\mathcal{U}_0}(y, z)|^2 dy dz. \quad (2.13)$$

### 2.5. Stability estimate

In previous works (see [7, 11, 37]), Lipschitz stability estimates have been established for piecewise constant and piecewise linear isotropic conductivities and a certain class of anisotropic conductivities respectively, in terms of the local D–N map. Here, we extend these results to the class of anisotropic conductivities defined in section 2.2. First, we determine a bound to the  $L^\infty$ -norm of the difference between two anisotropic conductivities in terms of the square root of the misfit functional introduced above. Then, we derive a Lipschitz stability result in terms of the local D–N map.

**Theorem 2.1.** *Let  $\Omega$  be a bounded domain as in assumptions (2.1). Let  $\sigma^{(1)}$  and  $\sigma^{(2)}$  be two anisotropic conductivities as in assumptions (2.2), i.e. of the form*

$$\sigma^{(i)}(x) = \sum_{m=1}^N \gamma_m^{(i)}(x) \chi_{D_m}(x) A(x), \quad \text{for any } x \in \Omega, \quad i = 1, 2, \quad (2.14)$$

where  $D = \{D_m\}_{m=1}^N$  is the partition of subdomains as in assumptions (2.1),  $A(x)$  is the known Lipschitz matrix and  $\gamma_m^{(i)}(x)$  are the piecewise-affine functions given by the formula

$$\gamma_m^{(i)}(x) = s_m^{(i)} + S_m^{(i)} \cdot x, \quad x \in D_m,$$

for  $s_m^{(i)} \in \mathbb{R}$  and  $S_m^{(i)} \in \mathbb{R}^n$ . Then there exists a positive constant  $C$  such that

$$\|\sigma^{(1)} - \sigma^{(2)}\|_{L^\infty(\Omega)} \leq C (\mathcal{J}(\sigma^{(1)}, \sigma^{(2)}))^{1/2}, \quad (2.15)$$

where  $C$  depends on the a priori data only.

From this result, it follows a Lipschitz stability estimate in terms of the local D–N maps.

**Corollary 2.2.** *Assume that the hypothesis of theorem 2.1 hold, then*

$$\|\sigma^{(1)} - \sigma^{(2)}\|_{L^\infty(\Omega)} \leq C \|\Lambda_{\sigma^{(1)}}^\Sigma - \Lambda_{\sigma^{(2)}}^\Sigma\|_*, \quad (2.16)$$

where  $C > 0$  is a constant depending on the a priori data only.

**Remark 2.3.** From now on, as we deal with two different anisotropic conductivities  $\sigma^{(i)}$ ,  $i = 1, 2$ , we will simply denote with the symbol  $\Lambda_i$  the local DN map  $\Lambda_{\sigma^{(i)}}^\Sigma$ .

## 3. Proof of the main result

The proof of theorem 2.1 is based on an argument that combines asymptotic estimates for the Green's function of the elliptic operator  $\text{div}(\sigma(\cdot)\nabla\cdot)$  (proposition 3.1), together with a result of unique continuation (proposition 3.2). In this section we introduce these technical results (proved in section 4), then we prove theorem 2.1 and corollary 2.2.

### 3.1. Technical tools

3.1.1. *Behaviour of Green's function near interfaces.* We shall denote with

$$\Gamma(x, y) = \frac{1}{n(2-n)\omega_n} |x-y|^{2-n}, \quad \omega_n = \frac{2\pi^{n/2}}{n\Gamma(n/2)}, \quad (3.1)$$

the fundamental solution for the Laplace operator (here  $\omega_n$  denotes the volume of the unit ball in  $\mathbb{R}^n$ ).

Let  $\{D_m\}_{m=0}^K$ ,  $K \in \{1, \dots, N\}$  be the chain of subdomains as in assumptions (2.1),  $\{\Sigma_m\}_{m=1}^K$  be the corresponding sequence of flat portions with special points  $P_1, \dots, P_K$ . Moreover, let  $\nu(P_{m+1})$  denotes the unit normal to  $\partial D_m$  at the point  $P_{m+1}$  pointing outside  $D_m$ .

**Proposition 3.1 (Asymptotic estimates).** *Fix an index  $m \in \{0, \dots, K-1\}$ , then there exist constants  $\alpha, \theta_1, \theta_2, 0 < \alpha, \theta_1, \theta_2 < 1$  and  $C_1, C_2, C_3 > 0$  depending on the a priori data only and a suitable constant  $C_4 > 1$  such that the following inequalities hold true for every  $x \in B_{\frac{r_0}{C_4}}(P_{m+1}) \cap D_{m+1}$  and every  $y = P_{m+1} - r\nu(P_{m+1})$ , where  $r \in (0, \frac{r_0}{C_4})$*

$$\left| G(x, y) - \frac{2}{\gamma_m(P_{m+1}) + \gamma_{m+1}(P_{m+1})} \Gamma(Jx, Jy) \right| \leq C_1 |x-y|^{3-n-\alpha}, \quad (3.2)$$

$$\left| \nabla_x G(x, y) - \frac{2}{\gamma_m(P_{m+1}) + \gamma_{m+1}(P_{m+1})} \nabla_x \Gamma(Jx, Jy) \right| \leq C_2 |x-y|^{1-n+\theta_1}, \quad (3.3)$$

$$\left| \nabla_y \nabla_x G(x, y) - \frac{2}{\gamma_m(P_{m+1}) + \gamma_{m+1}(P_{m+1})} \nabla_y \nabla_x \Gamma(Jx, Jy) \right| \leq C_3 |x-y|^{-n+\theta_2}, \quad (3.4)$$

where  $J$  is the positive definite matrix  $J = \sqrt{A(P_{m+1})}^{-1}$ .

3.1.2. *Quantitative estimates of propagation of smallness.* For any number  $b > 0$ , define the concave, non decreasing function  $\omega_b(t)$  on  $(0, +\infty)$  as

$$\omega_b(t) = \begin{cases} 2^b e^{-2} |\log t|^{-b}, & t \in (0, e^{-2}), \\ e^{-2}, & t \in [e^{-2}, +\infty). \end{cases}$$

We recall (see (4.34) and (4.35) in [11]) that

$$(0, +\infty) \ni t \rightarrow t\omega_b\left(\frac{1}{t}\right) \text{ is a non-decreasing function} \quad (3.5)$$

and for any  $\beta \in (0, 1)$  we have that

$$\omega_b\left(\frac{t}{\beta}\right) \leq |\log e\beta^{-1/2}|^b \omega_b(t), \quad \omega_b(t^\beta) \leq \left(\frac{1}{\beta}\right)^b \omega_b(t). \quad (3.6)$$

Furthermore, we shall denote the iterative compositions of  $\omega$  as

$$\omega_b^{(1)} = \omega_b, \quad \omega_b^{(j)} = \omega_b \circ \omega_b^{(j-1)} \quad j = 2, 3, \dots,$$

and we set  $\omega_b^{(0)}(t) = t^b$  for  $0 < b < 1$ .

Fix a chain of subdomains  $\{D_m\}_{m=0}^K$  as in assumptions (2.1) for the domain  $\tilde{\Omega}$ . Set

$$\mathcal{W}_k = \bigcup_{m=0}^k D_m, \quad \mathcal{U}_k = \tilde{\Omega} \setminus \overline{\mathcal{W}_k}, \quad \text{for } k = 0, \dots, K. \quad (3.7)$$

**Definition 3.1.** For any  $y, z \in \mathcal{W}_k$ , define the singular solution

$$S_{\mathcal{U}_k}(y, z) = \int_{\mathcal{U}_k} (\sigma^{(1)}(\cdot) - \sigma^{(2)}(\cdot)) \nabla G_1(\cdot, y) \cdot \nabla G_2(\cdot, z), \quad \text{for } k = 0, \dots, K.$$

The set  $\{S_{\mathcal{U}_k}(y, z)\}_{k=0}^K$  is a family of real-valued functions which satisfies the following inequality:

$$|S_{\mathcal{U}_k}(y, z)| \leq C \|\sigma^{(1)} - \sigma^{(2)}\|_{L^\infty(\Omega)} (d(y)d(z))^{1-\frac{n}{2}}, \quad \text{for every } y, z \in \mathcal{W}_k, \quad (3.8)$$

where  $d(y) = \text{dist}(y, \mathcal{U}_k)$  and  $C$  is a positive constant depending on  $\lambda$  and  $n$  only.

One can prove (see [11]) that for every  $y, z \in \mathcal{W}_k$  with  $k = 0, \dots, K$ , the functions  $S_{\mathcal{U}_k}(\cdot, z), S_{\mathcal{U}_k}(y, \cdot)$  belongs to  $H_{\text{loc}}^1(\mathcal{W}_k)$  and are weak solutions, respectively, to

$$\text{div}(\sigma^{(1)}(\cdot) \nabla S_{\mathcal{U}_k}(\cdot, z)) = 0, \quad \text{div}(\sigma^{(2)}(\cdot) \nabla S_{\mathcal{U}_k}(y, \cdot)) = 0 \text{ in } \mathcal{W}_k.$$

We introduce the following parameters:

$$\begin{aligned} \beta &= \arctan \frac{1}{L}, \quad \beta_1 = \arctan \left( \frac{\sin \beta}{4} \right), \quad \lambda_1 = \frac{r_0}{1 + \sin \beta_1}, \\ \rho_1 &= \lambda_1 \sin \beta_1, \quad a = \frac{1 - \sin \beta_1}{1 + \sin \beta_1}, \\ \lambda_m &= a \lambda_{m-1}, \quad \rho_m = a \rho_{m-1}, \quad \text{for every } m \geq 2, \\ d_m &= \lambda_m - \rho_m, \quad m \geq 1. \end{aligned} \quad (3.9)$$

Notice that  $d_m = r_0 a^m, 0 < a < 1$ .

Choose  $l \in \mathbb{N}$ , fix a point  $\bar{y} \in \Sigma_{m+1}$ , then define

$$w = w_l(\bar{y}) = \bar{y} - \lambda_l \nu(\bar{y}), \quad \text{for every } l \geq 1, \quad (3.10)$$

where  $w$  is a point into the domain  $D_m$  near the interface  $\Sigma_{m+1}$ . For a given  $r \in (0, d_1]$  define the function

$$\bar{h}(r) = \min\{l \in \mathbb{N} : d_l \leq r\}. \quad (3.11)$$

For successive estimates, it is important to point out the following inequality:

$$\log \left( \frac{r}{d_1} \right)^C \leq \bar{h}(r) - 1 \leq \log \left( \frac{r}{d_1} \right)^C + 1, \quad C = \frac{1}{|\log a|}. \quad (3.12)$$

The following estimate for  $S_{\mathcal{U}_k}(y, z)$  holds true, for any  $k = 1, \dots, K$ .

**Proposition 3.2 (Estimates of unique continuation).** *Suppose that for a positive number  $\varepsilon_0$  and  $r > 0$  we have*

$$|S_{U_k}(y, z)| \leq r_0^{2-n} \varepsilon_0, \quad \text{for every } (y, z) \in (D_0)_r \times (D_0)_r, \quad (3.13)$$

then the following inequalities hold true for every  $r \in (0, d_1]$

$$|S_{U_k}(w_{\bar{h}}(Q_{k+1}), w_{\bar{h}}(Q_{k+1}))| \leq C_1^{\bar{h}}(E + \varepsilon_0) \left( \omega_{1/C}^{(2k)} \left( \frac{\varepsilon_0}{E + \varepsilon_0} \right) \right)^{(1/C)^{\bar{h}}}, \quad (3.14)$$

$$|\partial_{y_j} \partial_{z_i} S_{U_k}(w_{\bar{h}}(Q_{k+1}), w_{\bar{h}}(Q_{k+1}))| \leq C_2^{\bar{h}}(E + \varepsilon_0) \left( \omega_{1/C}^{(2k)} \left( \frac{\varepsilon_0}{E + \varepsilon_0} \right) \right)^{(1/C)^{\bar{h}}}, \quad (3.15)$$

for any  $i, j = 1, \dots, n$ , where  $Q_{k+1} \in \Sigma_{k+1} \cap B_{\frac{r_0}{8}}(P_{k+1})$ ,  $w_{\bar{h}(r)}(Q_{k+1}) = Q_{k+1} - \lambda_{\bar{h}(r)} \nu(Q_{k+1})$ , with  $\lambda_{\bar{h}(r)}$  as above,  $\nu(Q_{k+1})$  is the exterior unit normal to  $\partial D_k$  at the point  $Q_{k+1}$  pointing outside  $D_k$  and  $C_1, C_2 > 0$  depend on the *a priori* data only.

### 3.2. Proof of theorem 2.1 and the corollary 2.2

**Proof of theorem 2.1.** First, notice that

$$\|\sigma^{(1)} - \sigma^{(2)}\|_{L^\infty(\Omega)} \leq \|\gamma^{(1)} - \gamma^{(2)}\|_{L^\infty(\Omega)} \bar{A},$$

where  $\bar{A}$  is the Lipschitz constant from assumptions (2.2). Let  $D_K$  be the subdomain of  $\Omega$  such that

$$\|\gamma^{(1)} - \gamma^{(2)}\|_{L^\infty(\Omega)} = \|\gamma_K^{(1)} - \gamma_K^{(2)}\|_{L^\infty(D_K)}.$$

Then, inequality (2.15) will follow from

$$\|\gamma_K^{(1)} - \gamma_K^{(2)}\|_{L^\infty(D_K)} \leq C(\mathcal{J}(\sigma^{(1)}, \sigma^{(2)}))^{1/2}, \quad (3.16)$$

for  $C > 1$  a positive constant depending on *a priori* estimates.

To prove (3.16), we find convenient, as previously stated, to work in the augmented domain  $\tilde{\Omega}$  as in (2.9), where  $D_0$  is the domain defined in (2.8), on which we have defined the extended conductivity  $\sigma^{(i)}$  for  $i = 1, 2$  by setting  $\sigma^{(i)}|_{D_0} = I$ . Recalling that  $D_K$  is the subdomain of  $\Omega$  where the maximum of  $|\gamma^{(1)} - \gamma^{(2)}|$  is reached, let  $D_0, D_1, \dots, D_K$  be the chain of subdomains as in section 2.1 and let  $\Sigma_1, \dots, \Sigma_K$  be the corresponding flat portions. Set

$$\varepsilon_0 = (\mathcal{J}(\sigma^{(1)}, \sigma^{(2)}))^{1/2}, \quad E = \|\gamma_K^{(1)} - \gamma_K^{(2)}\|_{L^\infty(D_K)}, \quad (3.17)$$

$$\delta_k = \|\gamma^{(1)} - \gamma^{(2)}\|_{L^\infty(W_k)}, \quad \text{for } k = 1, \dots, K. \quad (3.18)$$

Given a differentiable function  $f$  on a domain  $\Omega$ , we can split its differential as

$$Df(x) = D_T f(x) + \partial_\nu f(x), \quad \text{for } x \in \Sigma_k, \quad k = 1, \dots, K,$$

where  $D_T f$  is the  $n - 1$  dimensional vector of the tangential partial derivatives of  $f$  on  $\Sigma_k$  and  $\partial_\nu f$  denotes the normal partial derivative of  $f$  on  $\Sigma_k$ , respectively for  $k = 1, 2, \dots, K$ .

Fix  $0 < r_2 < r_1$  such that  $\Sigma_k \cap B_{r_1}(P_k) \neq \emptyset$  for  $k = 1, 2, \dots, K$ . We observe that the norm  $\|\gamma_k^{(1)} - \gamma_k^{(2)}\|_{L^\infty(D_k)}$  can be estimated in terms of the quantities

$$\|\gamma_k^{(1)} - \gamma_k^{(2)}\|_{L^\infty(\Sigma_k \cap B_{r_1}(P_k))} \quad \text{and} \quad \left| \partial_\nu(\gamma_k^{(1)} - \gamma_k^{(2)})(P_k) \right|. \tag{3.19}$$

In fact, fix an orthonormal basis  $\{e_j^k\}_{j=1, \dots, n-1}$  which generates the hyperplane containing the flat part  $\Sigma_k$ . Set

$$\alpha_k + \beta_k \cdot x = \left( \gamma_k^{(1)} - \gamma_k^{(2)} \right) (x), \quad x \in D_k.$$

If we evaluate  $\left( \gamma_k^{(1)} - \gamma_k^{(2)} \right)$  at the points  $P_k + r_2 e_j^k$ ,  $j = 1, \dots, n - 1$ , it follows that

$$\left| \alpha_k + \beta_k \cdot (P_k + r_2 e_j^k) \right| \leq |\alpha_k + \beta_k \cdot P_k| + r_1 \sum_{j=1}^{n-1} |\beta_k \cdot e_j^k| \leq C \|\gamma_k^{(1)} - \gamma_k^{(2)}\|_{L^\infty(\Sigma_k \cap B_{r_1}(P_k))}.$$

Next, notice that

$$|\beta_k \cdot \nu| = \left| \partial_\nu(\gamma_k^{(1)} - \gamma_k^{(2)})(P_k) \right|.$$

In conclusion, for  $k = 1, \dots, K$ ,

$$|\alpha_k| + |\beta_k| \leq C \left( \|\gamma_k^{(1)} - \gamma_k^{(2)}\|_{L^\infty(\Sigma_k \cap B_{r_1}(P_k))} + \left| \partial_\nu(\gamma_k^{(1)} - \gamma_k^{(2)})(P_k) \right| \right).$$

Hence, our task will be to estimate the quantities introduced in (3.19) for  $k = 1, \dots, K$  in terms of the function  $\omega_{1/C}$  introduced in section 3.1.2,  $\varepsilon_0$  and  $E$ .

**3.2.1. Boundary estimates.** Let us start from the case  $k = 1$ . We will prove the following estimate:

$$\|\gamma_1^{(1)} - \gamma_1^{(2)}\|_{L^\infty(\Sigma_1 \cap B_{r_1}(P_1))} + \left| \partial_\nu(\gamma_1^{(1)} - \gamma_1^{(2)})(P_1) \right| \leq C(\varepsilon_0 + E)\omega_{1/C}^{(0)} \left( \frac{\varepsilon_0}{\varepsilon_0 + E} \right). \tag{3.20}$$

For every  $y, z \in (D_0)_r$ , by Green formula the following equalities hold:

$$\begin{aligned} & \int_{\Sigma} [G_2(\cdot, z)\sigma^{(1)}(\cdot)\nabla G_1(\cdot, y) \cdot \nu - G_1(\cdot, y)\sigma^{(2)}(\cdot)\nabla G_2(\cdot, z) \cdot \nu] \, dS \\ &= \int_{\Omega} (\sigma^{(1)} - \sigma^{(2)})(\cdot)\nabla G_1(\cdot, y) \cdot \nabla G_2(\cdot, z), \end{aligned} \tag{3.21}$$

and

$$\begin{aligned} & \int_{\Sigma} [\partial_{z_n} G_2(\cdot, z)\sigma^{(1)}(\cdot)\nabla \partial_{y_n} G_1(\cdot, y) \cdot \nu - \partial_{y_n} G_1(\cdot, y)\sigma^{(2)}(\cdot)\nabla \partial_{z_n} G_2(\cdot, z) \cdot \nu] \, dS \\ &= \int_{\Omega} (\sigma^{(1)} - \sigma^{(2)})(\cdot)\nabla \partial_{y_n} G_1(\cdot, y) \cdot \nabla \partial_{z_n} G_2(\cdot, z), \end{aligned} \tag{3.22}$$

where  $G_1(\cdot, y)$  and  $G_2(\cdot, z)$  are weak solutions to the problem (2.10). Since  $S_{\mathcal{U}_0}(y, z)$  and  $\partial_{y_n} \partial_{z_n} S_{\mathcal{U}_0}(y, z)$  are weak solutions to the following equation

$$\operatorname{div} (\sigma^{(1)}(\cdot)\nabla S_{\mathcal{U}_0}(\cdot, z)) + \operatorname{div} (\sigma^{(2)}(\cdot)\nabla S_{\mathcal{U}_0}(y, \cdot)) = 0, \quad \text{in } D_y \times D_z,$$

we can apply a result of local boundedness for weak solutions of a uniformly elliptic operator (see [38, chapter 8, theorem 8.17]) that allows us to bound the supremum of  $S_{\mathcal{L}_0}(y, z)$  by its  $L^2$ -norm as follows:

$$\sup_{(y,z) \in (D_y)_r \times (D_z)_r} |S_{\mathcal{L}_0}(y, z)| \leq Cr^{-n} \left( \int_{D_y \times D_z} |S_{\mathcal{L}_0}(y, z)|^2 dy dz \right)^{1/2}, \tag{3.23}$$

where  $C$  depends on  $n, \lambda, |\Omega|$  and  $r \in (0, r_0/6)$ .

Let  $\rho_0 = r_0/C_4$ , where  $C_4$  is the constant introduced in proposition 3.1. Let  $r \in (0, d_2]$  and define the point  $w = w(P_1) = P_1 - \tau\nu(P_1)$  where  $\nu(P_1)$  is the unit outward normal of  $\partial D_1$  at the point  $P_1$  and  $\tau = \lambda_{\bar{h}(r)} = a^{\bar{h}-1}\lambda_1, \bar{h} = \bar{h}(r)$  is defined in (3.11).

Set  $y = z = w$ , split the right-hand side of (3.21) into the sum of two integrals  $I_1(w)$  and  $I_2(w)$ :

$$S_{\mathcal{L}_0}(w, w) = I_1(w) + I_2(w),$$

where

$$\begin{aligned} I_1(w) &= \int_{B_{\rho_0}(P_1) \cap D_1} (\gamma_1^{(1)} - \gamma_1^{(2)})(\cdot) A(\cdot) \nabla G_1(\cdot, w) \cdot \nabla G_2(\cdot, w), \\ I_2(w) &= \int_{\Omega \setminus (B_{\rho_0}(P_1) \cap D_1)} (\sigma^{(1)} - \sigma^{(2)})(\cdot) \nabla G_1(\cdot, w) \cdot \nabla G_2(\cdot, w). \end{aligned}$$

The integral  $I_2(w)$  can be easily estimate using [11, proposition 3.1] as

$$|I_2(w)| \leq CE\rho_0^{2-n}, \tag{3.24}$$

Let us estimate  $I_1(w)$  from below in terms of  $\|\gamma_1^{(1)} - \gamma_1^{(2)}\|_{L^\infty(\Sigma_1 \cap B_{r_1}(P_1))}$ . Let  $\bar{x} \in \overline{\Sigma_1 \cap B_{r_1}(P_1)}$  be such that

$$(\gamma_1^{(1)} - \gamma_1^{(2)})(\bar{x}) = \|\gamma_1^{(1)} - \gamma_1^{(2)}\|_{L^\infty(\Sigma_1 \cap B_{r_1}(P_1))}.$$

Since  $(\gamma_1^{(1)} - \gamma_1^{(2)})(x) = \alpha_1 + \beta_1 \cdot x$ ,

$$\begin{aligned} I_1(w) &= \int_{B_{\rho_0}(P_1) \cap D_1} (\gamma_1^{(1)} - \gamma_1^{(2)})(\bar{x}) A(x) \nabla G_1(x, w) \cdot \nabla G_2(x, w) dx \\ &\quad + \int_{B_{\rho_0}(P_1) \cap D_1} \beta_1 \cdot (x - \bar{x}) A(x) \nabla G_1(x, w) \cdot \nabla G_2(x, w) dx, \end{aligned} \tag{3.25}$$

which leads to

$$\begin{aligned} |I_1(w)| &\geq \left| \int_{B_{\rho_0}(P_1) \cap D_1} (\gamma_1^{(1)} - \gamma_1^{(2)})(\bar{x}) A(x) \nabla G_1(x, w) \cdot \nabla G_2(x, w) \right| \\ &\quad - \bar{A} \int_{B_{\rho_0}(P_1) \cap D_1} |\beta_1 \cdot (x - \bar{x})| |\nabla G_1(x, w)| |\nabla G_2(x, w)| dx. \end{aligned} \tag{3.26}$$

If we set  $\tilde{c}^{(1)} = \frac{2}{1+\gamma_1^{(1)}(P_1)}$  and  $\tilde{c}^{(2)} = \frac{2}{1+\gamma_1^{(2)}(P_1)}$ , by adding and subtracting the fundamental solution  $\tilde{c}^{(i)}\Gamma$  we have

$$\begin{aligned}
|I_1(w)| \geq & \left| \int_{B_{\rho_0}(P_1) \cap D_1} (\gamma_1^{(1)} - \gamma_1^{(2)})(\bar{x}) A(x) \tilde{c}^{(1)} \tilde{c}^{(2)} |\nabla \Gamma(Jx, Jw)|^2 \right. \\
& - \int_{B_{\rho_0}(P_1) \cap D_1} |(\gamma_1^{(1)} - \gamma_1^{(2)})(\bar{x})| |A(x) \nabla(G_1(x, w) - \tilde{c}^{(1)}\Gamma(Jx, Jw)) \\
& \cdot \nabla(G_2(x, w) - \tilde{c}^{(2)}\Gamma(Jx, Jw))| dx - \bar{A} \int_{B_{\rho_0}(P_1) \cap D_1} |(\gamma_1^{(1)} - \gamma_1^{(2)})(\bar{x})| \\
& \times |\nabla(G_1(x, w) - \tilde{c}^{(1)}\Gamma(Jx, Jw))| \tilde{c}^{(2)} |\nabla \Gamma(Jx, Jw)| dx \\
& - \bar{A} \int_{B_{\rho_0}(P_1) \cap D_1} |(\gamma_1^{(1)} - \gamma_1^{(2)})(\bar{x})| \tilde{c}^{(1)} |\nabla \Gamma(Jx, Jw)| |\nabla(G_2(x, w) \\
& - \tilde{c}^{(2)}\Gamma(Jx, Jw))| dx - \int_{B_{\rho_0}(P_1) \cap D_1} |\beta_1 \cdot (x - \bar{x})| |A(x) \nabla \Gamma(Jx, Jw)| \\
& \cdot \nabla \Gamma(Jx, Jw)| dx. \tag{3.27}
\end{aligned}$$

Now, up to a change of coordinate we can suppose that  $P_1$  is the origin  $O$ . Let us apply the asymptotic estimate (3.4) to (3.27) for  $J = \sqrt{A^{-1}(0)}$ , it follows that

$$\begin{aligned}
|I_1(w)| \geq & \|\gamma_1^{(1)} - \gamma_1^{(2)}\|_{L^\infty(\Sigma_1 \cap B_{r_1})} C \lambda^{-1} \int_{B_{\rho_0} \cap D_1} |\nabla_x \Gamma(Jx, Jw)|^2 dx \\
& - C E \int_{B_{\rho_0} \cap D_1} |\nabla_x \Gamma(Jx, Jw)| |x - w|^{\theta_1 + 1 - n} dx \\
& - C E \int_{B_{\rho_0} \cap D_1} |x - w|^{2\theta_1 + 2 - 2n} dx \\
& - C E \int_{B_{\rho_0} \cap D_1} |x - \bar{x}| |x - w|^{2 - 2n} dx,
\end{aligned}$$

where the  $C > 0$  depends on the *a priori* data only. By definition (3.1), we can express explicitly the fundamental solution  $\Gamma$  inside the integrals and obtain:

$$\begin{aligned}
|I_1(w)| \geq & \|\gamma_1^{(1)} - \gamma_1^{(2)}\|_{L^\infty(\Sigma_1 \cap B_{r_1})} C \lambda^{-1} \int_{B_{\rho_0} \cap D_1} \frac{|J^2(x - w)|^2}{|J(x - w)|^n} dx \\
& - C E \int_{B_{\rho_0} \cap D_1} \frac{|J^2(x - w)|}{|J(x - w)|^n} |x - w|^{\theta_1 + 1 - n} dx \\
& - C E \int_{B_{\rho_0} \cap D_1} |x - w|^{2\theta_1 + 2 - 2n} dx \\
& - C \int_{B_{\rho_0} \cap D_1} |\beta_1| |x - \bar{x}| |x - w|^{2 - 2n} dx. \tag{3.28}
\end{aligned}$$



By estimating the integrals in (3.28) with respect to the parameter  $\tau$ , we can bound  $|I_1(w)|$  from below as follows:

$$|I_1(w)| \geq \|\gamma_1^{(1)} - \gamma_1^{(2)}\|_{L^\infty(\Sigma_1 \cap B_{r_1})} C \tau^{2-n} - C E \tau^{2-n+\theta_1} - C E \tau^{2-n+2\theta_1} - C E \tau^{3-n}. \quad (3.29)$$

By (3.23) and (3.24), it follows that

$$|I_1(w)| \leq |S_{\mathcal{L}_0}(w, w)| + |I_2(w)| \leq C \varepsilon_0 \tau^{-n} + C E \rho_0^{2-n},$$

which leads to the following estimate for the conductivity:

$$\begin{aligned} \|\gamma_1^{(1)} - \gamma_1^{(2)}\|_{L^\infty(\Sigma_1 \cap B_{r_1}(P_1))} \tau^{(2-n)} &\leq C \varepsilon_0 \tau^{-n} + C E \rho_0^{2-n} + C E \tau^{2-n+\theta_1} \\ &\quad + C E \tau^{2-n+2\theta_1} + C E \tau^{3-n}. \end{aligned}$$

Dividing by  $\tau^{2-n}$  both sides and by minimising with respect to  $\tau$  we obtain

$$\|\gamma_1^{(1)} - \gamma_1^{(2)}\|_{L^\infty(\Sigma_1 \cap B_{r_1}(P_1))} \leq C(\varepsilon_0 + E) \left( \frac{\varepsilon_0}{\varepsilon_0 + E} \right)^{\frac{\theta_1}{\theta_1+2}}, \quad (3.30)$$

where  $C$  is a positive constant which depends on the *a priori* data only.

Let us estimate  $|\partial_\nu(\gamma_1^{(1)} - \gamma_1^{(2)})(P_1)|$ . From (3.22), for  $y = z = w$  as above, we split again the  $n$ th partial derivative of the singular solution as follows:

$$\partial_{y_n} \partial_{z_n} S_{\mathcal{L}_0}(w, w) = \bar{I}_1(w) + \bar{I}_2(w), \quad (3.31)$$

where

$$\bar{I}_1(w) = \int_{B_{\rho_0}(P_1) \cap D_1} (\gamma_1^{(1)} - \gamma_1^{(2)})(\cdot) A(\cdot) \nabla \partial_{y_n} G_1(\cdot, w) \cdot \nabla \partial_{z_n} G_2(\cdot, w),$$

$$\bar{I}_2(w) = \int_{\Omega \setminus (B_{\rho_0}(P_1) \cap D_1)} (\sigma^{(1)} - \sigma^{(2)})(\cdot) \nabla \partial_{y_n} G_1(\cdot, w) \cdot \nabla \partial_{z_n} G_2(\cdot, w).$$

With a similar argument as in (3.24) one can determine an upper bound for  $\bar{I}_2$  of the form

$$|\bar{I}_2(w)| \leq C E \rho_0^{-n}, \quad (3.32)$$

where  $C$  depends on the *a priori* data. Notice that for any point  $x \in B_{\rho_0}(P_1) \cap D_1$ , the following equality holds

$$\begin{aligned} (\gamma_1^{(1)} - \gamma_1^{(2)})(x) &= (\gamma_1^{(1)} - \gamma_1^{(2)})(P_1) + (D_T(\gamma_1^{(1)} - \gamma_1^{(2)})(P_1)) \cdot (x - P_1)' \\ &\quad + (\partial_\nu(\gamma_1^{(1)} - \gamma_1^{(2)})(P_1))(x - P_1)_n. \end{aligned}$$

Proceeding as in (3.25) and (3.26),

$$\begin{aligned} |\bar{I}_1(w)| &\geq \left| \int_{B_{\rho_0}(P_1) \cap D_1} (\partial_\nu(\gamma_1^{(1)} - \gamma_1^{(2)})(P_1))(x - P_1)_n A(\cdot) \nabla \partial_{y_n} G_1(\cdot, w) \cdot \nabla \partial_{z_n} G_2(\cdot, w) \right| \\ &\quad - \int_{B_{\rho_0}(P_1) \cap D_1} |(D_T(\gamma_1^{(1)} - \gamma_1^{(2)})(P_1)) \cdot (x - P_1)'| |A(\cdot) \nabla \partial_{y_n} G_1(\cdot, w) \cdot \nabla \partial_{z_n} G_2(\cdot, w)| \\ &\quad - \int_{B_{\rho_0}(P_1) \cap D_1} |(\gamma_1^{(1)} - \gamma_1^{(2)})(P_1)| |A(\cdot) \nabla \partial_{y_n} G_1(\cdot, w) \cdot \nabla \partial_{z_n} G_2(\cdot, w)|. \end{aligned}$$

Up to a rigid transformation, we can assume that  $P_1$  coincides with the origin  $O$  of the coordinate system. Using a similar technique as in (3.27) and by proposition 3.1, this leads to

$$\begin{aligned} |\bar{I}_1(w)| &\geq |\partial_\nu(\gamma_1^{(1)} - \gamma_1^{(2)})(O)| C \int_{B_{\rho_0} \cap D_1} |\nabla_x \partial_{y_n} \Gamma(Jx, Jw)|^2 |x_n| \\ &\quad - C \left\{ E \int_{B_{\rho_0} \cap D_1} |\partial_{y_n} \nabla_x \Gamma(Jx, Jw)| |x - w|^{\theta_2 - n} |x_n| \right. \\ &\quad \left. + E \int_{B_{\rho_0} \cap D_1} |x - w|^{\theta_2 - 2n} |x_n| \right\} \\ &\quad - \int_{B_{\rho_0} \cap D_1} |D_T(\gamma_1^{(1)} - \gamma_1^{(2)})| |x'| |\nabla \partial_{y_n} G_1(\cdot, w)| |\nabla \partial_{z_n} G_2(\cdot, w)| \\ &\quad - \int_{B_{\rho_0} \cap D_1} |(\gamma_1^{(1)} - \gamma_1^{(2)})(O)| |\nabla \partial_{y_n} G_1(\cdot, w)| |\nabla \partial_{z_n} G_2(\cdot, w)|. \quad (3.33) \end{aligned}$$

By (3.30), we derive the following lower bound:

$$\begin{aligned} |\bar{I}_1(w)| &\geq |\partial_\nu(\gamma_1^{(1)} - \gamma_1^{(2)})(O)| C \int_{B_{\rho_0}(P_1) \cap D_1} |x - w|^{1-2n} \\ &\quad - C \left\{ E \int_{B_{\rho_0} \cap D_1} |x - w|^{1-2n+\theta_2} - \int_{B_{\rho_0} \cap D_1} |x - w|^{2-2n+\theta_2} \right. \\ &\quad \left. - \varepsilon_0 \int_{B_{\rho_0} \cap D_1} |x - w|^{1-2n} - \varepsilon_0 \int_{B_{\rho_0} \cap D_1} |x - w|^{-2n} \right\}, \end{aligned}$$

which leads to

$$|\partial_\nu(\gamma_1^{(1)} - \gamma_1^{(2)})(O)| \tau^{1-n} \leq |I_1(w)| + C (\varepsilon_0 \tau^{-n} + E \tau^{1-n+\theta_2}). \quad (3.34)$$

By the quantitative estimate (3.15),

$$\begin{aligned} |\bar{I}_1(w)| &\leq |\partial_{y_n} \partial_{z_n} S_{\mathcal{U}_0}(w, w)| + |I_2(w)| \\ &\leq C \varepsilon_0 \tau^{-(2+n)} + C E \rho_0^{-n}. \end{aligned} \quad (3.35)$$

Thus, by combining together (3.34) and (3.35), it follows that

$$|\partial_\nu(\gamma_1^{(1)} - \gamma_1^{(2)})(O)|\tau^{1-n} \leq C (\varepsilon_0\tau^{-(2+n)} + E\rho_0^{-n} + \varepsilon_0\tau^{-n} + E\tau^{1-n+\theta_2}),$$

which leads to

$$|\partial_\nu(\gamma_1^{(1)} - \gamma_1^{(2)})(O)| \leq C (\varepsilon_0\tau^{-3} + E\tau^{\theta_2}).$$

Finally, optimizing the right-hand side with respect to  $\tau$ , the estimate is given by the following inequality

$$|\partial_\nu(\gamma_1^{(1)} - \gamma_1^{(2)})(O)| \leq C\varepsilon_0^{\frac{\theta_2}{\theta_2+3}}(E + \varepsilon_0)^{\frac{3}{3+\theta_2}},$$

so that (3.20) is proved.

**3.2.2. Interior estimates.** We show that from the case  $k = 1$  we obtain the following estimate for the case  $k = 2$ :

$$\|\sigma_2^{(1)} - \sigma_2^{(2)}\|_{L^\infty(\Sigma_2 \cap B_{r_1}(P_2))} \leq C(\varepsilon_0 + E) \left( \omega_{1/C}^{(3)} \left( \frac{\varepsilon_0}{\varepsilon_0 + E} \right) \right)^{\frac{1}{C}}, \quad (3.36)$$

$$\left| \partial_\nu(\sigma_2^{(1)} - \sigma_2^{(2)})(P_2) \right| \leq C(\varepsilon_0 + E) \left( \omega_{1/C}^{(4)} \left( \frac{\varepsilon_0}{\varepsilon_0 + E} \right) \right)^{\frac{1}{C}}. \quad (3.37)$$

Since the proofs of (3.36) and (3.37) are similar, we prove (3.37), assuming that (3.36) holds.

$$\begin{aligned} & \int_{\Sigma} [\partial_{z_n} G_2(\cdot, z) \sigma^{(1)}(\cdot) \nabla \partial_{y_n} G_1(\cdot, y) \cdot \nu - \partial_{y_n} G_1(\cdot, y) \sigma^{(2)}(\cdot) \nabla \partial_{z_n} G_2(\cdot, z) \cdot \nu] \, dS \\ &= \partial_{y_n} \partial_{z_n} S_{\mathcal{U}_1}(y, z) + \int_{\mathcal{W}_1} (\sigma^{(1)} - \sigma^{(2)})(\cdot) \partial_{y_n} \nabla G_1(\cdot, y) \cdot \partial_{z_n} \nabla G_2(\cdot, z). \end{aligned} \quad (3.38)$$

Let  $\rho_0 = r_0/C_4$ , where  $C_4$  is the constant introduced in proposition 3.1. Pick  $r \in (0, r_0/6)$ . Fix the point  $w = w(P_2) = P_2 - \tau\nu(P_2)$  where  $\tau = a^{\tilde{h}-1}\lambda_1$ . We split the integral solution into two parts:

$$\partial_{y_n} \partial_{z_n} S_{\mathcal{U}_1}(w, w) = I_1(w) + I_2(w), \quad (3.39)$$

where

$$\begin{aligned} I_1(w) &= \int_{B_{\rho_0}(P_2) \cap D_2} (\gamma_2^{(1)} - \gamma_2^{(2)})(\cdot) A(\cdot) \partial_{y_n} \nabla G_1(\cdot, w) \cdot \partial_{z_n} \nabla G_2(\cdot, w), \\ I_2(w) &= \int_{\mathcal{U}_2 \setminus (B_{\rho_0}(P_2) \cap D_2)} (\sigma^{(1)} - \sigma^{(2)})(\cdot) \partial_{y_n} \nabla G_1(\cdot, w) \cdot \partial_{z_n} \nabla G_2(\cdot, w). \end{aligned}$$

As in the boundary estimates, we can bound from above  $I_2(w)$  as follows:

$$|I_2(w)| \leq CE\rho_0^{-n}. \quad (3.40)$$

Now, let us estimate from below the integral  $I_1(w)$  in terms of the quantity  $|\partial_\nu(\sigma_2^{(1)} - \sigma_2^{(2)})(P_2)|$ . First, notice that for any  $x \in B_{\rho_0}(P_2) \cap \Sigma_2$  we can rewrite  $\gamma_2^{(i)}$  as

$$\gamma_2^{(i)}(x) = \gamma_2^{(i)}(P_2) + D_T \gamma_2^{(i)}(P_2) \cdot (x - P_2)' + \partial_\nu(\gamma_2^{(i)}(P_2))(x - P_2)_n. \quad (3.41)$$

By (3.41),

$$\begin{aligned} |I_1(w)| &\geq \left| \int_{B_{\rho_0}(P_1) \cap D_2} (\partial_\nu(\gamma_2^{(1)} - \gamma_2^{(2)})(P_2))(x - P_2)_n A(x) \partial_{y_n} \nabla G_1(\cdot, w) \cdot \partial_{z_n} \nabla G_2(\cdot, w) \right. \\ &\quad - \int_{B_{\rho_0}(P_2) \cap D_2} |(D_T(\gamma_2^{(1)} - \gamma_2^{(2)})(P_2)) \cdot (x - P_2)'| |A(x) \partial_{y_n} \nabla G_1(\cdot, w) \cdot \partial_{z_n} \nabla G_2(\cdot, w)| \\ &\quad \left. - \int_{B_{\rho_0}(P_2) \cap D_2} |(\gamma_2^{(1)} - \gamma_2^{(2)})(P_2)| |A(x) \partial_{y_n} \nabla G_1(\cdot, w) \cdot \partial_{z_n} \nabla G_2(\cdot, w)| \right|. \end{aligned}$$

Up to a rigid transformation of coordinates, we can assume that  $P_2$  coincides with the origin  $O$  of the coordinate system. By proposition 3.1,

$$\begin{aligned} |I_1(w)| &\geq |\partial_\nu(\gamma_2^{(1)} - \gamma_2^{(2)})(O)| C \int_{B_{\rho_0} \cap D_2} |\partial_{y_n} \nabla_x \Gamma(Jx, Jw)|^2 |x_n| \\ &\quad - CE \int_{B_{\rho_0} \cap D_2} |\partial_{y_n} \nabla_x \Gamma(Jx, Jw)| |x - w|^{\theta_2 - n} |x_n| \\ &\quad - CE \int_{B_{\rho_0} \cap D_2} |x - w|^{2\theta_2 - 2n} |x_n| \\ &\quad - \int_{B_{\rho_0} \cap D_2} |D_T(\gamma_2^{(1)} - \gamma_2^{(2)})(O)| |x'| |A(x) \partial_{y_n} \nabla G_1(\cdot, w) \cdot \partial_{z_n} \nabla G_2(\cdot, w)| \\ &\quad - \int_{B_{\rho_0} \cap D_2} |(\gamma_2^{(1)} - \gamma_2^{(2)})(O)| |A(x) \partial_{y_n} \nabla G_1(\cdot, w) \cdot \partial_{z_n} \nabla G_2(\cdot, w)|. \quad (3.42) \end{aligned}$$

We can estimate the two last terms of the right-hand side by (3.36). Then

$$\begin{aligned} |I_1(w)| &\geq |\partial_\nu(\gamma_2^{(1)} - \gamma_2^{(2)})(O)| C \int_{B_{\rho_0} \cap D_2} |x - w|^{1-2n} \\ &\quad - CE \int_{B_{\rho_0} \cap D_2} |x - w|^{\theta_2 + 1 - 2n} \\ &\quad - CE \int_{B_{\rho_0} \cap D_2} |x - w|^{2\theta_2 + 1 - 2n} \\ &\quad - (\varepsilon_0 + E) \left( \omega_{1/C}^{(3)} \left( \frac{\varepsilon_0}{\varepsilon_0 + E} \right) \right)^{1/C} \int_{B_{\rho_0} \cap D_2} |x - w|^{1-2n} \\ &\quad - (\varepsilon_0 + E) \left( \omega_{1/C}^{(3)} \left( \frac{\varepsilon_0}{\varepsilon_0 + E} \right) \right)^{1/C} \int_{B_{\rho_0} \cap D_2} |x - w|^{-2n}, \end{aligned}$$

where the constant  $C > 0$  depends on the *a priori* data and on  $J$ . This leads to

$$\left| \partial_\nu(\gamma_2^{(1)} - \gamma_2^{(2)})(O) \right| \bar{r}^{(1-n)} \leq |I_1(w)| + C \left\{ (\varepsilon_0 + E) \left( \omega_{1/C}^{(3)} \left( \frac{\varepsilon_0}{\varepsilon_0 + E} \right) \right)^{1/C} \tau^{-n} + E \frac{\tau^{1-n+\theta_2}}{\rho_0^{\theta_2}} \right\}.$$

Secondly, by (3.39) and (3.40),

$$|I_1(w)| \leq |\partial_{y_n} \partial_{z_n} S_{\mathcal{U}_1}(w, w)| + CE\rho_0^{-n}.$$

Combining the last two inequalities, it follows that

$$\begin{aligned} \left| \partial_\nu(\gamma_2^{(1)} - \gamma_2^{(2)}) \right| \tau^{(1-n)} &\leq |\partial_{y_n} \partial_{z_n} S_{\mathcal{U}_1}(w, w)| + C \left\{ E\rho_0^{-n} \right. \\ &\quad \left. + (\varepsilon_0 + E) \left( \omega_{1/C}^{(3)} \left( \frac{\varepsilon_0}{\varepsilon_0 + E} \right) \right)^{1/C} \tau^{-n} + E \frac{\tau^{1-n+\theta_2}}{\rho_0^{\theta_2}} \right\}. \end{aligned}$$

By quantitative estimates of propagation of smallness (proposition 3.2), we can estimate the integral solution as

$$|\partial_{y_j} \partial_{z_n} S_{\mathcal{U}_1}(w, w)| \leq r_0^{-n} C^{\bar{h}} (\varepsilon_0 + \delta_1 + E) \left( \omega_{1/C}^{(2)} \left( \frac{\varepsilon_0 + \delta_1}{E + \delta_1 + \varepsilon_0} \right) \right)^{(1/C)^{\bar{h}}},$$

so that

$$\begin{aligned} \left| \partial_\nu(\gamma_2^{(1)} - \gamma_2^{(2)})(O) \right| &\leq C^{\bar{h}} (\varepsilon_0 + \delta_1 + E) \left( \omega_{1/C}^{(2)} \left( \frac{\varepsilon_0 + \delta_1}{E + \delta_1 + \varepsilon_0} \right) \right)^{(1/C)^{\bar{h}}} \tau^{(n-1)} \\ &\quad + C\tau^{(-1)} (\varepsilon_0 + E) \left( \omega_{1/C}^{(3)} \left( \frac{\varepsilon_0}{\varepsilon_0 + E} \right) \right)^{1/C} + CE \frac{\tau^{\theta_2}}{\rho_0^{\theta_2}}. \end{aligned} \quad (3.43)$$

Since  $\bar{h}$  is a function of  $r$ , we have to estimate  $C^{\bar{h}}$  and  $(\frac{1}{C})^{\bar{h}}$  in terms of  $r$ . Recalling (3.12), it turns out that

$$\left( \frac{d_1}{r} \right)^{C_1} \leq C^{\bar{h}} \leq C_2 \left( \frac{d_1}{r} \right)^{C_1}.$$

Since  $\tau \leq \lambda_1 \cdot \frac{r}{d_1}$ ,

$$\begin{aligned} |\partial_\nu(\gamma_2^{(1)} - \gamma_2^{(2)})(O)| &\leq C(\varepsilon_0 + E) \left\{ \left( \frac{r}{d_1} \right)^{n-1-C} \left( \omega_{1/C}^{(2)} \left( \frac{\varepsilon_0 + \delta_1}{E + \delta_1 + \varepsilon_0} \right) \right)^{\left( \frac{r}{d_1} \right)^C} \right. \\ &\quad \left. + \left( \frac{r}{d_1} \right)^{-1} \left( \omega_{1/C}^{(3)} \left( \frac{\varepsilon_0}{\varepsilon_0 + E} \right) \right)^{1/C} + \left( \frac{r}{d_1} \right)^{\theta_2} \right\}. \end{aligned} \quad (3.44)$$

One can show that the following inequality holds:

$$\frac{\varepsilon_0 + \delta_1}{E + \delta_1 + \varepsilon_0} \leq C\omega_{1/C}^{(0)} \left( \frac{\varepsilon_0}{\varepsilon_0 + E} \right). \quad (3.45)$$

Then, combining (3.45) together with (3.44),

$$\left| \partial_\nu(\gamma_2^{(1)} - \gamma_2^{(2)})(O) \right| \leq C(\varepsilon_0 + E) \left\{ \left( \frac{r}{d_1} \right)^{n-1-C} \left( \omega_{1/C}^{(3)} \left( \frac{\varepsilon_0}{E + \varepsilon_0} \right) \right)^{\left( \frac{r}{d_1} \right)^C} + \left( \frac{r}{d_1} \right)^{\theta_2} \right\}.$$

Finally, optimizing with respect to  $r$ , (3.37) follows.

Proceeding as above, for  $k = 3, \dots, K$ , one can show that the following inequalities hold:

$$\|\gamma_k^{(1)} - \gamma_k^{(2)}\|_{L^\infty(\Sigma_k \cap B_{r_1}(P_k))} \leq C(\varepsilon_0 + E) \left( \omega_{1/C}^{(2k-1)} \left( \frac{\varepsilon_0}{\varepsilon_0 + E} \right) \right)^{\frac{1}{C}}, \quad (3.46)$$

$$\left| \partial_\nu(\gamma_k^{(1)} - \gamma_k^{(2)})(P_k) \right| \leq C(\varepsilon_0 + E) \left( \omega_{1/C}^{(2k)} \left( \frac{\varepsilon_0}{\varepsilon_0 + E} \right) \right)^{\frac{1}{C}}. \quad (3.47)$$

By reformulating (3.21) and (3.22) as

$$\begin{aligned} & \int_{\Sigma} [G_2(\cdot, z)\sigma^{(1)}(\cdot)\nabla G_1(\cdot, y) \cdot \nu - G_1(\cdot, y)\sigma^{(2)}(\cdot)\nabla G_2(\cdot, z) \cdot \nu] \, dS \\ &= S_{\mathcal{U}_{k-1}}(y, z) + \int_{\mathcal{W}_{k-1}} (\sigma^{(1)} - \sigma^{(2)})(\cdot)\nabla G_1(\cdot, y) \cdot \nabla G_2(\cdot, z) \end{aligned} \quad (3.48)$$

and

$$\begin{aligned} & \int_{\Sigma} [\partial_{z_n} G_2(\cdot, z) \sigma^{(1)}(\cdot)\nabla \partial_{y_n} G_1(\cdot, y) \cdot \nu - \partial_{y_n} G_1(\cdot, y)\sigma^{(2)}(\cdot)\nabla \partial_{z_n} G_2(\cdot, z) \cdot \nu] \, dS \\ &= \partial_{y_n} \partial_{z_n} S_{\mathcal{U}_{k-1}}(y, z) + \int_{\mathcal{W}_{k-1}} (\sigma^{(1)} - \sigma^{(2)})(\cdot)\nabla \partial_{y_n} G_1(\cdot, y) \cdot \nabla \partial_{z_n} G_2(\cdot, z), \end{aligned} \quad (3.49)$$

respectively, the procedure is similar to the one seen above. We just point out that, for  $(y, z) \in \mathcal{W}_k \times \mathcal{W}_k$ ,

$$|S_{\mathcal{U}_{k-1}}(y, z)| \leq Cr_0^{2-n}(\varepsilon_0 + \delta_{k-1}),$$

then we can bound from above the integral solution by unique continuation (3.14) and (3.15).

Notice that

$$\delta_k \leq \delta_{k-1} + \|\gamma_k^{(1)} - \gamma_k^{(2)}\|_{L^\infty(D_k)}.$$

From the property (3.5) it follows that

$$\omega_{1/C}^{(2k)}(1) \leq \frac{\varepsilon_0 + \delta_{k-1} + E}{\varepsilon_0 + \delta_{k-1}} \omega_{1/C}^{(2k)} \left( \frac{\varepsilon_0 + \delta_{k-1}}{\varepsilon_0 + \delta_{k-1} + E} \right)$$

and

$$\delta_{k-1} + \varepsilon_0 \leq (\omega_{1/C}^{(2k)}(1))^{-1}(\varepsilon_0 + \delta_{k-1} + E) \left( \omega_{1/C}^{(2k)} \left( \frac{\varepsilon_0 + \delta_{k-1}}{\varepsilon_0 + \delta_{k-1} + E} \right) \right).$$

By the estimates (3.46) and (3.47) it follows that

$$\delta_k + \varepsilon_0 \leq C(\varepsilon_0 + E) \left( \omega_{1/C}^{(2k)} \left( \frac{\varepsilon_0}{\varepsilon_0 + E} \right) \right)^{1/C}.$$

This leads to the following estimate for  $E = \delta_K$

$$E \leq C(\varepsilon_0 + E) \left( \omega_{1/C}^{(2K)} \left( \frac{\varepsilon_0}{\varepsilon_0 + E} \right) \right)^{\frac{1}{C}}.$$

Since the function  $\omega_{1/C}$  is invertible for  $\varepsilon_0 < e^{-2}E$  (otherwise the statement is proven), it follows that

$$E \leq \frac{C - \left( \omega_{1/C}^{(2K)} \left( \frac{1}{C} \right) \right)^{-1}}{\left( \omega_{1/C}^{(2K)} \left( \frac{1}{C} \right) \right)^{-1}} \varepsilon_0.$$

Hence, (3.50) is proven.  $\square$

**Proof of corollary 2.2.** Assume that the hypothesis of theorem 2.1 hold, then there exists a constant  $C > 1$  such that

$$\|\sigma^{(1)} - \sigma^{(2)}\|_{L^\infty(\Omega)} \leq C(\mathcal{J}(\sigma^{(1)}, \sigma^{(2)}))^{1/2}.$$

First, by the Alessandrini's identity,

$$S_{\mathcal{U}_0}(y, z) = \langle (\Lambda_1 - \Lambda_2)G_1(\cdot, y), G_2(\cdot, z) \rangle,$$

where  $G_1(\cdot, y), G_2(\cdot, z) \in H_{00}^{1/2}(\Sigma)$  for  $y, z \in D_0$  since they are weak solutions to the problem (2.10). Then, it follows that

$$|S_{\mathcal{U}_0}(y, z)| \leq C\|\Lambda_1 - \Lambda_2\|_*,$$

where

$$\|\Lambda_1 - \Lambda_2\|_* = \sup_{f, g \in H_{00}^{1/2}(\Sigma), \|g\| = \|\varphi\| = 1} |\langle (\Lambda_1 - \Lambda_2)g, \varphi \rangle|.$$

Then

$$(\mathcal{J}(\sigma^{(1)}, \sigma^{(2)}))^{1/2} \leq C\|\Lambda_1 - \Lambda_2\|_*, \quad (3.50)$$

where  $C > 0$  depends on the *a priori* data only. Then the inequality (2.16) trivially follows.  $\square$

#### 4. Proof of technical propositions

In this section we give the proof of the propositions needed for the proof of the main result (theorem 2.1).

4.1. Asymptotic estimates

Let  $0 < \mu < 1$  and  $B^+ \in C^\mu(Q_r^+)$ ,  $B^- \in C^\mu(Q_r^-)$  be symmetric, positive definite, matrix valued functions and define

$$B(x) = \begin{cases} B^+(x), & x \in Q_r^+, \\ B^-(x), & x \in Q_r^-, \end{cases}$$

such that  $B$  satisfies the uniform ellipticity condition

$$\lambda_0^{-1} |\xi|^2 \leq B(x)\xi \cdot \xi \leq \lambda_0 |\xi|^2, \quad \text{for a.e. } x \in Q_r, \text{ for every } \xi \in \mathbb{R}^n,$$

where  $\lambda_0 > 0$  is a constant. Let  $\bar{b} > 0$  and define

$$b(x) = \begin{cases} b^+ + B^+ \cdot x, & x \in Q_r^+, \\ b^- + B^- \cdot x, & x \in Q_r^-, \end{cases}$$

where  $b^+, b^- \in \mathbb{R}$ ,  $B^+, B^- \in \mathbb{R}^n$  and  $0 < \bar{b}^{-1} \leq b(x) \leq \bar{b}$ .

**Theorem 4.1.** *Let  $r > 0$  be a fixed number. Let  $b(x)$  and  $B(x)$  be as above. Let  $U \in H^1(Q_r)$  be a solution to*

$$\operatorname{div}(b(x)B(x)\nabla U) = 0, \quad \text{in } Q_r.$$

*Then, there exist positive constants  $0 < \alpha' \leq 1, C > 0$  depending on  $\bar{b}, r, \lambda_0$  and  $n$  only, such that for any  $\rho \leq \frac{r}{2}$  and for any  $x \in Q_{r-2\rho}$ , the following estimate holds*

$$\|\nabla U\|_{L^\infty(Q_\rho(x))} + \rho^{\alpha'} |\nabla U|_{\alpha', Q_\rho(x) \cap Q_r^+} + \rho^{\alpha'} |\nabla U|_{\alpha', Q_\rho(x) \cap Q_r^-} \leq \frac{C}{\rho^{1+n/2}} \|U\|_{L^2(Q_{2\rho}(x))}. \quad (4.1)$$

**Proof.** For the proof we refer to Li–Vogelius [52], where piecewise  $C^{1,\alpha'}$  estimates for solutions to elliptic equations in divergence form with piecewise Hölder continuous coefficients have been demonstrated. □

**Proof of theorem 3.1.** Let us consider a conductivity  $\sigma$  of the form

$$\sigma(x) = \sum_{k=1}^N \gamma_k(x) \chi_{D_k}(x) A(x).$$

First, fix  $k = 1, \dots, K$ . Up to a rigid transformation, we the point  $P_{k+1}$  can be identified with the origin and  $\gamma_k(0) = \gamma^-$  and  $\gamma_{k+1}(0) = \gamma^+$  for  $k \in \mathbb{R}$ . For any  $x = (x', x_n)$ , denote  $x^* = (x', -x_n)$ .

Let us introduce a linear change of coordinates

$$\begin{aligned} L : \mathbb{R}^n &\rightarrow \mathbb{R}^n \\ \xi &\mapsto L\xi := RJ\xi, \end{aligned}$$



where  $J = \sqrt{A^{-1}(0)}$  and the matrix  $R$  is orthogonal and represents the planar rotation in  $\mathbb{R}^n$  that rotates the unit vector  $\frac{v}{\|v\|}$ , where  $v = \sqrt{A(0)}e_n$  to the  $n$ th standard unit vector  $e_n$  and such that

$$R|_{(\pi)^\perp} = Id|_{(\pi)^\perp},$$

where  $\pi$  is the plane generated by  $e_n$  and  $v$  and  $(\pi)^\perp$  is the orthogonal complement of  $\pi$  (see [37]). Moreover, the following relations hold

- $A(0) = L^{-1} \cdot (L^{-1})^T$ ,
- $(L\xi) \cdot e_n = \frac{1}{\|v\|} \xi \cdot e_n$ ,
- $\sigma_{A(0)}(\xi) = L^{-1} \sigma_I(L\xi)(L^{-1})^T$ , where  $\sigma_I(L\xi) = \sigma_I(x) = (\gamma^- + (\gamma^+ - \gamma^-)\chi^+(x))I$ .

A fundamental solution of the operator  $\operatorname{div}_\xi((\gamma^- + (\gamma^+ - \gamma^-)\chi^+(\cdot))A(0)\nabla_\xi \cdot)$  has the following explicit form

$$H_{A(0)}(\xi, \eta) = \begin{cases} |J| \left( \frac{1}{\gamma^+} \Gamma(L\xi, L\eta) + \frac{\gamma^+ - \gamma^-}{\gamma^+(\gamma^+ + \gamma^-)} \Gamma(L\xi, L^*\eta) \right), & \text{if } \xi_n, \eta_n > 0, \\ |J| \left( \frac{2}{\gamma^+ + \gamma^-} \Gamma(L\xi, L\eta) \right), & \text{if } \xi_n \eta_n < 0, \\ |J| \left( \frac{1}{\gamma^-} \Gamma(L\xi, L\eta) + \frac{\gamma^- - \gamma^+}{\gamma^-(\gamma^+ + \gamma^-)} \Gamma(L\xi, L^*\eta) \right), & \text{if } \xi_n, \eta_n < 0, \end{cases} \quad (4.2)$$

where  $|J|$  denotes the determinant of the matrix  $J$  and  $L^*$  is the matrix whose coefficients follow the rule

$$l_{ij}^* = l_{ij}, \quad \text{for } i = 1, \dots, n-1, j = 1, \dots, n, \quad l_{nj}^* = -l_{nj} \quad \text{for } j = 1, \dots, n.$$

Set  $H(\xi, \eta) = H_{A(0)}(\xi, \eta)$ . Denote with  $\tilde{\Omega}$  the augmented domain obtained after having performed the change of coordinates  $L$ . Define the distribution

$$R(\xi, \eta) = G(\xi, \eta) - H(\xi, \eta), \quad (4.3)$$

where  $G(\cdot, \eta)$  is the weak solution to (2.10), then  $R(\xi, \eta)$  is a weak solution to the following boundary value problem

$$\begin{cases} \operatorname{div}_\xi (\sigma(\cdot) \nabla R(\cdot, \eta)) = -\operatorname{div}_\xi ((\sigma(\cdot) - \sigma_0(\cdot)) \nabla_\xi H(\cdot, \eta)), & \text{in } \tilde{\Omega}, \\ R(\cdot, \eta) = -H(\cdot, \eta), & \text{on } \partial\tilde{\Omega}, \end{cases}$$

where  $\sigma_0(\cdot) = (\gamma^- + (\gamma^+ - \gamma^-)\chi^+(\cdot))A(0)$ . By the representation formula over  $\tilde{\Omega}$ , it follows that  $R$  satisfies the following integral identity

$$R(\xi, \eta) = -\int_{\tilde{\Omega}} (\sigma(\zeta) - \sigma_0(\zeta)) \nabla_\zeta H(\zeta, \eta) \cdot \nabla_\zeta G(\zeta, \xi) d\zeta + \int_{\partial\tilde{\Omega}} \sigma(\zeta) \nabla G(\zeta, \xi) \cdot \nu H(\zeta, \eta) dS(\zeta). \quad (4.4)$$

The integral over  $\partial\tilde{\Omega}$  at the right-hand side of (4.4) can be easily bounded from above as in [7, equation (4.10)] by a constant  $C >$  which depends on the *a priori* data only.

Set  $\gamma_0(\cdot) = \gamma^- + (\gamma^+ - \gamma^-)\chi^+(\cdot)$ . Locally, in a neighbourhood of the origin, the following estimate holds

$$\begin{aligned} |\sigma(\zeta) - \sigma_0(\zeta)| &\leq |\gamma(\zeta)A(\zeta) - \gamma_0(\zeta)A(0)| \leq |\gamma(\zeta)| |A(\zeta) - A(0)| \\ &\quad + |\gamma(\zeta) - \gamma_0(\zeta)| |A(0)| \leq C |\zeta|, \end{aligned} \quad (4.5)$$

where  $C > 0$  depends on  $\bar{\gamma}, \bar{A}$  only. Moreover by (2.11) we find the following two pointwise bounds:

$$\begin{aligned} |\nabla_\zeta G(\zeta, \xi)| &\leq C |\zeta - \xi|^{1-n} \quad \text{for every } \zeta, \xi \in Q_{r_0}, \\ |\nabla_\zeta H(\zeta, \eta)| &\leq C |\zeta - \eta|^{1-n} \quad \text{for every } \zeta, \eta \in Q_{r_0}, \end{aligned}$$

which together with (4.5) leads to

$$\left| \int_{\Omega} (\sigma(\zeta) - \sigma_0(\zeta)) \nabla_\zeta H(\zeta, \eta) \cdot \nabla_\zeta G(\zeta, \xi) d\zeta \right| \leq C_1 |\xi - \eta|^{3-n-\alpha}, \quad (4.6)$$

for any  $0 < \alpha < 1$ . In conclusion, for  $\xi \in B_{r_0}^+$ ,  $\eta = \eta_n e_n$  with  $\eta_n \in (-r_0, 0)$ ,

$$|R(\xi, \eta)| \leq C |\xi - \eta|^{3-n-\alpha}. \quad (4.7)$$

We focus on the estimate for  $\nabla_\xi R(\xi, e_n \eta_n)$ . Fix  $\xi \in B_{r_0/4}^+$  and  $\eta_n \in (-r_0/4, 0)$ , consider the cylinder  $Q = B'_{h/4}(\xi') \times (\xi_n, \xi_n + \frac{h}{4})$  where  $h = |\xi - \eta|$ . Notice that  $Q \subset Q_{\frac{r_0}{2}}^+$ ,  $Q \subset Q_{\frac{h}{2}}(\xi)$  and  $\xi \in \partial Q$ .

By theorem 4.1 it follows that

$$|\nabla_\xi G(\cdot, e_n \eta_n)|_{\alpha', Q}, \quad |\nabla_\xi H(\cdot, e_n \eta_n)|_{\alpha', Q} \leq Ch^{-\alpha'+1-n}. \quad (4.8)$$

Hence by (4.3) and (4.8) we

$$|\nabla_\xi R(\cdot, e_n \eta_n)|_{\alpha', Q} \leq Ch^{-\alpha'+1-n}. \quad (4.9)$$

From the following interpolation inequality

$$\|\nabla_\xi R(\cdot, e_n \eta_n)\|_{L^\infty(Q)} \leq C \left( \|R(\cdot, e_n \eta_n)\|_{L^\infty(Q)}^{\alpha'/1+\alpha'} |\nabla_\xi R(\cdot, e_n \eta_n)|_{\alpha', Q}^{1/1+\alpha'} + \frac{1}{h} \|R(\cdot, \eta_n e_n)\|_{L^\infty(Q)} \right),$$

together with (4.7) we obtain

$$|\nabla_\xi R(\cdot, \eta_n e_n)| \leq Ch^{\theta_1+1-n},$$

where  $\theta_1 = \frac{\alpha'(1-\alpha)}{1+\alpha}$ .

Now, we look for a pointwise bound for  $\nabla_\eta \nabla_\xi R(\xi, \eta)$ . Define the cylinder  $\hat{Q} = B'_{\frac{h}{8}}(0) \times (\eta_n - \frac{h}{8}, \eta_n)$ . As before, we have that  $\hat{Q} \subset Q_{\frac{r_0}{4}}^-$ ,  $\hat{Q} \subset Q_{\frac{h}{4}}(\eta)$  and  $\xi \notin Q_{\frac{h}{4}}(\eta)$ .

Let  $k$  be an integer such that  $k \in \{1, \dots, n\}$ . Notice that  $\partial_{\xi_k} \Gamma(\xi, \cdot)$  is a weak solution to the Laplace equation

$$\Delta_\eta (\partial_{\xi_k} \Gamma(\xi, \cdot)) = 0 \quad \text{in } Q_{\frac{h}{4}}(\eta),$$

and  $\partial_{\xi_k} G(\xi, \cdot)$  is a weak solutions to the problem

$$\begin{cases} \operatorname{div}(\sigma(\cdot)\nabla\partial_{\xi_k}G_i(\xi, \cdot)) = -\delta(\xi - \cdot) & \text{in } Q_{\frac{h}{4}}(\eta), \\ G_i(\xi, \cdot) = 0 & \text{on } \partial\Omega. \end{cases}$$

By theorem 4.1, it follows that

$$|\nabla_{\eta}\partial_{\xi_k}G(\xi, \cdot)|_{\alpha', \hat{Q}} \leq Ch^{-\alpha'-1-\frac{n}{2}}\|\partial_{\xi_k}G(\xi, \cdot)\|_{L^2(Q_{\frac{h}{4}}(\eta))}. \quad (4.10)$$

Fix  $\bar{\eta} \in Q_{\frac{h}{4}}(\eta)$ , then  $\bar{\eta} \notin Q_{\frac{h}{16}}(\xi)$ . By theorem 4.1, it follows that

$$\|\nabla_{\xi}G(\cdot, \bar{\eta})\|_{L^{\infty}(Q_{\frac{h}{32}}(\xi))} \leq Ch^{-1-\frac{n}{2}}\|G(\cdot, \bar{\eta})\|_{L^{\infty}(Q_{\frac{h}{16}}(\xi))} \leq Ch^{1-n}. \quad (4.11)$$

From (4.10) and (4.11) it follows that

$$|\nabla_{\eta}\partial_{\xi_k}G(\xi, \cdot)|_{\alpha', \hat{Q}} \leq Ch^{-\alpha'-n}. \quad (4.12)$$

By the representation formula for  $\Gamma$ ,

$$|\nabla_{\eta}\partial_{\xi_k}\Gamma(\xi, \cdot)|_{\alpha', \hat{Q}} \leq Ch^{-\alpha'-n}, \quad (4.13)$$

and by (4.12) and (4.13),

$$|\nabla_{\eta}\partial_{\xi_k}R(\xi, \cdot)|_{\alpha', \hat{Q}} \leq Ch^{-\alpha'-n}. \quad (4.14)$$

Arguing as above, the following estimate holds:

$$\|\partial_{\xi_k}R(\xi, \cdot)\|_{L^{\infty}(\hat{Q})} \leq Ch^{\theta_1+1-n}. \quad (4.15)$$

By the following interpolation inequality

$$\|\nabla_{\eta}\partial_{\xi_k}R(\xi, \cdot)\|_{L^{\infty}(\hat{Q})} \leq C\|\partial_{\xi_k}R(\xi, \cdot)\|_{L^{\infty}(\hat{Q})}^{\frac{\alpha'}{\alpha'+1}}|\nabla_{\eta}\partial_{\xi_k}R(\xi, \cdot)|_{\alpha', \hat{Q}}^{\frac{1}{\alpha'+1}} \quad (4.16)$$

and by (4.15) and (4.14), we conclude that

$$|\nabla_{\eta}\partial_{\xi_k}R(\xi, \eta)| \leq Ch^{\theta_2-n}, \quad (4.17)$$

where  $\theta_2 = \frac{\theta_1\alpha'}{1+\alpha'}$ . □

#### 4.2. Propagation of smallness

In order to prove proposition 3.2, we state and prove a preliminary proposition 4.2, where we determine a pointwise bound for the weak solution to the conductivity equation in the interior of  $\tilde{\Omega}$ .

**Proposition 4.2.** *Let  $v \in H^1(\tilde{\Omega})$  be a weak solution to*

$$\operatorname{div}(\sigma \nabla v) = 0 \quad \text{in } \mathcal{W}_k, \quad (4.18)$$

where  $k \in \{0, \dots, K-1\}$ . Suppose there exist  $E, \epsilon > 0$  such that

$$|v(x)| \leq r_0^{2-n} \epsilon \quad \forall x \in D_0, \quad (4.19)$$

$$|v(x)| \leq E(r_0 d(x))^{1-(n/2)} \quad \forall x \in \mathcal{W}_k. \quad (4.20)$$

Then, for every  $r \in (0, d_1]$ ,

$$|v(w_{\bar{h}}(P_{k+1}))| \leq r_0^{2-n} C^{\bar{h}} (E + \epsilon) \left( \omega_{1/C}^{(K)} \left( \frac{\epsilon}{\epsilon + E} \right) \right)^{(1/C)^{\bar{h}}}, \quad (4.21)$$

where  $C > 1$  depends only on a priori data.

**Proof of proposition 4.2.** We adapt the proof in [11, proposition 4.4] to the case of the anisotropic conductivity.

To begin with, we introduce some parameters. Recall from (3.7) that  $\mathcal{W}_k = \bigcup_{m=0}^k D_m$ , then for the domain index  $m \in \{0, \dots, K-1\}$ ,

$$r_l = \frac{r_0}{l}, \quad \bar{\rho} = \frac{r_l}{32l\sqrt{1+L^2}}, \quad (4.22)$$

$$y_{m+1} = P_{m+1} - \frac{r_l}{32} \nu(P_{m+1}), \quad \tilde{y}_{m+1} = P_{m+1} + \frac{r_l}{32} \nu(P_{m+1}), \quad (4.23)$$

$$v_m = v|_{D_m}, \quad (4.24)$$

where  $P_{m+1}$  and  $\nu(P_{m+1})$  have been defined in subsection 3.1.1. We claim that for every  $m \in \{0, \dots, K-1\}$ ,

$$\|v\|_{L^\infty(B_{\bar{\rho}}(\tilde{y}_{m+1}))} \leq r_0^{2-n} C^{m+1} (E + \epsilon) \left( \omega_{1/C}^{(m+1)} \left( \frac{\epsilon}{\epsilon + E} \right) \right) \quad (4.25)$$

and prove (4.25) by induction as follows.

Case  $m = 0$ .

Up to a rigid transformation of coordinate, we can suppose that  $y_1 = -\frac{r_l}{32} e_n$ . From (4.20),

$$\|v\|_{L^\infty(D_0)} \leq E \left( r \sup_{x \in D_0} d(x) \right)^{1-n/2}. \quad (4.26)$$

Choose an arbitrary point  $\bar{y} \in \Sigma_1$ , possibly different from  $P_1$ . Let  $\phi$  be a Jordan curve joining  $y_1$  to  $w_1(\bar{y})$  such that  $\phi \subset (D_0)_{\bar{d}}$ , where  $\bar{d} = \min\{\text{dist}(y_1, \Sigma_1), \text{dist}(w_1(\bar{y}), \Sigma_1)\}$ , and  $(D_0)_{\bar{d}}$  is connected. Notice that  $w_1(\bar{y}) \in (D_0)_{\bar{d}}$ . Let us define a set of points  $\{\phi_i\}$ ,  $i = 1, \dots, s$  through the following process:

- $\phi_1 = \phi(0) = y_1$ ;
- For  $i > 1$ , set

$$\phi_{i+1} = \begin{cases} \phi(t_i), & \text{if } |\phi_i - w_1(\bar{y})| > 2r_l \text{ where } t_i = \max\{t_i : |\phi(t) - \phi_i| = 2r_l\}, \\ w_1(\bar{y}), & \text{if } |\phi_i - w_1(\bar{y})| < 2r_l \text{ and set } s = i + 1. \end{cases}$$

Apply the three sphere inequality in the case of pure principal part (see [10, theorem 2.1]) on spheres centred at  $\phi_1 = y_1$  for which estimates (4.26) and (4.19) hold, with suitable rays  $r, 3r, 4r$ :

$$\|v\|_{L^2(B_{3r}(y_1))} \leq Q \|v\|_{L^2(B_r(y_1))}^\delta \|v\|_{L^2(B_{4r}(y_1))}^{1-\delta} \leq Q r_0^{2-n} \epsilon^\delta E^{1-\delta},$$

where  $\delta = \frac{\log(\frac{4\lambda}{3})}{\log(\frac{4\lambda}{3}) + C \log(\frac{3}{\lambda})}$  and  $Q > 1$  is a constant which depends on  $\lambda, L, \max\{\frac{4r}{r_0}, 1\}$ .

Notice that  $B_r(\phi_2) \subset B_{3r}(\phi_1) = B_{3r}(y_1)$  so that the  $L^2$ -norm of  $v$  on  $B_r(\phi_2)$  can be easily estimated applying the three sphere inequality for the spheres of rays  $r, 3r, 4r$  centred at  $\phi_2$ . Moreover, by [38, theorem 8.17], since  $v$  is a weak solution to (4.18), it follows that

$$\|v\|_{L^\infty(B_{R/2}(y))} \leq C \rho^{n/2} \|v\|_{L^2(B_R(y))},$$

where  $C$  depends on  $n, \lambda$  and  $|\Omega|$ . By iterating this process, we can estimate the  $L^\infty$ -norm of  $v$  along the chain of spheres centred at points  $\phi_i$  of the curve  $\phi$ . In conclusion,

$$\|v\|_{L^\infty(B_r(w_1(\bar{y})))} \leq \|v\|_{L^\infty(B_{3r}(\phi_{s-1}))} \leq C r^{2-n} \epsilon^{\delta s} E^{1-\delta s}. \quad (4.27)$$

Fix  $r \in (0, d_1]$ . Recalling the parameters introduced in (3.9), the following inclusions hold:

$$B_{\rho_{k+1}}(w_{k+1}(\bar{y})) \subset B_{3\rho_k}(w_k(\bar{y})) \subset B_{4\rho_k}(w_k(\bar{y})) \subset C(\bar{y}, \nu(\bar{y}), \beta_1, r_0/3),$$

for any  $k = 1, 2, \dots$ . Notice that  $\rho_1 < r_1$  for a suitable  $l$ , then  $B_{\rho_1}(w_1(\bar{y})) \subset B_{r_1}(w_1(\bar{y}))$ . We proceed by moving from one centre to the successive one along the axis of the cone  $C(\bar{y}, \nu(\bar{y}), \beta_1, r_0/3)$  allowing to get closer and closer to the vertex  $\bar{y}$  and stop this process when we reach the sphere of radius  $\rho_{\bar{h}}$ . Then, from (4.27),

$$\|v\|_{L^\infty(B_{\rho_{\bar{h}}}(w_{\bar{h}}(\bar{y})))} \leq C \epsilon^{\delta s + \bar{h} - 1} E^{1 - \delta s + \bar{h} - 1}. \quad (4.28)$$

By the triangular inequality,

$$|v(\bar{y})| \leq |v(\bar{y}) - v(\bar{y} - r\nu(\bar{y}))| + |v(\bar{y} - r\nu(\bar{y}))|. \quad (4.29)$$

First, we estimate the second term on the right-hand side of (4.29). Since  $\bar{y} - r e_n \in B_{\rho_{\bar{h}}}(w_{\bar{h}}(\bar{y}))$ ,

$$|v(\bar{y} - r\nu(\bar{y}))| \leq C r_0^{2-n} \epsilon^{\delta s + \bar{h} - 1} E^{1 - \delta s + \bar{h} - 1} \leq C r^{2-n} (\epsilon + E) \left( \frac{\epsilon}{E + \epsilon} \right)^{1 - \delta s + \bar{h} - 1}.$$

Secondly, we estimate the first term on the right-hand side of (4.29). Since  $\bar{y} \in \mathcal{W}_k$ , by (4.20),

$$|v(\bar{y})| \leq C E \left( r_0 \sup_{x \in D_0} d(x) \right)^{1 - (n/2)} \leq C r_0^{2-n} E.$$

Hence, by theorem 4.1,

$$|v(\bar{y}) - v(\bar{y} - r\nu(\bar{y}))| \leq \|\nabla v\|_{L^\infty(Q_{r_0/3})} r \leq \frac{C}{r_0^{1+n/2}} \|v\|_{L^2(Q_{2r_0/3})} r \leq C r_0^{2-n} (E + \epsilon) \left( \frac{r}{r_0} \right).$$

Therefore,

$$|v(\bar{y})| \leq C r_0^{2-n} (E + \epsilon) \left( \frac{r}{r_0} + \left( \frac{\epsilon}{E + \epsilon} \right)^{\delta s + \bar{h} - 1} \right).$$

Minimizing the right-hand side of the last inequality with respect to  $r$ , the following inequality holds:

$$|v(\bar{y})| \leq Cr_0^{2-n}(E + \epsilon) \left| \log \left( \frac{\epsilon}{E + \epsilon} \right) \right|^{\delta^s} \left|^{-\frac{c}{2|\log \delta|}} \right|,$$

for a suitable constant  $C > 0$ . Set  $\tilde{\Sigma}_1 = \Sigma_1 \cap Q_{r_1}(P_1)$ . By the arbitrariness of  $\bar{y}$ , we obtain

$$\|v\|_{L^\infty(\tilde{\Sigma}_1)} \leq Cr_0^{2-n}(E + \epsilon)\omega_{1/C} \left( \frac{\epsilon}{\epsilon + E} \right). \quad (4.30)$$

In order to prove our claim, we need to estimate the gradient of  $v$ . Recalling that  $v_0 = v|_{D_0}$  and  $v_1 = v|_{D_1}$  and  $v_0$  is harmonic in  $D_0$ , from the three sphere inequality applied to  $\nabla v_0$  and the results of [52], one can recover the following estimates:

$$\|\nabla v_0\|_{L^\infty(\tilde{\Sigma}_1)} \leq Cr_0^{2-n}(E + \epsilon)\omega_{1/C} \left( \frac{\epsilon}{\epsilon + E} \right), \quad (4.31)$$

and

$$\|\nabla_T v_1\|_{L^\infty(\tilde{\Sigma}_1)} = \|\nabla_T v_0\|_{L^\infty(\tilde{\Sigma}_1)} \leq \|\nabla v_0\|_{L^\infty(\tilde{\Sigma}_1)} \leq Cr_0^{2-n}(E + \epsilon)\omega_{1/C} \left( \frac{\epsilon}{\epsilon + E} \right). \quad (4.32)$$

Now we can apply the following estimate due to Trytten [62]:

$$\begin{aligned} \int_{D_1 \cap B_{3r_1/8}(P_1)} |\nabla v_1|^2 &\leq \frac{c}{r_0} \left( \int_{\tilde{\Sigma}_1} v_1^2 + r_0^2 \int_{\tilde{\Sigma}_1} |\nabla v_1|^2 \right)^{\delta_1} \\ &\times \left( \int_{\tilde{\Sigma}_1} v_1^2 + r_0^2 \int_{\tilde{\Sigma}_1} |\nabla v_1|^2 + r_0 \int_{D_1 \cap B_{r_1/4}(P_1)} A |\nabla v_1|^2 \right)^{1-\delta_1}. \end{aligned} \quad (4.33)$$

In order to bound the left-hand side of (4.33), we have to estimate the following quantities:

- (a)  $\int_{\tilde{\Sigma}_1} v_1^2$ ;
- (b)  $\int_{\tilde{\Sigma}_1} |\nabla v_1|^2$ ;
- (c)  $\int_{D_1 \cap B_{r_1/4}(P_1)} A |\nabla v_1|^2$ .

For (a), we can just use (4.30). For (b), since  $\nabla v_1 = \nabla_T v_1 + (\nabla v_1 \cdot \nu)\nu$ ,

$$\int_{\tilde{\Sigma}_1} |\nabla v_1|^2 \leq \int_{\tilde{\Sigma}_1} |\nabla_T v_1|^2 + \int_{\tilde{\Sigma}_1} |(\nabla v_1 \cdot \nu)\nu|^2.$$

The first integral on the right-hand side can be estimated using (4.32). For the other term, one uses the transmission conditions

$$A(x)\nabla v_0 \cdot \nu = A(x)\nabla v_1 \cdot \nu, \quad \text{on } \Sigma_1. \quad (4.34)$$

Then,

$$\|\nabla v_1\|_{L^\infty(\tilde{\Sigma}_1)} \leq Cr_0^{1-n}(E + \epsilon)\omega_{1/C} \left( \frac{\epsilon}{\epsilon + E} \right). \quad (4.35)$$

Finally, (c) follows from standard energy estimates.

From the following trace estimate

$$\int_{D_1 \cap B_{3r_l/16}(P_1)} v_1^2 \leq C \left( r_0 \int_{\bar{\Sigma}_1} v_1^2 + r_0^2 \int_{D_1 \cap B_{3r_l/8}(P_1)} |\nabla v_1|^2 \right), \quad (4.36)$$

(4.30), (4.33), (4.35) and (4.36) it follows that

$$\|v_1\|_{L^\infty(B_{\bar{\gamma}_1})} \leq Cr_0^{1-n}(E + \epsilon)\omega_{1/C} \left( \frac{\epsilon}{\epsilon + E} \right). \quad (4.37)$$

Case  $m \implies m + 1$ . Set

$$\epsilon_m = C^{m+1}r_0^{2-n}(E + \epsilon) \left( \omega_{1/C}^{(m+1)} \left( \frac{\epsilon}{\epsilon + E} \right) \right).$$

By proceeding as above, we end up with the following inequality

$$\|v_1\|_{L^\infty(B_{\bar{\gamma}_{m+1}})} \leq Cr_0^{1-n}(E + \epsilon_m)\omega_{1/C} \left( \frac{\epsilon_m}{\epsilon_m + E} \right). \quad (4.38)$$

By the properties (3.5) and (3.6) of  $\omega_{1/C}$ , the claim follows. To summarise it, we have proved that for any point close enough to the interface, the  $L^\infty$ -norm of  $v$  on a small ball can be bound in terms of the quantities the right-hand side of (4.19) and (4.20).

For  $m < K - 1$  the thesis follows by the inequality (4.28), choosing  $\bar{y} = P_{m+1}$ .

For  $m = K - 1$ , by condition (4.20), arguing as in the inequality (4.28) and applying the claim, it follows that

$$\begin{aligned} |v(w_{\bar{h}}(P_K))| &\leq C(r_0^{2-n}\epsilon_K)^{\delta^{s+\bar{h}-1}}(r_0d_1a^{\bar{h}-1}E)^{1-\delta^{s+\bar{h}-1}} \leq C^{\bar{h}}r_0^{2-n}(\epsilon_K + E)\omega_{1/C} \left( \frac{\epsilon_K}{\epsilon_K + E} \right) \\ &\leq C^{\bar{h}}r_0^{2-n}(\epsilon + E)\omega_{1/C}^{(K)} \left( \frac{\epsilon}{\epsilon + E} \right)^{(1/C^{\bar{h}})}. \end{aligned}$$

**Proof of proposition 3.2.** To begin with, recall that for any  $(y, z) \in (D_0)_r \times (D_0)_r$ , for  $r \in (0, d_1]$ , the following bound holds:

$$|S_{\mathcal{U}_k}(y, z)| \leq \|\sigma^{(1)} - \sigma^{(2)}\|_{L^\infty(\Omega)}(\text{dist}(y, \mathcal{U}_k) \text{dist}(z, \mathcal{U}_k))^{1-n/2}.$$

For any  $y, z \in B_{\rho_{\bar{h}(r)}}(w_{\bar{h}(r)}(Q_{k+1}))$ , we apply proposition 4.2 once to  $v = S_{\mathcal{U}_k}(\cdot, z)$  and then to  $v = S_{\mathcal{U}_k}(y, \cdot)$  to obtain

$$|S_{\mathcal{U}_k}(y, z)| \leq r_0^{2-n}C^{\bar{h}(r)}(E + \epsilon_0) \left( \omega_{1/C}^{(2k)} \left( \frac{\epsilon_0}{E + \epsilon_0} \right) \right)^{(1/C)^{\bar{h}(r)}}. \quad (4.39)$$

Hence (3.14) follows from (4.39).

Since  $S_{\mathcal{U}_k}(y_1, \dots, y_n, z_1, \dots, z_n)$  is a weak solution in  $D_k \times D_k$  of the elliptic equation

$$\text{div}_y(\sigma^{(1)}(y)\nabla_y S_{\mathcal{U}_k}(y, z)) + \text{div}_z(\sigma^{(2)}(z)\nabla_z S_{\mathcal{U}_k}(y, z)) = 0, \quad (4.40)$$

for any  $i, j = 1, \dots, n$  it follows that

$$\begin{aligned} & \|\partial_{x_i} \partial_{x_j} \mathcal{S} \mathcal{U}_k(x_1, \dots, x_n, x_{n+1}, \dots, x_{2n})\|_{L^\infty \left( B_{\frac{\rho_{\bar{h}(r)}}{2}}(w_{\bar{h}(r)}(\mathcal{Q}_{k+1})) \times B_{\frac{\rho_{\bar{h}(r)}}{2}}(w_{\bar{h}(r)}(\mathcal{Q}_{k+1})) \right)} \\ & \leq \frac{C}{\rho_{\bar{h}(r)-1}^2} \|\mathcal{S} \mathcal{U}_k(x_1, \dots, x_n, x_{n+1}, \dots, x_{2n})\|_{L^\infty(B_{\frac{\rho_{\bar{h}(r)}}{2}}(w_{\bar{h}(r)}(\mathcal{Q}_{k+1})) \times B_{\frac{\rho_{\bar{h}(r)}}{2}}(w_{\bar{h}(r)}(\mathcal{Q}_{k+1})))}, \end{aligned} \quad (4.41)$$

where  $x_i = y_i, x_{i+n} = z_i$  for  $i = 1, \dots, n$ .

Moreover, since  $d_{\bar{h}(r)-1} > r$ , it follows that  $r < \frac{d_0}{a\rho_0} \rho_{\bar{h}(r)}$ , which in turn leads to

$$\|\partial_{x_i} \partial_{x_j} \mathcal{S} \mathcal{U}_k(x_1, \dots, x_{2n})\|_{L^\infty(\bar{\mathcal{Q}}_{\frac{\rho_{\bar{h}(r)}}{2}}(w_{\bar{h}(r)}(\mathcal{Q}_{k+1})))} \leq \frac{C}{r^2} \|\mathcal{S} \mathcal{U}_k(x_1, \dots, x_{2n})\|_{L^\infty(\bar{\mathcal{Q}}_{\rho_{\bar{h}(r)}}(w_{\bar{h}(r)}(\mathcal{Q}_{k+1})))}. \quad (4.42)$$

By (3.12), it follows that  $r^{-2} \leq \left(\frac{a}{r_0}\right)^2 \left(\frac{1}{a^2}\right)^{\bar{h}(r)}$ , and by combining (4.42) and the above inequality we get the desired estimate.  $\square$

## Acknowledgments

The authors wish to thank the anonymous referees for useful comments and remarks which improved the presentation of the paper. RG would like to acknowledge the support of the Dipartimento di matematica e geoscienze, Università degli Studi di Trieste, where the research of this paper was initiated during her sabbatical leave in 2020. The work of ES was performed under the PRIN Grant No. 201758MTR2-007.

## Data availability statement

No new data were created or analysed in this study.

## ORCID iDs

Sonia Foschiatti  <https://orcid.org/0000-0002-3505-2295>

Romina Gaburro  <https://orcid.org/0000-0002-6186-0712>

Eva Sincich  <https://orcid.org/0000-0002-7405-1842>

## References

- [1] Alessandrini G 1988 Stable determination of conductivity by boundary measurements *Appl. Anal.* **27** 153–72
- [2] Alessandrini G 1990 Singular solutions of elliptic equations and the determination of conductivity by boundary measurements *J. Differ. Equ.* **84** 252–72
- [3] Alessandrini G, Beretta E, Rosset E and Vessella S 2000 Optimal stability for inverse elliptic boundary value problems with unknown boundaries *Ann. Scuola Norm. Sup. Pisa, Cl. Sci.* **XXXIX** 755–806



- [4] Alessandrini G, de Hoop M V and Gaburro R 2017 Uniqueness for the electrostatic inverse boundary value problem with piecewise constant anisotropic conductivities *Inverse Problems* **33** 125013
- [5] Alessandrini G and Gaburro R 2001 Determining conductivity with special anisotropy by boundary measurements *SIAM J. Math. Anal.* **33** 153–71
- [6] Alessandrini G and Gaburro R 2009 The local Calderón problem and the determination at the boundary of the conductivity *Commun. PDE* **34** 918–36
- [7] Alessandrini G, de Hoop M V, Gaburro R and Sincich E 2017 Lipschitz stability for the electrostatic inverse boundary value problem with piecewise linear conductivities *J. Math. Pure. Appl.* **107** 638–64
- [8] Alessandrini G, de Hoop M V, Gaburro R and Sincich E 2018 Lipschitz stability for a piecewise linear Schrödinger potential from local Cauchy data *Asympt. Anal.* **108** 115–49
- [9] Alessandrini G, de Hoop M V, Faucher F, Gaburro R and Sincich E 2019 Inverse problem for the Helmholtz equation with Cauchy data: reconstruction with conditional well-posedness driven iterative regularization *ESAIM: Math. Modelling Numer. Anal.* **53** 1005–30
- [10] Alessandrini G, Rondi L, Rosset E and Vessella S 2009 The stability for the Cauchy problem for elliptic equations *Inverse Problems* **25** 123004
- [11] Alessandrini G and Vessella S 2005 Lipschitz stability for the inverse conductivity problem *Adv. Appl. Math.* **35** 207–41
- [12] Astala K and Päivärinta L 2006 Calderón’s inverse conductivity problem in the plane *Ann. Math.* **163** 265–99
- [13] Astala K, Päivärinta L and Lassas M 2005 Calderón’s inverse problem for anisotropic conductivity in the plane *Commun. PDE* **30** 207–24
- [14] Barceló J A, Barceló T and Ruiz A 2001 Stability of the inverse conductivity problem in the plane for less regular conductivities *J. Differ. Equ.* **173** 231–70
- [15] Barceló T, Faraco D and Ruiz A 2007 Stability of Calderón inverse conductivity problem in the plane *J. Math. Pure. Appl.* **88** 522–56
- [16] Belishev M I 2003 The Calderon problem for two-dimensional manifolds by the BC-method *SIAM J. Math. Anal.* **35** 172–82
- [17] Beretta E, De Hoop M V and Qiu L 2013 Lipschitz stability of an inverse boundary value problem for a Schrödinger-type equation *SIAM J. Math. Anal.* **45** 679–99
- [18] Beretta E, De Hoop M V, Faucher F and Scherzer O 2016 Inverse boundary value problem for the Helmholtz equation: quantitative conditional Lipschitz stability estimates *SIAM J. Math. Anal.* **48** 3962–83
- [19] Beretta E and Francini E 2011 Lipschitz stability for the electrical impedance tomography problem: the complex case *Commun. PDE* **36** 1723–49
- [20] Beretta E, Francini E, Francini E and Vessella S 2014 Uniqueness and Lipschitz stability for the identification of Lamé parameters from boundary measurements *Inverse Problems Imaging* **8** 611–44
- [21] Beretta E, Francini E, Morassi A, Rosset E and Vessella S 2014 Lipschitz continuous dependence of piecewise constant Lamé coefficients from boundary data: the case of non-flat interfaces *Inverse Problems* **30** 125005
- [22] Borcea L 2002 Electrical impedance tomography *Inverse Problems* **18** R99–136
- [23] Calderón A P 1980 On an inverse boundary value problem *Seminar on Numerical Analysis and its Applications to Continuum Physics* (Rio de Janeiro) pp 65–73 Soc. Brasil. Mat. reprinted in *Comput. Appl. Math.* 25 (2006) 133–138
- [24] Cheney M, Isaacson D and Newell J C 1999 Electrical impedance tomography *SIAM Rev.* **41** 85–101
- [25] Clop A, Faraco D, Faraco D and Ruiz A 2010 Stability of Calderón’s inverse conductivity problem in the plane for discontinuous conductivities *Inverse Problems Imaging* **4** 49–91
- [26] Dos Santos Ferreira D, Kenig C E, Salo M and Uhlmann G 2009 Limiting Carleman weights and anisotropic inverse problems *Invent. Math.* **178** 119–71
- [27] Dos Santos Ferreira D, Kurylev Y, Lassas M and Salo M 2016 The Calderón problem in transversally anisotropic geometries *J. Eur. Math. Soc.* **18** 2579–626
- [28] Druskin V 1982 The unique solution of the inverse problem of electrical surveying and electrical well-logging for piecewise-continuous conductivity *Izv. Earth Phys.* **18** 51–3 (in Russian)
- [29] Druskin V 1985 On uniqueness of the determination of the three-dimensional underground structures from surface measurements with variously positioned steady-state or monochromatic field sources *Sov. Phys.-Solid Earth* **21** 210–4 (in Russian)

- [30] Druskin V 1998 On the uniqueness of inverse problems from incomplete boundary data *SIAM J. Appl. Math.* **58** 1591–603
- [31] De Hoop M V, Qiu L and Scherzer O 2012 Local analysis of inverse problems: Hölder stability and iterative reconstruction *Inverse Problems* **28** 045001
- [32] De Hoop M V, Qiu L and Scherzer O 2015 An analysis of a multi-level projected steepest descent iteration for nonlinear inverse problems in Banach spaces subject to stability constraints *Numer. Math.* **129** 127–48
- [33] Faucher F, Alessandrini G, Barucq H, de Hoop M V, Gaburro R and Sincich E 2020 Full reciprocity-gap waveform inversion enabling sparse-source acquisition *Geophysics* **85** R461–76
- [34] Faraco D, Kurylev Y and Ruiz A 2013 G-convergence, Dirichlet to Neumann maps and invisibility *J. Funct. Anal.* **267** 2478–506
- [35] Faucher F, de Hoop M V and Scherzer O 2021 Reciprocity-gap misfit functional for distributed acoustic sensing, combining data from passive and active sources *Geophysics* **86** R211
- [36] Gaburro R and Lionheart W R B 2009 Recovering Riemannian metrics in monotone families from boundary data *Inverse Problems* **25** 045004
- [37] Gaburro R and Sincich E 2015 Lipschitz stability for the inverse conductivity problem for a conformal class of anisotropic conductivities *Inverse Problems* **31** 015008
- [38] Gilbarg D and Trudinger N S 1983 *Elliptic Partial Differential Equations of Second Order* 2nd edn (Berlin: Springer)
- [39] Garde H and Hyvönen N 2020 Optimal depth-dependent distinguishability bounds for electrical impedance tomography in arbitrary dimension *SIAM J. Appl. Math.* **80** 20–43
- [40] Greenleaf A, Lassas M and Uhlmann G 2003 Anisotropic conductivities that cannot be detected by EIT *Physiol. Meas.* **24** 413
- [41] Greenleaf A, Lassas M and Uhlmann G 2003 On nonuniqueness for Calderón’s inverse problem *Math. Res. Lett.* **10** 685–93
- [42] Kohn R and Vogelius M 1984 Identification of an unknown conductivity by means of measurements at the boundary *SIAM-AMS Proc.* **14** 113–23
- [43] Kohn R and Vogelius M 1984 Determining conductivity by boundary measurements *Commun. Pure Appl. Math.* **37** 289–98
- [44] Kohn R V and Vogelius M 1985 Determining conductivity by boundary measurements: II. Interior results *Commun. Pure Appl. Math.* **38** 643–67
- [45] Lassas M and Uhlmann G 2001 On determining a Riemannian manifold from the Dirichlet-to-Neumann map *Ann. Sci. École Norm. Sup.* **34** 771–87
- [46] Lassas M, Taylor M and Uhlmann G 2003 The Dirichlet-to-Neumann map for complete Riemannian manifolds with boundary *Commun. Anal. Geom.* **11** 207–21
- [47] Lee J M and Uhlmann G 1989 Determining anisotropic real-analytic conductivities by boundary measurements *Commun. Pure Appl. Math.* **42** 1097–112
- [48] Lionheart W R B 1997 Conformal uniqueness results in anisotropic electrical impedance imaging *Inverse Problems* **13** 125
- [49] Lions J L and Magenes E 1972 *Non-homogeneous Boundary Value Problems and Applications I (Die Grundlehren der mathematischen Wissenschaften vol 181)* (New York: Springer)
- [50] Liu L 1997 Stability estimates for the two-dimensional inverse conductivity problem *PhD Thesis* University of Rochester, New York
- [51] Littman W, Stampacchia G and Weinberger H W 1963 Regular points for elliptic equations with discontinuous coefficients *Ann. Scuola Norm. Pisa Cl. Sci.* **3** 43–77
- [52] Li Y Y and Vogelius M 2000 Gradient estimates for solutions to divergence form elliptic equations with discontinuous coefficients *Arch. Ration. Mech. Anal.* **153** 91–151
- [53] Mandache N 2001 Exponential instability in an inverse problem for the Schrödinger equation *Inverse Problems* **17** 1435–44
- [54] Nachman A I 1996 Global Uniqueness for a two-dimensional inverse boundary value problem *Ann. Math.* **143** 71–96
- [55] Nagayasu S, Uhlmann G and Wang J-N 2009 A depth-dependent stability estimate in electrical impedance tomography *Inverse Problems* **25** 075001
- [56] Rondi L 2006 A remark on a paper by Alessandrini and Vessella *Adv. Appl. Math.* **36** 67–9
- [57] Rüländ A and Sincich E 2019 Lipschitz stability for the finite dimensional fractional Calderón problem with finite Cauchy data *Inverse Problems Imaging* **13** 1023–44
- [58] Rüländ A and Sincich E 2020 On Runge approximation and Lipschitz stability for a finite-dimensional Schrödinger inverse problem *Appl. Anal.* 1–12

- [59] Schlumberger C 1920 *Etude sur la prospection électrique du sous-sol* (Paris: Gauthier Villars)
- [60] Sylvester J 1990 An anisotropic inverse boundary value problem *Commun. Pure Appl. Math.* **43** 201–32
- [61] Sylvester J and Uhlmann G 1987 A global uniqueness Theorem for an inverse boundary value problem *Ann. Math.* **125** 153–69
- [62] Trytten G N 1963 Pointwise bounds for solutions of the Cauchy problem for elliptic equations *Arch. Ration. Mech. Anal.* **13** 222–44
- [63] Uhlmann G 2009 Electrical impedance tomography and Calderón’s problem *Inverse Problems* **25** 123011